Title: Hyperornithinemia-Hyperammonemia-Homocitrullinuria Syndrome GeneReview

Table 5

Authors: Camacho J, Rioseco-Camacho N

Date: May 2012

Note: The following information is provided by the authors listed above and has not

been reviewed by GeneReviews staff.

Table 5. Selected SLC25A15 Pathologic Allelic Variants

DNA Nucleotide Change	Protein Amino Acid Change	Reference
c.562_564delTTC	p.Phe188del ^{1, 2}	Camacho et al [1999]
c.535C>T	p.Arg179ter ^{1, 3}	Tsujino et al [2000]
c.538G>A	p.Glu180Lys 3, 4	Camacho et al [1999]
c.569G>A	p.Gly190Asp ^{1, 2}	Salvi et al [2001]
c.564C>G	p.Phe188Leu	Tessa et al [2009]
c.95C>G	p.Thr32Arg ^{2, 4}	Camacho et al [2006]
c.79G>A	p.Gly27Arg ^{1, 3}	Salvi et al [2001]
c.658G>A	p.Gly220Arg ^{3, 4}	Al-Hassan et al [2008]
c.44C>A	p.Ala15Glu ¹	Shih & Ficicioglu [2000]
c.823C>T	p.Arg275ter ³	Torisu et al [2006]
c.824G>A	p.Arg275Gln ³	Salvi et al [2001]
c.110T>G	p.Met37Arg ³	Tessa et al [2009]
c.818T>A	p.Met273Lys ²	Fecarotta et al [2006]
c.337G>T	p.Gly113Cys ²	Fecarotta et al [2006]
c.847C>T	p.Leu283Phe 1, 2	Tessa et al [2009]
c.815C>T	p.Thr272Iso ²	Tessa et al [2009]
c.212T>A	p.Leu71Gln ^{1, 2}	Tessa et al [2009]

^{1.} SLC25A15 (ORNT1) mutations associated with neonatal onset

^{2.} Residual function

^{3.} No residual function

^{4.} Targets normally to mitochondria

References

Al-Hassnan ZN, Rashed MS, Al-Dirbashi OY, et al. Hyperornithinemia-Hyperammonemia-Homocitrullinuria syndrome with stroke-like imaging presentation: Clinical, biochemical and molecular analysis. J Neurol Sci. 2008;264:187-94.

Camacho JA, Mardach R, Rioseco-Camacho N, et al. Clinical and functional characterization of a human ornt1 mutation (T32R) in the hyperornithinemia-hyperammonemia-homocitrullinuria (HHH) syndrome. Pediatr Res. 2006;60:423-9.

Camacho JA, Obie C, Biery B, et al. Hyperornithinaemia-Hyperammonaemia-Homocitrullinuria syndrome is caused by mutations in a gene encoding a mitochondrial ornithine transporter. Nat Genet. 1999;22:151-8.

Fecarotta S, Parenti G, Vajro P, et al. HHH syndrome (Hyperornithinaemia, Hyperammonaemia, Homocitrullinuria), with fulminant hepatitis-like presentation. J Inherit Metab Dis. 2006;29:186-9.

Salvi S, Santorelli FM, Bertini E, et al. Clinical and molecular findings in Hyperornithinemia-Hyperammonemia-Homocitrullinuria syndrome. Neurology. 2001;57:911-4.

Shih VE, Laframboise R, Mandell R, et al. Neonatal form of the Hyperornithinaemia, Hyperammonaemia, and Homocitrullinuria (HHH) syndrome and prenatal diagnosis. Prenat Diagn. 1992;12:717-23.

Tessa A, Fiermonte G, Dionisi-Vici C, et al Identification of novel mutations in the SLC25A15 gene in Hyperornithinemia-Hyperammonemia-Homocitrullinuria (HHH) syndrome: A clinical, molecular, and functional study. Hum Mutat. 2009;30:741-8.

Torisu H, Kira R, Kanazawa N, Takemoto M, Sanefuji M, Sakai Y, Tsujino S, Hara T. A novel R275X mutation of the SLC25A15 gene in a Japanese patient with the HHH syndrome. Brain Dev. 2006;28:332-5.

Tsujino S, Kanazawa N, Ohashi T, et al. Three novel mutations (G27E, insAAC, R179X) in the ornt1 gene of Japanese patients with Hyperornithinemia, Hyperammonemia, and Homocitrullinuria syndrome. Ann Neurol. 2000;47:625-31.