

Title: Pancreatitis Overview *GeneReview* – Genetic Risk Factors that Predispose to Pancreatitis

Authors: LaRusch J, Solomon S, Whitcomb DC

Date: March 2014

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

## Genetic Risk Factors that Predispose to Pancreatitis

It is critical to note that the phenotype of individuals with the same allelic variant(s) may differ and the majority of individuals heterozygous for a single mutation in one of these genes (other than *PRSS1*) do not have pancreatitis.

**Table 1. Molecular Genetic Risk Factors that Predispose to Pancreatitis**

Degree of Risk	Inheritance Pattern	Frequency in Control Population	Gene (Variant)	Molecular Consequence of Variant	Other
<b>High</b>	Autosomal dominant	<0.1% (frequency same for all 3 alleles?)	<i>PRSS1</i> (gene duplication, R122H,N29I)	-50% increase in protein levels -Protein resistant to degradation	See Table 2
	Autosomal dominant with low penetrance or multigenic		<i>PRSS1</i> (A16V)	Protein partially resistant to degradation	
<b>Moderate</b>	Multigenic or autosomal dominant with low penetrance	<0.1%	<i>SPINK1</i> (c.27delC, p.Tyr54His)	Severe loss of protein function	See Table 3
	Multigenic or autosomal dominant with low penetrance	<0.1% 0.4% <0.1%	<i>CTRC</i> p.K247_R254del24 p.R254W p.Ala73Thr		
	Autosomal recessive or multigenic	3.0% <0.1% <0.1%	<i>CFTR</i> p.F508del c.del_exon2, 3 p.G551D		

Degree of Risk	Inheritance Pattern	Frequency in Control Population	Gene (Variant)	Molecular Consequence of Variant	Other
<b>Mild</b>	Autosomal recessive or multigenic	5-10%	<i>CFTR</i> (p.R75Q)	Selective loss of protein function	See Table 4
	Autosomal recessive or multigenic	3-5%	<i>SPINK1</i> (p.N34S, no known change)		
<b>Modifiers</b>	Complex	70%	<i>PRSS1</i> (c.-408T>C)	Altered expression of protein	See Table 5
	Complex	26%	<i>CLDN2</i> (c.-275-1293G>A)		
	Complex	20%	<i>CTRC</i> (c.180C>T)	No known change	

1. This table is provided by the authors of this GeneReview and has not been reviewed by GeneReviews staff.

2. Allelic variants may be described in [www.CFTR2.org](http://www.CFTR2.org) and [www.pancreasgenetics.org/](http://www.pancreasgenetics.org/)

**Table 2. *PRSS1* Allelic Variants**

Class of Variant Allele	DNA Nucleotide Change	Protein Amino Acid Change	Reference Sequences
<b>Pathogenic</b>		p.Arg122His	<a href="#">NM_002769.4</a> <a href="#">NP_002760.1</a>
		p.Arg122Cys	
		p.Ala16Val	
		p.Cys139Ser	
		p.Val39Ala	
		p.Asn29Ile	

**Table 3. *SPINK1* Allelic Variants**

Class of Variant Allele	DNA Nucleotide Change	Protein Amino Acid Change	Reference Sequences
<b>Pathogenic</b>	c.101A>G	p.Asn34Ser	<a href="#">NM_003122.3</a> <a href="#">NP_003113.2</a>
	c.194+2T>C 1		
	c.150T>G	p.Asp50Glu	
	c.160T>C	p.Tyr54His	
	c.199C>T	p.Arg67Cys	
	c.163C>T	p.Pro55Ser	

1. A T to C mutation at the +2 position of intron 3 causes exon 3 skipping [Kume et al 2006]

**Table 4. *CFTR* Allelic Variants**

Class of Variant Allele	DNA Nucleotide Change (Alias <sup>1</sup> )	Protein Amino Acid Change	Reference Sequences
<b>Benign</b>	c.1408A>G	p.Met470Val	<a href="#">NM_000492.3</a> <a href="#">NP_000483.3</a>
	c.3870A>G	p.Pro1290Pro	
	c.1584G>A	p.Glu528Glu	
<b>Pathogenic</b>	c.1521_1523delCTT	p.Phe508del	
	c.224G>A	p.Arg75Gln	
	c.1652G>A	p.Glu551Asp	
	c.1624G>T	p.Glu542Ter	
	(del_exon 2,3) <sup>2</sup>		
	(del_exon 22,23) <sup>2</sup>		

1. Variant designation that does not conform to current naming conventions

2. Large deletion mutations of multiple exons in *CFTR* c.54-5940\_273+10250del21kb (del\_exon2,3) and c.3963-78\_4242+577del (del\_exon22,23)

**Table 5. *CTRC* Allelic Variants Discussed in This *GeneReview***

Class of Variant Allele	DNA Nucleotide Change	Protein Amino Acid Change	Reference Sequences
<b>Pathogenic</b>	c.760C>T	p.Arg254Trp	<a href="#">NM_007272.2</a> <a href="#">NP_009203.2</a>
	c.738_761del24	p.Lys247_Arg254del	
	c.164G>A	p.Trp55Ter	

**Other genes** in which mutations or common variants have been implicated as risk factors for chronic pancreatitis include:

- *CASR*: Mutations have either a small effect or have not been studied and/or confirmed in all populations [Felderbauer et al 2003, Muddana et al 2008, Murugaian et al 2008]. *CASR* loss-of-function variants appear to increase the risk of pancreatitis in individuals with *CFTR* mutations, while loss-of-function mutations appear to increase the risk of pancreatitis in heavy users of alcohol.
- *GGT1*: Variation in the gene encoding γ-glutamyltransferase 1 has been recently associated with both pancreatic cancer and chronic pancreatitis [Brand et al 2013].
- Non-coding variants in the *CLDN2* locus and *PRSS1-PRSS2* locus: Common variants that do not directly alter amino acid sequence have recently been identified as risk modifiers for chronic pancreatitis in North America, but have not been investigated in families with hereditary pancreatitis [Whitcomb et al 2012]. *CLDN2*-locus haplotypes on the X chromosome is an X-linked-recessive risk factor for alcoholic pancreatitis, while a *PRSS1-PRSS2*-locus haplotype appears

to reduce trypsinogen gene expression and reduces the risk for pancreatitis from multiple causes.

- CPA1: carboxypeptidase 1 is the second most abundant pancreatic digestive enzyme produced by pancreatic acinar cells. Multiple rare mutations increase risk of chronic pancreatitis through protein missfolding and increased endoplasmic reticulum stress response, especially in children [Witt et al 2013].
- Note: Genes in which mutations do not appear to be associated with chronic pancreatitis include: *PRSS3*, *TGFB1*, *TNF*, *TNFRSF1A*, *IL10*, *UGT1A7*, *PDGFB*, *EPHX1*, *HFE* (in which mutations cause [HFE-related hereditary hemochromatosis](#)), *GP2* [Masson et al 2010] and *SERPINA1* (in which mutations cause [alpha1 antitrypsin deficiency](#)).

**Variants in the secondary trypsinogens.** Anionic trypsinogen encoded by *PRSS2* and mesotrypsin encoded by *PRSS3* are generally not considered major risk factors in non-*PRSS1* hereditary pancreatitis, but isolated instances have been described of mutations associated with either disease or healthy controls. For example:

- **Damaging variants.** A double gain-of-function mutation was reported in which a hybrid duplication of exons 1 and 2 of *PRSS2* and exons 3 to 5 of *PRSS1* resulted in a copy number variant (CNV) of trypsin with a damaging Isoleucine at the 29th amino acid [Masson et al 2008].
- **Protective variants**
  - It has been suggested that a rare protective loss-of-function *PRSS2* mutation- p.Gly191Arg -may mitigate risk alleles in other genes [Witt et al 2006, Santhosh et al 2008].
  - Protective mitigation is also seen in the frequent low expression haplotype *PRSS1* characterized by the intronic SNP rs10273639, which is significantly less frequent in persons with pancreatitis than healthy controls [Whitcomb et al 2012].

## References

- Brand H, Diergaarde B, O'Connell MR, Whitcomb DC, Brand RE. Variation in the  $\gamma$ -glutamyltransferase 1 gene and risk of chronic pancreatitis. *Pancreas*. 2013;42:836-40.
- Felderbauer P, Hoffmann P, Einwächter H, Bulut K, Ansorge N, Schmitz F, Schmidt WE. A novel mutation of the calcium sensing receptor gene is associated with chronic pancreatitis in a family with heterozygous SPINK1 mutations. *BMC Gastroenterol*. 2003;3:34.
- Kume K, Masamune A, Kikuta K, Shimosegawa T. [-215G>A; IVS3+2T>C] mutation in the SPINK1 gene causes exon 3 skipping and loss of the trypsin binding site. *Gut*. 2006;55:1214.
- Masson E, Le Maréchal C, Chandak GR, Lamoril J, Bezieau S, Mahurkar S, Bhaskar S, Reddy DN, Chen JM, Férec C. Trypsinogen copy number mutations in patients with idiopathic chronic pancreatitis. *Clin Gastroenterol Hepatol*. 2008;6:82-8
- Masson E, Paliwal S, Bhaskar S, Prakash S, Scotet V, Reddy DN, Le Maréchal C, Ratan Chandak G, Chen JM, Férec C. Genetic analysis of the glycoprotein 2 gene in patients with chronic pancreatitis. *Pancreas*. 2010;39:353-8.
- Muddana V, Lamb J, Greer JB, Elinoff B, Hawes RH, Cotton PB, Anderson MA, Brand RE, Slivka A, Whitcomb DC. Association between calcium sensing receptor gene polymorphisms and chronic

pancreatitis in a US population: role of serine protease inhibitor Kazal 1 type and alcohol. *World J Gastroenterol.* 2008;14:4486-91.

Murugaian EE, Premkumar RM, Radhakrishnan L, Vallath B. Novel mutations in the calcium sensing receptor gene in tropical chronic pancreatitis in India. *Scand J Gastroenterol.* 2008;43:117-21.

Santhosh S, Witt H, te Morsche RH, Nemoda Z, Molnár T, Pap A, Jansen JB, Drenth JP. A loss of function polymorphism (G191R) of anionic trypsinogen (PRSS2) confers protection against chronic pancreatitis. *Pancreas.* 2008;36:317-20.

Whitcomb DC, LaRusch J, Krasinskas AM, Klei L, Smith JP, Brand RE, Neoptolemos JP, Lerch MM, Tector M, Sandhu BS, Guda NM, Orlichenko L; Alzheimer's Disease Genetics Consortium, Alkaade S, Amann ST, Anderson MA, Baillie J, Banks PA, Conwell D, Côté GA, Cotton PB, DiSario J, Farrer LA, Forsmark CE, Johnstone M, Gardner TB, Gelrud A, Greenhalf W, Haines JL, Hartman DJ, Hawes RA, Lawrence C, Lewis M, Mayerle J, Mayeux R, Melhem NM, Money ME, Muniraj T, Papachristou GI, Pericak-Vance MA, Romagnuolo J, Schellenberg GD, Sherman S, Simon P, Singh VP, Slivka A, Stolz D, Sutton R, Weiss FU, Wilcox CM, Zarnescu NO, Wisniewski SR, O'Connell MR, Kienholz ML, Roeder K, Barmada MM, Yadav D, Devlin B. Common genetic variants in the CLDN2 and PRSS1-PRSS2 loci alter risk for alcohol-related and sporadic pancreatitis. *Nat Genet.* 2012;44:1349-54.

Witt H, Beer S, Rosendahl J, Chen JM, Chandak GR, Masamune A, Bence M, Szmola R, Oracz G, Macek M Jr, Bhatia E, Steigenberger S, Lasher D, Bühler F, Delaporte C, Tebbing J, Ludwig M, Pilsak C, Saum K, Bugert P, Masson E, Paliwal S, Bhaskar S, Sobczynska-Tomaszewska A, Bak D, Balascak I, Choudhuri G, Nageshwar Reddy D, Rao GV, Thomas V, Kume K, Nakano E, Kakuta Y, Shimosegawa T, Durko L, Szabó A, Schnúr A, Hegyi P, Rakonczay Z Jr, Pfützer R, Schneider A, Groneberg DA, Braun M, Schmidt H, Witt U, Friess H, Algül H, Landt O, Schuelke M, Krüger R, Wiedenmann B, Schmidt F, Zimmer KP, Kovacs P, Stumvoll M, Blüher M, Müller T, Janecke A, Teich N, Grützmann R, Schulz HU, Mössner J, Keim V, Löhr M, Férec C, Sahin-Tóth M. Variants in CPA1 are strongly associated with early onset chronic pancreatitis. *Nat Genet.* 2013;45:1216-20.

Witt H, Sahin-Tóth M, Landt O, Chen JM, Kähne T, Drenth JP, Kukor Z, Szepessy E, Halangk W, Dahm S, Rohde K, Schulz HU, Le Maréchal C, Akar N, Ammann RW, Truninger K, Bargetzi M, Bhatia E, Castellani C, Cavestro GM, Cerny M, Destro-Bisol G, Spedini G, Eiberg H, Jansen JB, Koudova M, Rausova E, Macek M Jr, Malats N, Real FX, Menzel HJ, Moral P, Galavotti R, Pignatti PF, Rickards O, Spicak J, Zarnescu NO, Böck W, Gress TM, Friess H, Ockenga J, Schmidt H, Pfützer R, Löhr M, Simon P, Weiss FU, Lerch MM, Teich N, Keim V, Berg T, Wiedenmann B, Luck W, Groneberg DA, Becker M, Keil T, Kage A, Bernardova J, Braun M, Güldner C, Halangk J, Rosendahl J, Witt U, Treiber M, Nickel R, Férec C. A degradation-sensitive anionic trypsinogen (PRSS2) variant protects against chronic pancreatitis. *Nat Genet.* 2006;38:668-73.