

Title: Hutchinson-Gilford Progeria Syndromes *GeneReview* Preclinical Studies

Authors: Gordon LB, Brown WT, Collins FS

Date: December 2014

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

Preclinical Studies in HGPS Cells and Murine Models

The inability to release progerin from the nuclear membrane results in structural stress on the nucleus. Immunofluorescence of HGPS fibroblasts with antibodies directed against lamin A reveals a visible abnormality, an irregular shape of the nuclear envelope, in 40%-50% of cells [Eriksson et al 2003]. It is hypothesized that this permanently farnesylated mutant form of prelamin A (progerin) acts in a dominant-negative fashion and leads to the progressive defects in nuclear architecture that are seen in HGPS [Goldman et al 2004]. Progerin also mislocalizes into insoluble cytoplasmic aggregates and membranes during mitosis, causing abnormal chromosome segregation, binucleation, and a lag in progression out of late mitosis [Cao et al 2007].

Functionally, patient cells exhibit loss of plasticity and increased stiffness [Dahl et al 2010], which may be in part why tissues exposed to mechanical stress (vasculature, bone, joints) are clinically affected. Nuclear function is globally abnormal in patient cells, with widespread alterations of chromatin structure, such as the loss of heterochromatin domains, changes in epigenetic markers maintenance of genomic integrity [Pegoraro et al 2009], DNA repair defects [Musich & Zou 2009, Benson et al 2010], replication fork stalling [Liu et al 2006], and telomere dysfunction [Benson et al 2010, Cao et al 2011].

Broad abnormalities in cellular stress-response pathways involved in HGPS include hyperactivation of p53 signaling [Varela et al 2005, Liu et al 2006, Kudlow et al 2008], decreased numbers of adult stem cells [Espada et al 2008, Scaffidi & Misteli 2008, Rosengardten et al 2011], dysregulation of the somatotrophic axis and several miRNA-controlled circuits [Marino et al 2010, Ugalde et al 2011], and profound changes in glucose and lipid metabolism [Marino et al 2008].

Induced pluripotent stem cells (iPSC) and their potential. Patient-derived iPSC currently provide a unique tool to study HGPS cells and pathways normally unavailable for study, such as mesenchymal stem cells and differentiation pathways of tissues affected by the disease such as vascular smooth muscle cells (VSMCs), adipose tissue and bone [Zhang et al 2011]. In addition, they have been used to develop strategies that can successfully correct the HGPS mutation [Liu et al 2011].

References

- Benson EK, Lee SW, Aaronson SA. Role of progerin-induced telomere dysfunction in HGPS premature cellular senescence. *J Cell Sci.* 2010;123:2605-12.
- Cao K, Blair CD, Faddah DA, Kieckhaefer JE, Olive M, Erdos MR, Nabel EG, Collins FS. Progerin and telomere dysfunction collaborate to trigger cellular senescence in normal human fibroblasts. *J Clin Invest.* 2011;121:2833-44.
- Cao K, Capell BC, Erdos MR, Djabali K, Collins FS. A lamin A protein isoform overexpressed in Hutchinson-Gilford progeria syndrome interferes with mitosis in progeria and normal cells. *Proc Natl Acad Sci U S A.* 2007;104:4949-54.
- Dahl KN, Kalinowski A, Pekkan K. Mechanobiology and the microcirculation: cellular, nuclear and fluid mechanics. *Microcirculation.* 2010;17:179-91.
- Eriksson M, Brown WT, Gordon LB, Glynn MW, Singer J, Scott L, Erdos MR, Robbins CM, Moses TY, Berglund P, Dutra A, Pak E, Durkin S, Csoka AB, Boehnke M, Glover TW, Collins FS. Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. *Nature.* 2003;423:293-8.
- Espada J, Varela I, Flores I, Ugalde AP, Cadiñanos J, Pendás AM, Stewart CL, Tryggvason K, Blasco MA, Freije JM, López-Otín C. Nuclear envelope defects cause stem cell dysfunction in premature-aging mice. *J Cell Biol.* 2008;181:27-35.
- Goldman RD, Shumaker DK, Erdos MR, Eriksson M, Goldman AE, Gordon LB, Gruenbaum Y, Khuon S, Mendez M, Varga R, Collins FS. Accumulation of mutant lamin A causes progressive changes in nuclear architecture in Hutchinson-Gilford progeria syndrome. *Proc Natl Acad Sci U S A.* 2004;101:8963-8.
- Kudlow BA, Stanfel MN, Burtner CR, Johnston ED, Kennedy BK. Suppression of proliferative defects associated with processing-defective lamin A mutants by hTERT or inactivation of p53. *Mol Biol Cell.* 2008;19:5238-48.
- Liu Y, Rusinol A, Sinensky M, Wang Y, Zou Y. DNA damage responses in progeroid syndromes arise from defective maturation of prelamin A. *J Cell Sci.* 2006;119:4644-9.
- Mariño G, Ugalde AP, Fernández AF, Osorio FG, Fueyo A, Freije JM, López-Otín C. Insulin-like growth factor 1 treatment extends longevity in a mouse model of human premature aging by restoring somatotroph axis function. *Proc Natl Acad Sci U S A.* 2010;107:16268-73.
- Mariño G, Ugalde AP, Salvador-Montoliu N, Varela I, Quirós PM, Cadiñanos J, van der Pluijm I, Freije JM, López-Otín C. Premature aging in mice activates a systemic metabolic response involving autophagy induction. *Hum Mol Genet.* 2008;17:2196-211.
- Musich PR, Zou Y. Genomic instability and DNA damage responses in progeria arising from defective maturation of prelamin A. *Aging (Albany NY).* 2009;1:28-37.
- Pegoraro G, Kubben N, Wickert U, Göhler H, Hoffmann K, Misteli T. Ageing-related chromatin defects through loss of the NURD complex. *Nat Cell Biol.* 2009;11:1261-7.
- Rosengardten Y, McKenna T, Grochová D, Eriksson M. Stem cell depletion in Hutchinson-Gilford progeria syndrome. *Aging Cell.* 2011;10:1011-20.
- Scaffidi P, Misteli T. Lamin A-dependent misregulation of adult stem cells associated with accelerated ageing. *Nat Cell Biol.* 2008;10:452-9.
- Ugalde AP, Ramsay AJ, de la Rosa J, Varela I, Mariño G, Cadiñanos J, Lu J, Freije JM, López-Otín C. Aging and chronic DNA damage response activate a regulatory pathway involving miR-29 and p53. *EMBO J.* 2011;30:2219-32.
- Varela I, Cadiñanos J, Pendás AM, Gutiérrez-Fernández A, Folgueras AR, Sánchez LM, Zhou Z, Rodríguez FJ, Stewart CL, Vega JA, Tryggvason K, Freije JM, López-Otín C. Accelerated ageing in mice deficient in Zmpste24 protease is linked to p53 signalling activation. *Nature.* 2005;437:564-8.
- Zhang J, Lian Q, Zhu G, Zhou F, Sui L, Tan C, Mutalif RA, Navasankari R, Zhang Y, Tse HF, Stewart CL, Colman A. A human iPSC model of Hutchinson Gilford Progeria reveals vascular smooth muscle and mesenchymal stem cell defects. *Cell Stem Cell.* 2011;8:31-45.