

Title: *PROP1*-Related Combined Pituitary Hormone Deficiency *GeneReview* – *POU1F1* Variants

Author: de Graaff L

Date: August 2014

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

### Specific variants in *POU1F1*

Many *POU1F1* variants have been described to date. For a comprehensive review, see Pfäffle & Klammt [2011].

Recently, several new *POU1F1* mutations were described:

- In 2011, Tenenbaum et al reported the follow up of a patient with a nonsense mutation, in whom central hypothyroidism was diagnosed at the age of 2 months and GH and PRL deficiencies were documented at 9 months. MRI at 14 years revealed a hypoplastic adenohypophysis. The patient underwent spontaneous but delayed puberty. A novel disease-causing mutation (c.502insT) was identified in the homozygous state in exon 4 of *POU1F1*. This insertion results in a frameshift introducing an early termination codon at position 174 (p.Thr168IlefsX7), leading to a severely truncated protein lacking the entire homeodomain. This mutation abolishes *POU1F1*'s transactivation properties on three target promoters [Tenenbaum-Rakover et al 2011].
- In 2012, Turton et al described a male patient with extreme short stature, learning difficulties, anterior pituitary hypoplasia, secondary hypothyroidism and undetectable prolactin, growth hormone (GH) and insulin-like growth factor 1 (IGF1), with normal random cortisol. In this patient, combined pituitary hormone deficiency in this patient was caused by loss of *POU1F1* function by two novel mechanisms, namely aberrant splicing (IVS1+3nt (A>G) and protein instability (R265W).
- Inoue et al 2012 described a novel heterozygous splice site mutation (Ex2 + 1G>T; c.214 + 1G>T) in identical twin brothers with mild CPHD. In vitro splicing studies suggested this mutation to result in an in-frame skipping of exon 2, thus producing an internally deleted protein lacking most of the R2 transactivation subdomain (TAD-R2). Heterologous expression studies of the mutated *POU1F1* protein showed modest reductions in its transactivation activities in HEK293T cells, while acting as a dominant-negative inhibitor of the endogenous activities of *POU1F1* in pituitary GH3 cells.

## References

Inoue H, Mukai T, Sakamoto Y, Kimura C, Kangawa N, Itakura M, Ogata T, Ito Y, Fujieda K; Japan Growth Genome Consortium (2012) Identification of a novel mutation in the exon 2 splice donor site of the POU1F1/PIT-1 gene in Japanese identical twins with mild combined pituitary hormone deficiency. (Clin Endocrinol (Oxf). 2012;76:78-87.

Pfäffle R, Klammt J. Pituitary transcription factors in the aetiology of combined pituitary hormone deficiency. Best Pract Res Clin Endocrinol Metab. 2011;25:43-60.

Tenenbaum-Rakover Y, Sobrier ML, Amselem S. A novel POU1F1 mutation (p.Thr168IlefsX7) associated with an early and severe form of combined pituitary hormone deficiency: functional analysis and follow-up from infancy to adulthood. Clin Endocrinol (Oxf). 2011;75:214-9.

Turton JP, Strom M, Langham S, Dattani MT, Le Tissier P. Two novel mutations in the POU1F1 gene generate null alleles through different mechanisms leading to combined pituitary hormone deficiency. (Clin Endocrinol (Oxf). 2012;76:387-93.