Title: *ATP1A3*-Related Neurologic Disorders *GeneReview* Table 2 Authors: Brashear A, Sweadner KJ, Cook JF, Swoboda KJ, Ozelius L

Date: November 2014

Note: The following information is provided by the authors and has not been reviewed by

GeneReviews staff.

Table 2. ATP1A3 Genotypes as They Correlate with RDP, AHC, and CAPOS Phenotypes

Nucleotide Change	Protein Change	Phenotype	Inherited or <i>de novo</i> Mutation	Recurrent (Yes/No), Frequency
410C>A	Ser137Tyr	AHC	de novo	Yes, infrequent
410C>T	Ser137Phe	AHC	de novo	No
419A>T	Gln140Leu	AHC	de novo	No
821T>A	Ile274Asn	AHC	de novo Familial (dominant)	Yes, infrequent
821T>C	lle274Thr	RDP	de novo	No
829G>A	Glu277Lys	RDP	de novo	Yes, infrequent
965T>A	Val322Asp	AHC	de novo	Yes, infrequent
979_981delCTG	Leu327del	RDP	de novo	No
998G>T	Cys333Phe	AHC	de novo	Yes, infrequent
1003A>C	Thr335Pro	AHC	de novo	No
1072G>T	Gly358Cys	AHC	de novo	No
1109C>A	Thr370Asn	RDP	de novo	No
1112T>C	Leu371Pro	AHC	de novo	No
1144T>C	Trp382Arg	RDP	de novo	No
1250T>C	Leu417Pro	RDP	de novo	No
1838C>T	Thr613Met	RDP	de novo Familial (dominant)	Yes, intermediate N = 6
2051C>T	Ser684Phe	RDP	de novo	No
2264G>C	Gly755Ala	AHC	de novo	No
2263G>A	Gly755Ser	AHC	de novo	Yes, infrequent
2263G>T	Gly755Cys	AHC	de novo	Yes, infrequent
2267G>A	Arg756His	RDP	de novo	No
2270T>C	Leu757Pro	AHC	de novo	No
2273T>G	lle758Ser	RDP	Familial (dominant)	No
2312C>A	Thr771Asn	AHC	de novo	No
2316C>A	Ser772Arg	AHC	de novo	No
2318A>G	Asn773Ser	AHC	de novo	No
2318A>T	Asn773lle	AHC	de novo	No
2338T>C	Phe780Leu	RDP	Familial (dominant)	No
Not reported	Asp801Glu	AHC	de novo /familial	No (twins)
2401G>A	Asp801Asn	AHC	de novo	Yes, frequent N >60 reported

Nucleotide Change	Protein Change	Phenotype	Inherited or <i>de novo</i> Mutation	Recurrent (Yes/No), Frequency
2401G>T	Asp801Tyr	RDP	Familial	No
2411C>T	Thr804lle	AHC	de novo	Yes, infrequent
2415C>G	Asp805Glu	AHC	de novo	No
2417T>G	Met806Arg	AHC	de novo	No
2428A>T	Ile810Phe	AHC	de novo	No
2429T>G	lle810Ser	AHC	de novo	No
2431T>C	Ser811Pro	AHC	de novo	Yes, infrequent
2443G>A	Glu815Lys	AHC	de novo	Yes, frequent N >40 reported
2452G>A	Glu818Lys	Capos	de novo Familial (dominant)	Yes, low frequency N = 3 families; 1 <i>de novo</i>
2542+1G>A	Splice site	AHC	de novo	Yes, infrequent
2600G>A	Gly867Asn	RDP/AHC		No
2755_2757delGTC	Val919del	AHC	de novo	No
2767G>A	Asp923Asn	AHC/RDP	Familial (AHC) de novo & familial (RDP)	Yes, low frequency N = 5
2767G>T	Asp923Thr	AHC	de novo	No
2780G>A	Cys927Tyr	AHC	de novo	No
2780G>T	Cys927Phe	AHC	de novo	Yes, infrequent
2781C>G	Cys927Trp	AHC	de novo	No
2839G>A	Gly947Arg	AHC	de novo	Yes, intermediate N >10
2839G>C	Gly947Arg	AHC	de novo	Yes, low frequency N = 4
2864C>A	Ala955Asp	AHC	de novo	No
2974G>T	Asp992Tyr	AHC	de novo	Yes, infrequent
3038_3040dupACT	Tyr1013dup	RDP	de novo	No

AHC = alternating hemiplegia of childhood

RDP = rapid-onset dystonia parkinsonism

CAPOS = cerebellar ataxia, areflexia, pes cavus, optic atrophy, and sensorineural hearing loss

Updated May 4, 2014

Reference Sequences: <u>NM 152296.3</u>; <u>NP 689509.1</u>

RDP. Fourteen variants have been described to date to cause RDP (reviewed in Heinzen et al 2014, Rosewich et al 2014).

- Three are recurrent:
 - c.829G>A encoding p.Glu277Lys (~13% all de novo) [de Carvalho Aguiar et al 2004, Brashear et al 2007, Tarsy et al 2010];
 - c.1838C>T resulting in p.Thr613Met (~26% both *de novo* and familial occurrence) [de Carvalho Aguiar et al 2004, Brashear et al 2007, Lee et al 2007, McKeon et al 2007; Barbano 2012];
 - o c.2767G>A encoding p.Asp923Asn (~8% all *de nov*o) [Zanotti-Fregonara et al 2008, Anselm et al 2009].
- The following mutations have been reported at least once: c.821T>C (p.lle274Thr);
 c.979_981delCTG (p.Leu327del); c.1109C>A (p.Thr370Asn); c.1144T>C (p.Trp382Arg);
 c.1250T>C (p.Leu417Pro); c.2051C>T (p.Ser684Phe); c.2267G>A (p.Arg756His); c.2273T>G
 (p.lle758Ser); c.2338T>C (p.Phe780Leu); c.2401G>T (p.Asp801Tyr); c.3038_3040dupACT (p.Tyr1013dup).

AHC. More than 30 pathogenic variants have been reported to result in an AHC phenotype in more than 150 individuals.

- The following mutations are recurrent and occur at a reasonably high frequency; together they
 account for ~ 2/3 of simplex cases reported to date:
 - o c.2401G>A in exon17 resulting in p.Asp801Asn (~40%)
 - o c.2443G>A in exon 17 resulting in p.Gly815Lys (~24%)
 - o c.2839G>A and c.2839G>C in exon 21 resulting in p.Gly947Arg (~8%)
- The following mutations have been reported more than once, but at a much lower frequency: c.410C>A (p.Ser137Tyr); c.821T>A (p.Ile274Asn); c.965T>A (p.Val322Asp); c.998G>T (p.Cys333Phe); c.2263G>A (p.Gly755Ala); c.2263G>A (p.Gly755Ser); c.2263G>T (p.Gly755Cys); c.2411C>T (p.Thr804lle); c.2431T>C (p.Ser811Pro); splice site change c.2452+1G>A; c.2780G>T (p.Cys927Phe); c.2974G>T (p.Asp992Tyr).
- The following mutations have been reported at least once: c.410C>T (p.Ser137Phe); c.419A>T (p.Gln140Leu); c.1003A>C (p.Thr335Pro); c.1072G>T (p.Gly358Cys); c.1112T>C (p.Leu371Pro); c.2264G>C (p.Gly755Ala); c.2270T>C (p.Leu757Pro; c.2312C>A (p.Thr771Asn); c.2316C>A (p.Ser772Arg); c.2318A>G (p.Asn773Ser); c.2318A>T (p.Asn773lle); c.2415C>G (p.Asp805Glu); c.2417T>G (p.Met806Arg); c.2428A>T (p.Ile810Phe); c.2429T>G (p.Ile810Ser); c.2600G>A (p.Gly867Asn); c.2755_2757delGTC (p.Val919del); c.2767G>A (p.Asp923Asn); c.2767G>T (p.Asp923Thr); c.2780G>A (p.Cys927Thr); c.2781C>G (p.Cys927Trp); c.2864C>A (p.Ala955Asp).

References

Anselm IA, Sweadner KJ, Gollamudi S, Ozelius LJ, Darras BT. Rapid-onset dystonia-parkinsonism in a child with a novel atp1a3 gene mutation. Neurology. 2009;73:400-1.

Barbano RL, Hill DF, Snively BM, Light LS, Boggs N, McCall WV, Stacy M, Ozelius L, Sweadner KJ, Brashear A. New triggers and non-motor findings in a family with rapid-onset dystonia-parkinsonism. Parkinsonism Relat Disord. 2012;18:737-41.

Brashear A, Dobyns WB, de Carvalho Aguiar P, Borg M, Frijns CJ, Gollamudi S, Green A, Guimaraes J, Haake BC, Klein C, Linazasoro G, Munchau A, Raymond D, Riley D, Saunders-Pullman R, Tijssen MA, Webb D, Zaremba J, Bressman SB, Ozelius LJ. The phenotypic spectrum of rapid-onset dystonia-parkinsonism (RDP) and mutations in the ATP1A3 gene. Brain. 2007;130:828-35.

de Carvalho Aguiar P, Sweadner KJ, Penniston JT, Zaremba J, Liu L, Caton M, Linazasoro G, Borg M, Tijssen MA, Bressman SB, Dobyns WB, Brashear A, Ozelius LJ. Mutations in the Na+/K+ -ATPase alpha3 gene ATP1A3 are associated with rapid-onset dystonia parkinsonism. Neuron 2004;43:169-75.

Heinzen EL, Arzimanoglou A, Brashear A, Clapcote SJ, Gurrieri F, Goldstein DB, Jóhannesson SH, Mikati MA, Neville B, Nicole S, Ozelius LJ, Poulsen H, Schyns T, Sweadner KJ, van den Maagdenberg A, Vilsen Bfor the ATP1A3 Working Group. Distinct neurological disorders with ATP1A3 mutations. Lancet Neurol. 2014;13:503-14.

Lee JY, Gollamudi S, Ozelius LJ, Kim JY, Jeon BS. ATP1A3 mutation in the first Asian case of rapid-onset dystonia-parkinsonism. Mov Disord. 2007;22:1808-9.

McKeon A, Ozelius LJ, Hardiman O, Greenway MJ, Pittock SJ. Heterogeneity of presentation and outcome in the Irish rapid-onset dystonia-Parkinsonism kindred. Mov Disord. 2007;22:1325-7.

Rosewich H, Ohlenbusch A, Huppke P, Schlotawa L, Baethmann M, Carrilho I, Fiori S, Lourenço CM, Sawyer S, Steinfeld R, Gärtner J, Brockmann K. The expanding clinical and genetic spectrum of ATP1A3-related disorder. Neurology. 2014;82:945-55.

Tarsy D, Sweadner KJ, Song PC. Case records of the Massachusetts General Hospital. Case 17-2010 - A 29-year-old woman with flexion of the left hand and foot and difficulty speaking. N Engl J Med. 2010;362:2213-9.

Zanotti-Fregonara P,Vidailhet M, Kas A, Ozelius LJ, Clot F, Hindié E, Ravasi L, Devaux JY, Roze E. [123I]-FP-CIT and [99mTc]-HMPAO single photon emission computed tomography in a new sporadic case of rapid-onset dystonia-parkinsonism. J Neurol Sci. 2008;273:148-51.