

Heme/Onc Literature Surveillance -- Cycle 2026-07-02

- **Surveillance window:** 2026-06-02 to 2026-07-02
 - **Disease groups covered:** Acute Myeloid Leukemia, Myelodysplastic Syndromes, Myeloproliferative Neoplasms, Chronic Lymphocytic Leukemia, Diffuse Large B-Cell Lymphoma, Follicular Lymphoma, Mantle Cell Lymphoma, Marginal Zone Lymphoma, Hodgkin Lymphoma, Multiple Myeloma, Acute Lymphoblastic Leukemia, Chronic Myeloid Leukemia, Aplastic Anemia, Clonal Hematopoiesis of Indeterminate Potential, Sickle Cell Disease
 - **Total new items surfaced:** 15
 - **Flagged for retraction/concern:** 0
 - **Needs human review:** 0
 - **Mode:** abstracts only — no AI summarization, extraction, or appraisal. Each item shows its identification info and the verbatim source abstract.
-

Acute Myeloid Leukemia

Trials

A randomized trial of GVHD prophylaxis in haploidentical PBSC transplantation: ATG, PTCy, and low-dose combination therapy.

Journal: Blood | **Date:** 2026-Jun-25 | **PMID:** 41849227 | **DOI:** 10.1182/blood.2025032569

Publication types: Journal Article, Randomized Controlled Trial, Clinical Trial, Phase III, Multicenter Study

Abstract:

The optimal graft-versus-host disease (GVHD) prophylaxis strategy in haploidentical peripheral blood stem cell transplantation remains controversial. In this open-label, phase 3 study, patients aged 14 to 70 years with acute myeloid leukemia or myelodysplastic syndromes with excess blasts ■ or ■ were randomized (2:1:1) to receive low-dose antithymocyte globulin (ATG; 5 mg/kg) plus posttransplant cyclophosphamide (PTCy; 50 mg/kg; referred to as ATG/PTCy), standard-dose ATG (total dose, 10 mg/kg), or a PTCy-based (total dose, 100 mg/kg) regimen for GVHD prophylaxis. The coprimary end points were the cumulative incidence (CI) of grade 2 to 4 acute GVHD (aGVHD) by day 100 and GVHD-free, relapse-free survival at 1 year after transplant. A total of 407 patients were randomized to receive an ATG/PTCy (185 patients), ATG (113 patients), or PTCy (109 patients) regimen for GVHD prophylaxis. By day +100, the CI of grade 2 to 4 aGVHD did not differ significantly among the 3 groups ($P = .210$). Although the overall incidence of chronic GVHD (cGVHD) was comparable across all groups ($P = .110$), the 2-year CI of moderate-to-severe cGVHD was numerically lower in the ATG/PTCy (17.4%) and ATG (17.3%) groups than the PTCy group (28.3%), without reaching statistical significance ($P = .095$). No significant differences were observed in survival outcomes among the 3 groups. Notably, the CI of neutrophil and platelet recovery was significantly higher in the ATG/PTCy group than in the other groups ($P < .001$). This trial suggested that the 3 GVHD prophylaxis strategies presented similar efficacy in preventing grade 2 to 4 aGVHD and yielded comparable survival. This trial was registered at www.clinicaltrials.gov as NCT03608059.

Cladribine With Low-Dose Cytarabine and Venetoclax Alternating With Azacitidine and Venetoclax for Newly Diagnosed Acute Myeloid Leukemia.

Journal: American journal of hematology | **Date:** 2026-Jul | **PMID:** 42035228 | **DOI:** 10.1002/ajh.70328

Publication types: Clinical Trial, Phase II, Journal Article

Abstract:

Venetoclax-based low-intensity regimens have improved the outcomes of older or unfit patients with acute myeloid leukemia (AML). This phase II study investigated the combination of cladribine plus low-dose cytarabine and venetoclax alternating with azacitidine plus venetoclax for older or unfit patients with newly diagnosed AML. A total of 190 patients were included; the median age was 68 years (range, 47-84 years; 13% $\geq$ 75 years). By the European LeukemiaNet 2022 classification, 16%, 20%, and 64% were stratified as favorable, intermediate, and adverse risk, respectively. The rates of complete remission (CR)/CR with incomplete blood count recovery (CRi) and minimal residual disease (MRD) negative CR/CRi were 84% and 75% overall and 91% and 77% among patients with TP53-wild type AML, respectively. The 4- and 8-week mortality rates were 1% and 3%, respectively. Among responders, 44% proceeded to allogeneic hematopoietic stem cell transplantation. The median overall survival (OS) and event free survival (EFS) were 52 and 50 months, respectively. The 2- and 5-year OS rates were 60% and 45%, respectively. The 2- and 5-year EFS rates were 56% and 43%, respectively. Patients achieving MRD-negative CR had a median OS not reached and a 2-year OS rate of 70%. The median time to absolute neutrophil count recovery ($> 1 \times 10^9$)

Myelodysplastic Syndromes

Trials

A randomized trial of GVHD prophylaxis in haploidentical PBSC transplantation: ATG, PTCy, and low-dose combination therapy.

Journal: Blood | **Date:** 2026-Jun-25 | **PMID:** 41849227 | **DOI:** 10.1182/blood.2025032569

Publication types: Journal Article, Randomized Controlled Trial, Clinical Trial, Phase III, Multicenter Study

Abstract:

The optimal graft-versus-host disease (GVHD) prophylaxis strategy in haploidentical peripheral blood stem cell transplantation remains controversial. In this open-label, phase 3 study, patients aged 14 to 70 years with acute myeloid leukemia or myelodysplastic syndromes with excess blasts $\blacksquare$ or $\blacksquare$ were randomized (2:1:1) to receive low-dose antithymocyte globulin (ATG; 5 mg/kg) plus posttransplant cyclophosphamide (PTCy; 50 mg/kg; referred to as ATG/PTCy), standard-dose ATG (total dose, 10 mg/kg), or a PTCy-based (total dose, 100 mg/kg) regimen for GVHD prophylaxis. The coprimary end points were the cumulative incidence (CI) of grade 2 to 4 acute GVHD (aGVHD) by day 100 and GVHD-free, relapse-free survival at 1 year after transplant. A total of 407 patients were randomized to receive an ATG/PTCy (185 patients), ATG (113 patients), or PTCy (109 patients) regimen for GVHD prophylaxis. By day +100, the CI of grade 2 to 4 aGVHD did not differ significantly among the 3 groups ($P = .210$). Although the overall incidence of chronic GVHD (cGVHD) was comparable across all groups ($P = .110$), the 2-year CI of moderate-to-severe cGVHD was numerically lower in the ATG/PTCy (17.4%) and ATG (17.3%) groups than the PTCy group (28.3%), without reaching statistical significance ($P = .095$). No significant differences were observed in survival outcomes among the 3 groups. Notably, the CI of neutrophil and platelet recovery was significantly higher in the ATG/PTCy group than in the other groups ($P < .001$). This trial suggested that the 3 GVHD prophylaxis strategies presented similar efficacy in preventing grade 2 to 4 aGVHD and yielded comparable survival. This trial was registered at www.clinicaltrials.gov as NCT03608059.

Imetelstat improves patient-reported outcomes and quality of life in lower-risk myelodysplastic syndromes: results from the phase III IMerge study.

Journal: Haematologica | **Date:** 2026-Jul-01 | **PMID:** 41568521 | **DOI:** 10.3324/haematol.2025.288956

Publication types: Journal Article, Clinical Trial, Phase III, Randomized Controlled Trial, Multicenter Study

Abstract:

Red blood cell (RBC) transfusions for anemia associated with lower-risk myelodysplastic syndromes/neoplasms (LR-MDS) often contribute to reduced quality of life (QOL). Thus, reduction in RBC transfusion dependency (TD) is a primary therapeutic goal. Imetelstat is a first-in-class, competitive telomerase inhibitor approved to treat certain adult

patients with LR-MDS with RBC-TD anemia who have not responded to, have lost response to, or are ineligible for erythropoiesis-stimulating agents. In the phase III IMerge study (clinicaltrials.gov identifier: NCT02598661), treatment with imetelstat resulted in clinically meaningful, statistically significant increases in the primary endpoint of ≥ 8 -week RBC transfusion independence (TI) versus placebo. Because patients with LR-MDS experience detrimental effects on numerous facets of QOL (physical, emotional, social, and functional), these exploratory analyses assessed patient-reported outcomes using the Functional Assessment of Chronic Illness Therapy-Fatigue, Quality of Life in Myelodysplasia Scale, and Functional Assessment of Cancer Therapy-Anemia questionnaires as part of the phase III IMerge study. Nominal P values were reported. Fewer imetelstat-treated patients experienced deterioration in fatigue levels and more imetelstat-treated patients experienced sustained improvement in fatigue levels and QOL versus placebo. In the imetelstat group, 8-week, 24-week, and 1-year RBC-TI responders had sustained improvements in predefined significance thresholds versus non-responders for fatigue (70%, 73%, and 88%, respectively, vs. 37%, 41%, and 44%, respectively; $P < 0.001$, $P = 0.004$, and $P = 0.002$) and QOL across different measures of response (43-53% vs. 21-30%; $P \leq 0.0126$). These results suggest that treatment with imetelstat may be associated with improvement in QOL beyond fatigue while sustaining RBC-TI in patients with LR-MDS with RBC-TD anemia.

Myeloproliferative Neoplasms

No new qualifying items this cycle.

Chronic Lymphocytic Leukemia

Guidelines

Updated consensus guidelines for the diagnosis and management of patients with HCL and HCL variant.

Journal: Blood | **Date:** 2026-Jul-02 | **PMID:** 41980020 | **DOI:** 10.1182/blood.2025032757

Publication types: Journal Article, Review, Practice Guideline

Abstract:

Hairy cell leukemia (HCL) and HCL variant (HCLv) are distinct, rare, and chronic splenic B-cell lymphomas/leukemias that partially overlap in clinicopathologic presentation but differ in genetic basis, prognosis, and management. HCL is caused by the BRAF-V600E kinase-activating mutation in $>95\%$ of the patients, usually has excellent responses to chemotherapy with purine analogues, and is also amenable to BRAF inhibitor-based targeted treatments. In contrast, HCLv lacks BRAFV600E mutation, requires combined therapy with purine analogues in addition to rituximab, and generally shows less durable responses. Here, an international team of hematologists, experts on these rare diseases, was convened by the Hairy Cell Leukemia Foundation to update the previous guidelines (published in 2017) by providing a summary of current methods to diagnose and manage patients with HCL and HCLv as well as a prospective on newer targeted therapies to further improve outcomes.

Trials

Venetoclax combinations in untreated CLL: 5-year results and patient-reported outcomes analysis of the CLL13/GAIA trial.

Journal: Blood | **Date:** 2026-Jun-18 | **PMID:** 41911073 | **DOI:** 10.1182/blood.2025032160

Publication types: Journal Article, Randomized Controlled Trial, Clinical Trial, Phase III, Multicenter Study

Abstract:

Fixed-duration venetoclax combinations have become a standard first-line treatment in chronic lymphocytic leukemia (CLL). The phase 3 CLL13/GAIA trial assesses 3 time-limited combinations: venetoclax-rituximab (RV),

venetoclax-obinutuzumab (GV), and venetoclax-obinutuzumab-ibrutinib (GIV), in comparison with chemoimmunotherapy (CIT). Fit patients with CLL without TP53 aberrations were randomized between 6 cycles of CIT (fludarabine-cyclophosphamide-rituximab [FCR] or bendamustine-rituximab [BR]) or 12 cycles of RV, GV, or GIV (GIV: ibrutinib continuation until cycle 36 if measurable residual disease at months 12/15). In total, 926 patients were randomized (GIV: 231, GV: 229, RV: 237, and CIT: 229 [FCR: 150, BR: 79]). With a median observation time of 63.8 months, 5-year progression-free survival (PFS) rates were 81.3% (GIV), 69.8% (GV), 57.4% (RV), and 50.7% (CIT). PFS was superior for GV and GIV compared with CIT and RV ($P < .001$ in each case). In addition, GIV showed longer PFS than GV ($P = .0046$). Venetoclax-based re-treatment after venetoclax-based first-line regimens was efficacious, with 2-year treatment-free survival $>80\%$ from second-line treatment. No differences in overall survival were observed between treatment arms (5-year rates: GIV, 94.3%; GV, 93.6%; RV, 94.7%; and CIT, 90.7%). The incidence rates of severe infections were highest with CIT, whereas cardiac events were most frequent with GIV. Compared with patients treated with CIT, those treated with GV or RV reported rapid and significantly greater quality-of-life (QoL) improvements. In the GIV arm, clinically relevant QoL improvements occurred later (month 15, after the end of treatment in most patients) than with GV/RV, likely due to a higher treatment-related symptom burden. This trial was registered at www.clinicaltrials.gov as NCT02950051.

Diffuse Large B-Cell Lymphoma

Trials

Ten-Year Outcomes after CAR T-Cell Therapy for B-Cell Lymphomas.

Journal: The New England journal of medicine | **Date:** 2026-Jun-25 | **PMID:** 42341302 | **DOI:** 10.1056/NEJMoa2518035

Publication types: Clinical Trial, Phase II, Journal Article, Randomized Controlled Trial

Abstract:

BACKGROUND: Anti-CD19 chimeric antigen receptor (CAR) T-cell therapy is a standard treatment for relapsed or refractory B-cell non-Hodgkin lymphomas. Long-term results and curative potential remain uncertain.

METHODS: We evaluated long-term outcomes in 38 patients with relapsed or refractory B-cell non-Hodgkin lymphomas (24 patients with large B-cell lymphoma and 14 with follicular lymphoma) who had been treated with CTL019 (now called tisagenlecleucel) - autologous T cells expressing CD19-directed, 4-1BB-costimulated chimeric receptors.

Lymphoma-free survival was defined as the time from the tisagenlecleucel infusion to relapse or lymphoma-related death. The incidence of non-relapse-related death and second primary cancer was estimated with the Aalen-Johansen method. The data-cutoff date was October 1, 2025.

RESULTS: At a median follow-up of 10.1 years (range, 7.9 to 11.5), no relapses had occurred beyond 5.4 years. The 10-year lymphoma-free survival was 32% (95% confidence interval [CI], 14 to 51) among patients with large B-cell lymphoma and 47% (95% CI, 20 to 71) among those with follicular lymphoma. In an analysis that included deaths from any cause, the 10-year progression-free survival was 17% (95% CI, 5 to 34) among patients with large B-cell lymphoma and 29% (95% CI, 9 to 52) among those with follicular lymphoma; the 10-year overall survival was 17% (95% CI, 5 to 34) and 50% (95% CI, 23 to 72), respectively. Persistent grade 2 or 3 neutropenia occurred in 2 patients (5%); no late anemia or thrombocytopenia was observed. A second primary cancer developed in 9 patients (10-year cumulative incidence, 21%). The 10-year non-relapse-related mortality was 18% (14% when deaths related to coronavirus disease 2019 were excluded). Higher CAR-transgene persistence appeared to be associated with long-term response. B-cell aplasia persisted in 44% of patients with a long-term response.

CONCLUSIONS: Among patients with heavily pretreated B-cell non-Hodgkin lymphoma, a single infusion of tisagenlecleucel led to decade-long remissions (lymphoma-free survival) in approximately one third of the patients with large B-cell lymphomas and in nearly one half of those with follicular lymphoma. (Funded by the Richard Berman Family Innovations Center in CLL and Lymphomas and others; ClinicalTrials.gov number, NCT02030834.).

Tafasitamab plus lenalidomide and R-CHOP versus R-CHOP for first-line treatment of patients with high-risk diffuse large B-cell lymphoma (frontMIND): a global, phase 3, randomised, double-blind, placebo-controlled trial.

Journal: Lancet (London, England) | **Date:** 2026-Jun-20 | **PMID:** 42217458 | **DOI:** 10.1016/S0140-6736(26)00866-4

Publication types: Journal Article, Randomized Controlled Trial, Clinical Trial, Phase III, Multicenter Study

Abstract:

BACKGROUND: Approximately 40% of patients with high-risk diffuse large B-cell lymphoma (DLBCL) are not cured with first-line R-CHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone or prednisolone). We aimed to investigate the addition of tafasitamab (an Fc-enhanced anti-CD19 monoclonal antibody) and lenalidomide to R-CHOP (tafa-len-R-CHOP) in patients with high-risk aggressive B-cell lymphomas.

METHODS: frontMIND is a phase 3, randomised, double-blind, placebo-controlled study conducted at 298 centres in North America, South America, Europe, and the Asia-Pacific region. Patients aged 18-80 years with previously untreated, high-intermediate-risk or high-risk DLBCL or high-grade B-cell lymphoma (HGBL) were randomly allocated (1:1), stratified by International Prognostic Index (IPI) or age-adjusted IPI and geographical region, to receive six 21-day cycles of standard R-CHOP (rituximab 375 mg/m

FINDINGS: Between May 11, 2021, and March 2, 2023, 1229 patients were screened, among whom 899 were randomly allocated: 448 (50%) to the tafa-len-R-CHOP group and 451 (50%) to the R-CHOP group. At the time of primary analysis (median follow-up 35.2 months [95% CI 35.0-35.4]), progression-free survival was improved in the tafa-len-R-CHOP group versus the R-CHOP group (hazard ratio [HR] 0.75 [95% CI 0.59-0.96]; $p=0.0194$), with 2-year progression-free survival rates of 71.1% (66.3-75.4) with tafa-len-R-CHOP versus 62.9% (57.9-67.5) with R-CHOP. Interim HR for overall survival was 0.85 (0.63-1.14). The overall rate of grade 3 or higher treatment-emergent adverse events was higher with tafa-len-R-CHOP (384 [87%] of 443) than with R-CHOP (340 [76%] of 447). Additionally, a higher rate of fatal treatment-emergent adverse events was observed with tafa-len-R-CHOP (26 [6%]) than with R-CHOP (17 [4%]).

However, the number of overall deaths in the study was lower with tafa-len-R-CHOP than with R-CHOP (82 [19%] vs 97 [22%]). Based on disposition data, rates of premature discontinuation of all study drugs were similar in the tafa-len-R-CHOP group (71 [16%] of 443) and R-CHOP group (66 [15%] of 447).

INTERPRETATION: Progression-free survival was significantly improved with tafa-len-R-CHOP versus R-CHOP; however, the safety profile indicated increases in adverse events, including treatment-emergent adverse events leading to death, with the addition of tafasitamab and lenalidomide. Overall survival data are immature; follow-up is ongoing. Further analyses, including of circulating tumor DNA, will help to assess whether deeper molecular responses are contributing to the progression-free survival benefit observed with tafa-len-R-CHOP. Tafa-len-R-CHOP might represent a potential new first-line treatment for patients with high-risk DLBCL or HGBL.

FUNDING: Incyte Corporation.

Follicular Lymphoma

Trials

Ten-Year Outcomes after CAR T-Cell Therapy for B-Cell Lymphomas.

Journal: The New England journal of medicine | **Date:** 2026-Jun-25 | **PMID:** 42341302 | **DOI:** 10.1056/NEJMoa2518035

Publication types: Clinical Trial, Phase II, Journal Article, Randomized Controlled Trial

Abstract:

BACKGROUND: Anti-CD19 chimeric antigen receptor (CAR) T-cell therapy is a standard treatment for relapsed or refractory B-cell non-Hodgkin lymphomas. Long-term results and curative potential remain uncertain.

METHODS: We evaluated long-term outcomes in 38 patients with relapsed or refractory B-cell non-Hodgkin lymphomas (24 patients with large B-cell lymphoma and 14 with follicular lymphoma) who had been treated with CTL019 (now called tisagenlecleucel) - autologous T cells expressing CD19-directed, 4-1BB-costimulated chimeric receptors.

Lymphoma-free survival was defined as the time from the tisagenlecleucel infusion to relapse or lymphoma-related death. The incidence of non-relapse-related death and second primary cancer was estimated with the Aalen-Johansen method. The data-cutoff date was October 1, 2025.

RESULTS: At a median follow-up of 10.1 years (range, 7.9 to 11.5), no relapses had occurred beyond 5.4 years. The 10-year lymphoma-free survival was 32% (95% confidence interval [CI], 14 to 51) among patients with large B-cell lymphoma and 47% (95% CI, 20 to 71) among those with follicular lymphoma. In an analysis that included deaths from any cause, the 10-year progression-free survival was 17% (95% CI, 5 to 34) among patients with large B-cell lymphoma and 29% (95% CI, 9 to 52) among those with follicular lymphoma; the 10-year overall survival was 17% (95% CI, 5 to 34) and 50% (95% CI, 23 to 72), respectively. Persistent grade 2 or 3 neutropenia occurred in 2 patients (5%); no late anemia or thrombocytopenia was observed. A second primary cancer developed in 9 patients (10-year cumulative incidence, 21%). The 10-year non-relapse-related mortality was 18% (14% when deaths related to coronavirus disease 2019 were excluded). Higher CAR-transgene persistence appeared to be associated with long-term response. B-cell aplasia persisted in 44% of patients with a long-term response.

CONCLUSIONS: Among patients with heavily pretreated B-cell non-Hodgkin lymphoma, a single infusion of tisagenlecleucel led to decade-long remissions (lymphoma-free survival) in approximately one third of the patients with large B-cell lymphomas and in nearly one half of those with follicular lymphoma. (Funded by the Richard Berman Family Innovations Center in CLL and Lymphomas and others; ClinicalTrials.gov number, NCT02030834.).

Lenalidomide plus rituximab for previously untreated advanced follicular lymphoma: the 10-year RELEVANCE trial analysis.

Journal: Blood | **Date:** 2026-Jun-18 | **PMID:** 41915772 | **DOI:** 10.1182/blood.2026033126

Publication types: Journal Article, Randomized Controlled Trial, Clinical Trial, Phase III, Multicenter Study

Abstract:

In the multinational, phase 3 RELEVANCE trial, 1030 patients with previously untreated follicular lymphoma were randomized to receive rituximab + lenalidomide (R2; n = 513) or rituximab-based immunochemotherapy (R-Chemo; n = 517). In the final analysis, at 120 months of follow-up, median progression-free survival (PFS) was comparable between the treatment groups: 110.6 months with R2 vs 102.8 months with R-Chemo, according to the independent review committee assessment. The 10-year PFS rates were 46.4% for R2 and 46.6% for R-chemo. Median overall survival (OS) and time-to-next lymphoma treatment (TTNLT) were not reached in either arm; 10-year OS rates were 82.4% for R2 and 81.1% for R-chemo, and 10-year TTNLT rates were 62.2% for R2 and 66.3% for R-chemo. Overall, patients with progression of disease within 24 months (POD24) had a poorer prognosis than those without POD24 (hazard ratio, 6.215; $P < .0001$); however, no difference was observed between the study groups. The incidence of second primary malignancies (SPMs) was 2.11 cases per 100 patient-years (95% confidence interval, 1.80-2.46). Only 9 transformations occurred after 24 months (3 with R2 vs 6 with R-chemo). In each study group, 87 patients died, mainly because of lymphoma progression and SPMs. This long-term follow-up of RELEVANCE confirmed that R2 provides a chemotherapy-free alternative to immunochemotherapy in this patient population. This trial was registered at www.clinicaltrials.gov as NCT01476787 and NCT01650701 and at EudraCT as 2011-002792-42.

Mantle Cell Lymphoma

No new qualifying items this cycle.

Marginal Zone Lymphoma

No new qualifying items this cycle.

Hodgkin Lymphoma

Trials

Gonadal function and fertility after Hodgkin lymphoma treatment with nivolumab-AVD in the phase 2 GHSG NIVAHL trial.

Journal: Blood | **Date:** 2026-Jun-18 | **PMID:** 41926569 | **DOI:** 10.1182/blood.2025031809

Publication types: Clinical Trial, Phase II, Journal Article, Randomized Controlled Trial

Abstract:

In 82 patients with classic Hodgkin lymphoma, stable gonadal hormone levels were observed up to 24 months after nivolumab in combination with doxorubicin, vinblastine, and dacarbazine (N-AVD) first-line treatment. These findings suggest preserved gonadal function and fertility after 4×N-AVD. The trial was registered at www.clinicaltrials.gov as NCT03004833.

Multiple Myeloma

No new qualifying items this cycle.

Acute Lymphoblastic Leukemia

No new qualifying items this cycle.

Chronic Myeloid Leukemia

No new qualifying items this cycle.

Aplastic Anemia

No new qualifying items this cycle.

Clonal Hematopoiesis of Indeterminate Potential

No new qualifying items this cycle.

Sickle Cell Disease

Guidelines

Gene Therapy for Sickle Cell Disease: Practice Recommendations from the American Society for Transplantation and Cellular Therapy and the International Society for Cell & Gene Therapy.

Journal: Transplantation and cellular therapy | **Date:** 2026-Jul | **PMID:** 41905703 | **DOI:** 10.1016/j.jtct.2026.03.019

Publication types: Journal Article, Review, Practice Guideline

Abstract:

Gene therapy has emerged as a transformative treatment option for individuals with sickle cell disease (SCD), with recent regulatory approvals marking a pivotal shift in clinical care. However, the complexity of patient selection, stem cell mobilization, manufacturing, conditioning, and long-term follow-up underscores the need for standardized, evidence-informed guidance. This consensus document, derived from early clinical experience and developed by the American Society for Transplantation and Cellular Therapy and the International Society for Cell & Gene Therapy, provides practical recommendations for the clinical implementation of ex vivo gene therapies for SCD. Key areas addressed include eligibility assessment, comparative considerations with allogeneic hematopoietic cell transplantation, mobilization and apheresis strategies, conditioning with myeloablative chemotherapy, fertility preservation, psychosocial care, manufacturing quality attributes, and lifelong surveillance for late effects. Collectively, these recommendations aim to harmonize clinical practice; support shared decision-making; and promote safe, equitable, and durable delivery of gene therapy for individuals with SCD.

Trials

Nonmyeloablative HLA-identical sibling transplant for sickle cell disease in children: a multicenter prospective study.

Journal: Blood advances | **Date:** 2026-Jun-23 | **PMID:** 41945757 | **DOI:** 10.1182/bloodadvances.2025019567

Publication types: Clinical Trial, Phase II, Journal Article, Multicenter Study

Abstract:

HLA-identical sibling donor hematopoietic cell transplant (HCT) is an established curative therapy for sickle cell disease (SCD). Although myeloablative HCT conditioning achieves a high event-free survival (EFS) in children, it carries substantial toxicity, including risks of graft-versus-host disease (GVHD), transplant-related mortality, and infertility. Nonmyeloablative conditioning with alemtuzumab, 300 cGy total body irradiation, and sirolimus has demonstrated tolerability as a curative approach in adults with SCD. We conducted a multicenter, prospective clinical trial to evaluate this regimen in children and adolescents with SCD. Thirty-eight patients (aged 2.9-21.9 years; median age, 13.5 years) underwent HCT. All achieved initial engraftment; however, 7 patients (18.4%) experienced secondary graft failure between day +39 and day +391 with autologous hematologic reconstitution. One additional patient received an unconditioned second HCT. At 2 years after HCT, overall survival was 100%, and EFS was 78.9%. Among patients with engraftment, 2-year median donor myeloid chimerism was 95%, and T-cell chimerism was 75%. No patients developed acute or chronic GVHD. The median number of platelet transfusions received was 0 (interquartile range, 0-3). Patient-reported health-related quality of life remained stable early after HCT and significantly improved at 1 year. After year 2, 3 additional patients received a reduced-intensity second HCT to avert secondary graft failure. Although regimen modifications are needed to reduce graft failure risk, this nonmyeloablative approach offers a relatively low-toxicity curative option for children with SCD who have an HLA-identical sibling donor. This trial was registered at www.clinicaltrials.gov as NCT03587272.

Increasing daily step counts improves physical fitness, and reduces pain and arterial stiffness in sickle cell patients.

Journal: Haematologica | **Date:** 2026-Jul-01 | **PMID:** 41641647 | **DOI:** 10.3324/haematol.2025.300290

Publication types: Journal Article, Randomized Controlled Trial

Abstract:

Patients with sickle cell anemia (SCA) have long been discouraged from physical activity (PA). The aim of the present study was to assess the impact of increasing daily step counts on physical fitness, pain and vascular function in patients with SCA. Thirty-eight patients with SCA were recruited and equipped with a Fitbit wrist-worn accelerometer-based PA tracker for five weeks to objectively quantify their baseline daily step counts. Patients were then randomly assigned to one of three groups: 1) control group - no specific information regarding PA was given for eight weeks (N=12); 2) PA1 group - daily step counts increased by 25% of baseline for eight weeks (N=12); 3) PA2 group - daily step counts increased by 25% for four weeks, then by 50% for an additional four weeks (N=14). Pain intensity and frequency

decreased after the intervention in the PA1 and PA2 groups. In addition, patients from these two groups increased the distance walked in six minutes. Arterial stiffness decreased in both PA1 and PA2 groups, without any change in the autonomic nervous system activity. Several inflammatory markers slightly decreased in the PA2 group. Incubation of cultured endothelial cells with patient plasma showed a decrease in the percentage of ICAM-1 positive cells in the PA2 group. This study is the first to show that adopting a simple approach to increase daily PA (i.e., increasing daily step count by 25-50%) for eight weeks is sufficient to decrease pain, and improve physical condition and vascular function of patients with SCA.

HLA-haploidentical hematopoietic stem cell transplantation in patients with sickle cell disease: results from the phase II DREP-HAPLO trial.

Journal: Haematologica | **Date:** 2026-Jul-01 | **PMID:** 41376570 | **DOI:** 10.3324/haematol.2025.300013

Publication types: Journal Article, Clinical Trial, Phase II, Multicenter Study

Abstract:

We report the final analysis of the DREP-HAPLO multicenter phase II protocol, which evaluated haploidentical transplantation after a reduced-intensity conditioning regimen incorporating thiotepa, in children and adults with severe sickle cell disease (SCD). Twenty-two patients (median age, 17 years; range, 12-40) received a conditioning regimen consisting of 2 Gy total body irradiation, thymoglobulin, cyclophosphamide, fludarabine, and thiotepa, followed by T-cell-replete bone marrow infusion and graft-versus-host disease (GvHD) prophylaxis based on post-transplant cyclophosphamide. The primary endpoint, event-free survival, defined as survival without graft failure and without moderate-to-severe chronic GvHD, was 68.18% (95% confidence interval [95% CI]: 51.25%-90.70%) and 61.98% (95% CI: 44.07%-87.19%) at 1 and 4 years, respectively. Overall survival and rejection-free survival at 4 years were 90.15% (95% CI: 78.03%-100%) and 85.56% (95% CI: 71.63%- 100%), respectively. Six patients (27%) developed moderate-to-severe GVHD. In most cases (4/6) GvHD resolved, leaving only two patients (9%) with persistent moderate-to-severe GvHD at their last follow-up. Eighty-five grade 2 to 4 infectious episodes were reported in 21 patients during the 24 months of follow-up, most of which were bacterial (38 cases). These data strengthen existing evidence supporting the feasibility of haploidentical transplantation in both pediatric and adult patients with severe SCD, demonstrating a very low rejection rate when thiotepa is incorporated into the conditioning regimen. Future efforts should focus on reducing chronic GvHD and infection rates.

Needs human review

None this cycle.

Flagged: retraction / expression of concern

None this cycle.

Footer: exact queries used

```
{ "aml": { "pubmed_term": "(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND (\"Leukemia, Myeloid, Acute\"[MeSH Terms]) AND humans[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"Journal of Hematology & Oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology & Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal])", "pubmed_querytranslation": "(\"clinical trial, phase iii\"[Publication Type] OR \"clinical trial, phase ii\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND \"leukemia, myeloid, acute\"[MeSH Terms] AND \"humans\"[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", "pubmed_count": 2, "window": [ "2026-06-18", "2026-07-02" ] }, "mds": { "pubmed_term": "(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND (\"Myelodysplastic Syndromes\"[MeSH Terms]) AND humans[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"Journal of Hematology & Oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology & Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal])", "pubmed_querytranslation": "(\"clinical trial, phase iii\"[Publication Type] OR \"clinical trial, phase ii\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND \"Myelodysplastic Syndromes\"[MeSH Terms] AND \"humans\"[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", "pubmed_count": 2, "window": [ "2026-06-18", "2026-07-02" ] }, "mpn": { "pubmed_term": "(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND (\"Myeloproliferative Disorders\"[MeSH Terms]) AND humans[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR
```

"Leukemia\"[Journal] OR \"Journal of Hematology & Oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology & Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\"[Publication Type] OR \\\"Practice Guideline\"[Publication Type]) AND \\\"Myeloproliferative Disorders\"[MeSH Terms] AND \\\"humans\"[MeSH Terms] AND (\\\"N Engl J Med\"[Journal] OR \\\"Lancet\"[Journal] OR \\\"JAMA\"[Journal] OR \\\"Nature Medicine\"[Journal] OR \\\"J Clin Oncol\"[Journal] OR \\\"The Lancet Oncology\"[Journal] OR \\\"JAMA Oncology\"[Journal] OR \\\"Ann Oncol\"[Journal] OR \\\"Blood\"[Journal] OR \\\"The Lancet Haematology\"[Journal] OR \\\"Blood Advances\"[Journal] OR \\\"Haematologica\"[Journal] OR \\\"Leukemia\"[Journal] OR \\\"Journal of hematology oncology\"[Journal] OR \\\"J Thromb Haemost\"[Journal] OR \\\"American Journal of Hematology\"[Journal] OR \\\"Blood Cancer Journal\"[Journal] OR \\\"British Journal of Haematology\"[Journal] OR \\\"experimental hematology oncology\"[Journal] OR \\\"Bone Marrow Transplantation\"[Journal] OR \\\"HemaSphere\"[Journal] OR \\\"Blood Reviews\"[Journal] OR \\\"Thrombosis and Haemostasis\"[Journal] OR \\\"Transplantation and Cellular Therapy\"[Journal] OR \\\"leukemia lymphoma\"[Journal] OR \\\"Annals of Hematology\"[Journal] OR \\\"Transfusion\"[Journal] OR \\\"Seminars in Hematology\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", \"pubmed_count\": 0, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"c1l\": { \"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication Type] OR \\\"Guideline\"[Publication Type] OR \\\"Practice Guideline\"[Publication Type]) AND (\\\"Leukemia, Lymphocytic, Chronic, B-Cell\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\"[Journal] OR \\\"Lancet\"[Journal] OR \\\"JAMA\"[Journal] OR \\\"Nature Medicine\"[Journal] OR \\\"J Clin Oncol\"[Journal] OR \\\"The Lancet Oncology\"[Journal] OR \\\"JAMA Oncology\"[Journal] OR \\\"Ann Oncol\"[Journal] OR \\\"Blood\"[Journal] OR \\\"The Lancet Haematology\"[Journal] OR \\\"Blood Advances\"[Journal] OR \\\"Haematologica\"[Journal] OR \\\"Leukemia\"[Journal] OR \\\"Journal of Hematology & Oncology\"[Journal] OR \\\"J Thromb Haemost\"[Journal] OR \\\"American Journal of Hematology\"[Journal] OR \\\"Blood Cancer Journal\"[Journal] OR \\\"British Journal of Haematology\"[Journal] OR \\\"Experimental Hematology & Oncology\"[Journal] OR \\\"Bone Marrow Transplantation\"[Journal] OR \\\"HemaSphere\"[Journal] OR \\\"Blood Reviews\"[Journal] OR \\\"Thrombosis and Haemostasis\"[Journal] OR \\\"Transplantation and Cellular Therapy\"[Journal] OR \\\"Leukemia & Lymphoma\"[Journal] OR \\\"Annals of Hematology\"[Journal] OR \\\"Transfusion\"[Journal] OR \\\"Seminars in Hematology\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\"[Publication Type] OR \\\"Practice Guideline\"[Publication Type]) AND \\\"leukemia, lymphocytic, chronic, b cell\"[MeSH Terms] AND \\\"humans\"[MeSH Terms] AND (\\\"N Engl J Med\"[Journal] OR \\\"Lancet\"[Journal] OR \\\"JAMA\"[Journal] OR \\\"Nature Medicine\"[Journal] OR \\\"J Clin Oncol\"[Journal] OR \\\"The Lancet Oncology\"[Journal] OR \\\"JAMA Oncology\"[Journal] OR \\\"Ann Oncol\"[Journal] OR \\\"Blood\"[Journal] OR \\\"The Lancet Haematology\"[Journal] OR \\\"Blood Advances\"[Journal] OR \\\"Haematologica\"[Journal] OR \\\"Leukemia\"[Journal] OR \\\"Journal of hematology oncology\"[Journal] OR \\\"J Thromb Haemost\"[Journal] OR \\\"American Journal of Hematology\"[Journal] OR \\\"Blood Cancer Journal\"[Journal] OR \\\"British Journal of Haematology\"[Journal] OR \\\"experimental hematology oncology\"[Journal] OR \\\"Bone Marrow Transplantation\"[Journal] OR \\\"HemaSphere\"[Journal] OR \\\"Blood Reviews\"[Journal] OR \\\"Thrombosis and Haemostasis\"[Journal] OR \\\"Transplantation and Cellular Therapy\"[Journal] OR \\\"leukemia lymphoma\"[Journal] OR \\\"Annals of Hematology\"[Journal] OR \\\"Transfusion\"[Journal] OR \\\"Seminars in Hematology\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", \"pubmed_count\": 2, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"dlbcl\": { \"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication Type] OR \\\"Guideline\"[Publication Type] OR \\\"Practice Guideline\"[Publication Type]) AND (\\\"Lymphoma, Large B-Cell, Diffuse\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\"[Journal] OR \\\"Lancet\"[Journal] OR \\\"JAMA\"[Journal] OR \\\"Nature Medicine\"[Journal] OR \\\"J Clin Oncol\"[Journal] OR \\\"The Lancet Oncology\"[Journal] OR \\\"JAMA Oncology\"[Journal] OR \\\"Ann Oncol\"[Journal] OR \\\"Blood\"[Journal] OR \\\"The Lancet Haematology\"[Journal] OR \\\"Blood Advances\"[Journal] OR \\\"Haematologica\"[Journal] OR \\\"Leukemia\"[Journal] OR \\\"Journal of Hematology & Oncology\"[Journal] OR \\\"J Thromb Haemost\"[Journal] OR \\\"American Journal of Hematology\"[Journal] OR \\\"Blood Cancer Journal\"[Journal] OR \\\"British Journal of Haematology\"[Journal] OR \\\"Experimental Hematology & Oncology\"[Journal] OR \\\"Bone Marrow Transplantation\"[Journal] OR \\\"HemaSphere\"[Journal] OR \\\"Blood Reviews\"[Journal] OR \\\"Thrombosis and Haemostasis\"[Journal] OR \\\"Transplantation and Cellular Therapy\"[Journal] OR \\\"Leukemia & Lymphoma\"[Journal] OR \\\"Annals of Hematology\"[Journal] OR \\\"Transfusion\"[Journal] OR \\\"Seminars in Hematology\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\"[Publication Type] OR \\\"Practice Guideline\"[Publication Type]) AND

"lymphoma, large b cell, diffuse"[MeSH Terms] AND "humans"[MeSH Terms] AND ("N Engl J Med"[Journal] OR "Lancet"[Journal] OR "JAMA"[Journal] OR "Nature Medicine"[Journal] OR "J Clin Oncol"[Journal] OR "The Lancet Oncology"[Journal] OR "JAMA Oncology"[Journal] OR "Ann Oncol"[Journal] OR "Blood"[Journal] OR "The Lancet Haematology"[Journal] OR "Blood Advances"[Journal] OR "Haematologica"[Journal] OR "Leukemia"[Journal] OR "Journal of hematology oncology"[Journal] OR "J Thromb Haemost"[Journal] OR "American Journal of Hematology"[Journal] OR "Blood Cancer Journal"[Journal] OR "British Journal of Haematology"[Journal] OR "experimental hematology oncology"[Journal] OR "Bone Marrow Transplantation"[Journal] OR "HemaSphere"[Journal] OR "Blood Reviews"[Journal] OR "Thrombosis and Haemostasis"[Journal] OR "Transplantation and Cellular Therapy"[Journal] OR "leukemia lymphoma"[Journal] OR "Annals of Hematology"[Journal] OR "Transfusion"[Journal] OR "Seminars in Hematology"[Journal] OR "Research and Practice in Thrombosis and Haemostasis"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]", "pubmed_count": 2, "window": ["2026-06-18", "2026-07-02"] }, "follicular_lymphoma": { "pubmed_term": "(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND (\"Lymphoma, Follicular\"[MeSH Terms]) AND humans[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"Journal of Hematology & Oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology & Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal])", "pubmed_querytranslation": "(\"clinical trial, phase iii\"[Publication Type] OR \"clinical trial, phase ii\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND \"lymphoma, follicular\"[MeSH Terms] AND \"humans\"[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]", "pubmed_count": 2, "window": ["2026-06-18", "2026-07-02"] }, "mantle_cell_lymphoma": { "pubmed_term": "(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND (\"Lymphoma, Mantle-Cell\"[MeSH Terms]) AND humans[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology & Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal])", "pubmed_querytranslation": "(\"clinical trial, phase iii\"[Publication Type] OR \"clinical trial, phase ii\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type]) AND \"lymphoma, mantle cell\"[MeSH Terms] AND \"humans\"[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal]

OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR
\"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date -
Publication]\", \"pubmed_count\": 0, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"marginal_zone_lymphoma\": {
\"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication
Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND (\\\"Lymphoma, B-Cell,
Marginal Zone\\\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR
\\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet
Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The
Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR
\\\"Leukemia\\\"[Journal] OR \\\"Journal of Hematology & Oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR
\\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of
Haematology\\\"[Journal] OR \\\"Experimental Hematology & Oncology\\\"[Journal] OR \\\"Bone Marrow
Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and
Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"Leukemia &
Lymphoma\\\"[Journal] OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in
Hematology\\\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal])\",
\"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase
ii\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND
\\\"lymphoma, b cell, marginal zone\\\"[MeSH Terms] AND \\\"humans\\\"[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR
\\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The
Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR
\\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR
\\\"Leukemia\\\"[Journal] OR \\\"journal of hematology oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR
\\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of
Haematology\\\"[Journal] OR \\\"experimental hematology oncology\\\"[Journal] OR \\\"Bone Marrow
Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and
Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"leukemia lymphoma\\\"[Journal]
OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR
\\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal]) AND 2026/06/18:2026/07/02[Date -
Publication]\", \"pubmed_count\": 0, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"hodgkin\": { \"pubmed_term\":
\"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication Type] OR
\\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND (\\\"Hodgkin Disease\\\"[MeSH
Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR
\\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA
Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR
\\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"Journal of Hematology
& Oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR
\\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"Experimental Hematology
& Oncology\\\"[Journal] OR \\\"Bone Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood
Reviews\\\"[Journal] OR \\\"Thrombosis and Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular
Therapy\\\"[Journal] OR \\\"Leukemia & Lymphoma\\\"[Journal] OR \\\"Annals of Hematology\\\"[Journal] OR
\\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR \\\"Research and Practice in Thrombosis and
Haemostasis\\\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR
\\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice
Guideline\\\"[Publication Type]) AND \\\"Hodgkin Disease\\\"[MeSH Terms] AND \\\"humans\\\"[MeSH Terms] AND (\\\"N Engl J
Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin
Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal]
OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR
\\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"journal of hematology oncology\\\"[Journal] OR \\\"J
Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal]
OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"experimental hematology oncology\\\"[Journal] OR \\\"Bone
Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and
Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"leukemia lymphoma\\\"[Journal]
OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR
\\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal]) AND 2026/06/18:2026/07/02[Date -
Publication]\", \"pubmed_count\": 1, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"multiple_myeloma\": {
\"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication
Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND (\\\"Multiple
Myeloma\\\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR
\\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet
Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The
Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR
\\\"Leukemia\\\"[Journal] OR \\\"Journal of Hematology & Oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR

\ "American Journal of Hematology\"[Journal] OR \ "Blood Cancer Journal\"[Journal] OR \ "British Journal of Haematology\"[Journal] OR \ "Experimental Hematology & Oncology\"[Journal] OR \ "Bone Marrow Transplantation\"[Journal] OR \ "HemaSphere\"[Journal] OR \ "Blood Reviews\"[Journal] OR \ "Thrombosis and Haemostasis\"[Journal] OR \ "Transplantation and Cellular Therapy\"[Journal] OR \ "Leukemia & Lymphoma\"[Journal] OR \ "Annals of Hematology\"[Journal] OR \ "Transfusion\"[Journal] OR \ "Seminars in Hematology\"[Journal] OR \ "Research and Practice in Thrombosis and Haemostasis\"[Journal])",
"pubmed_querytranslation": "(\\ \"clinical trial, phase iii\"[Publication Type] OR \\ \"clinical trial, phase ii\"[Publication Type] OR \\ \"Guideline\"[Publication Type] OR \\ \"Practice Guideline\"[Publication Type]) AND \\ \"Multiple Myeloma\"[MeSH Terms] AND \\ \"humans\"[MeSH Terms] AND (\\ \"N Engl J Med\"[Journal] OR \\ \"Lancet\"[Journal] OR \\ \"JAMA\"[Journal] OR \\ \"Nature Medicine\"[Journal] OR \\ \"J Clin Oncol\"[Journal] OR \\ \"The Lancet Oncology\"[Journal] OR \\ \"JAMA Oncology\"[Journal] OR \\ \"Ann Oncol\"[Journal] OR \\ \"Blood\"[Journal] OR \\ \"The Lancet Haematology\"[Journal] OR \\ \"Blood Advances\"[Journal] OR \\ \"Haematologica\"[Journal] OR \\ \"Leukemia\"[Journal] OR \\ \"Journal of hematology oncology\"[Journal] OR \\ \"J Thromb Haemost\"[Journal] OR \\ \"American Journal of Hematology\"[Journal] OR \\ \"Blood Cancer Journal\"[Journal] OR \\ \"British Journal of Haematology\"[Journal] OR \\ \"Experimental Hematology & Oncology\"[Journal] OR \\ \"Bone Marrow Transplantation\"[Journal] OR \\ \"HemaSphere\"[Journal] OR \\ \"Blood Reviews\"[Journal] OR \\ \"Thrombosis and Haemostasis\"[Journal] OR \\ \"Transplantation and Cellular Therapy\"[Journal] OR \\ \"leukemia lymphoma\"[Journal] OR \\ \"Annals of Hematology\"[Journal] OR \\ \"Transfusion\"[Journal] OR \\ \"Seminars in Hematology\"[Journal] OR \\ \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", "pubmed_count": 0, "window": ["2026-06-18", "2026-07-02"] }, "all": { "pubmed_term": "(\\ \"Clinical Trial, Phase III\"[Publication Type] OR \\ \"Clinical Trial, Phase II\"[Publication Type] OR \\ \"Guideline\"[Publication Type] OR \\ \"Practice Guideline\"[Publication Type]) AND (\\ \"Precursor Cell Lymphoblastic Leukemia-Lymphoma\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\ \"N Engl J Med\"[Journal] OR \\ \"Lancet\"[Journal] OR \\ \"JAMA\"[Journal] OR \\ \"Nature Medicine\"[Journal] OR \\ \"J Clin Oncol\"[Journal] OR \\ \"The Lancet Oncology\"[Journal] OR \\ \"JAMA Oncology\"[Journal] OR \\ \"Ann Oncol\"[Journal] OR \\ \"Blood\"[Journal] OR \\ \"The Lancet Haematology\"[Journal] OR \\ \"Blood Advances\"[Journal] OR \\ \"Haematologica\"[Journal] OR \\ \"Leukemia\"[Journal] OR \\ \"Journal of Hematology & Oncology\"[Journal] OR \\ \"J Thromb Haemost\"[Journal] OR \\ \"American Journal of Hematology\"[Journal] OR \\ \"Blood Cancer Journal\"[Journal] OR \\ \"British Journal of Haematology\"[Journal] OR \\ \"Experimental Hematology & Oncology\"[Journal] OR \\ \"Bone Marrow Transplantation\"[Journal] OR \\ \"HemaSphere\"[Journal] OR \\ \"Blood Reviews\"[Journal] OR \\ \"Thrombosis and Haemostasis\"[Journal] OR \\ \"Transplantation and Cellular Therapy\"[Journal] OR \\ \"Leukemia & Lymphoma\"[Journal] OR \\ \"Annals of Hematology\"[Journal] OR \\ \"Transfusion\"[Journal] OR \\ \"Seminars in Hematology\"[Journal] OR \\ \"Research and Practice in Thrombosis and Haemostasis\"[Journal])",
"pubmed_querytranslation": "(\\ \"clinical trial, phase iii\"[Publication Type] OR \\ \"clinical trial, phase ii\"[Publication Type] OR \\ \"Guideline\"[Publication Type] OR \\ \"Practice Guideline\"[Publication Type]) AND \\ \"Precursor Cell Lymphoblastic Leukemia-Lymphoma\"[MeSH Terms] AND \\ \"humans\"[MeSH Terms] AND (\\ \"N Engl J Med\"[Journal] OR \\ \"Lancet\"[Journal] OR \\ \"JAMA\"[Journal] OR \\ \"Nature Medicine\"[Journal] OR \\ \"J Clin Oncol\"[Journal] OR \\ \"The Lancet Oncology\"[Journal] OR \\ \"JAMA Oncology\"[Journal] OR \\ \"Ann Oncol\"[Journal] OR \\ \"Blood\"[Journal] OR \\ \"The Lancet Haematology\"[Journal] OR \\ \"Blood Advances\"[Journal] OR \\ \"Haematologica\"[Journal] OR \\ \"Leukemia\"[Journal] OR \\ \"journal of hematology oncology\"[Journal] OR \\ \"J Thromb Haemost\"[Journal] OR \\ \"American Journal of Hematology\"[Journal] OR \\ \"Blood Cancer Journal\"[Journal] OR \\ \"British Journal of Haematology\"[Journal] OR \\ \"experimental hematology oncology\"[Journal] OR \\ \"Bone Marrow Transplantation\"[Journal] OR \\ \"HemaSphere\"[Journal] OR \\ \"Blood Reviews\"[Journal] OR \\ \"Thrombosis and Haemostasis\"[Journal] OR \\ \"Transplantation and Cellular Therapy\"[Journal] OR \\ \"leukemia lymphoma\"[Journal] OR \\ \"Annals of Hematology\"[Journal] OR \\ \"Transfusion\"[Journal] OR \\ \"Seminars in Hematology\"[Journal] OR \\ \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", "pubmed_count": 0, "window": ["2026-06-18", "2026-07-02"] }, "cml": { "pubmed_term": "(\\ \"Clinical Trial, Phase III\"[Publication Type] OR \\ \"Clinical Trial, Phase II\"[Publication Type] OR \\ \"Guideline\"[Publication Type] OR \\ \"Practice Guideline\"[Publication Type]) AND (\\ \"Leukemia, Myelogenous, Chronic, BCR-ABL Positive\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\ \"N Engl J Med\"[Journal] OR \\ \"Lancet\"[Journal] OR \\ \"JAMA\"[Journal] OR \\ \"Nature Medicine\"[Journal] OR \\ \"J Clin Oncol\"[Journal] OR \\ \"The Lancet Oncology\"[Journal] OR \\ \"JAMA Oncology\"[Journal] OR \\ \"Ann Oncol\"[Journal] OR \\ \"Blood\"[Journal] OR \\ \"The Lancet Haematology\"[Journal] OR \\ \"Blood Advances\"[Journal] OR \\ \"Haematologica\"[Journal] OR \\ \"Leukemia\"[Journal] OR \\ \"journal of hematology oncology\"[Journal] OR \\ \"J Thromb Haemost\"[Journal] OR \\ \"American Journal of Hematology\"[Journal] OR \\ \"Blood Cancer Journal\"[Journal] OR \\ \"British Journal of Haematology\"[Journal] OR \\ \"Experimental Hematology & Oncology\"[Journal] OR \\ \"Bone Marrow Transplantation\"[Journal] OR \\ \"HemaSphere\"[Journal] OR \\ \"Blood Reviews\"[Journal] OR \\ \"Thrombosis and Haemostasis\"[Journal] OR \\ \"Transplantation and Cellular Therapy\"[Journal] OR \\ \"Leukemia & Lymphoma\"[Journal] OR \\ \"Annals of Hematology\"[Journal] OR \\ \"Transfusion\"[Journal] OR \\ \"Seminars in Hematology\"[Journal] OR \\ \"Research and Practice in Thrombosis and Haemostasis\"[Journal])",
"pubmed_querytranslation": "(\\ \"clinical trial, phase iii\"[Publication Type] OR \\ \"clinical trial, phase ii\"[Publication Type] OR \\ \"Guideline\"[Publication Type] OR \\ \"Practice Guideline\"[Publication Type]) AND \\ \"leukemia, myelogenous, chronic, bcr abl positive\"[MeSH Terms] AND \\ \"humans\"[MeSH Terms] AND (\\ \"N Engl J

Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", \"pubmed_count\": 0, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"aplastic_anemia\": { \"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND (\\\"Anemia, Aplastic\\\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"Journal of Hematology & Oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"Experimental Hematology & Oncology\\\"[Journal] OR \\\"Bone Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"Leukemia & Lymphoma\\\"[Journal] OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type]) AND \\\"anemia, aplastic\\\"[MeSH Terms] AND \\\"humans\\\"[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"journal of hematology oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"experimental hematology oncology\\\"[Journal] OR \\\"Bone Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"leukemia lymphoma\\\"[Journal] OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal]) AND 2026/06/18:2026/07/02[Date - Publication]\", \"pubmed_count\": 0, \"window\": [\"2026-06-18\", \"2026-07-02\"] }, \"chip\": { \"pubmed_term\": \"(\\\"Clinical Trial, Phase III\\\"[Publication Type] OR \\\"Clinical Trial, Phase II\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type] OR \\\"Observational Study\\\"[Publication Type] OR \\\"Comparative Study\\\"[Publication Type] OR \\\"Multicenter Study\\\"[Publication Type]) AND (\\\"Clonal Hematopoiesis\\\"[MeSH Terms]) AND humans[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"Journal of Hematology & Oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"Experimental Hematology & Oncology\\\"[Journal] OR \\\"Bone Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and Haemostasis\\\"[Journal] OR \\\"Transplantation and Cellular Therapy\\\"[Journal] OR \\\"Leukemia & Lymphoma\\\"[Journal] OR \\\"Annals of Hematology\\\"[Journal] OR \\\"Transfusion\\\"[Journal] OR \\\"Seminars in Hematology\\\"[Journal] OR \\\"Research and Practice in Thrombosis and Haemostasis\\\"[Journal])\", \"pubmed_querytranslation\": \"(\\\"clinical trial, phase iii\\\"[Publication Type] OR \\\"clinical trial, phase ii\\\"[Publication Type] OR \\\"Guideline\\\"[Publication Type] OR \\\"Practice Guideline\\\"[Publication Type] OR \\\"Observational Study\\\"[Publication Type] OR \\\"Comparative Study\\\"[Publication Type] OR \\\"Multicenter Study\\\"[Publication Type]) AND \\\"Clonal Hematopoiesis\\\"[MeSH Terms] AND \\\"humans\\\"[MeSH Terms] AND (\\\"N Engl J Med\\\"[Journal] OR \\\"Lancet\\\"[Journal] OR \\\"JAMA\\\"[Journal] OR \\\"Nature Medicine\\\"[Journal] OR \\\"J Clin Oncol\\\"[Journal] OR \\\"The Lancet Oncology\\\"[Journal] OR \\\"JAMA Oncology\\\"[Journal] OR \\\"Ann Oncol\\\"[Journal] OR \\\"Blood\\\"[Journal] OR \\\"The Lancet Haematology\\\"[Journal] OR \\\"Blood Advances\\\"[Journal] OR \\\"Haematologica\\\"[Journal] OR \\\"Leukemia\\\"[Journal] OR \\\"journal of hematology oncology\\\"[Journal] OR \\\"J Thromb Haemost\\\"[Journal] OR \\\"American Journal of Hematology\\\"[Journal] OR \\\"Blood Cancer Journal\\\"[Journal] OR \\\"British Journal of Haematology\\\"[Journal] OR \\\"experimental hematology oncology\\\"[Journal] OR \\\"Bone Marrow Transplantation\\\"[Journal] OR \\\"HemaSphere\\\"[Journal] OR \\\"Blood Reviews\\\"[Journal] OR \\\"Thrombosis and

```

Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal]
OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR
\"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/02:2026/07/02[Date -
Publication]\", \"pubmed_count\": 0, \"window\": [ \"2026-06-02\", \"2026-07-02\" ] }, \"sickle_cell\": { \"pubmed_term\":
\"(\"Clinical Trial, Phase III\"[Publication Type] OR \"Clinical Trial, Phase II\"[Publication Type] OR
\"Guideline\"[Publication Type] OR \"Practice Guideline\"[Publication Type] OR \"Clinical Trial\"[Publication
Type] OR \"Observational Study\"[Publication Type]) AND (\"Anemia, Sickle Cell\"[MeSH Terms]) AND humans[MeSH
Terms] AND (\"N Engl J Med\"[Journal] OR \"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature
Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The Lancet Oncology\"[Journal] OR \"JAMA
Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR \"The Lancet Haematology\"[Journal] OR
\"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR \"Leukemia\"[Journal] OR \"Journal of Hematology
& Oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR \"American Journal of Hematology\"[Journal] OR
\"Blood Cancer Journal\"[Journal] OR \"British Journal of Haematology\"[Journal] OR \"Experimental Hematology
& Oncology\"[Journal] OR \"Bone Marrow Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood
Reviews\"[Journal] OR \"Thrombosis and Haemostasis\"[Journal] OR \"Transplantation and Cellular
Therapy\"[Journal] OR \"Leukemia & Lymphoma\"[Journal] OR \"Annals of Hematology\"[Journal] OR
\"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR \"Research and Practice in Thrombosis and
Haemostasis\"[Journal])\", \"pubmed_querytranslation\": \"(\"clinical trial, phase iii\"[Publication Type] OR
\"clinical trial, phase ii\"[Publication Type] OR \"Guideline\"[Publication Type] OR \"Practice
Guideline\"[Publication Type] OR \"Clinical Trial\"[Publication Type] OR \"Observational Study\"[Publication
Type]) AND \"anemia, sickle cell\"[MeSH Terms] AND \"humans\"[MeSH Terms] AND (\"N Engl J Med\"[Journal] OR
\"Lancet\"[Journal] OR \"JAMA\"[Journal] OR \"Nature Medicine\"[Journal] OR \"J Clin Oncol\"[Journal] OR \"The
Lancet Oncology\"[Journal] OR \"JAMA Oncology\"[Journal] OR \"Ann Oncol\"[Journal] OR \"Blood\"[Journal] OR
\"The Lancet Haematology\"[Journal] OR \"Blood Advances\"[Journal] OR \"Haematologica\"[Journal] OR
\"Leukemia\"[Journal] OR \"journal of hematology oncology\"[Journal] OR \"J Thromb Haemost\"[Journal] OR
\"American Journal of Hematology\"[Journal] OR \"Blood Cancer Journal\"[Journal] OR \"British Journal of
Haematology\"[Journal] OR \"experimental hematology oncology\"[Journal] OR \"Bone Marrow
Transplantation\"[Journal] OR \"HemaSphere\"[Journal] OR \"Blood Reviews\"[Journal] OR \"Thrombosis and
Haemostasis\"[Journal] OR \"Transplantation and Cellular Therapy\"[Journal] OR \"leukemia lymphoma\"[Journal]
OR \"Annals of Hematology\"[Journal] OR \"Transfusion\"[Journal] OR \"Seminars in Hematology\"[Journal] OR
\"Research and Practice in Thrombosis and Haemostasis\"[Journal]) AND 2026/06/18:2026/07/02[Date -
Publication]\", \"pubmed_count\": 4, \"window\": [ \"2026-06-18\", \"2026-07-02\" ] } }

```

"Surveillance aid only. Verify every item against the primary source and apply clinical judgment and institutional protocols before any clinical use. Automated extraction can misread numbers, endpoints, and populations."