

Krylov Linear Solvers and Quasi Monte Carlo Methods for Transport Simulations

Sam Pasmann,^a C. T. Kelley,^b and Ryan McClarren^{*,a}

*^aDepartment of Aerospace and Mechanical Engineering
University of Notre Dame
Fitzpatrick Hall, Notre Dame, IN 46556*

*^bNorth Carolina State University, Department of Mathematics
3234 SAS Hall, Box 8205
Raleigh NC 27695-8205*

*Email: rmcclarr@nd.edu

Number of pages: 7

Number of tables: 2

Number of figures: 2

Abstract

QMC + Krylov

Keywords — Quasi Monte Carlo Methods, Krylov Linear Solvers

I. INTRODUCTION

II. COMPUTATIONAL RESULTS

In this section we consider an example from [1]. The formulation of the transport problem is taken from [2]. The equation for the angular flux ψ is

$$\mu \frac{\partial \psi}{\partial x}(x, \mu) + \Sigma_t(x) \psi(x, \mu) = \frac{1}{2} \left[\Sigma_s(x) \int_{-1}^1 \psi(x, \mu') d\mu' + q(x) \right] \text{ for } 0 \leq x \leq \tau \quad (1)$$

The boundary conditions are

$$\psi(0, \mu) = \psi_l(\mu), \mu > 0; \psi(\tau, \mu) = \psi_r(\mu), \mu < 0.$$

In the computations we use

$$\tau = 5, \Sigma_s(x) = \omega_0 e^{-x/s}, \Sigma_t(x) = 1, q(x) = 0, \psi_l(\mu) = 1, \psi_r(\mu) = 0,$$

and consider two cases $s = 1$ and $s = \infty$

II.A. Source Iteration and Linear Solvers

II.B. QMC and Krylov Linear Solvers

We use two krylov methods [3], GMRES [4] and Bi-CGSTAB [5].

The linear and nonlinear solvers come from the Julia package [SIAMFANLEQ.jl](#) [6]. The documentation for these codes is in the [Juila notebooks](#) [7] and the book [8] that accompany the package.

I can solve the QMC linear problem with both Krylov methods now and, just like the classical case, I'm seeing fewer than half of the number of transport sweeps. Herewith the results for $N=1000$ and $N_x=100$.

II.C. Validation and calibration study

We conclude this section with a validation study. We compare the QMC results with the results from [1]. The results in [1] are exit distributions and are accurate to six figures. We have




Fig. 2. $s = \infty$

duplicated those results with an Sn computation on a fine angular and spatial mesh.

Sam, Ryan, should we use more or different values of N and Nx ?

For $N = 1000$ and $Nx = 100$ we obtain the cell-average fluxes from the QMC approximation. We then use a single Sn transport sweep to recover the exit distributions from the QMC cell-average fluxes. We report the results and the corresponding results from [1] in Tables I and II.

The exit distributions, as is clear from Table I can vary by five orders of magnitude. Even so, the results from QMC agree with the benchmarks to roughly two figures.

III. CONCLUSION

TABLE I
Exit Distributiions: $s = 1$

μ	Garcia/Siewert		QMC	
	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$
5.00e-02	5.89664e-01	6.07488e-06	5.71197e-01	5.85487e-06
1.00e-01	5.31120e-01	6.92516e-06	5.22137e-01	6.66741e-06
2.00e-01	4.43280e-01	9.64232e-06	4.41567e-01	9.25261e-06
3.00e-01	3.80306e-01	1.62339e-05	3.81029e-01	1.54416e-05
4.00e-01	3.32964e-01	4.38580e-05	3.34673e-01	4.09691e-05
5.00e-01	2.96090e-01	1.69372e-04	2.98224e-01	1.57373e-04
6.00e-01	2.66563e-01	5.73465e-04	2.68871e-01	5.35989e-04
7.00e-01	2.42390e-01	1.51282e-03	2.44749e-01	1.42448e-03
8.00e-01	2.22235e-01	3.24369e-03	2.24583e-01	3.07431e-03
9.00e-01	2.05174e-01	5.96036e-03	2.07478e-01	5.67991e-03
1.00e+00	1.90546e-01	9.77123e-03	1.92789e-01	9.35351e-03

TABLE II
Exit Distributiions: $s = \infty$

μ	Garcia/Siewert		QMC	
	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$
5.00e-02	8.97798e-01	1.02202e-01	8.47454e-01	1.00663e-01
1.00e-01	8.87836e-01	1.12164e-01	8.52822e-01	1.10325e-01
2.00e-01	8.69581e-01	1.30419e-01	8.47710e-01	1.29064e-01
3.00e-01	8.52299e-01	1.47701e-01	8.35879e-01	1.46849e-01
4.00e-01	8.35503e-01	1.64497e-01	8.22291e-01	1.64034e-01
5.00e-01	8.18996e-01	1.81004e-01	8.08044e-01	1.80827e-01
6.00e-01	8.02676e-01	1.97324e-01	7.93459e-01	1.97336e-01
7.00e-01	7.86493e-01	2.13507e-01	7.78672e-01	2.13625e-01
8.00e-01	7.70429e-01	2.29571e-01	7.63768e-01	2.29725e-01
9.00e-01	7.54496e-01	2.45504e-01	7.48818e-01	2.45642e-01
1.00e+00	7.38721e-01	2.61279e-01	7.33889e-01	2.61361e-01

ACKNOWLEDGMENTS

The research of CTK was supported by Department of Energy grant DE-NA003967, and National Science Foundation Grants DMS-1745654, and DMS-1906446.

REFERENCES

- [1] R. GARCIA and C. SIEWERT, “Radiative transfer in finite inhomogeneous plane-parallel atmospheres,” *J. Quant. Spectrosc. Radiat. Transfer*, **27**, 141 (1982).
- [2] J. WILLERT, C. T. KELLEY, D. A. KNOLL, and H. K. PARK, “Hybrid Deterministic/Monte Carlo Neutronics,” *SIAM J. Sci. Comp.*, **35**, S62 (2013).
- [3] C. T. KELLEY, *Iterative Methods for Linear and Nonlinear Equations*, no. 16 in Frontiers in Applied Mathematics, SIAM, Philadelphia (1995).
- [4] Y. SAAD and M. SCHULTZ, “GMRES a generalized minimal residual algorithm for solving nonsymmetric linear systems,” *SIAM J. Sci. Stat. Comp.*, **7**, 856 (1986).
- [5] H. A. VAN DER VORST, “Bi-CGSTAB: A fast and smoothly converging variant to Bi-CG for the solution of nonsymmetric systems,” *SIAM J. Sci. Statist. Comput.*, **13**, 631 (1992).
- [6] C. T. KELLEY, “SIAMFANLEquations.jl,” <https://github.com/ctkelley/SIAMFANLEquations.jl> (2020); 10.5281/zenodo.4284807., URL <https://github.com/ctkelley/SIAMFANLEquations.jl>, julia Package.
- [7] C. T. KELLEY, “Notebook for Solving Nonlinear Equations with Iterative Methods: Solvers and Examples in Julia,” <https://github.com/ctkelley/NotebookSIAMFANL> (2020); 10.5281/zenodo.4284687., URL <https://github.com/ctkelley/NotebookSIAMFANL>, iJulia Notebook.
- [8] C. T. KELLEY, “Solving Nonlinear Equations with Iterative Methods: Solvers and Examples in Julia,” (2020) Unpublished book ms, under contract with SIAM.