

Krylov Linear Solvers and Quasi Monte Carlo Methods for Transport Simulations

C. T. Kelley; Ryan McClarren, Sam Pasmann, Ilham Variansyah

The Center for Exascale Monte Carlo Neutron Transport (CEMeNT)
North Carolina State University and University of Notre Dame

September 30, 2021



NC STATE



UNIVERSITY OF
NOTRE DAME



Oregon State
University

1. Our Place in Project Roadmap
2. Model Problem and QMC
 - Model Problem
 - Formulation as Linear System
 - QMC overview
3. Preliminary Results
4. Conclusions
5. Appendix
 - Links
 - References

Our Place in Project Roadmap

CEMENT

$$\begin{aligned}\mu \frac{\partial \psi}{\partial x}(x, \mu) + \Sigma_t(x) \psi(x, \mu) &= \frac{1}{2} \left[\Sigma_s(x) \int_{-1}^1 \psi(x, \mu') d\mu' + q(x) \right] \\ &\equiv \frac{1}{2} (\phi(x) + q(x)),\end{aligned}$$

where $0 \leq x \leq \tau$ and

$$\phi(x) = \int_{-1}^1 \psi(x, \mu') d\mu'.$$

Subject to boundary conditions

$$\psi(0, \mu) = \psi_l(\mu), \mu > 0; \psi(\tau, \mu) = \psi_r(\mu), \mu < 0.$$

Given an iteration n and $\phi_n(x)$ Compute ϕ_{n+1} by solving the fixed source problem

$$\mu \frac{\partial \psi_{n+1}}{\partial x}(x, \mu) + \Sigma_t(x) \psi(x, \mu) = \frac{1}{2} [\Sigma_s(x) \phi_n(x) + q(x)],$$

and then compute

$$\phi_{n+1}(x) = \int_{-1}^1 \psi_{n+1}(x, \mu') d\mu' \equiv \mathcal{S}(\phi_n, q, \psi_l, \psi_r)$$

Source iteration is Picard iteration for the fixed point problem

$$\phi = \mathcal{S}(\phi, q, \psi_l, \psi_r)$$

To use other solvers we must convert to a linear system via

$$\mathcal{K}(\phi) = \mathcal{S}(\phi, 0, 0, 0) \text{ and } f = \mathcal{S}(0, q, \psi_l, \psi_r)$$

to get



$$A\phi \equiv (I - \mathcal{K})\phi = f,$$

which we can send to a linear solver.

- Evaluating the Source iteration map
 - Conventional deterministic methods (SN ...)
 - Monte Carlo
 - Quasi Monte Carlo (this talk)
- Solving the linear problem
 - Source (Picard) iteration (in most codes)
 - Krylov methods: matrix-free
 - needs **CONSISTENT** evaluation of \mathcal{S} to get $A\phi$
MC has trouble with this
 - but no matrix assembly

- Initialization
 - \mathbf{x}_0 : initial iterate
 - $\mathbf{r}_0 = \mathbf{b}_0 - \mathbf{Ax}_0$: initial residual
- Find iteration \mathbf{x}_k in
 $\mathcal{K}_k = \text{span}(\mathbf{r}_0, \mathbf{Ar}_0, \dots, \mathbf{A}^{k-1}\mathbf{r}_0)$
- Example GMRES (Sadd/Schultz 1986): \mathbf{x}_k minimizes $\|\mathbf{b} - \mathbf{Ax}\|$ over \mathcal{K}_k
- All is well if matrix-vector product is consistent.
 - \mathbf{Ax} returns the same result with every call.
 - MC is not deterministic and hence not consistent.

QMC (red) and MC (blue)



A few facts

- MC and QMC are continuous in space/angle/energy
- With N samples/particles
 - QMC error $O(1/N)$
 - MC variance $O(1/\sqrt{N})$
- MC randomness
 - can break Krylov solvers
 - Willert et al 2013, 2014, 2015
 - Simoncini et al, 2003
- QMC gives consistent matrix-vector products

- All codes in Julia on github
<https://github.com/ctkelley/NDA-QMC.jl>
- Model problem from Garcia/Siewert 1982

$$\tau = 5, \Sigma_s(x) = \omega_0 e^{-x/s}, \Sigma_t(x) = 1, q(x) = 0, \psi_l(\mu) = 1, \psi_r(\mu) = 0.$$

- Two cases: $s = 1$ (easy) and $s = \infty$ (hard)
- Evaluate source iteration map with QMC
- Compare GMRES and Bi-CGSTAB (van der Vorst 1992) to Picard
- Validation with Garcia/Siewert exit distributions
 - QMC: N samples, N_x cells
 - Two figure accuracy with $N = 1000$ and $N_x = 100$


Validation: Easy case, $N = 1000$, $N_x = 100$

μ	Garcia/Siewert		QMC	
	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$
5.00e-02	5.89664e-01	6.07488e-06	5.71197e-01	5.85487e-06
1.00e-01	5.31120e-01	6.92516e-06	5.22137e-01	6.66741e-06
2.00e-01	4.43280e-01	9.64232e-06	4.41567e-01	9.25261e-06
3.00e-01	3.80306e-01	1.62339e-05	3.81029e-01	1.54416e-05
4.00e-01	3.32964e-01	4.38580e-05	3.34673e-01	4.09691e-05
5.00e-01	2.96090e-01	1.69372e-04	2.98224e-01	1.57373e-04
6.00e-01	2.66563e-01	5.73465e-04	2.68871e-01	5.35989e-04
7.00e-01	2.42390e-01	1.51282e-03	2.44749e-01	1.42448e-03
8.00e-01	2.22235e-01	3.24369e-03	2.24583e-01	3.07431e-03
9.00e-01	2.05174e-01	5.96036e-03	2.07478e-01	5.67991e-03
1.00e+00	1.90546e-01	9.77123e-03	1.92789e-01	9.35351e-03


Validation: Hard case, $N = 1000$, $N_x = 100$

μ	Garcia/Siewert		QMC	
	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$	$\psi(0, -\mu)$	$\psi(\tau, -\mu)$
5.00e-02	8.97798e-01	1.02202e-01	8.47454e-01	1.00663e-01
1.00e-01	8.87836e-01	1.12164e-01	8.52822e-01	1.10325e-01
2.00e-01	8.69581e-01	1.30419e-01	8.47710e-01	1.29064e-01
3.00e-01	8.52299e-01	1.47701e-01	8.35879e-01	1.46849e-01
4.00e-01	8.35503e-01	1.64497e-01	8.22291e-01	1.64034e-01
5.00e-01	8.18996e-01	1.81004e-01	8.08044e-01	1.80827e-01
6.00e-01	8.02676e-01	1.97324e-01	7.93459e-01	1.97336e-01
7.00e-01	7.86493e-01	2.13507e-01	7.78672e-01	2.13625e-01
8.00e-01	7.70429e-01	2.29571e-01	7.63768e-01	2.29725e-01
9.00e-01	7.54496e-01	2.45504e-01	7.48818e-01	2.45642e-01
1.00e+00	7.38721e-01	2.61279e-01	7.33889e-01	2.61361e-01

Convergence: Easy Case



Convergence: Hard Case



- QMC for neutron transport simulations
 - Completely deterministic
 - Avoids problems with MC and Krylov methods
 - Krylov solvers very promising
- Test on benchmark problem
- Next up: NDA

- Jupyter notebook draft of paper:
https://github.com/ctkelley/NDA_QMC.jl
- Julia package: <https://github.com/ctkelley/SIAMFANLEquations.jl>
- Jupyter notebook draft of solver book:
<https://github.com/ctkelley/NotebookSIAMFANL>

- SAMUEL PASMANN, ILHAM VARIANSYAH, AND RYAN G. MCCLARREN, *Convergent Transport Source Iteration Calculations with Quasi-Monte Carlo*, in 2021 ANS Virtual Annual Meeting, 2021, pp. 192–195.
- R.D.M. GARCIA AND C.E. SIEWERT, *Radiative transfer in finite inhomogeneous plane-parallel atmospheres*, J. Quant. Spectrosc. Radiat. Transfer, 27 (1982), pp. 141–148.
- C. T. KELLEY, *Solving Nonlinear Equations with Iterative Methods: Solvers and Examples in Julia*, 2020. Unpublished book ms, under contract with SIAM.
- C. T. KELLEY, *Notebook for Solving Nonlinear Equations with Iterative Methods: Solvers and Examples in Julia*. <https://github.com/ctkelley/NotebookSIAMFANL>, 2020.

- J. WILLERT, XIAOJUN CHEN, AND C. T. KELLEY, *Newton's method for Monte Carlo-based residuals*, SIAM J. Numer. Anal., 53 (2015), pp. 1738–1757.
- J. WILLERT, C. T. KELLEY, D. A. KNOLL, AND H. K. PARK, *Hybrid deterministic/Monte Carlo neutronics*, SIAM J. Sci. Comp., 35 (2013), pp. S62–S83.
- J. WILLERT, C. T. KELLEY, D. A. KNOLL, AND H. K. PARK, *A hybrid deterministic/Monte Carlo method for solving the k -eigenvalue problem with a comparison to analog Monte Carlo solutions*, J. Comp. Th. Transport, 43 (2014), pp. 50–67.
- V. SIMONCINI AND D. B. SZYLD, *Theory of inexact Krylov subspace methods and applications to scientific computing*, SIAM J. Sci. Comput., 25 (2003), pp. 454–477.

- C. T. KELLEY, *SIAMFANLEquations.jl*. <https://github.com/ctkelley/SIAMFANLEquations.jl>, 2020. Julia Package.
- C. T. KELLEY, *Iterative Methods for Linear and Nonlinear Equations*, no. 16 in Frontiers in Applied Mathematics, SIAM, Philadelphia, 1995.
- C. T. KELLEY, *Solving Nonlinear Equations with Newton's Method*, no. 1 in Fundamentals of Algorithms, SIAM, Philadelphia, 2003.
- Y. SAAD AND M.H. SCHULTZ, *GMRES a generalized minimal residual algorithm for solving nonsymmetric linear systems*, SIAM J. Sci. Stat. Comp., 7 (1986), pp. 856–869.
- H. A. VAN DER VORST, *Bi-CGSTAB: A fast and smoothly converging variant to Bi-CG for the solution of nonsymmetric systems*, SIAM J. Sci. Statist. Comput., 13 (1992), pp. 631–644.