



Unsecured attractants, collisions, and high mortality strain coexistence between grizzly bears and people in the Elk Valley, southeast British Columbia

Journal:	<i>Journal of Wildlife Management and Wildlife Monographs</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Lamb, Clayton; Biodiversity Pathways Ltd. Smit, Laura; Ministry of Forests, Lands, Natural Resource Operations and Rural Development, Research Program Mowat, Garth; Ministry of Forests, Lands, Natural Resource Operations and Rural Development, Research Program McLellan, Bruce; Ministry of Forests, Lands & Natural Resource Operations, Proctor, Michael
Keywords:	carnivore, demography, genetic capture recapture, reproduction, roadkill, Ursus
Abstract:	Historical persecution of grizzly bears in North America reduced the species range by 55%. Today, dedicated recovery efforts and shifting societal perceptions have supported the recovery and expansion of grizzly bear populations in many areas. With increasing overlap between people and bears, conservation actions and scientific inquiry are now shifting efforts towards supporting coexistence with bears. Here we assessed the demography and behaviour of grizzly bears in a coexistence landscape in southeast British Columbia, Canada, where abundant grizzly bear populations occur among busy, human-settled valleys. Between 2016 and 2022 we captured 76 individual grizzly bears and monitored their conflict behaviour, survival, and reproduction for 160 animal-years. The cause of death for fourteen animals with a functioning collar was human-wildlife conflict (n=6), road or rail collision (n=6), unknown but human suspected (n=1), and natural (n=1). Subadult survival was the lowest recorded in North America, while adult survival was similar to other studies, suggesting an intense demographic filter for young animals. We estimate that human-caused mortality is underreported in government databases by 65%, or for every recorded mortality there are ~2 that go unreported. Reporting was especially low for road and rail mortalities. Grizzly bear mortality in the Elk Valley due to collisions and conflicts with people is an order of magnitude greater than elsewhere in British Columbia. Combining DNA- and collar-based estimates of population growth we show that grizzly bear abundance is stable due to source-sink dynamics, whereby ~7 immigrant bears per year offset the high mortality rates in the area. Grizzly bears dispersing into the valley are often young and more conflict-naïve, creating a

conflict spiral that can be interrupted by reducing mortality of young animals. Creating a self-sustaining population of bears within the study area will require targeted efforts to reduce or secure attractants on private property and strategies to minimize collisions with trains and vehicles.

SCHOLARONE™
Manuscripts

1 30 Nov 2022

2 RH: Lamb et al. • Straining Coexistence

3 **Unsecured attractants, collisions, and high mortality strain coexistence between grizzly**
4 **bears and people in the Elk Valley, southeast British Columbia**

5 Clayton T. Lamb, Biodiversity Pathways Ltd., Department of Biology, Kelowna, BC, Canada

6 V1V 1V7

7 Laura Smit, Ministry of Forests, Fish and Wildlife Branch, Nelson, BC, Canada V1L 4K3

8 Garth Mowat, Ministry of Forests, Fish and Wildlife Branch, Nelson, BC, Canada V1L 4K3 and
9 University of British Columbia, Department of Earth, Environmental and Geographic
10 Sciences, Kelowna, BC

11 Bruce McLellan, International Union for the Conservation of Nature Bear Specialist Group,
12 D'Arcy, BC, Canada V0N 1L0

13 Michael Proctor, International Union for the Conservation of Nature Bear Specialist Group,
14 Kaslo, BC, Canada V0G 1M0;

15

16 **Correspondence:** Clayton T. Lamb Biodiversity Pathways Ltd., Department of Biology,
17 Kelowna, BC, Canada V1V 1V7. Email: ctlamb@ualberta.ca

18

19 **ABSTRACT** Historical persecution of grizzly bears in North America reduced the species range
20 by 55%. Today, dedicated recovery efforts and shifting societal perceptions have supported the
21 recovery and expansion of grizzly bear populations in many areas. With increasing overlap
22 between people and bears, conservation actions and scientific inquiry are now shifting efforts
23 towards supporting coexistence with bears. Here we assessed the demography and behaviour of

24 grizzly bears in a coexistence landscape in southeast British Columbia, Canada, where abundant
25 grizzly bear populations occur among busy, human-settled valleys. Between 2016 and 2022 we
26 captured 76 individual grizzly bears and monitored their conflict behaviour, survival, and
27 reproduction for 160 animal-years. The cause of death for fourteen animals with a functioning
28 collar was human-wildlife conflict (n=6), road or rail collision (n=6), unknown but human
29 suspected (n=1), and natural (n=1). Subadult survival was the lowest recorded in North America,
30 while adult survival was similar to other studies, suggesting an intense demographic filter for
31 young animals. We estimate that human-caused mortality is underreported in government
32 databases by 65%, or for every recorded mortality there are ~2 that go unreported. Reporting was
33 especially low for road and rail mortalities. Grizzly bear mortality in the Elk Valley due to
34 collisions and conflicts with people is an order of magnitude greater than elsewhere in British
35 Columbia. Combining DNA- and collar-based estimates of population growth we show that
36 grizzly bear abundance is stable due to source-sink dynamics, whereby ~7 immigrant bears per
37 year offset the high mortality rates in the area. Grizzly bears dispersing into the valley are often
38 young and more conflict-naïve, creating a conflict spiral that can be interrupted by reducing
39 mortality of young animals. Creating a self-sustaining population of bears within the study area
40 will require targeted efforts to reduce or secure attractants on private property and strategies to
41 minimize collisions with trains and vehicles.

42 **KEYWORDS** carnivore, demography, genetic capture recapture, reproduction, roadkill, *Ursus*

43 During the early to mid-twentieth century, grizzly bear (*Ursus arctos*) populations were
44 dramatically reduced in North America (Mattson and Merrill 2002). The species was considered
45 a 'dangerous impediment to progress' by many settlers (Peek et al. 2003), and due to shooting,
46 trapping, and poisoning across much of the continent, the species range contracted by 53%

47 (Laliberte and Ripple 2004). As the environmental movement grew in the second half of the
48 twentieth century and societal views of peoples' place in nature shifted from a perspective of
49 dominion to mutualism (Manfredo et al. 2020), the persecution of grizzly bears slowed
50 (Bruskotter et al. 2017). In 1975 after a century of persecution, the grizzly bear was listed as a
51 threatened species in the contiguous United States under the Endangered Species Act. For over
52 thirty years, efforts have been made to reduce human-caused mortality of grizzly bears and
53 increase population connectivity in the United States and Canada (Schwartz et al. 2006,
54 Hebblewhite et al. 2022). For example, significant changes to policy and regulation in British
55 Columbia between 1964 and 1996 restricted the hunter kill and secured persistent attractants
56 such as open garbage dumps, reflecting the shifting public attitudes towards grizzly bears.
57 Several populations have since recovered, some of which were once small, isolated, and in peril.
58 Grizzly bear populations are now increasing in many areas in and around the defined US
59 Recovery Zones in four U.S states, and in portions of southern Canada, such as in central British
60 Columbia (McLellan 1989, Apps et al. 2014, Lamb et al. 2018, Hatter et al. 2018, McLellan et
61 al. 2021) or expanding eastward into portions of their historic range in Alberta (Morehouse and
62 Boyce 2016). The grizzly bear population in the Greater Yellowstone Ecosystem that was
63 estimated at 175 individuals in 1975 has since increased 5-fold, and more than 750 grizzlies now
64 range into landscapes that have been dramatically transformed by people since the animals last
65 walked there a century ago (Schwartz et al. 2006). The Yellowstone example highlights a
66 situation that is unfolding across much of the southern distribution of grizzly bears in both
67 Canada and the United States; successful conservation efforts have allowed the species to
68 increase in their current range and expand into portions of their historic range. During this range
69 recolonization grizzly bears are dispersing across, or living in, human-dominated landscapes,

70 ushering in a new era of large carnivore conservation focused better understanding human-bear
71 interactions and applying innovative programs to support both parties and promote coexistence
72 (Morehouse and Boyce 2016, Proctor et al. 2018, Morehouse et al. 2020).

73 Coexistence between people and wildlife is a state where both exist in shared landscapes
74 and conduct activities necessary to life within tolerable levels of risk (Frank 2016, Lute and
75 Carter 2020). Importantly, the version of coexistence that we subscribe to does not imply the
76 situation is always peaceful, rather the situation needs to be at least demographically sustainable
77 and without excessive burdens on either party (Lamb et al. 2020). While this seems like an
78 achievable goal, plentiful conflicts still occur between grizzly bears and people as bear
79 populations increase and expand, challenging the viability of coexistence when bears pose risks
80 to human safety and property. In response, some grizzly bears have altered their behaviour to
81 more nocturnal patterns to avoid conflicts with people and associated mortality. Despite this
82 behavioural avoidance of risk, grizzly bear populations in most human-dominated landscapes are
83 not self-sustaining. Due to high mortality rates the presence of bears in human-dominated
84 landscapes is reliant on immigration from less disturbed areas (Lamb et al. 2020). In these
85 emerging landscapes of coexistence, the viability of coexistence teeters on our ability to provide
86 the necessary tools to keep people and their property safe while allowing bears to move across
87 landscapes, survive, and reproduce at rates that support stable populations.

88 The southeast corner of British Columbia, Canada, is a landscape that presents both
89 opportunity and challenges for human-bear coexistence. Here abundant grizzly bear populations
90 occur among busy, human-settled valleys. While the area is perhaps unsurprisingly a hotspot of
91 human-bear conflicts, the persistence of grizzly bears suggests there is much to learn about how
92 grizzly bears coexist with transportation corridors, towns, intensive resource extraction,

93 agriculture, and expanding recreation. Previous investigations into the demography of grizzly
94 bears in the Elk Valley of southeast British Columbia used composite metrics of growth derived
95 from DNA capture-recapture data and revealed high mortality rates were contributing to source-
96 sink dynamics (Lamb et al. 2017). However, because DNA data does not provide known fates or
97 age-related information, the specific demographic mechanisms facilitating persistence remained
98 hazy. Here we sought to understand the demographics of individual grizzly bears in the Elk
99 Valley, identify what was killing them, and determine whether those mortalities were being
100 reported. At the population level, we investigate the number of immigrant bears currently
101 subsidizing the persistence of bears in the Elk Valley and ultimately propose solutions to
102 operationalize coexistence between people and grizzly bears.

103

104 **STUDY AREA**

105 The 5,073 km² study area is in the Rocky Mountains of southeast British Columbia, Canada
106 (Figure 1). We initially defined a general study area based on the ecological trap area in Lamb et
107 al. (2017) to guide collaring efforts but then refined the area post-hoc as the 99th percentile of a
108 utilization distribution generated by pooling locations from all collared grizzly bears. We refer to
109 the study area as the “Elk Valley” although the upper headwaters of the Elk River are not
110 included (Figure 1). The study area stands out as a unique area of grizzly bear coexistence and
111 conflict due to the moderate density of grizzly bears (15-56 grizzly bears / 1,000 km² (McLellan
112 2015, Lamb et al. 2019)) living in close proximity to three towns of >5,000 people each, a
113 highway with >10,000 vehicles per day, an active railway, five large open pit coal mines, and
114 abundant recreation including off-road vehicle use, mountain biking, hiking, hunting, and fishing
115 occurring across the landscape.

116

117 **METHODS**118 **Capture, handling, and collaring**

119 Grizzly bears were captured using multiple methods throughout their active season (April to
120 November) between 2016 and 2022. Some bears were darted from a helicopter, but this method
121 was not viable in all portions of the study area due to human settlement in the valley bottom. In
122 more human-dominated areas, we captured bears in culvert traps and leg-restraining snares,
123 which allowed us to choose captures sites based on safety concerns. Our capture effort was
124 primarily directed toward the valley bottom and tributaries of the Elk Valley and therefore our
125 inference primarily pertains to the areas that correspond to the clusters of telemetry locations
126 (Figure 1).

127 Bears involved in human-wildlife conflict were sometimes captured by members of the
128 British Columbia Conservation Officer Service (COS). When their capture did not end in
129 euthanasia, we often collared these animals and included them in our sample. Although other
130 studies have separated the demography of conflict bears from the study population, at least until
131 a conflict bear is captured in a research trap and becomes a research animal for the rest of its life
132 (Schwartz et al. 2006), we chose to pool all captured animals together. Unlike other studies that
133 captured bears across large areas, both near and far from human settlements, our study focussed
134 on bears in human-dominated landscapes and thus all the bears in our sample were at least
135 potentially conflict animals. None of the bears first captured by the COS died while collared so
136 their inclusion did not appear to bias our sample.

137 We used Vectronic VERTEX Lite collars (VECTRONIC Aerospace, Berlin, Germany)
138 and Followit Geos collars (Followit AB, Lindesberg, Sweden), each of which took between 1

139 and 12 relocations a day and was equipped with a VHF beacon for real time manual relocation.
140 All collars were fitted with a cotton belt break away of varying thickness that was designed to rot
141 within 1-5 years. In addition to the cotton belt break away, most collars were equipped with a
142 remote blow off within the collar that was pre-programmed to activate within 2-4 years
143 (depending on the bear's age) that could also be activated remotely by satellite at any time. We
144 provide additional details on traps, drug information, and handling procedures in Supporting
145 Information A.

146

147 **Demographic monitoring**

148 Mortality was primarily monitored via a 12-hour inactivity switch within the collars. In addition,
149 we opportunistically recovered dead ear-tagged animals that were no longer collared. We
150 generally responded to mortality notifications from collars within 12 hours. Cause of death was
151 often apparent (for example where the carcass was on a railway or highway with excessive blunt-
152 force trauma), but it was ascertained by necropsy when cause was less clear. We assessed animal
153 body condition either subjectively or measured accumulated fat depth over the rump. We
154 censored collars that rotted off, were blown off, or failed. Early in the study we had poor collar
155 performance and the GPS and VHF transmitters often stopped working prematurely. We
156 assessed the potential for these instances to be mortalities and the collar destroyed (i.e., cryptic
157 poaching of collared animals) by collating known outcomes determined through other means
158 such as DNA sampling, subsequent live capture, or confirmed mortality after collar failure. All
159 grizzly bears killed by people must be reported to a wildlife officer in BC. During this
160 compulsory inspection, data and samples are collected and the data is stored in the BC

161 Compulsory Inspection (CI) database. We genotyped all CI samples as part of a larger genetic
162 monitoring program in this area (Mowat et al. 2020).

163 We monitored the reproduction of females via annual aerial cub surveys in May, as well
164 as ancillary observations at subsequent captures or via remote cameras. For each observation we
165 recorded the female identity, and the number of offspring observed and their age (cub of year,
166 yearling, or two-year-old, etc.). In cases where we did not observe offspring as cubs of the year,
167 it was sometimes difficult to discriminate between yearling and two-year-old bears in the field,
168 so we used a combination of body size, mother's age, and observations in subsequent years to
169 estimate the age of offspring.

170

171 **Estimating demographic parameters**

172 We estimated survival parameters for males and females separately for three age classes: 1)
173 dependent cubs and yearlings (0-1 years old), 2) independent subadult animals (2-6 years old),
174 and 3) adults (>6 years old). The youngest females (n=2) to produce a litter of cubs in our study
175 were five years old and generally younger animals appeared to produce fewer cubs until they
176 were 7, so we estimated reproduction for females in two age classes: 5-6 and >6.

177 Annualized survival rates for collared animals (subadults and adults) were estimated
178 using the Kaplan-Meier known fate and staggered entry approach over monthly time periods.
179 Annualized survival for dependent animals (cubs and yearlings) was estimated by following the
180 fate of litters from collared females. We estimated dependent survival as the proportion of
181 individuals that were observed the following year with their mother. We did not include two-
182 year-olds in this estimate because many of them are not seen with their mother as three-year-olds
183 due to family breakup which often occurs in spring (McLellan 2015).

184 Annual reproduction for subadult and adult females was estimated as the total number of
185 cubs of the year observed with collared females of each age class divided by the total number of
186 collared females monitored in each age class (Garshelis et al. 2005). We estimated the average
187 age of primiparity following the approach described in Garshelis et al. (1998), wherein we
188 calculated the number of cubs produced per nulliparous female aged 5-9. We weighted these
189 results by the proportion of the population available to produce cubs (i.e., those animals that
190 were not currently with offspring and still alive/monitored). We were not able to calculate birth
191 intervals due to small sample sizes (n=4).

192 We estimated the intrinsic population growth rate using a deterministic Leslie matrix,
193 which represents the growth rate of grizzly bears without the influence of immigration and
194 emigration and assuming a stable age distribution. The Leslie matrix included demographic
195 transitions for animals 0-27 years old, which we populated with the age class specific vital rates
196 calculated above. We set reproductive senescence at 27 years of age (Schwartz et al. 2003). We
197 compared this intrinsic growth rate from collared individuals to the observed population growth
198 calculated using genetic tags and spatial capture-recapture (SCR) (Mowat et al. 2020). The
199 primary difference between these two measures of population growth is that intrinsic population
200 growth only considers the influence of reproduction and survival, while observed population
201 growth also includes immigration and emigration and thus represents the observed change in
202 abundance through time. By calculating the difference between observed and intrinsic growth
203 rates, immigration rates can be directly estimated; a demographic parameter that is challenging to
204 estimate in other systems (Kokko 2006, Lamb et al. 2020). We estimated uncertainty for each
205 parameter by resampling individuals with replacement (bootstrapping) 5,000 times, estimating
206 demographic parameters with each bootstrapped sample, and extracting the standard error and

207 90% confidence intervals of the resulting distribution. All analyses were conducted in Program R
208 (R Core Team 2021). To ensure reproducibility, our analysis code and data have been posted on
209 GitHub (https://github.com/ctlamb/ElkValley_Grizzly_Demography_22).

210 The long-term genetic capture-recapture dataset encompassed 4,059 detections of 849
211 grizzly bears across 12,000 km² in the southern Rocky Mountains of BC between 2006 and
212 2021. To estimate demographic parameters for our study area and account for SCR analysis
213 which use home range centers as the parameter of interest, we subset the genetic data to our
214 study area (Figure 1) and reduced its size to 3,210 km² using an interior buffer of 5 km to
215 encompass the home range centers of bears in our study but not additional area (Figure S1,
216 available in Supporting Information). The reduced study area excluded genetically tagged bears
217 whose home range centers were towards the edge of the study area and thus experienced less risk
218 than our collared sample. The subset of genetic data encompassed 1,462 detections of 291
219 grizzly bears. We fit two types of SCR models to these data: 1) closed models which estimated
220 density for each year using the 'secr' package, and 2) open models which estimated population
221 trend by following individuals entering and leaving the population across years using the
222 'openCR' package. For both models we included covariates for sex and trap type (bait site or rub
223 tree) as detection covariates. We included all years (2006-2021) of data to maximize the number
224 of individuals and recaptures and thus improve precision in both the closed and open models, but
225 we focus on the demographic estimates for the 2016-2020 period to align with our period of
226 monitoring the collared bears.

227 We compiled grizzly conflict reports and mortality records by source and location across
228 BC using the publicly available Wildlife Alert Reporting Program data
229 (<https://warp.wildsafebc.com/>) and CI data to assess the degree to which the Elk Valley study

230 area has disproportionately high levels of human-bear conflict and mortality than elsewhere in
231 the province.

232

233 **Estimating unreported mortality**

234 We estimated unrecorded mortality using three methods. Because people may be more
235 likely to report the death of a collared bear than an uncollared bear, and because sample sizes
236 were small, we felt it was important to calculate the unreported rate in multiple ways to assess
237 the robustness of estimates across methods. For each method we provide an overall unreported
238 mortality rate and, where possible, a cause-specific rate.

239 The first method, hereafter termed the “collar fates” approach, used collar fates only. For
240 each bear that died while wearing a functioning radiocollar, we noted whether the animal’s death
241 was reported and recorded in the CI database. We calculated the underreporting rate by dividing
242 the number of collared bear mortalities that were unreported by the total number of collared bear
243 mortalities.

244 For the second method, hereafter called the “CI ratio” method, we replicated the
245 approach of McLellan et al. (2018) and compared the number of bears killed by COS to the
246 number killed by other sources, both for bears wearing functioning radiocollars and for
247 uncollared bears recorded in the CI database. This second approach assumes all mortalities that
248 involve a Conservation Officer are recorded, which is reasonable because Conservation Officers
249 do extensive reporting on each conflict, especially when it results in a mortality, and all
250 mortalities are required to be recorded in the Compulsory Inspection database. To calculate the
251 underreporting rate using the CI ratio method, we first estimated the number of grizzlies killed
252 by human causes but not reported in the CI database ($HC_{unreported}$, eq. 1), where COS_{ci} is the

253 number of bears killed by COS in the CI data, $HCci$ is the number of non-COS human-caused
254 kills in the CI database, COS is the number of collared bears killed by COS, and HC is the
255 number of collared bears killed by non-COS human causes. To get an underreporting rate, we
256 divided $HCunreported$ by the sum of $HCunreported$ plus $HCci$.

257

$$HCunreported = \left(\frac{COSci}{\frac{COS}{HC}} \right) - HCci$$

258 Eq.1 Estimating unreported mortality using the Compulsory Inspection method from McLellan et al.
259 (2018)
260

261 For the third method, hereafter called the “ear tag ratio” method, we took the ratio of
262 animals with functioning radiocollars killed by COS to those killed by other human sources
263 (described in the CI ratio method above) and compared it to the ratio expected based on returned
264 ear tags. We ear-tagged 76 individual bears, but at any one time only 10-20 bears had
265 functioning radiocollars. Some ear-tagged bears died, but there were many uncollared ear-tagged
266 bears on the landscape, and this number increased through the study. This third approach is
267 based on the assumptions that 1) the collared sample is a subset of the larger sample of animals
268 that were marked with permanent ear tags, and 2) the ear tags are reported when a mortality is
269 reported. Ear-tagged bears were reported by Conservation Officers and highway crews and
270 detected via remote cameras in the area (Emily Chow, pers. comm. April 12, 2022), and it is
271 reasonable to assume all ear tags would be reported with reported mortalities because each bear
272 that dies and ends up in the CI database is handled by either a CO or biologist, and all check for
273 tags.

274 To calculate the underreporting rate with the ear tag ratio method, we first estimated the
275 expected return ratio of ear tags for COS kills ($TRco$) by dividing the number of COS-killed ear-

276 tagged animals wearing functioning radiocollars by the total number of ear-tagged animals (both
277 collared and uncollared) killed by COS. Because bears euthanized by the COS had perfect
278 recording for both radiocollared and uncollared but ear-tagged bears, this $TRco$ ratio was the
279 expected ear tag return rate for other mortality sources, if they also had perfect reporting.
280 However, other causes of mortality that we can detect in the radiocollared sample, such as train
281 and highway kills or poaching, are unlikely to have perfect reporting of uncollared bears. For
282 these causes of mortality, we expect fewer ear tags to be returned for uncollared but ear-tagged
283 animals due to either cryptic poaching or animals dying a short distance from the right of way
284 after a road or rail collision. To calculate the expected number of returned ear tags for mortality
285 sources with imperfect reporting ($TRex$, eq. 2), we divided the number of radiocollared bears
286 deaths for a given mortality source ($CDms$) by the expected ear tag return rate for COS kills
287 ($TRco$). To get an underreporting rate for the ear tag ratio method, we divided the total reported
288 number of ear-tagged bears killed ($TRrep$) by $TRex$ and subtracted it from 1.

$$TRex = \left(\frac{CDms}{TRco} \right)$$

290 Eq.2 Estimating the expected number of recovered ear tags, which is then compared to the actual number
291 of recovered ear tags to estimate the unreported rate with the ear tag ratio method.

292 To contextualize the ear tag method, consider an example where for every radiocollared
293 bear killed by COS, the COS also euthanized 2 uncollared bears with ear tags. We assume the
294 COS report all ear-tagged animals and thus our collared bear represents 1/3 of the total ear-
295 tagged animals killed by COS ($TRco = 0.33$). Now consider a source of mortality with imperfect
296 reporting, such as highway collisions. In this example, 4 deaths of radiocollared animals were
297 reported on the highway ($CDms = 4$), and 2 mortalities of uncollared ear-tagged bears were also
298 reported ($TRrep = 6$). Assuming perfect reporting, we expect one third (1/3) of ear-tagged
299 animals to be wearing functioning collars; however, for the highway sample two thirds (4/6) of
300 animals to be wearing functioning collars; however, for the highway sample two thirds (4/6) of

301 ear-tagged animals were wearing a collar, suggesting a lack of ear tag recovery. To calculate the
302 expected number of returned ear tags (TR_{ex}), we divided 4 (CD_{ms}) by 0.33 (TR_{co}) and found
303 that 12 ear-tagged animals likely died in highway collisions. To get an underreporting rate using
304 the ear tag ratio method for this example, we divided 6 (TR_{rep}) by 12 (TR_{ex}) and subtracted it
305 from 1 to find that an estimated 50% of highway mortalities were unreported.

306 Finally, we integrated the estimates from all three methods (collar fates, CI ratio, and ear
307 tag ratio) into a single ensemble estimate. To do this, we compiled the bootstrapped results from
308 all methods, calculated a mean result for each bootstrap iteration across methods, and reported
309 this ensemble estimate along with its error.

310

311 **RESULTS**

312 **Capture, handling, and collaring**

313 Between 2016 and 2022 we radiocollared 70 individuals (110 capture events) and 6 bears were
314 marked but not radiocollared. Researchers were responsible for ~92% of the captures while the
315 remaining ~8% were caught by Conservation Officers. Bears were captured in culvert traps
316 ($n=12$), free-range darting from the ground ($n=6$), free-range darting from a helicopter ($n=15$),
317 and in leg restraints ($n=77$). The collared animals were captured mostly as adults (>6 years old):
318 $n=27$ males; $n=30$ females) and subadults (2-6 years old: $n=21$ males; $n=23$ females), and one
319 male was collared at 1.5 years old. Capture effort was concentrated in seasonal habitats, which
320 was generally in the valley bottom of the Elk Valley in the fall. Once collared, bears ranged well
321 beyond the valley bottom into adjacent valleys and inter-provincially (Figure 2).

322 Males were consistently heavier than females, and this difference increased as they aged
323 (Figure 2). The average age of captured adults was 12 for males and 11 for females, while the

324 oldest male was 27 and the oldest female was estimated at approximately 20 years old based on
325 tooth wear (Table S1, available in Supporting Information). Fat levels were similar across age
326 classes and sexes but differed through the year with increasing fat levels in the fall. As a
327 percentage of body weight, the maximum fat level recorded was 38.6% for a female and 39.2%
328 for a male. Bears captured due to conflicts with people were in good body condition and
329 appeared to be as fat as, or fatter than, bears captured for research purposes (Figure 2). Bears
330 killed due to conflicts with people had an average of 2.4 cm (n=8, range=1-4 cm) of rump fat,
331 and those killed in road/rail collisions had 4.2 cm (n=3, range=3.5-5 cm) of rump fat, indicating
332 generally healthy animals in both cases.

333

334 **Demographic monitoring**

335 We recorded mortality of 22 of the 76 marked animals (Figure 5). Of the 76 marked animals, 70
336 were radiocollared, and 14 died while their collar was functioning (Table 1). The other 8 marked
337 animals that died were either never collared (only ear tagged) or were not wearing a functional
338 collar when they died. We monitored the survival of 70 individual collared animals across 160
339 animal-years. The cause of death for the 14 animals with a functioning collar was human-
340 wildlife conflict (n=6), road collision (n=2), railway collision (n=3), road or rail collision (n=1),
341 unknown but human suspected (n=1), and natural (n=1). The human-wildlife conflict kills
342 generally stemmed from unsecured attractants and subsequent conflicts at private residences
343 (n=4), but one animal was killed due to habituated behaviour on a coal mine, and another was
344 shot and killed ~2 km from town and motive of the shooter was unknown because the mortality
345 was not reported. We suspected human causes for the one mortality of unknown cause because
346 the animal was a 5-year-old female in good health (25% body fat) when she was captured just

347 over a month earlier. She was found dead ~50 meters from a gravel road and ~500 meters from a
348 highway, but due to delayed transmission of the collar's mortality signal, the carcass was too
349 decomposed to assess whether blunt force trauma from a collision had occurred or if she had
350 been shot. The natural mortality was a female that died in a clifffy area near the top of a
351 mountain. Telemetry data showed she had gone up into the cliffs and stayed there for a week
352 before she died. When found, she was emaciated with no signs of trauma. Toxicology results
353 suggested she was not poisoned.

354 All the human-caused mortalities occurred in the valley bottom, which made up less than
355 half of the area the bears ranged across (Figure 2). Three of the mortalities occurred while
356 collared females were with dependent offspring. In one case all three cubs and their mother were
357 struck and killed by a train, and in another case one of two yearlings were killed with their
358 mother in an unreported conflict mortality. In the third case we detected one of two cubs alive for
359 the following four years after its collared mother had died and the cub (now a subadult) is
360 currently still alive and collared. Five of the 70 radiocollared bears in our study were initially
361 captured by Conservation Officers, but none of the 14 animals that died while collared had been
362 involved in a conflict situation at first capture.

363 Of the 101 capture events where collars were deployed, the fate of the animal was known
364 in 95 cases and unknown in 6 cases. Known fates included death (n=14), the animal was alive
365 but had dropped its collar (n=47), or the animal was still wearing a functioning collar at the time
366 of writing (n=17). In the remaining 23 instances, we lost connection with collars; however, we
367 know the animals were alive in 17 of these instances due to subsequent recapture or DNA
368 detection. In the 6 cases where the bears' fate remained unknown, it is possible the collar was
369 destroyed during a human-caused mortality (i.e., unreported conflict kill, poaching, or collision),

370 but we know the majority of the connection failures were not mortalities but rather collar
371 failures. Of the 6 unknown fates, 4 animals had last collar locations >1.5 km from a road,
372 railway, or human settlement, suggesting the connection loss was unlikely due to a human-
373 caused mortality. Of the remaining 2 animals with unknown fates, the last relocation for one was
374 0.5-1.5 km from a road, railway, or human settlement, and the other was <0.5 km. Indeed, collars
375 involved in road and rail collisions were often severely damaged, impairing their normal
376 function. Thus, it's possible some of these unknown fates were undetected mortalities. However,
377 it is also important to note that many of the collars with connection failures that were eventually
378 confirmed to be simply collar failures and not bear mortalities had also stopped working close to
379 roads and people. For this analysis we assume the 6 unknown fates are also censored fates and
380 not deaths while acknowledging that this assumption means we are estimating a conservative
381 mortality rate which may be slightly higher if some of these unknown fates were deaths.

382 We monitored reproduction of 36 subadult and adult females across 94 animal-years and
383 detected 23 litters of various aged offspring. Females spent 54 animal-years alone, 18 with cubs,
384 13 with yearlings, 7 with two-year-olds, and 2 with three-year-olds. There was an average of 1.9
385 cubs per litter, 1.5 yearlings, 1.4 two-year-olds, and 1.5 three-year-olds. We observed a total of
386 41 dependent offspring, of which 28 were monitored for more than one year. Of these 28, we
387 observed 26 as cubs and 19 were observed with their mother the following spring while 7
388 presumably died. We observed 15 offspring as yearlings, of which 11 were observed with their
389 mother the following spring at two years of age; the 4 undetected two-year-olds may have died,
390 or they were simply not with their mother during our flight in May either due to dispersal or
391 temporary displacement during breeding season.

392 We monitored the reproductive status of 16 females between the ages of 5 and 9. Two
393 animals were known to have had a litter at 5, and one animal had a litter at 6. These were the
394 only animals to have a litter before the age of 7. Most females were with cubs when aged 7-9
395 (Figure S2, available in Supporting Information). The age of first parturition was estimated at 7.2
396 years including all 16 females, and 7.5 years when we excluded two females that were only
397 monitored at 9+ years old, and we could not be sure they had not had cubs previously.

398

399 **Estimating demographic parameters**

400 Annual survival of dependent young, 0-1 years old, was 0.73 (90% CI: 0.61-0.83) for both sexes
401 combined, 0.60 (90% CI: 0.38-0.82) for subadult males, 0.71 (90% CI: 0.54-0.88) for subadult
402 females, 1.0 (90% CI: 0.83-1.00) for adult males, and 0.96 (90% CI: 0.91-1.0) for adult females.
403 Annual reproduction (female cubs/female/year) by females aged 5-6 was 0.15 (90% CI: 0.00-
404 0.31), and 0.24 (90% CI: 0.15-0.33) for females over 6 years old. When combined in the Leslie
405 matrix, these vital rates suggested the intrinsic population growth rate for Elk Valley grizzly
406 bears was 0.94 (90% CI: 0.86-1.01), with 93% of bootstrapped estimates <1 (Figure 6).

407 Open spatial capture-recapture modelling suggested the abundance of grizzly bears in the
408 Elk Valley study area has been stable 2006-2021 with an observed population growth rate of
409 1.01 (90% CI: 0.99-1.03). We tested whether this overall stable trend was different during our
410 period of study (2016-2022) compared to pre-2016 and found no evidence for the more complex
411 model structure (delta AIC=0.4). The density of grizzly bears in the Elk Valley study area
412 between 2016 and 2021 averaged 32.0 bears/1,000 km² (90% CI: 28.9-35.0), or 103 individuals
413 (90% CI: 92.7-112.0). Using a stable observed population growth rate of 1.01, and the intrinsic
414 population growth rate calculated above from radiocollared animals, we estimated the viability

415 of grizzly bears in the Elk Valley is subsidized through immigration which annually adds 6.9%
416 (90% CI: 0-15) of the population, or ~7 bears, into the study area to maintain stable abundance
417 (Figure 6). Indeed, we observed 3 examples of radiocollared male bears immigrating into the Elk
418 Valley study area from 77-95 km away (Figure 6). All three of these bears were eventually
419 killed, highlighting the spatial extent of the source-sink dynamics in the Elk Valley study area
420 and the risk immigrant bears are exposed to once settled.

421 Recorded conflicts and mortality were higher in the Elk Valley study area than the rest of
422 BC. There was an average of 65.3 conflict reports per 10,000 sq.km/year in the Elk Valley
423 compared to only 5.8 per 10,000 sq.km/year across the rest of the province (Figure 4). Hunting, a
424 regulated source of mortality, showed a similar prevalence (mortality per unit area) in the Elk
425 Valley compared to the rest of the province. In contrast, conflicts with people and road/rail
426 mortalities were one to two orders of magnitude more prevalent in the Elk Valley than elsewhere
427 (Table 2). The Elk Valley study area, which accounts for less than 1% of the grizzly bear range
428 in BC, but encompassed 33% and 42% of the provincially reported road and rail mortalities,
429 respectively.

430

431 **Estimating unreported mortality**

432 Of the 13 grizzly bears killed by people that were wearing functioning radiocollars, 7 were not
433 reported to authorities. The unreported mortalities were from road or rail collisions (n=4),
434 conflicts at private property (n=1), shot and left (n=1), and of unknown cause but where humans
435 were suspected (n=1) (Table 1). When estimating the unreported rate, we classified the shot and
436 left bear as a conflict kill. We estimated the unreported rate of human-caused mortality using the
437 rate of reporting from collared bears at 0.54 (90% CI: 0.31-0.77). Although sample sizes were

438 small, we calculated cause-specific unreported rate rates to identify any obvious differences in
439 rates between sources. Two of 4 mortalities that resulted from conflicts with people but without
440 CO involvement were not reported (0.50), 4 of 6 road and rail mortalities were not reported
441 (0.67), and the unknown but human suspected mortality was not reported (1).

442 Using the CI ratio method, we estimated the unreported rate at 0.64 (90% CI: 0.0-0.9).
443 Using the ear tag ratio method, we estimated the unreported rate at 0.76 (90% CI: 0.54-1.0).
444 Putting all estimates together, we estimated an ensemble unreported rate at 0.65 (90% CI: 0.35-
445 0.81). We calculated cause-specific unreported rates using both the CI and ear tag ratio methods
446 (Table 1).

447

448 **DISCUSSION**

449 Grizzly bears in the Elk Valley provide unique insights into how human-dominated landscapes
450 shape grizzly bear behaviour and demography, and how grizzly bears in turn are slowly
451 reshaping the behaviour of people who are adopting coexistence solutions. Grizzly bears are
452 currently abundant in the Elk Valley despite living among 15,000 people, major highways and
453 railways, extensive resource extraction, and widespread recreation. The Elk Valley hosts more
454 than twice the grizzly bear density (32 bears/1,000 km²) compared to 100 km the north in Banff
455 National Park (12 bears/1,000 km², (Whittington et al. 2018))—Canada’s flagship protected area.
456 A desire to understand the demographic mechanisms that allowed grizzly bears to persist and
457 apparently thrive in the Elk Valley motivated this work.

458 We show that young grizzly bears in the Elk Valley are surviving poorly, with up to 40%
459 annual mortality (Figure 5A). Adult animals, however, had survival rates over 95% which is as
460 high as, or higher than, survival rates seen in other studies such as those done in Banff (Garshelis

461 et al. 2005), Flathead Valley (McLellan 2015), northwest Montana (Mace et al. 2012), and
462 Yellowstone ((Schwartz et al. 2006); Fig. 5C). Consistent with other studies, we show that
463 people caused most mortalities (93%, 13/14). The primary cause of death was conflicts with
464 people due to unsecured attractants on private property and collisions with vehicles or trains. No
465 collared bears were killed by hunters, but the grizzly bear hunting season was closed a year after
466 our study began. Despite many people living throughout the study area, and the Conservation
467 Officer headquarters being in the study area, we estimate that only about one-third of the human-
468 caused mortalities that did not involve Conservation Officers were reported to authorities.

469 Although this is a slightly higher reporting rate than seen in more remote areas (McLellan et al.
470 2018), the low reporting rate means that the Compulsory Inspection data currently under-
471 represents the severity of conflict, road, and rail mortalities in the Elk Valley and likely
472 elsewhere in BC. The stark discrepancy in survival between subadults and adults in the Elk
473 Valley highlights the intense demographic filter (*sensu* Ford et al. 2017) that essentially provides
474 two options for a young bear: 1) learn how to avoid conflicts and stay safe near transportation
475 corridors, or 2) likely die before adulthood.

476 High mortality rates were not offset by reproduction in our study population (Fig. 5B).
477 The low intrinsic population growth rate suggested bear density in the lower Elk Valley would
478 decrease by approximately 7% a year without immigration. Without being buoyed by
479 immigration, the bears that spend time in the lower Elk valley bottom would decline rapidly
480 (Figure 6A and B). However, such a decline has not been observed and bear density has been
481 relatively stable for the past 15 years. According to local observations and population
482 reconstructions, grizzly bear numbers had also been increasing in the area prior to our study
483 (Hatter et al. 2018, Lamb et al. 2019, Mowat et al. 2020). The source-sink dynamic observed

484 here appears to be currently sustainable at the broader landscape scale beyond the Elk Valley and
485 us supported by the current level of connectivity between the Elk Valley and adjacent secure
486 habitats. We do not know how fragile the source-sink dynamic is, and whether habitat alteration
487 in adjacent habitats could disrupt this dynamic and impede the flow of bears needed to sustain
488 the Elk Valley in the future.

489 Grizzly bears can be a challenging species for people to have living nearby. Along with
490 the Terrace-Kitimat and Bella Coola valleys, the Elk Valley is a provincial hotspot for human-
491 grizzly bear conflict, as evidenced by the multitude of conflicts reported each year (Figure 4C).
492 In addition to conflicts between people and bears over unsecured attractants, grizzly bears
493 occasionally cause physical harm to people. In the last ten years, at least six people have been
494 attacked by grizzly bears in the Elk Valley and adjacent Kootenay valley outside Cranbrook; this
495 accounts for approximately half the grizzly-caused human injuries in the entire province during
496 that period. In each case, the victims were either actively hunting or scouting for animals before
497 hunting season. Victims often defended themselves by shooting at the bear, or in one case by
498 stabbing the bear with an arrow. While many people live and recreate in the valley without ever
499 having a conflict with a grizzly bear—many have never even seen a grizzly bear due to their
500 nocturnal behavior—the consistent flurry of conflicts in the spring and fall, as well as infrequent
501 but consistent physical confrontations, indicate human-grizzly coexistence in the Elk Valley
502 remains challenging.

503 Collisions between vehicles or trains and wildlife were common in our study. Like other
504 challenges to human-wildlife coexistence, collisions are lose-lose situations where neither party
505 benefits. Collisions with wildlife often result in dead animals, human injury or death, damaged
506 vehicles, and the interruption of the flow of goods and people along transportation corridors.

507 While collisions with bears are less frequent than with other species such as deer, elk, moose, or
508 sheep—largely due to their relative abundance on the landscape—we show here that collisions
509 between grizzly bears and vehicles or trains are a leading cause of death contributing to
510 unsustainable mortality rates for grizzly bears in the Elk Valley. About one third of British
511 Columbia's recorded grizzly bear road collisions occur in the Elk Valley. Rail collisions with
512 grizzly bears only occur in a few areas of the province, but nearly half the recorded mortalities
513 occur in the Elk Valley. Rail mortality through the Highway 1 corridor is a leading mortality
514 factor for grizzly bears in Banff National Park (St. Clair et al. 2019), which is the only other
515 place in Canada where train collisions with grizzly bears are regularly reported.

516 Although grizzly bears in the Elk Valley are clearly exposed to high levels of risk from
517 various human activities on the landscape, many adult grizzly bears in our study lived near
518 people without reported conflict. We followed multiple adult female bears, some of which also
519 had offspring, that spent most of their active season living in the valley bottom where their daily
520 movements involved crossing railways, highways, and spending time near residential properties.
521 These bears were often strictly nocturnal (Lamb et al. 2020), allowing them to spend time near
522 residences and even access human-sourced foods such as apples, without ever being detected by
523 people. In contrast, subadult animals in our study often accessed human foods during the day,
524 increasing the likelihood that they would be detected by people and be killed. Because offspring
525 generally separate from their mothers before they are old enough to safely wear a collar, we were
526 not able to determine if cubs raised by a savvy mother also had higher survival. However,
527 (Morehouse et al. 2016) found conflict behaviour of mothers dictated the conflict behaviour of
528 offspring, suggesting behaviours that reduce or promote conflicts can be learned. Currently many
529 young bears in the Elk Valley are immigrants from areas without human settlement or

530 transportation corridors (Figure 6), and they are likely more naïve to these risks and more prone
531 to conflict. We thus expect conflicts in the Elk Valley could be reduced by ensuring high
532 survival of resident adult female bears who know how to coexist and can continue teaching their
533 offspring these habitats.

534 While the abundance of grizzly bears appears stable in the Elk Valley, does stability
535 subsidized through immigration, recurring seasonal damage to private property, and occasional
536 physical confrontations signal coexistence? Coexistence likely falls along a spectrum. Take for
537 example areas where grizzly bears have been extirpated, such as the Okanagan Valley, Peace
538 River Valley, Lower Fraser Valley, or the prairies. Coexistence is not happening in these
539 landscapes because grizzly bears are not present, and grizzly bears that disperse into the human-
540 dominated portions of these areas are often killed or relocated. On the other extreme would be an
541 environment where thousands of people and abundant grizzly bears can share the same
542 landscapes with little risk to life or property, likely with significant behavioural adjustments from
543 both parties. Such a landscape doesn't yet exist, but some are trending in that direction (Proctor
544 et al. 2018, Morehouse et al. 2020). The Elk Valley fits somewhere in the middle of these two
545 scenarios, with an abundant and stable grizzly bear population sharing a valley with people, but
546 conflicts and grizzly bear mortality remain high in portions of the valley. We view this as a form
547 of coexistence due to the consistently high number of grizzly bears that share space with people;
548 however, the situation is far from perfect and is not "peaceful coexistence", especially for the
549 dead bears and injured people. Future efforts should focus on finding ways to keep people and
550 bears safer in the valley, with a goal of reducing the risk to people and property, grizzly
551 mortality, and ultimately the reliance on immigration to sustain this population.

552

553 MANAGEMENT IMPLICATIONS

554 Here we provide evidence that grizzly bear mortality and conflicts need to be reduced in
555 the Elk Valley study area to facilitate human-bear coexistence and a self-sustaining bear
556 population. Tools are increasingly available to improve the safety of bears and people, such as
557 bear aware training and improved technologies for personal and property safety. A
558 comprehensive review from Alaska demonstrated that bear spray improves personal safety by
559 stopping brown bear charges at least 90% of the time, and leaves 98% of the people uninjured
560 who deploy the spray on a bear (Smith et al. 2008). Electric fencing has been shown to be one of
561 the most effective tools to repel grizzly bears from attractants such as livestock or fruit trees,
562 reducing property damage by 80-100% (Johnson 2018, Khorozyan and Waltert 2020). Lethal
563 removal of problem bears generally provides short-term relief but does not address the
564 underlying causes of conflict, and thus is not effective long term unless lethal removal is done
565 continuously (Khorozyan and Waltert 2020). Programs that provide bear spray training and help
566 landowners eliminate access to attractants, such as cost shared electric fencing or removing and
567 replacing fruit trees, have made a positive difference for coexistence when applied at a landscape
568 scale (Proctor et al. 2018, Eneas 2020). In British Columbia there are efforts to reduce conflicts,
569 supported by a government-private partnership called WildsafeBC, conservation groups, private
570 funders, and some municipalities, but the lack of dedicated funds for cost share programs limits
571 the long-term success of these solutions. However, creative solutions to reduce attractants are
572 being trialed locally. For example, the BC Ministry of Transportation and Infrastructure
573 implemented a program to move road killed animal carcasses in the Elk Valley to an electric
574 fenced compound where the carcasses are not accessible to bears. Highway strikes of ungulates
575 are very common in the Elk Valley, and previously the carcasses were often dumped in gravel

576 pits and commonly fed on by bears (Figure 3F), so this effort removed a large bear attractant
577 from the valley bottom. Further efforts to reduce bears' access to unsecured attractants such as
578 livestock, fruit trees, and garbage are needed at a landscape to meaningfully reduce conflicts and
579 mortality.

580 The Province of British Columbia is supporting a different coexistence solution—a collision
581 reduction system composed of wildlife crossing structures and fencing along Highway 3—that
582 will keep wildlife and people safer in our study area. An ambitious, grassroots project broke
583 ground in 2020 that aims to fence 27 kilometers of highway and build (n=3) or retrofit existing
584 (n=7) structures to serve as wildlife crossings. This section of highway encompasses multiple
585 collision hotspots (Lee et al. 2019), including the sites where one collared bear was killed on the
586 highway and where another was known to be struck and killed by either a vehicle or train. The
587 project is focused on a significant wildlife corridor connecting Canada and the USA, making it
588 an ideal location for mitigation (Proctor et al. 2015, Lee et al. 2019, Poole and Lamb 2020).

589 Crossing structures are used by bears regularly in Banff National Park (Sawaya et al. 2014, Ford
590 et al. 2017), and when combined with fencing that excludes wildlife from the roadway, can
591 reduce wildlife mortality by up to 96% (Ford et al. 2022). Currently, the only collision reduction
592 system within the core range of grizzly bears occurs in Banff National Park, but the low density
593 of bears in Banff limits sample sizes to measure the systems' effectiveness on grizzly bears (Ford
594 et al. 2022). In the Elk Valley the comparatively higher density of grizzly bears, collisions (Table
595 1), and the comprehensive "before" data provided here should eventually provide a robust
596 before-after comparison of the Highway 3 projects' effectiveness.

597 Several emerging trends in human behaviour and stewardship practices suggest the future
598 could be brighter in terms of reduced human-bear conflicts if such practices are adopted at scale.

599 We believe that creating programs to support local people and the bears who have learned to
600 navigate these challenging areas will encourage coexistence in the Elk Valley and help redefine
601 what the upper spectrum of coexistence could look like.

602

603 **ACKNOWLEDGMENTS**

604 Our work was conducted within ?amak?is Ktunaxa, the homelands of Ktunaxa people. We thank
605 the Ktunaxa Nation Council for their support of our k‡aw‡a (grizzly bear) work. Thanks to R.
606 Gordon, P. Burley, F. Deboon, P. Stent, E. Chow, S. Clow, and the South Rockies Grizzly Bear
607 Project volunteers for supporting our research project and helping us out in the field. These data
608 were collected with support from the Government of British Columbia, British Columbia
609 Conservation Officer Service, Habitat Conservation Trust Foundation, Teck Coal, Vanier
610 Canada Graduate Scholarship, Liber Ero Fellowship Program, National Science Engineering and
611 Research Council, Fish and Wildlife Compensation Program, Forest Enhancement Society of
612 British Columbia, Columbia Basin Trust, Counter Assault, Wildsight, Nature Conservancy of
613 Canada, Wildlife Conservation Society, Yellowstone to Yukon Conservation Initiative, Safari
614 Club International, Sparwood and District Fish and Wildlife Association, Elkford Rod and Gun
615 Club, Ministry of Transport and Infrastructure, Outdoor Research, and Patagonia. The authors
616 have no conflicts of interest to declare.

617

618 **ETHICS STATEMENT**

619 Captures were in accordance with University of Alberta Animal Ethics #AUP00002181 and
620 Province of British Columbia Capture Permit #CB17-264200.

621

622 **REFERENCES**

- 623 Apps, C., D. Paetkau, B. N. McLellan, A. N. Hamilton, and B. Bateman. 2014. Grizzly bear
624 population abundance, distribution, and connectivity across British Columbia's southern
625 Coast Ranges. Page 125. Aspen Wildlife Research and Ministry of Environment,
626 Victoria, British Columbia.
- 627 Bruskotter, J. T., J. A. Vucetich, M. J. Manfredo, G. R. Karns, C. Wolf, K. Ard, N. H. Carter, J.
628 V. López-Bao, G. Chapron, S. D. Gehrt, and W. J. Ripple. 2017. Modernization, Risk,
629 and Conservation of the World's Largest Carnivores. *BioScience* 67:646–655.
- 630 Cattet, M. R. L., N. A. Caulkett, and G. B. Stenhouse. 2003. Anesthesia of Grizzly Bears Using
631 Xylazine-Zolazepam-Tiletamine or Zolazepam-Tiletamine. *Ursus* 14:88–93.
- 632 Eneas, K. L. 2020. Influence of Livestock and Electrified Fences on Livestock Depredation and
633 Habitat Selection by Grizzly Bears in the Mission Valley, Montana. University of
634 Montana.
- 635 Farley, S. D., and C. T. Robbins. 1994. Development of two methods to estimate body
636 composition of bears. *Canadian Journal of Zoology* 72:220–226.
- 637 Flaa, J. P., S. B. Michel, and C. Borstad. 2009. Building a reliable snare cable for capturing
638 grizzly and American black bears. *Ursus* 20:50–55.
- 639 Ford, A. T., M. Barrueto, and A. P. Clevenger. 2017. Road mitigation is a demographic filter for
640 grizzly bears: Road Crossing Behavior in Grizzly Bears. *Wildlife Society Bulletin*
641 41:712–719.
- 642 Ford, A. T., B. Dorsey, T. S. Lee, and A. P. Clevenger. 2022. A before-after-control-impact
643 study of wildlife fencing along a highway in the Canadian Rocky Mountains. *Frontiers in*
644 *Conservation Science* 3:935420.

- 645 Frank, B. 2016. Human–Wildlife Conflicts and the Need to Include Tolerance and Coexistence:
646 An Introductory Comment. *Society & Natural Resources* 29:738–743.
- 647 Garshelis, D. L., M. L. Gibeau, and S. Herrero. 2005. Grizzly Bear Demographics in and around
648 Banff National Park and Kananaskis Country, Alberta. *The Journal of Wildlife
649 Management* 69:277–297.
- 650 Garshelis, D. L., K. V. Noyce, and P. L. Coy. 1998. Calculating Average Age of First
651 Reproduction Free of the Biases Prevalent in Bear Studies. *Ursus* 10:437–447.
- 652 Hatter, I. W., G. Mowat, and B. N. McLellan. 2018. Statistical population reconstruction to
653 evaluate grizzly bear trends in British Columbia, Canada. *Ursus* 29:1.
- 654 Hebblewhite, M., J. A. Hilty, S. Williams, H. Locke, C. Chester, D. Johns, G. Kehm, and W. L.
655 Francis. 2022. Can a large-landscape conservation vision contribute to achieving
656 biodiversity targets? *Conservation Science and Practice* 4:e588.
- 657 Johnson, B. J. 2018. Permeability of three-strand electric fences by black bears and grizzly bears.
658 Montana State University, Bozeman, Montana.
- 659 Khorozyan, I., and M. Waltert. 2020. Variation and conservation implications of the
660 effectiveness of anti-bear interventions. *Scientific Reports* 10:15341.
- 661 Kokko, H. 2006. From Individual Dispersal to Species Ranges: Perspectives for a Changing
662 World. *Science* 313:789–791.
- 663 Laliberte, A. S., and W. J. Ripple. 2004. Range Contractions of North American Carnivores and
664 Ungulates. *BioScience* 54:123.
- 665 Lamb, C. T., A. T. Ford, B. N. McLellan, M. F. Proctor, G. Mowat, L. Ciarniello, S. E. Nielsen,
666 and S. Boutin. 2020. The ecology of human–carnivore coexistence. *Proceedings of the
667 National Academy of Sciences* 117:17876–17883.

- 668 Lamb, C. T., A. T. Ford, M. F. Proctor, J. A. Royle, G. Mowat, and S. Boutin. 2019. Genetic
669 tagging in the Anthropocene: scaling ecology from alleles to ecosystems. *Ecological
670 Applications* 29:e01876.
- 671 Lamb, C. T., G. Mowat, B. N. McLellan, S. E. Nielsen, and S. Boutin. 2017. Forbidden fruit:
672 human settlement and abundant fruit create an ecological trap for an apex omnivore.
673 *Journal of Animal Ecology* 86:55–65.
- 674 Lamb, C. T., G. Mowat, A. Reid, L. Smit, M. Proctor, B. N. McLellan, S. E. Nielsen, and S.
675 Boutin. 2018. Effects of habitat quality and access management on the density of a
676 recovering grizzly bear population. *Journal of Applied Ecology* 55:1406–1417.
- 677 Lee, T., D. A. P. Clevenger, and D. C. Lamb. 2019. Amendment: Highway 3 Transportation
678 Mitigation for Wildlife and Connectivity in Elk Valley of British Columbia. Page 60.
679 Miistikis Institute, Calgary, Alberta.
- 680 Lute, M. L., and N. H. Carter. 2020. Are We Coexisting With Carnivores in the American West?
681 *Frontiers in Ecology and Evolution* 8:48.
- 682 Mace, R. D., D. W. Carney, T. Chilton-Radandt, S. A. Courville, M. A. Haroldson, R. B. Harris,
683 J. Jonkel, B. Mclellan, M. Madel, T. L. Manley, C. C. Schwartz, C. Servheen, G.
684 Stenhouse, J. S. Waller, and E. Wenum. 2012. Grizzly bear population vital rates and
685 trend in the Northern Continental Divide Ecosystem, Montana: NCDE Grizzly
686 Population Trend. *The Journal of Wildlife Management* 76:119–128.
- 687 Manfredo, M. J., E. G. Urquiza-Haas, A. W. Don Carlos, J. T. Bruskotter, and A. M. Dietsch.
688 2020. How anthropomorphism is changing the social context of modern wildlife
689 conservation. *Biological Conservation* 241:108297.

- 690 Mattson, D. J., and T. Merrill. 2002. Extirpations of Grizzly Bears in the Contiguous United
691 States, 1850-2000. *Conservation Biology* 16:1123–1136.
- 692 McLellan, B. N. 1989. Dynamics of a grizzly bear population during a period of industrial
693 resource extraction. III. Natality and rate of increase. *Canadian Journal of Zoology*
694 67:1865–1868.
- 695 McLellan, B. N. 2015. Some mechanisms underlying variation in vital rates of grizzly bears on a
696 multiple use landscape: Variation in Grizzly Bear Vital Rates. *The Journal of Wildlife
697 Management* 79:749–765.
- 698 McLellan, B. N., G. Mowat, and C. T. Lamb. 2018. Estimating unrecorded human-caused
699 mortalities of grizzly bears in the Flathead Valley, British Columbia, Canada. *PeerJ*
700 6:e5781.
- 701 McLellan, M. L., B. N. McLellan, R. Sollmann, and H. U. Wittmer. 2021. Vital rates of two
702 small populations of brown bears in Canada and range-wide relationship between
703 population size and trend. *Ecology and Evolution* 11:3422–3434.
- 704 Morehouse, A. T., and M. S. Boyce. 2016. Grizzly bears without borders: Spatially explicit
705 capture-recapture in southwestern Alberta. *The Journal of Wildlife Management*
706 80:1152–1166.
- 707 Morehouse, A. T., T. A. Graves, N. Mikle, and M. S. Boyce. 2016. Nature vs. Nurture: Evidence
708 for Social Learning of Conflict Behaviour in Grizzly Bears. *PLOS ONE* 11:e0165425.
- 709 Morehouse, A. T., C. Hughes, N. Manners, J. Bectell, and T. Bruder. 2020. Carnivores and
710 Communities: A Case Study of Human-Carnivore Conflict Mitigation in Southwestern
711 Alberta. *Frontiers in Ecology and Evolution* 8:2.

- 712 Mowat, G., L. Smit, C. Lamb, and N. Faught. 2020. South Rockies grizzly bear inventory:
713 progress report 2006-2019. Page 25. Province of BC, Nelson, BC.
- 714 Peek, J., J. Beecham, D. Garshelis, F. Messier, S. Miller, and D. Strickland. 2003. Management
715 Of Grizzly Bears In British Columbia: A Review By An Independent Scientific Panel.
716 Page 90. Minister of Water, Land and Air Protection Government of British Columbia,
717 Victoria, B.C.
- 718 Poole, K., and C. Lamb. 2020. Migration, movements and survival in a partially migratory elk
719 population in southeast British Columbia. Page 56. Sparwood & District Fish & Wildlife
720 Association.
- 721 Proctor, M. F., S. E. Nielsen, W. F. Kasworm, C. Servheen, T. G. Radandt, A. G. Machutchon,
722 and M. S. Boyce. 2015. Grizzly bear connectivity mapping in the Canada-United States
723 trans-border region: Grizzly Bear Connectivity Mapping. The Journal of Wildlife
724 Management 79:544–558.
- 725 Proctor, M., W. Kasworm, K. Annis, A. MacHutchon, J. Teisberg, T. Radandt, and C. Servheen.
726 2018. Conservation of Threatened Canada-USA Trans-border Grizzly Bears Linked to
727 Comprehensive Conflict Reduction. Human–Wildlife Interactions 12.
- 728 R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
729 Statistical Computing, Vienna, Austria.
- 730 Sawaya, M. A., S. T. Kalinowski, and A. P. Clevenger. 2014. Genetic connectivity for two bear
731 species at wildlife crossing structures in Banff National Park. Proceedings of the Royal
732 Society B: Biological Sciences 281:20131705.
- 733 Schwartz, C. C., M. A. Haroldson, G. C. White, R. B. Harris, S. Cherry, K. A. Keating, D.
734 Moody, and C. Servheen. 2006. Temporal, Spatial, and Environmental Influences on the

- 735 Demographics of Grizzly Bears in the Greater Yellowstone Ecosystem. *Wildlife*
736 *Monographs* 161:1–68.
- 737 Schwartz, C. C., K. A. Keating, H. V. Reynolds, V. G. Barnes, R. A. Sellers, J. E. Swenson, S.
738 D. Miller, B. N. McLellan, J. Keay, R. McCann, M. Gibeau, W. F. Wakkinen, R. D.
739 Mace, W. Kasworm, R. Smith, and S. Herrero. 2003. Reproductive maturation and
740 senescence in the female brown bear:11.
- 741 Smith, T. S., S. Herrero, T. D. Debruyn, and J. M. Wilder. 2008. Efficacy of Bear Deterrent
742 Spray in Alaska. *Journal of Wildlife Management* 72:640–645.
- 743 St. Clair, C. C., J. Backs, A. Friesen, A. Gangadharan, P. Gilhooly, M. Murray, and S. Pollock.
744 2019. Animal learning may contribute to both problems and solutions for wildlife–train
745 collisions. *Philosophical Transactions of the Royal Society B: Biological Sciences*
746 374:20180050.
- 747 Whittington, J., M. Hebblewhite, and R. B. Chandler. 2018. Generalized spatial mark-resight
748 models with an application to grizzly bears. *Journal of Applied Ecology* 55:157–168.
- 749
- 750 Figure Captions
- 751 Figure 1. Study area in the Elk Valley of southeast British Columbia between 2016 and 2022 is
752 enclosed by the white line and is the 99% Utilization Distribution of collared bear relocations.
753 Inset map shows the southern range of grizzly bears (dark shaded area) across western North
754 America.
- 755
- 756 Figure 2. Elk valley grizzly bear capture, telemetry, and mortality data collected between 2016
757 and 2022. A) capture locations, B) telemetry and mortality locations, C) capture weight (kg) by

758 sex and age with trend line fitted using locally weighted smoothing (LOESS), and D) percent
759 body fat at capture, measured using bioimpedance.

760

761 Figure 3. Collage depicting the life, death, and conflict of grizzly bears in the Elk Valley
762 between 2016 and 2022. A) A grizzly bear in the upper Elk River, BC. B) An adult female
763 grizzly bear (EVGF97) killed by a train; her three cubs were also killed in the same collision. C)
764 A subadult female grizzly bear (EVGF54) in the back of a BC Conservation Officer truck with
765 two dead pigs. EVGF54 was shot by a landowner while she was attacking their pigs. The
766 landowner had an electric fence, but it was not maintained and had shorted out due to long
767 vegetation against the fence, rendering it ineffective. D) A young male grizzly bear killed on
768 Highway 3 near Hosmer, BC. E) The cost of conflict to landowners. EVGF73 and her cubs'
769 paws can be seen on the door of this chicken coop that she opened. She and one of her yearling
770 cubs were illegally killed, and not reported, on an adjacent property one year later. F) A subadult
771 grizzly bear in an unpicked crab apple tree in Elkford, BC. G) A grizzly bear eating a road killed
772 deer in the valley bottom.

773

774 Figure 4. Reported human-bear conflicts as recorded in the Wildlife Alert Reporting System in
775 A) the Elk Valley study area between 2016 and 2021, B) seasonally within (A) per year, and C)
776 across the province between 2016 and 2021. The Elk Valley study area in southeast BC has the
777 highest rate of reported human-bear conflicts in the province (~65.3 conflict reports per 10,000
778 sq.km/year). The mean number of conflicts per 10,000 sq.km/year is 5.8 across the province. The

779 Lower Skeena valley near Kitimat and Terrace in west-central BC has a similar rate (64.8) to the
780 Elk Valley.

781
782 Figure 5. Elk Valley grizzly bear demographic data collected between 2016 and 2022.
783 Distributions represent the density of bootstrapped samples. A) Annual survival rates with
784 standard error bars. B) Reproductive rates with standard error bars. C) Comparison between Elk
785 Valley survival rates and published rates from across North America; error bars are 95% CIs. D)
786 Estimated unrecorded mortality; thick error bars cover 66% of the bootstrapped samples, and
787 thin error bars cover 95%.

788
789 Figure 6. Source-sink dynamics in the Elk Valley. A) Known immigrants from Alberta, Canada
790 and Montana, USA into the Elk Valley between 2016 and 2022. These immigrants were all
791 young (≤ 4) males and came from 77-95 km away. B) Intrinsic population growth rate of Elk
792 Valley grizzly bear population (i.e., without immigration and emigration). Thick error bars cover
793 66% of the bootstrapped samples, and thin error bars cover 95%. C) Abundance of grizzly bears
794 in the Elk Valley estimated from genetic spatial capture-recapture analysis between 2016 and
795 2021 and predicted from collar-based intrinsic population growth rate from (B). Projected
796 population trends to 2040 shown based on observed stable abundance, and abundance without
797 immigration subsidy.

798
799
800
801
802
803

804 Tables

805 Table 1. Mortality of collared animals while monitored, as well as human-caused grizzly
 806 mortalities reported in the CI database, and the number of ear-tagged animals known to have
 807 died by each mortality source between 2016 and 2022. The collar fates method estimates
 808 underreporting using unreported/monitored for actively collared animals killed by human causes
 809 (n=13). Unreported but monitored animals are shown in brackets. The CI ratio method uses the
 810 approach of McLellan et al. (2018) and compares the number of bears killed by COS with
 811 functioning radiocollars to those killed by other sources, for bears wearing functioning
 812 radiocollars and for uncollared bears recorded in the CI database. The ear tag ratio method uses
 813 the return ratio of ear tags of previously live-captured animals to radiocollar-monitored animals
 814 for COS kills and creates an expected number of tags returned for each mortality source, which
 815 is then used to calculate an unreported rate via 1-(returned/expected).

Cause	monitored	CI reporte d	tagged returned (reported)	tagged expected	unreporte d (collar method)	unreporte d (CI method)	unreporte d (eartag method)
Conflict	4 (2)	14	2	18	0.5	0.5	0.89
Conflict-COS	2 (0)	14	9	9	0	0	0
Road/Rail	6 (4)	11	3	27	0.67	0.74	0.89
Unk-human suspected	1 (1)	0	0	4.5	1	0	1
Hunter	0 (0)	3	0	0	0	0	0
Total	13 (7)	42	14	58.5	0.54	0.64	0.76


816

817 Table 2. Reported human-caused grizzly bear mortalities 2001-2021 within the Elk Valley study
 818 area compared to the rest of BC's grizzly bear range. Density [dead bears per 1,000 km²] shown
 819 for each area, with the total number of mortalities shown in brackets. Mortality data are from the
 820 British Columbia Compulsory Inspection database. The Elk Valley study area encompasses
 821 5,074 km² (0.66%) of the 764,330 km² BC grizzly bear range. "Excess" is how many times

822 higher the mortality density is than the rest of the province. The Elk Valley has
823 disproportionately high mortality for most sources. Note a total is not calculated because the
824 reporting rates differ within each source, so cannot be accurately totalled. Hunter kills are always
825 reported while the other sources are often underreported, as we show in Table 1.


Cause	Elk Valley	Rest of BC	Elk Valley share (%)	Excess (x higher than expected)
Human-bear conflict	13.44 (69)	1.25 (947)	7	11
Hunter	14.22 (73)	5.72 (4340)	2	2
Rail	3.7 (19)	0.03 (26)	42	108
Road	3.51 (18)	0.05 (37)	33	72
Human-bear conflict	13.44 (69)	1.25 (947)	7	11

826



Study area in the Elk Valley of southeast British Columbia between 2016 and 2022 is enclosed by the white line and is the 99% Utilization Distribution of collared bear relocations. Inset map shows the southern range of grizzly bears (dark shaded area) across western North America.

846x1058mm (72 x 72 DPI)




Elk valley grizzly bear capture, telemetry, and mortality data collected between 2016 and 2022. A) capture locations, B) telemetry and mortality locations, C) capture weight (kg) by sex and age with trend line fitted using locally weighted smoothing (LOESS), and D) percent body fat at capture, measured using bioimpedance.

1587x1111mm (72 x 72 DPI)




Collage depicting the life, death, and conflict of grizzly bears in the Elk Valley between 2016 and 2022. A) A grizzly bear in the upper Elk River, BC. B) An adult female grizzly bear (EVGF97) killed by a train; her three cubs were also killed in the same collision. C) A subadult female grizzly bear (EVGF54) in the back of a BC Conservation Officer truck with two dead pigs. EVGF54 was shot by a landowner while she was attacking their pigs. The landowner had an electric fence, but it was not maintained and had shorted out due to long vegetation against the fence, rendering it ineffective. D) A young male grizzly bear killed on Highway 3 near Hosmer, BC. E) The cost of conflict to landowners. EVGF73 and her cubs' paws can be seen on the door of this chicken coop that she opened. She and one of her yearling cubs were illegally killed, and not reported, on an adjacent property one year later. F) A subadult grizzly bear in an unpicked crab apple tree in Elkford, BC. G) A grizzly bear eating a road killed deer in the valley bottom.

338x314mm (150 x 150 DPI)



Reported human-bear conflicts as recorded in the Wildlife Alert Reporting System in A) the Elk Valley study area between 2016 and 2021, B) seasonally within (A) per year, and C) across the province between 2016 and 2021. The Elk Valley study area in southeast BC has the highest rate of reported human-bear conflicts in the province (~65.3 conflict reports per 10,000 sq.km/year). The mean number of conflicts per 10,000 sq.km/year is 5.8 across the province. The Lower Skeena valley near Kitimat and Terrace in west-central BC has a similar rate (64.8) to the Elk Valley.

1375x1111mm (72 x 72 DPI)




Elk Valley grizzly bear demographic data collected between 2016 and 2022. Distributions represent the density of bootstrapped samples. A) Annual survival rates with standard error bars. B) Reproductive rates with standard error bars. C) Comparison between Elk Valley survival rates and published rates from across North America; error bars are 95% CIs. D) Estimated unrecorded mortality; thick error bars cover 66% of the bootstrapped samples, and thin error bars cover 95%.


1375x1270mm (72 x 72 DPI)

A) Known immigrants into study area


3 male grizzly bears, 77-95 km displacements

**B) Intrinsic Population Growth Rate**

i.e., without immigration; 5000 bootstrapped samples

**C) Abundance**

Genetic capture recapture shows abundance is stable, collars show population would decline without immigration



Source-sink dynamics in the Elk Valley. A) Known immigrants from Alberta, Canada and Montana, USA into the Elk Valley between 2016 and 2022. These immigrants were all young (≤ 4) males and came from 77-95 km away. B) Intrinsic population growth rate of Elk Valley grizzly bear population (i.e., without immigration and emigration). Thick error bars cover 66% of the bootstrapped samples, and thin error bars cover 95%.

C) Abundance of grizzly bears in the Elk Valley estimated from genetic spatial capture-recapture analysis between 2016 and 2021 and predicted from collar-based intrinsic population growth rate from (B). Projected population trends to 2040 shown based on observed stable abundance, and abundance without immigration subsidy.

1270x1164mm (72 x 72 DPI)