

# WAHIDA NOORZAD ATTORNEY AT LAW

Ph: 925.321.7760 | Fax: 888.797.7040  
39111 Paseo Padre Pkwy | Suite 217 | Fremont, CA 94538

---

May 14, 2025

USCIS  
San Francisco Asylum Office  
P.O. Box 77530  
San Francisco, CA 94107

**Re: Request to Reschedule or Extend I-589 Asylum Interview for Ms. Jielian Yao  
(A# 209-788-393)**

Dear USCIS Officer,

I am Wahida Noorzad, the legal representative for Ms. Jielian Yao (A# 209-788-393) in connection with her I-589 application, which I have been handling since 2022.

I am writing in regard to the upcoming asylum interview for our client, Ms. Jielian Yao (A# 209-788-393), which is currently scheduled for **June 9, 2025, at 8:30 AM** at the **San Francisco Asylum Office**, located at 75 Hawthorne Street, 7th Floor, San Francisco, CA 94105.

Our office received a notice confirming this scheduled interview; however, due to the applicant's current mental health condition, I respectfully request a **rescheduling or extension** of this interview. Ms. Yao is currently suffering from what appears to be **Major Depressive Disorder with suicidal risk**. Additional mental health concerns, such as **delusional disorder or psychotic symptoms**, cannot be ruled out at this time. A physician has recommended **inpatient psychiatric treatment**, which further underscores the severity of her condition.

Ms. Yao is currently prescribed the following medications:

- Mirtazapine
- Olanzapine
- Pantoprazole
- Polyethylene Glycol
- Senna

She also has a scheduled follow-up medical appointment on **May 19, 2025**.

Given her fragile mental state and ongoing treatment, I respectfully request a postponement of the I-589 asylum interview to allow her time to stabilize and properly prepare for this important legal process.

# WAHIDA NOORZAD ATTORNEY AT LAW

Ph: 925.321.7760 | Fax: 888.797.7040

39111 Paseo Padre Pkwy | Suite 217 | Fremont, CA 94538

---


**Enclosed with this letter are the following supporting documents:**

- Copy of the I-589 Interview Notice for Ms. Jielian Yao (Page 1)
- Copies of medical records dated May 7, 2025, to May 13, 2025 (Page 2 - 8)
- Medical documentation dated May 8, 2025 (Page 9 - 11)
- Copies of current prescription medications (Page 12 – 37)

We sincerely apologize for any inconvenience this request may cause and deeply appreciate your understanding and assistance in this matter.

Thank you for your time and consideration.

Sincerely,



Wahida Noorzad  
Wahida Noorzad Attorney at Law  
39111 Paseo Padre Pkwy, Suite 217  
Tel: 925-321-7760

**THIS NOTICE DOES NOT GRANT ANY IMMIGRATION STATUS OR BENEFIT.**



|                                 |                |                                                                           |
|---------------------------------|----------------|---------------------------------------------------------------------------|
| Receipt Number<br>ZLA1600264580 |                | Case Type<br>I589 - APPLICATION FOR ASYLUM AND FOR WITHHOLDING OF REMOVAL |
| Received Date<br>12/19/2016     | Priority Date  | Applicant A209 788 393<br>YAO, JIELIAN                                    |
| Notice Date<br>04/26/2025       | Page<br>1 of 1 |                                                                           |

WAHIDA NOORZAD  
39111 PASEO PADRE PKWY  
217  
FREMONT CA 94538

**Please come to:**

75 Hawthorne St., 7th Floor  
San Francisco, CA 94105

**On (Date):** Monday, June 09, 2025  
**At (Time):** 08:30AM

We have mailed an official notice about this case (and any relevant documentation) according to the mailing preferences you chose on Form G-28, Notice of Entry of Appearance as Attorney or Accredited Representative. **This is a courtesy copy, not the official notice.**

**What the Official Notice Said**

**\*\*\* INTERVIEW RESCHEDULE NOTICE \*\*\***

As shown above, at your request we have rescheduled the asylum interview for you (and any family members included as dependents on your Form I-589, Application for Asylum and for Withholding of Removal). Your request to reschedule your asylum interview results in an applicant-caused delay for purposes of employment authorization based on your pending Form I-589. To RESOLVE this applicant-caused delay, you must appear for your rescheduled asylum interview. You may file a Form I-765, Application for Employment Authorization, based your pending asylum application 150 days after you filed your asylum application. You are not eligible to receive an Employment Authorization Document (EAD) until your asylum application has been pending for at least another 30 days, for a total of 180 days. 8 CFR 208.7(a)(1). The 150-day waiting period and the 180-day eligibility period, commonly referred to as the 180-Day Asylum EAD Clock, do not include delays that you request or cause while your asylum application is pending with an asylum office or with the Immigration Court. 8 CFR 208.7(a)(2).

If you fail to appear for this rescheduled asylum interview without good cause, or fail without good cause to provide a competent interpreter if you are required to do so, we may refer your Form I-589 to an immigration judge, and you will be ineligible for employment authorization based on your pending Form I-589. For additional information regarding establishing good cause, please review [uscis.gov/humanitarian/refugees-and-asylum/asylum/establishing-good-cause-or-exceptional-circumstances](https://uscis.gov/humanitarian/refugees-and-asylum/asylum/establishing-good-cause-or-exceptional-circumstances).

Please see the additional information on the back. You will be notified separately about any other cases you filed.

USCIS encourages you to sign up for a USCIS online account. To learn more about creating an account and the benefits, go to <https://www.uscis.gov/file-online>.

San Francisco Asylum Office  
U.S. CITIZENSHIP & IMMIGRATION SVC  
P.O. Box 77530  
San Francisco CA 94107

USCIS Contact Center: [www.uscis.gov/contactcenter](https://www.uscis.gov/contactcenter)





## 诊后摘要

Jielian Yao 病历号: 200451291 Mood problem 5/7/2025 - 5/13/2025 ZSFG 7C ACUTE PSYCHIATRIC B5

## Instructions



### 您的药物已更改

- 开始服用:
- mirtazapine (REMERON)
  - OLANZapine (ZyPREXA)
  - pantoprazole (PROTONIX)
  - polyethylene glycol (GLYCOLAX)
  - senna (senna)

请复核下方您更新后的药物列表。

## 您的后续步骤

### 请完成

- ☐ 在ZSFG OUTPATIENT PHARMACY领取下列药物
- mirtazapine
  - OLANZapine
  - pantoprazole
  - polyethylene glycol
  - senna



### Other instructions

#### Substance Use

If you smoke or chew tobacco, consider quitting. Call 1-800-No-Butts for smoking cessation counseling.

Do not use non-prescribed drugs. If you think you might have a problem with alcohol or drugs and would like to access to Substance Use Treatment, please contact TAP.

Treatment Access Program: 1-800-750-2727.

You may drop-in for information regarding substance abuse programs and services from Monday to Friday at 8:00 a.m. to 4:30 p.m. at 1380 Howard St, First Floor; San Francisco.

#### Discharge Question

If you have questions or concerns regarding your inpatient stay, or would like to obtain results from studies pending discharge, please contact the Charge Nurse at Psychiatric Emergency Services at Zuckerberg San Francisco General Hospital & Trauma Center (1001 Potrero Avenue, San Francisco, CA 94110 and phone number 415-206-8125.)

## Other instructions (续)

### Walk-in Mental Healthcare Support

- Dore Urgent Care Clinic, 52 Dore Street, 415-553-3100. They are open 24/7.
- Westside Crisis Clinic, 245 11th St, 415-355-0311. Drop-in hours are Mon-Fri from 8am-3pm.

**Hummingbird Places** offers drop-in support for food, shower, laundry, and rest. They are open 7 days a week, from 10am-6pm. \*Walk-ins welcomed, and you do not need a referral to receive the above services.

HUMMINGBIRD PLACE-POTRERO  
887 Potrero Ave, San Francisco 94110  
The facility is on ZSFGH grounds.  
Telephone: 628-206-2855

HUMMINGBIRD PLACE-VALENCIA  
1156 Valencia St, San Francisco 94110  
Telephone: 415-329-2005

### Suicide/Crisis Hotline

If you have an emergency, call 911.

You may also call:

- Mobile Crisis: 415-355-8300
- ZSFGH Psychiatric Emergency Services 415-206-8125
- Suicide Prevention/Crisis Hotline 415-781-0500

If you feel that you may hurt yourself, please call:

- Local: 415-781-0500
- National: 1-800-273-8255

You will be discharging home today to 17321 Via San Ardo San Lorenzo CA 94580-3527

## Procedure Information

No upcoming procedures

## 后续步骤

您当前没有即将到来的预约



### Additional Information

You have been connected to the following clinic for after care.

**Your follow up appointment is as follows:**

**5/19/2025 at 10:30 am with Jan Burch** (Please arrive 15 minutes prior to appointment)

### Pathways to Wellness

1970 Broadway STE 250  
Oakland, CA 94612  
1-800-975-7284

## 药物列表

|                                                                                                                                                                                                                                                                                                 | 上午                                                                                            | 中午左右 | 晚上 | 睡前                                                                                                 | 按需服用                                                                                            |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|------|----|----------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|
|  <b>mirtazapine 30 mg tablet</b><br>常用名: REMERON<br>Take 1 tablet (30 mg total) by mouth at bedtime for 14 days.<br>签署人: Colin Buzza                                                                           |                                                                                               |      |    | <br>1 tablet    |                                                                                                 |
|  <b>OLANZapine 10 mg tablet</b><br>常用名: ZyPREXA<br>每天口服一次 1 片。 (剂量 = 10 毫克)<br>签署人: Colin Buzza                                                                                                                | <br>1 tablet |      |    |                                                                                                    |                                                                                                 |
|  <b>pantoprazole 40 mg EC tablet</b><br>常用名: PROTONIX<br>每天早餐前口服一次 1 片。 (剂量 = 40 毫克) 不要压碎、咀嚼或掰开。<br>签署人: Colin Buzza                                                                                           | <br>1 tablet |      |    |                                                                                                    |                                                                                                 |
|  <b>polyethylene glycol 17 gram packet</b><br>常用名: GLYCOLAX<br>Stir and dissolve 1 (one) packet in 4 to 8 ounces of beverage and drink by mouth 1 time each day if needed for constipation<br>签署人: Colin Buzza |                                                                                               |      |    |                                                                                                    | <br>1 packet |
|  <b>senna 8.6 mg tablet</b><br>每天睡前一次口服 2 片。 (剂量 = 17.2 毫克)<br>签署人: Colin Buzza<br>药物通用名: senna                                                                                                              |                                                                                               |      |    | <br>2 tablets |                                                                                                 |

 如果您有任何疑问, 请咨询您的医生或其他医疗人员。

## 药物领取地点



在 **ZSFG OUTPATIENT PHARMACY** 领取下列药物

- mirtazapine 30 mg tablet
- OLANZapine 10 mg tablet
- pantoprazole 40 mg EC tablet
- polyethylene glycol 17 gram packet
- senna 8.6 mg tablet

地址: 1001 Potrero Avenue, SAN FRANCISCO CA 94110  
 小时: Monday-Friday 900AM-800PM Saturday 900AM - 100PM  
 电话: 628-206-8107

---

## MyChart注册

给您的医生发送信息，查看您的测试结果，更新您的处方，安排预约等。

前往<https://mychart.sfdph.org/MyChart/>，点击“现在注册”，然后输入您的激活码:2BR8T-N3WPG。激活码过期日期5/27/2025。

---

# 说明

## Additional Discharge Instructions

**Events leading to your inpatient hospitalization:** paranoid thought content and suicidal ideation w/ no plan

**Your diagnoses are:** Principal Problem:

Unspecified mood (affective) disorder (CMS/HCC)

Active Problems:

Unspecified psychosis not due to a substance or known physiological condition (CMS/HCC)

**Language preference:** Cantonese

## Important test results:

### Visit Vitals

|           |                                                               |
|-----------|---------------------------------------------------------------|
| BP        | 143/87 (BP Location: Right arm,<br>Patient Position: Sitting) |
| Pulse     | 77                                                            |
| Temp      | 36.7 °C (98.1 °F) (Temporal)                                  |
| Resp      | 17                                                            |
| Ht        | 1.5 m (4' 11.06")                                             |
| Wt        | 45.7 kg (100 lb 12.8 oz)                                      |
| SpO2      | 98%                                                           |
| BMI       | 20.32 kg/m <sup>2</sup>                                       |
| OB Status | Postmenopausal                                                |
| BSA       | 1.38 m <sup>2</sup>                                           |

Body mass index is 20.32 kg/m<sup>2</sup>.

### Lab Results

| Component | Value        | Date       |
|-----------|--------------|------------|
| COVID19   | NOT detected | 05/09/2025 |

### Lab Results

| Component | Value   | Date       |
|-----------|---------|------------|
| GLU       | 160     | 05/06/2025 |
| CALCIUM   | 9.5     | 05/06/2025 |
| NA        | 142     | 05/06/2025 |
| K         | 3.4 (L) | 05/06/2025 |
| CO2       | 29      | 05/06/2025 |
| CL        | 108 (H) | 05/06/2025 |
| BUN       | 11      | 05/06/2025 |
| LABCREA   | 0.64    | 05/06/2025 |

### Lab Results

| Component | Value | Date       |
|-----------|-------|------------|
| CHOL      | 154   | 05/08/2025 |

### Lab Results

| Component | Value | Date       |
|-----------|-------|------------|
| HDL       | 50    | 05/08/2025 |

### Lab Results



## 说明 (续)

| Component | Value | Date       |
|-----------|-------|------------|
| LDLCALC   | 87    | 05/08/2025 |

## Lab Results

| Component | Value | Date       |
|-----------|-------|------------|
| TRIG      | 89    | 05/08/2025 |

**Pending tests:** You have no pending tests at discharge.

**For your Medical Advance Directive,** the completion of an advanced directive is contraindicated due to your medical condition.

**For your Psychiatric Advance Directive,** the completion of an advanced directive is contraindicated due to your legal status.

The 11 element in this transitions record discussed with and provided to patient or caregiver at discharge, and a verbal endorsement of understanding was received by the discharging clinician.

**You will take the following medication upon discharge:**

- Zyprexa 10 mg once daily
- Mirtazapine 30 mg once at bedtime
- Protonix 40 mg once daily before breakfast
- Senna 8.6 once at bedtime as needed
- glycolax powder 17g packet once as needed

**You are to follow up with:**

Pathways to Wellness - Jan Burch 5/19 at 10:30 am

## 額外嘅出院指示

導致你住院嘅事件：偏激嘅思想內容同埋有計劃嘅自殺意念

你嘅診斷係：主要問題：

未明確嘅情緒(情緒)障礙( CMS / HCC )

活躍問題：

未指明嘅精神病，唔係因為某種物質或者已知嘅生理狀況( CMS / HCC )

待處理嘅檢查：你出院嗰陣有任何待處理嘅檢查。

對於你嘅醫療預先指令，由於你嘅醫療狀況，完成預先指令係禁忌。

對於你嘅精神科預先指令，由於你嘅法律地位，完成預先指令係禁忌。

呢個過渡期嘅11個元素記錄咗出院時同病人或照顧者討論同提供畀病人或照顧者，而出院嘅臨床醫生亦都收到咗口頭認可嘅理解。

你出院時會服用以下藥物：

- Zyprexa 10毫克每日一次
- 米塔扎平30毫克，睡前一次
- Protonix 40毫克，每日早餐前一次
- 根據需要，睡前一次服用塞納 8.6
- 甘草粉 17g 包一次按需要

## 说明 (续)

你要跟進：

健康之路 - Jan Burch 5/19上晝十點半

**Yao, Jielian**

MRN: 200451291

**Ridhaa F Sachidanandan, MD**

Resident

Emergency Medicine

Date of Service: 5/8/2025 1:30 PM

Psychiatric Legal Document    
Signed

State of California

Department of Health Care Services

Health and Human Services Agency

**NOTICE OF CERTIFICATION FOR  
INTENSIVE TREATMENT PURSUANT****CONFIDENTIAL PATIENT  
INFORMATION****TO SECTION 5250 (14 DAYS  
INTENSIVE TREATMENT) OR 5270.15  
(ADDITIONAL 30 DAYS INTENSIVE  
TREATMENT) OF THE WELFARE AND  
INSTITUTIONS CODE**☒ 14 day hold  
☐ 30 day holdThe authorized agency providing 14-day intensive treatment, County of **San Francisco** has custody of:Name **Jielian Yao**Address **17321 Via San Ardo****San Lorenzo CA 94580-3527**Marital Status **divorced**Date of birth **3/19/1963**Sex **female**

We, the undersigned, allege that the above-named person is, as a result of a mental disorder or

impairment by chronic alcoholism (mark all that apply):

☐ A danger to others ☒ A danger to himself or herself ☒ Gravely disabled as defined in subdivision (h) of Section 5008 of the Welfare and Institutions Code

The specific facts which form the basis for our opinion that the above-named person meets one or more of the classifications indicated above are as follows:

**Jielian endorses feeling targeted by people who are monitoring her asylum status, she is worried she will "die early" because of this, she has described her food as "rotten" and has not been eating well at home because of this, also states she will harm herself if she is kept in the hospital, has endorses thoughts of wanting to die to family.**

The above-named person has been informed of this evaluation, and has been advised of the need for, but has not been able or willing to accept treatment on a voluntary basis, or to accept referral to, the following services

**"Inpatient psychiatric hospitalization and treatment"**We, therefore, certify the above-named person to receive intensive treatment related to the mental disorder or impairment by chronic alcoholism beginning 05/08/25 in the intensive treatment facility herein named: **Zuckerberg San Francisco General Hospital**(Electronically signed by) Signature **Ridhaa F Sachidanandan,**  
**MD**

Date: 5/8/2025

Date: **5/8/2025**(Electronically signed by typing name) Signature: **Colin D Buzza,**  
**MD**

I hereby state that I delivered a copy of this notice this day to the above-named person and that I informed him or her that unless judicial review is requested a certification review hearing will be

held within four days of the date on which the person is certified for a period of intensive treatment and that an attorney or advocate will visit him or her to provide assistance in preparing for the hearing or to answer questions regarding his or her commitment or to provide other assistance. The court has been notified of this certification on this day.

(Electronically signed by) Signature **Ridhaa F Sachidanandan,** Date: **5/8/2025**  
**MD**

Original: Superior Court  
Copies: Person Certified -

Personally delivered

Person's Attorney  
Public Defender  
District Attorney  
Intensive Treatment

DHCS 1808 (Revised 12/2019)  
Facility

|                                                                                                                                                                          |                              |                                                                                               |                                          |  |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-----------------------------------------------------------------------------------------------|------------------------------------------|--|
| <b>SUPERIOR COURT OF CALIFORNIA, COUNTY OF<br/>SAN FRANCISCO</b><br>CIVIC CENTER COURTHOUSE, MENTAL HEALTH<br>DIVISION<br>400 McAllister Street, San Francisco, CA 94102 |                              |                                                                                               | <b>PATIENT NUMBER:</b><br>MRN: 200451291 |  |
| <b>CERTIFICATION REVIEW HEARING RECORD AND DECISION</b>                                                                                                                  |                              |                                                                                               |                                          |  |
| <input checked="" type="checkbox"/> <b>14 Day Certification</b><br>(Welf. & Inst. Code §5250)                                                                            |                              | <input type="checkbox"/> <b>Additional 30-Day Certification</b><br>(Welf. & Inst. Code §5270) |                                          |  |
| (PERSON CERTIFIED/PATIENT)                                                                                                                                               |                              |                                                                                               |                                          |  |
| FIRST NAME                                                                                                                                                               | MIDDLE NAME                  | LAST NAME                                                                                     | DATE OF BIRTH<br>(MM/DD/YYYY)            |  |
| Jiellian                                                                                                                                                                 |                              | Yao                                                                                           | 3/19/1963                                |  |
| <b>HEARING FACILITY</b>                                                                                                                                                  |                              | <b>CERTIFICATION FACILITY</b><br>(If different from Hearing Facility)                         | <b>WARD/UNIT:</b>                        |  |
| Zuckerberg San Francisco General<br>Hospital                                                                                                                             |                              |                                                                                               | PES                                      |  |
| INITIAL DETENTION<br>DATE                                                                                                                                                | 14-DAY<br>CERTIFICATION DATE | ADDITIONAL 30-<br>DAY CERTIFICATION<br>DATE                                                   | CERTIFICATION TYPE: (Use code<br>below)  |  |
| 05/05/25                                                                                                                                                                 | 05/08/25                     |                                                                                               | 05                                       |  |
| FACILITY REP:                                                                                                                                                            |                              | PATIENT REP:                                                                                  | ADVOCATE/OTHER:                          |  |
| TITLE:                                                                                                                                                                   |                              | TITLE:                                                                                        |                                          |  |

The above named patient was certified for ☒ **14 Days** / ☐ **Additional 30-Days** involuntary treatment pursuant to Welf. & Inst.

Code §5250 or §5270 et. seq.. The Certification was on the basis of one or more of the criteria specified by law: **(Mark all that apply)**

☐ A danger to others    ☒ A danger to self    ☒ Gravely disabled

A Certification Review Hearing was held this date to determine whether probable cause exists to continue to involuntarily detain the

10

patient for 14 days/30 additional days intensive treatment on the basis that as a result of mental disorder the patient is presently as specified, a danger to self, a danger to others, or gravely disabled (unable to provide for basic personal needs for food, clothing or shelter).

**THE PERSON CERTIFIED/PATIENT: (COURT USE ONLY)**

- ☐ Was present
 ☐ Was **NOT** present for other reasons:  
☐ Waived presence in accordance with Welf. & Inst. Code §5256
 ☐ Discharged
 ☐ Voluntarily admitted
 ☐ Hearing postponed
 ☐

**IT WAS DETERMINED THAT: (COURT USE ONLY)**
**CERTIFICATION CODE #:** (Use Code Below)

- ☐ Probable cause does not exist by Hearing Officer – no hearing held
 ☐ Patient went from 5150 to T-Con. No hearing held  
☐ Probable cause does not exist by findings on consensus
 ☐ **Probable cause does exist:** A danger to  
☐ Others ☐ Self ☐ Gravely disabled

The reason for continued involuntary detention and the evidence relied upon to form this determination (or evidence relied upon at the time this was filled out) is cited below:

| HEARING CONDUCTED BY | SIGNATURE | TITLE                         | DATE | TIME |
|----------------------|-----------|-------------------------------|------|------|
|                      |           | Mental Health Hearing Officer |      |      |

| CERTIFICATION CODES:    |                           |                      |                               |                                         |                                           |                                                    |
|-------------------------|---------------------------|----------------------|-------------------------------|-----------------------------------------|-------------------------------------------|----------------------------------------------------|
| 01) Danger to Self Only | 02) Danger to Others only | 03) Gravely Disabled | 04) Danger to Self and Others | 05) Danger to Self and Gravely Disabled | 06) Danger to Others and Gravely Disabled | 07) Danger to Self and Others and Gravely Disabled |

Original: Patient Chart Mail Copies: 1) SF Superior Court Mental Health Division 2) Person Certified (Personally deliver)

3) Person's Attorney/Public Defender 4) District Attorney 5) Facility Medical Director 6) Hearing Officer  
Electronically signed by Ridhaa F Sachidanandan, MD at 5/8/2025 1:49 PM

ED to Hosp-Admission (Current) on 5/5/2025 Note shared with patient

**Care Timeline**

05/07 Admitted from ED 2122  
2122

## Medication Monograph

---

**IMPORTANT: HOW TO USE THIS INFORMATION:** This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

### MIRTAZAPINE - ORAL

(mer-TAZE-uh-peen)

Remeron

Antidepressant medications are used to treat a variety of conditions, including depression and other mental/mood disorders. These medications can help prevent suicidal thoughts/attempts and provide other important benefits. However, studies have shown that a small number of people (especially people younger than 25) who take antidepressants for any condition may experience worsening depression, other mental/mood symptoms, or suicidal thoughts/attempts. It is very important to talk with the doctor about the risks and benefits of antidepressant medication (especially for people younger than 25), even if treatment is not for a mental/mood condition. Tell the doctor right away if you notice worsening depression/other psychiatric conditions, unusual behavior changes (including possible suicidal thoughts/attempts), or other mental/mood changes (including new/worsening anxiety, panic attacks, trouble sleeping, irritability, hostile/angry feelings, impulsive actions, severe restlessness, very rapid speech). Be especially watchful for these symptoms when a new antidepressant is started or when the dose is changed.

This medication is used to treat depression.

Mirtazapine works by helping to restore the balance of certain natural substances in the brain.

Read the Medication Guide provided by your pharmacist before you start using mirtazapine and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once daily at bedtime. The dosage is based on your medical condition and response to treatment. Use this medication regularly in order to get the most benefit from it. To help you remember, take it at the same time each day. It may take 1 to 4 weeks to notice improvement in your symptoms. Do not increase your dose or take it more often than prescribed. Keep taking this medication even if you feel well. Do not stop taking this medication without consulting your doctor. Some conditions may become worse when you suddenly stop this drug. Your dose may need to be gradually decreased. Tell your doctor if your condition does not get better or if it gets worse.

See also the Warning section. Dizziness, drowsiness, lightheadedness, increased appetite, weight gain, dry mouth, or constipation may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To relieve dry mouth, suck on (sugarless) hard candy or ice chips, chew (sugarless) gum, drink water or use a saliva substitute. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: swelling of the hands/feet, shaking (tremor), confusion. Get medical help right away if you have any very serious side effects, including: fast/irregular heartbeat, severe dizziness, fainting, eye pain/

---

**Medication Monograph**

---

swelling/redness, widened pupils, vision changes (such as seeing rainbows around lights at night, blurred vision). This medication may increase serotonin and rarely cause a very serious condition called serotonin syndrome/toxicity. The risk increases if you are also taking other drugs that increase serotonin, so tell your doctor or pharmacist of all the drugs you take (see Drug Interactions section). Get medical help right away if you develop some of the following symptoms: fast heartbeat, hallucinations, loss of coordination, severe dizziness, severe nausea/vomiting/diarrhea, twitching muscles, unexplained fever, unusual agitation/restlessness. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Before taking this medication, tell your doctor or pharmacist if you are allergic to it, or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: history or family history of psychiatric disorders (such as bipolar/manic-depressive disorder), history or family history of suicide attempts, liver disease, kidney disease, seizures, high blood cholesterol or triglyceride levels, heart disease (such as recent heart attack, angina), stroke, severe loss of body fluids (dehydration), low blood pressure,

personal or family history of glaucoma (angle-closure type). Mirtazapine may cause a condition that affects the heart rhythm (QT prolongation). QT prolongation can rarely cause serious (rarely fatal) fast/irregular heartbeat and other symptoms (such as severe dizziness, fainting) that need medical attention right away. The risk of QT prolongation may be increased if you have certain medical conditions or are taking other drugs that may cause QT prolongation. Before using mirtazapine, tell your doctor or pharmacist of all the drugs you take and if you have any of the following conditions: certain heart problems (heart failure, slow heartbeat, QT prolongation in the EKG), family history of certain heart problems (QT prolongation in the EKG, sudden cardiac death). Low levels of potassium or magnesium in the blood may also increase your risk of QT prolongation. This risk may increase if you use certain drugs (such as diuretics/"water pills") or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about using mirtazapine safely. This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Older adults may be more sensitive to the side effects of this drug, especially drowsiness and QT prolongation (see above). During pregnancy, this medication should be used only when clearly needed. If this medication is used during the last 3 months of pregnancy, rarely your newborn may develop symptoms including feeding or breathing difficulties, seizures, muscle stiffness, jitteriness or constant crying. Report any such symptoms to your doctor promptly. However, since untreated mental/mood disorders (such as depression) can be a serious condition, do not stop taking this medication unless your doctor directs you to do so. If you are planning pregnancy, become pregnant, or think you may be pregnant, discuss with your doctor right away the benefits and risks

## Medication Monograph

---

of using this medication during pregnancy. This drug passes into breast milk. Consult your doctor before breastfeeding.

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Taking MAO inhibitors with this medication may cause a serious (possibly fatal) drug interaction. Avoid taking MAO inhibitors (isocarboxazid, linezolid, metaxalone, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, tranylcypromine) during treatment with this medication. Most MAO inhibitors should also not be taken for two weeks before and after treatment with this medication. Ask your doctor when to start or stop taking this medication. The risk of serotonin syndrome/toxicity increases if you are also taking other drugs that increase serotonin. Examples include street drugs such as MDMA/"ecstasy," St. John's wort, certain antidepressants (including SSRIs such as fluoxetine/paroxetine, SNRIs such as duloxetine/venlafaxine), tryptophan, among others. The risk of serotonin syndrome/toxicity may be more likely when you start or increase the dose of these drugs. Tell your doctor or pharmacist if you are taking other products that cause drowsiness including alcohol, marijuana (cannabis), antihistamines (such as cetirizine, diphenhydramine), drugs for sleep or anxiety (such as alprazolam, diazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), and opioid pain relievers (such as codeine, hydrocodone). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your

pharmacist about using those products safely.

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: very fast/irregular heartbeat, severe dizziness, fainting.

Lab and/or medical tests should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. Do not share this medication with others.

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up.

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised May 2024. Copyright(c) 2025 First Databank, Inc.



## Medication Guide

### Mirtazapine Tablets, USP

(mir taz' a peen)

**What is the most important information I should know about mirtazapine tablets?**

**Mirtazapine tablets may cause serious side effects, including:**

- **Increased risk of suicidal thoughts or actions in some children and young adults.** Mirtazapine tablets and other antidepressant medicines may increase suicidal thoughts or actions in some people 24 years of age and younger, **especially within the first few months of treatment or when the dose is changed.** Mirtazapine tablets are not for use in children.
  - **Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.**

#### **How can I watch for and try to prevent suicidal thoughts and actions?**

- Pay close attention to any changes, especially sudden changes in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions. This is very important when an antidepressant medicine is started or when the dose is changed.
- Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.

**Call your healthcare provider or get emergency medical help right away if you or your family member have any of the following symptoms, especially if they are new, worse, or worry you:**

- attempts to commit suicide
- acting aggressive, being angry or violent
- new or worse depression
- panic attacks
- new or worse irritability
- an extreme increase in activity or talking (mania)
- acting on dangerous impulses
- thoughts about suicide or dying
- new or worse anxiety
- feeling very agitated or restless
- trouble sleeping
- other unusual changes in behavior or mood

### **What are mirtazapine tablets?**

Mirtazapine tablets are prescription medicines used to treat a certain type of depression called Major Depressive Disorder (MDD) in adults.

It is not known if mirtazapine tablets are safe and effective for use to treat MDD in children.

### **Who should not take mirtazapine tablets?**

**Do not take mirtazapine tablets if you:**

- take a Monoamine Oxidase Inhibitor (MAOI)
- have stopped taking an MAOI in the last 14 days
- are being treated with the antibiotic linezolid or intravenous methylene blue
- if you are allergic to mirtazapine or any of the ingredients in mirtazapine tablets. See the end of this Medication Guide for a complete list of ingredients in mirtazapine tablets.

Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid or intravenous methylene blue.

**Do not start taking an MAOI for at least 14 days after you stop treatment with mirtazapine tablets.**

**Before taking mirtazapine tablets, tell your healthcare provider about all of your medical conditions, including if you:**

- have a history of suicide or depression
- have a history or family history of bipolar disorder, mania or hypomania
- have a low white blood cell count
- have glaucoma (high pressure in the eye)

- have or had heart problems or stroke
- have an abnormal heart beat called QT prolongation or a family history of QT prolongation
- have seizures
- have high cholesterol or triglyceride levels
- have low sodium levels in your blood
- have or had kidney or liver problems
- have low blood pressure
- are pregnant or plan to become pregnant. It is not known if mirtazapine tablets will harm your unborn baby.
  - Talk to your healthcare provider if you become pregnant or think you may be pregnant during treatment with mirtazapine tablets.
  - If you become pregnant while taking mirtazapine tablets, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants. You can register by calling 1-844-405-6185 or visiting online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>. The purpose of this registry is to monitor the pregnancy outcomes in women who have been treated with mirtazapine tablets at any time during pregnancy.
- are breastfeeding or plan to breastfeed. Mirtazapine may pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with mirtazapine tablets.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Mirtazapine tablets and other medicines may affect each other causing possible serious side effects.

Mirtazapine tablets may affect the way other medicines work and other medicines may affect the way mirtazapine tablets work.

**Especially tell your healthcare provider if you take:**

- MAOIs
- medicines to treat migraine headaches known as triptans
- tricyclic antidepressants
- fentanyl
- lithium
- tramadol
- tryptophan
- buspirone
- amphetamines
- benzodiazepines
- St. John's Wort
- medicines used to treat mood, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs)
- medicines that may affect your heart rhythm (such as certain antibiotics and some antipsychotics)

Ask your healthcare provider if you are not sure if you are taking any of these medicines. Your healthcare provider can tell you if it is safe to take mirtazapine tablets with your other medicines.

Do not start or stop any other medicines during treatment with mirtazapine tablets without talking to your healthcare provider first. Stopping mirtazapine tablets suddenly may cause you to have serious side effects. See, **"What are the possible side effects of mirtazapine tablets?"**

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

**How should I take mirtazapine tablets?**

- Take mirtazapine tablets exactly as your healthcare provider tells you to. Do not change your dose or stop taking mirtazapine tablets without first talking to your healthcare provider.
- Your healthcare provider may need to change the dose of mirtazapine tablets until it is the right dose for you.
- Take mirtazapine tablets 1 time each day, preferably in the evening at bedtime.
- If you take too many mirtazapine tablets call your healthcare provider or poison control center at 1-800-222-1222 right away or go to the nearest hospital emergency room.

**What should I avoid while taking mirtazapine tablets?**

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how mirtazapine tablets affect you. Mirtazapine tablets can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly.
- Avoid drinking alcohol during treatment with mirtazapine tablets.
- Avoid taking medicines used to treat anxiety, insomnia, and seizures, called benzodiazepines, during treatment with mirtazapine tablets. Ask your healthcare provider if you are not sure if you take one of these medicines.

#### What are the possible side effects of mirtazapine tablets?

Mirtazapine tablets may cause serious side effects, including:

- See, **"What is the most important information I should know about mirtazapine tablets?"**
- **Low white blood cell count.** Tell your healthcare provider right away if you develop any signs or symptoms of a low white blood cell count, including:
  - o fever
  - o sore throat
  - o flu-like symptoms
  - o chills
  - o mouth or nose sores
  - o infections
- **Serotonin syndrome.** A potentially life-threatening problem called serotonin syndrome can happen when you take mirtazapine tablets with certain other medicines. See, **"Who should not take mirtazapine tablets?"** Stop taking mirtazapine tablets and call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the following signs and symptoms of serotonin syndrome:
  - o agitation
  - o confusion
  - o fast heart beat
  - o dizziness
  - o flushing
  - o tremors, stiff muscles, or muscle twitching
  - o seizures
  - o seeing or hearing things that are not real (hallucinations)
  - o coma
  - o blood pressure changes
  - o sweating
  - o high body temperature (hyperthermia)
  - o loss of coordination
  - o nausea, vomiting, diarrhea
- **Eye problems (angle-closure glaucoma).** Mirtazapine tablets may cause a certain type of eye problem called angle-closure glaucoma. Call your healthcare provider if you have eye pain, changes in your vision, or swelling or redness in or around the eye. Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.
- **Heart rhythm problems.**
- **Severe skin reaction.** Mirtazapine tablets may cause a severe skin reaction that may include rash, fever, swollen glands, and other organ involvement such as liver, kidney, lung and heart. The reaction may sometimes be fatal. Tell your healthcare provider right away if you experience any of these signs.
- **Increased appetite and weight gain.**
- **Sleepiness.** See, **"What should I avoid while taking mirtazapine tablets?"**
- **Mania or hypomania (manic episodes)** in people who have a history of bipolar disorder. Symptoms may include:
  - o greatly increased energy
  - o racing thoughts
  - o unusually grand ideas
  - o talking more or faster than usual
  - o severe trouble sleeping
  - o reckless behavior
  - o excessive happiness or irritability
- **Seizures (convulsions).**
- **Increased fat levels (cholesterol and triglycerides) in your blood.**
- **Low sodium levels in your blood (hyponatremia).** Low sodium levels in your blood may be serious and may cause death. Elderly people may be at greater risk for this. Signs and Symptoms of low sodium levels in your blood may include:
  - o headache
  - o memory changes
  - o difficulty concentrating
  - o confusion

- o weakness and unsteadiness on your feet which can lead to falls

**In severe or more sudden cases, signs and symptoms include:**

- o hallucinations (seeing or hearing things that are not real)
- o seizures
- o respiratory arrest
- **Changes in liver function tests.**
- **Discontinuation syndrome.** Suddenly stopping mirtazapine tablets may cause you to have serious side effects. Your healthcare provider may want to decrease your dose slowly. Symptoms may include:
  - o dizziness
  - o irritability and agitation
  - o anxiety
  - o sweating
  - o seizures
  - o ringing in your ears (tinnitus)
  - o nausea and vomiting
  - o problems sleeping
  - o tiredness
  - o confusion
  - o electric shock sensation (paresthesia)
  - o shaking (tremor)
  - o headache
  - o abnormal dreams
  - o changes in your mood
  - o hypomania

**The most common side effects of mirtazapine tablets include:**

- sleepiness
- increased appetite
- weight gain
- dizziness

These are not all the possible side effects of mirtazapine tablets.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**How should I store mirtazapine tablets?**

- Store mirtazapine tablets at room temperature between 20° to 25°C (68° to 77°F).
- Keep mirtazapine tablets away from light and moisture.

**Keep mirtazapine tablets and all medicines out of the reach of children.**

**General information about the safe and effective use of mirtazapine tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use mirtazapine tablets for a condition for which they were not prescribed. Do not give mirtazapine tablets to other people, even if they have the same symptoms that you have. They may harm them. You can ask your healthcare provider or pharmacist for information about mirtazapine tablets that is written for healthcare professionals.

**What are the ingredients in mirtazapine tablets?**

**Active ingredient:** mirtazapine

**Inactive ingredients:** anhydrous lactose, colloidal silicon dioxide, croscarmellose sodium, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch (corn), sodium lauryl sulfate, titanium dioxide and triacetin.

**For more information about mirtazapine tablets, call Mylan at 1-877-446-3679 (1-877-4-INFO-RX).**

The brands listed are trademarks of their respective owners.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured for:

**Mylan Pharmaceuticals Inc.**

Morgantown, WV 26505 U.S.A.

Manufactured by:

**ALPHAPHARM PTY LTD**

15 Garnet Street  
Carole Park QLD 4300  
Australia

Revised: 10/2022  
ALP:MTZP:R23mh/ALP:MG:MTZP:R7mh  
PM 2847/8

## Medication Monograph

---

**IMPORTANT: HOW TO USE THIS INFORMATION:** This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

### SENNOSIDES - ORAL

(SEN-oh-sides)

Sennosides are used to treat constipation. They may also be used to clean out the intestines before a bowel examination/surgery. Sennosides are known as stimulant laxatives. They work by keeping water in the intestines, which causes movement of the intestines.

Take this medication by mouth as directed by your doctor. If you are self-treating, follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. If you are using the chewable tablets, chew each tablet well and swallow. If you are using the dissolving strips, allow each strip to dissolve on the tongue and then swallow, with or without water. If you are using the liquid form of this medication, measure the dose carefully using a special measuring device/cup. Do not use a household spoon because you may not get the correct dose. Shake the container well before each dose. Dosage is based on your age, medical condition, and response to treatment. Do not increase your dose or take this drug more often than directed. Do not take this medication for more than 7 days unless directed by your doctor. Serious side effects may occur with overuse of this medication (see also Side Effects section). It may take 6 to 12 hours before this medication causes a bowel movement. Tell your doctor if

your condition lasts or gets worse, or if bleeding from the rectum occurs. If you think you may have a serious medical problem, get medical help right away.

Stomach/abdominal pain or cramping, nausea, diarrhea, or weakness may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. This medication may cause your urine to turn reddish-brown. This effect is harmless and will disappear when the medication is stopped. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: nausea/vomiting/diarrhea that doesn't stop, muscle cramps/weakness, irregular heartbeat, dizziness, decreased urination, mental/mood changes (such as confusion). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Before taking this medication, tell your doctor or pharmacist if you are allergic to it; or to senna; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: appendicitis or

## Medication Monograph

---

symptoms of appendicitis (such as nausea/vomiting, sudden or unexplained stomach/abdominal pain), a sudden change in bowel habits that lasts for longer than 2 weeks, bleeding from the rectum, intestinal blockage. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Chewable tablets, liquid products, or dissolving strips may contain sugar. Caution is advised if you have diabetes or any other condition that requires you to limit/avoid sugar in your diet. Ask your doctor or pharmacist about using this product safely. Tell your doctor if you are pregnant before using this medication. Using it for long periods during pregnancy is not recommended. Consult your doctor for more details. This medication passes into breast milk, but is unlikely to harm a nursing infant. Consult your doctor before breastfeeding.

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: nausea/vomiting/diarrhea that doesn't stop, severe stomach/abdominal pain or cramping.

Keep all medical and lab appointments. Lifestyle changes that may prevent or relieve constipation include exercising, drinking enough water, and

eating a proper diet with fiber-rich foods such as bran, fresh fruits/vegetables. Talk to your doctor or pharmacist about lifestyle changes that might benefit you.

If your doctor has directed you to take this medication on a regular schedule and you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised June 2024. Copyright(c) 2025 First Databank, Inc.



## Medication Monograph

---

**IMPORTANT: HOW TO USE THIS INFORMATION:** This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

### POLYETHYLENE GLYCOL (PEG) 3350 - ORAL

(paw-lee-ETH-ih-leen GLY-coll)

Miralax

This medication is used to treat occasional constipation. It works by holding water in the stool to soften the stool and increases the number of bowel movements. It is known as an osmotic-type laxative. This medication is also available without a prescription. If you are taking this medication for self-treatment, it is important to read the manufacturer's package instructions carefully so you know when to consult your doctor or pharmacist. (See also Precautions.)

Take this medication by mouth usually once daily, or as directed by your doctor or the directions on the product package. If you are prescribed the individual packets, mix the powder as directed by your doctor or pharmacist. If you are using the bulk bottle, use the provided cap to measure the prescribed dose. Mix the powder with a glass (4-8 ounces/120-240 milliliters) of liquid such as water, juice, soda, coffee, or tea. Before drinking the solution, stir the powder well until it completely dissolves. If you have any questions, ask your pharmacist. This medication may cause certain liquid thickeners (starch-based products) to not work well, which may increase the risk of choking. If you are using liquid thickeners with this

medication, talk to your pharmacist to select a liquid thickener which will not interact with this medication. It may take 2 to 4 days after you have started this medication to have a bowel movement. Do not increase your dose or take it more often than prescribed. Also, do not use this medication for more than 2 weeks unless directed by your doctor. Tell your doctor if your condition lasts or gets worse. Extended use or overuse may result in dependence on laxatives and chronic constipation. Overuse may cause diarrhea, dehydration, and mineral imbalance (such as low sodium). If you are using nonprescription polyethylene glycol for self-treatment of occasional constipation and this treatment has not worked after 7 days, talk to your doctor for additional treatment advice.

Nausea, abdominal cramping, or gas may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have diarrhea that doesn't stop, stomach/abdominal pain that is severe or doesn't go away, an excessive number of bowel movements, bloody stools, or rectal bleeding while using this medication. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.



## Medication Monograph

---

Before taking this medication, tell your doctor or pharmacist if you are allergic to polyethylene glycol; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: nausea/vomiting/abdominal pain that doesn't go away, stomach/intestinal problems (such as bowel obstruction, irritable bowel syndrome), kidney disease. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially diarrhea. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breastfeeding.

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: severe diarrhea, dizziness, decrease in the amount of urine.

Do not share this medication with others. To

prevent constipation, remember to exercise regularly, drink adequate fluids, and eat fiber-rich foods such as bran, whole grain, fresh fruits, and vegetables. Talk to your doctor or pharmacist about lifestyle changes that might benefit you.

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up.

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised May 2024. Copyright(c) 2025 First Databank, Inc.

## Medication Monograph

---

**IMPORTANT: HOW TO USE THIS INFORMATION:** This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

### PANTOPRAZOLE DELAYED-RELEASE - ORAL

(pan-TOE-pruh-zole)

Protonix

Pantoprazole is used to treat certain stomach and esophagus problems (such as acid reflux). It works by decreasing the amount of acid your stomach makes. This medication relieves symptoms such as heartburn, difficulty swallowing, and cough. It helps heal acid damage to the stomach and esophagus, helps prevent ulcers, and may help prevent cancer of the esophagus. Pantoprazole belongs to a class of drugs known as proton pump inhibitors (PPIs).

This section contains uses of this drug that are not listed in the approved US professional labeling for the drug but that may be prescribed by your health care professional. Use this drug for a condition that is listed in this section only if it has been so prescribed by your health care professional. This drug may also be used for stomach and intestinal ulcers.

Read the Medication Guide provided by your pharmacist before you start taking pantoprazole and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, usually once daily. The dosage and length of treatment are based on your medical condition

and response to treatment. Children's dosage is also based on weight. If you are taking the tablets, you may take them with or without food. Swallow the tablets whole. Do not split, crush, or chew the medication. Doing so may destroy the drug. If you are taking the granules, take your dose 30 minutes before a meal. To take it by mouth, open the packet and mix the granules in applesauce or apple juice. Do not mix with other foods or liquids. Do not crush or chew the granules. Sprinkle the granules on 1 teaspoon (5 milliliters) of applesauce and swallow all of the mixture right away (within 10 minutes). Follow with sips of water. Or you can mix the granules with 1 teaspoon (5 milliliters) of apple juice in a small cup, stir for 5 seconds, and swallow all of the mixture right away. To make sure you take the entire dose, rinse the cup once or twice with apple juice to mix any remaining granules, and swallow the juice. Do not prepare the mixture ahead of time for later use. If you are giving the granules through a tube into the stomach (nasogastric or gastric tube), ask your health care professional for detailed instructions on how to properly mix and give it. If needed, antacids may be taken along with this medication. If you are also taking sucralfate, take pantoprazole at least 30 minutes before sucralfate. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Continue to take this medication for the prescribed length of treatment even if you are feeling better. Tell your doctor if your condition lasts or gets worse. The risk of side effects goes up over time. Ask your doctor how long you should take this medication.

Headache or diarrhea may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have

## Medication Monograph

---

any serious side effects, including: symptoms of a low magnesium blood level (such as muscle spasms, irregular heartbeat, seizures), signs of lupus (such as rash on nose and cheeks, new or worsening joint pain), signs of kidney problems (such as change in the amount of urine, pink/bloody urine). This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Rarely, proton pump inhibitors (such as pantoprazole) have caused vitamin B-12 deficiency. The risk is increased if they are taken every day for a long time (3 years or longer). Tell your doctor right away if you develop symptoms of vitamin B-12 deficiency (such as unusual weakness, sore tongue, or numbness/tingling of the hands/feet). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Before taking pantoprazole, tell your doctor or pharmacist if you are allergic to it; or to similar drugs (such as lansoprazole, omeprazole); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your

pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, lupus. Some symptoms may actually be signs of a more serious condition. Get medical help right away if you have: heartburn with lightheadedness/sweating/dizziness, chest/jaw/arm/shoulder pain (especially with shortness of breath, unusual sweating), unexplained weight loss. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Proton pump inhibitors (such as pantoprazole) may increase your risk for bone fractures, especially with longer use, higher doses, and in older adults. Talk with your doctor or pharmacist about ways to prevent bone loss/fracture, such as by taking calcium (such as calcium citrate) and vitamin D supplements. Older adults may be more sensitive to the side effects of this drug, especially bone loss and fractures (see above), and *C. difficile* infection (see Side Effects section). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication passes into breast milk. Consult your doctor before breastfeeding.

See also How to Use section. Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: methotrexate (especially high-dose treatment). Some products need stomach acid so that the body can absorb them properly. Pantoprazole decreases stomach acid, so it may change how well these products work. Some affected products include ampicillin, atazanavir, erlotinib, levoketoconazole, nelfinavir,

## Medication Guide

### Pantoprazole Sodium Delayed-Release Tablets, USP (pan toe' pra zole soe'dee um)

#### What is the most important information I should know about pantoprazole sodium delayed-release tablets?

You should take pantoprazole sodium delayed-release tablets exactly as prescribed, at the lowest dose possible and for the shortest time needed.

Pantoprazole sodium delayed-release tablets may help your acid-related symptoms, but you could still have serious stomach problems. Talk with your doctor.

#### Pantoprazole sodium delayed-release tablets can cause serious side effects, including:

- **A type of kidney problem (acute tubulointerstitial nephritis).** Some people who take proton pump inhibitor (PPI) medicines, including pantoprazole sodium delayed-release tablets, may develop a kidney problem called acute tubulointerstitial nephritis that can happen at any time during treatment with pantoprazole sodium delayed-release tablets. Call your doctor right away if you have a decrease in the amount that you urinate or if you have blood in your urine.
- **Diarrhea caused by an infection (*Clostridium difficile*) in your intestines.** Call your doctor right away if you have watery stools or stomach pain that does not go away. You may or may not have a fever.
- **Bone fractures (hip, wrist, or spine).** Bone fractures in the hip, wrist, or spine may happen in people who take multiple daily doses of PPI medicines and for a long period of time (a year or longer). Tell your doctor if you have a bone fracture, especially in the hip, wrist, or spine.
- **Certain types of lupus erythematosus.** Lupus erythematosus is an autoimmune disorder (the body's immune cells attack other cells or organs in the body). Some people who take PPI medicines, including pantoprazole sodium delayed-release tablets, may develop certain types of lupus erythematosus or have worsening of the lupus they already have. Call your doctor right away if you have new or worsening joint pain or a rash on your cheeks or arms that gets worse in the sun.
- **Low magnesium and other mineral levels in your body** can happen in people who have taken pantoprazole sodium delayed-release tablets for at least 3 months. Tell your doctor if you have symptoms of low magnesium levels, including seizures, dizziness, irregular heartbeat, jitteriness, muscle aches or weakness, and spasms of hands, feet or voice.

Talk to your doctor about your risk of these serious side effects.

Pantoprazole sodium delayed-release tablets can have other serious side effects. See **"What are the possible side effects of pantoprazole sodium delayed-release tablets?"**

#### What are pantoprazole sodium delayed-release tablets?

A prescription medicine called a proton pump inhibitor (PPI) used to reduce the amount of acid in your stomach.

#### In adults, pantoprazole sodium delayed-release tablets are used for:

- up to 8 weeks for the healing and symptom relief of acid-related damage to the lining of the esophagus (called erosive esophagitis or EE). Your doctor may prescribe another 8 weeks of pantoprazole sodium delayed-release tablets in patients whose EE does not heal.
- maintaining healing of EE and to help prevent the return of heartburn symptoms caused by GERD. It is not known if pantoprazole sodium delayed-release tablets are safe and effective when used for longer than 12 months for this purpose.
- the long-term treatment of conditions where your stomach makes too much acid. This includes a rare condition called Zollinger-Ellison Syndrome.

#### In children 5 years of age and older, pantoprazole sodium delayed-release tablets are used for:

- up to 8 weeks for the healing and symptom relief of EE.

It is not known if pantoprazole sodium delayed-release tablets are safe if used longer than 8 weeks in children.

Pantoprazole sodium delayed-release tablets are not for use in children under 5 years of age.

It is not known if pantoprazole sodium delayed-release tablets are safe and effective in children for treatment other than EE.

**Do not take pantoprazole sodium delayed-release tablets if you are:**

- allergic to pantoprazole sodium, any other PPI medicine, or any of the ingredients in pantoprazole sodium delayed-release tablets. See the end of this Medication Guide for a complete list of ingredients.
- taking a medicine that contains rilpivirine (EDURANT<sup>®</sup>, COMPLERA<sup>®</sup>, ODEFSEY<sup>®</sup>, JULUCA<sup>®</sup>) used to treat HIV-1 (Human Immunodeficiency Virus).

**Before taking pantoprazole sodium delayed-release tablets, tell your doctor about all of your medical conditions, including if you:**

- have low magnesium levels, low calcium levels and low potassium levels in your blood.
- are pregnant or plan to become pregnant. Pantoprazole sodium delayed-release tablets may harm your unborn baby. Tell your doctor if you become pregnant or think you may be pregnant during treatment with pantoprazole sodium delayed-release tablets.
- are breastfeeding or plan to breastfeed. Pantoprazole can pass into your breast milk. Talk with your doctor about the best way to feed your baby if you take pantoprazole sodium delayed-release tablets.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins and herbal supplements. **Especially tell your doctor if you take** methotrexate (Otrexup<sup>®</sup>, Rasuvo<sup>®</sup>, Trexall<sup>®</sup>, XATMEP<sup>®</sup>), digoxin (LANOXIN<sup>®</sup>), or a water pill (diuretic).

**How should I take pantoprazole sodium delayed-release tablets?**

- Take pantoprazole sodium delayed-release tablets exactly as prescribed by your doctor.  
**Pantoprazole sodium delayed-release tablets:**
  - **Do not split, chew, or crush pantoprazole sodium delayed-release tablets.**
  - Swallow pantoprazole sodium delayed-release tablets whole, with or without food.
  - Tell your doctor if you are not able to swallow your pantoprazole sodium delayed-release tablet.
  - You may use antacids while taking pantoprazole sodium delayed-release tablets.
- If you miss a dose of pantoprazole sodium delayed-release tablets, take it as soon as possible. If it is almost time for your next dose, do not take the missed dose. Take the next dose at your regular time. Do not take 2 doses at the same time.
- If you take too many pantoprazole sodium delayed-release tablets, call your doctor or your poison control center at 1-800-222-1222 right away or go to the nearest emergency room.

**What are the possible side effects of pantoprazole sodium delayed-release tablets?**

**Pantoprazole sodium delayed-release tablets can cause serious side effects, including:**

- See "What is the most important information I should know about pantoprazole sodium delayed-release tablets?"
- **Low vitamin B-12 levels** in your body can happen in people who have taken pantoprazole sodium delayed-release tablets for a long time (more than 3 years). Tell your doctor if you have symptoms of low vitamin B-12 levels, including shortness of breath, lightheadedness, irregular heartbeat, muscle weakness, pale skin, feeling tired, mood changes, and tingling or numbness in the arms and legs.
- **Stomach growths (fundic gland polyps).** People who take PPI medicines for a long time have an increased risk of developing a certain type of stomach growths called fundic gland polyps, especially after taking PPI medicines for more than 1 year.
- **Severe skin reactions.** Pantoprazole sodium delayed-release tablets can cause rare but severe skin reactions that may affect any part of your body. These serious skin reactions may need to be treated in a hospital and may be life threatening:
  - Skin rash which may have blistering, peeling or bleeding on any part of your skin (including your lips, eyes, mouth, nose, genitals, hands or feet).
  - You may also have fever, chills, body aches, shortness of breath, or enlarged lymph nodes.

Stop taking pantoprazole sodium delayed-release tablets and call your doctor right away. These symptoms may be the first sign of a severe skin reaction.

#### Medication Monograph

---

pazopanib, rilpivirine, sparsentan, certain azole antifungals (itraconazole, ketoconazole, posaconazole), among others. This medication may interfere with certain lab tests (such as urine test for tetrahydrocannabinol-THC, blood test to find certain tumors), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669.

Do not share this medication with others. If your doctor instructs you to use this medication regularly for a long time, lab and/or medical tests (such as a magnesium blood test, vitamin B-12 levels) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details.

If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised August 2024. Copyright (c) 2025 First Databank, Inc.

**The most common side effects of pantoprazole sodium delayed-release tablets in adults include:** headache, diarrhea, nausea, stomach-area (abdominal) pain, vomiting, gas, dizziness, and joint pain.

**The most common side effects of pantoprazole sodium delayed-release tablets in children include:** upper respiratory infection, headache, fever, diarrhea, vomiting, rash, and stomach-area (abdominal) pain.

These are not all the possible side effects of pantoprazole sodium delayed-release tablets. **Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**How should I store pantoprazole sodium delayed-release tablets?**

Store pantoprazole sodium delayed-release tablets at room temperature between 20° to 25°C (68° to 77°F).

**Keep pantoprazole sodium delayed-release tablets and all medicines out of the reach of children.**

**General information about the safe and effective use of pantoprazole sodium delayed-release tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use pantoprazole sodium delayed-release tablets for a condition for which they were not prescribed. Do not give pantoprazole sodium delayed-release tablets to other people, even if they have the same symptoms that you have. They may harm them. You can ask your doctor or pharmacist for information about pantoprazole sodium delayed-release tablets that is written for health professionals.

**What are the ingredients in pantoprazole sodium delayed-release tablets?**

**Active ingredient:** pantoprazole sodium sesquihydrate

**Inactive ingredients in pantoprazole sodium delayed-release tablets:** anhydrous sodium carbonate, calcium stearate, crospovidone, hypromellose, mannitol, methacrylic acid and ethyl acrylate copolymer dispersion, povidone, propylene glycol, titanium dioxide, triethyl citrate and yellow iron oxide. The black imprinting ink contains black iron oxide, propylene glycol and shellac glaze.

For more information call Mylan at 1-877-446-3679 (1-877-4-INFO-RX).

This Medication Guide has been approved by the U.S. Food and Drug Administration.

The brands listed are trademarks of their respective owners.

Manufactured for:

**Mylan Pharmaceuticals Inc.**

Morgantown, WV 26505 U.S.A.

Manufactured by:

**Mylan Laboratories Limited**

Hyderabad — 500 096, India

75098854

Revised: 8/2023

MX:PANTDR:R17mmh/MX:MG:PANTDR:R12m/MX:MG:PANTDR:R11mh



## Medication Monograph

---

**IMPORTANT: HOW TO USE THIS INFORMATION:** This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

### OLANZAPINE - ORAL

(oh-LAN-za-peen)

Zyprexa

There may be a slightly increased risk of serious, possibly fatal side effects (such as stroke, heart failure, fast/irregular heartbeat, pneumonia) when this medication is used by older adults with dementia. This medication is not approved for the treatment of dementia-related behavior problems. Discuss the risks and benefits of this medication, as well as other effective and possibly safer treatments for dementia-related behavior problems, with the doctor. If you are using olanzapine in combination with other medication to treat depression, also carefully read the drug information for the other medication.

This medication is used to treat certain mental/mood disorders (such as schizophrenia, bipolar disorder). It may also be used in combination with other medication to treat depression. Olanzapine belongs to a class of drugs known as atypical antipsychotics. It works by helping to restore the balance of certain natural substances in the brain. Talk to the doctor about the risks and benefits of treatment (especially when used by teenagers). See also Precautions section.

This section contains uses of this drug that are not listed in the approved professional labeling

for the drug but that may be prescribed by your health care professional. Use this drug for a condition that is listed in this section only if it has been so prescribed by your health care professional. This drug may also be used to reduce the risk of nausea and vomiting caused by cancer drug treatment (chemotherapy).

Read the Medication Guide provided by your pharmacist before you start taking olanzapine and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once daily. The dosage is based on your medical condition and response to treatment. To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose and gradually increase your dose. Follow your doctor's instructions carefully. Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication even if you feel well. Do not stop taking this medication without consulting your doctor. Tell your doctor if your condition lasts or gets worse.

Drowsiness, dizziness, lightheadedness, stomach upset, dry mouth, constipation, increased appetite, or weight gain may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Dizziness and lightheadedness can increase the risk of falling. Get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: difficulty swallowing, shaking (tremor), slow heartbeat, fainting, mental/mood changes (such as confusion, restlessness), numbness/tingling of arms/legs, yellowing eyes/skin, severe stomach/abdominal pain, trouble



**Medication Monograph**

---

urinating, interrupted breathing during sleep. This drug may rarely make your blood sugar rise, which can cause or worsen diabetes. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. This drug may also cause significant weight gain and a rise in your blood cholesterol (or triglyceride) levels, especially in teenagers. These effects, along with diabetes, may increase your risk for developing heart disease. Discuss the risks and benefits of treatment with your doctor. (See also Notes section.) Olanzapine may rarely cause a condition known as tardive dyskinesia. In some cases, this condition may be permanent. Tell your doctor right away if you develop any unusual/uncontrolled movements (especially of the face, lips, mouth, tongue, arms or legs). This medication may increase a certain natural substance (prolactin) made by your body. For females, this increase in prolactin may result in unwanted breast milk, missed/stopped periods, or difficulty becoming pregnant. For males, it may result in decreased sexual ability, inability to produce sperm, or enlarged breasts. If you develop any of these symptoms, tell your doctor right away. Get medical help right away if you have any very serious side effects, including: seizures. This medication may rarely cause a very serious condition called neuroleptic malignant syndrome (NMS). Get medical help right away if you have any of the following symptoms: fever, muscle stiffness/pain/tenderness/weakness, severe tiredness, severe confusion, sweating, fast/irregular heartbeat, dark urine, signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble

breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Before taking olanzapine, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver problems, seizures, difficulty swallowing, low white blood cell count, dementia, difficulty urinating (for example, due to enlarged prostate), glaucoma (narrow angle), stomach/intestinal problems (such as blockage, chronic constipation, paralytic ileus), smoking, personal or family history of diabetes, heart disease, high cholesterol/triglyceride levels, breathing trouble during sleep (sleep apnea). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may make you sweat less, making you more likely to get heat stroke. Avoid doing things that may cause you to overheat, such as hard work or exercise in hot weather, or using hot tubs. When the weather is hot, drink a lot of fluids and dress lightly. If you overheat, quickly look for a place to cool down and rest. Get medical help right away if you have a fever that does not go away, mental/mood changes,



**Medication Monograph**

---

headache, or dizziness. Teenagers may be more sensitive to the side effects of this drug, especially weight gain, and also increased amounts of cholesterol, triglycerides, and prolactin. See also Side Effects section for more details. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, constipation, trouble urinating, confusion, dizziness, and lightheadedness. Drowsiness, confusion, dizziness, and lightheadedness can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Babies born to mothers who have used this drug during the last 3 months of pregnancy may rarely develop symptoms including muscle stiffness or shakiness, drowsiness, feeding/breathing difficulties, or constant crying. If you notice any of these symptoms in your newborn especially during their first month, tell the doctor right away. Since untreated mental/mood problems (such as schizophrenia, bipolar disorder, depression) can be a serious condition, do not stop taking this medication unless directed by your doctor. If you are planning pregnancy, become pregnant, or think you may be pregnant, immediately discuss with your doctor the benefits and risks of using this medication during pregnancy. This drug passes into breast milk and may have undesirable effects on a nursing infant. Tell the doctor right away if your baby develops symptoms such as muscle stiffness or shakiness, unusual sleepiness, or difficulty feeding. Consult your doctor before breastfeeding.

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: metoclopramide. Tell your doctor or

pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: severe drowsiness/dizziness, fast/irregular heartbeat, unusual/uncontrolled movements, seizures.

Do not share this medication with others. Lab and/or medical tests (such as blood sugar, weight, blood pressure, blood cholesterol/triglyceride levels, liver function) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details.

If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

OLANzapine 10 mg tablet

Call ZSFG OUTPATIENT PHARMACY TO REQUEST A PHARMACIST TELEPHONE CONSULTATION AT 628-206-8107  
Monday-Friday 900AM-800PM Saturday 900AM - 100PM

**Medication Monograph**

---

Information last revised May 2024. Copyright(c)  
2025 First Databank, Inc.

Dispense with Medication Guide available at:

[www.drreddys.com/medguide/olanzapinetabs.pdf](http://www.drreddys.com/medguide/olanzapinetabs.pdf)

## Medication Guide

### Olanzapine Tablets USP

(oh lan' za peen)

Read the Medication Guide that comes with olanzapine tablets before you start taking them and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about your medical condition or treatment. Talk with your doctor or pharmacist if there is something you do not understand or you want to learn more about olanzapine tablets.

#### What is the most important information I should know about olanzapine tablets?

Olanzapine tablets may cause serious side effects, including:

1. Increased risk of death in elderly people who are confused, have memory loss and have lost touch with reality (dementia-related psychosis).
2. High blood sugar (hyperglycemia).
3. High fat levels in your blood (increased cholesterol and triglycerides), especially in teenagers age 13 to 17, or when used in combination with fluoxetine in children age 10 to 17.
4. Weight gain, especially in teenagers age 13 to 17, or when used in combination with fluoxetine in children age 10 to 17.

These serious side effects are described below.

**1. Increased risk of death in elderly people who are confused, have memory loss and have lost touch with reality (dementia-related psychosis).** Olanzapine tablets are not approved for treating psychosis in elderly people with dementia.

**2. High blood sugar (hyperglycemia).** High blood sugar can happen if you have diabetes already or if you have never had diabetes. High blood sugar could lead to:

- a build up of acid in your blood due to ketones (ketoacidosis)
- coma
- death

Your doctor should do tests to check your blood sugar before you start taking olanzapine tablets and during treatment. In people who do not have diabetes, sometimes high blood sugar goes away when olanzapine tablets are stopped. People with diabetes and some people who did not have diabetes before taking olanzapine tablets need to take medicine for high blood sugar even after they stop taking olanzapine tablets.

If you have diabetes, follow your doctor's instructions about how often to check your blood sugar while taking olanzapine tablets.

**Call your doctor** if you have any of these symptoms of high blood sugar (hyperglycemia) while taking olanzapine tablets:

- feel very thirsty
- need to urinate more than usual
- feel very hungry
- feel weak or tired
- feel sick to your stomach



- feel confused or your breath smells fruity

**3. High fat levels in your blood (cholesterol and triglycerides).** High fat levels may happen in people treated with olanzapine tablets, especially in teenagers (13 to 17 years old), or when used in combination with fluoxetine in children (10 to 17 years old). You may not have any symptoms, so your doctor should do blood tests to check your cholesterol and triglyceride levels before you start taking olanzapine tablets and during treatment.

**4. Weight gain.** Weight gain is very common in people who take olanzapine tablets. Teenagers (13 to 17 years old) are more likely to gain weight and to gain more weight than adults. Children (10 to 17 years old) are also more likely to gain weight and to gain more weight than adults when olanzapine is used in combination with fluoxetine. Some people may gain a lot of weight while taking olanzapine tablets, so you and your doctor should check your weight regularly. Talk to your doctor about ways to control weight gain, such as eating a healthy, balanced diet, and exercising.

### **What are Olanzapine tablets?**

Olanzapine tablets are prescription medicine used to treat:

- schizophrenia in people age 13 or older.
- bipolar disorder, including:
  - manic or mixed episodes that happen with bipolar I disorder in people age 13 or older.
  - manic or mixed episodes that happen with bipolar I disorder, when used with the medicine lithium or valproate, in adults.
- long-term treatment of bipolar I disorder in adults.
- episodes of depression that happen with bipolar I disorder, when used with the medicine fluoxetine (Prozac®), in people age 10 or older.
- Episodes of depression that do not get better after 2 other medicines, also called treatment resistant depression, when used with the medicine fluoxetine (Prozac), in adults.

Olanzapine tablets have not been approved for use in children under 13 years of age. Olanzapine in combination with fluoxetine has not been approved for use in children under 10 years of age.

The symptoms of schizophrenia include hearing voices, seeing things that are not there, having beliefs that are not true, and being suspicious or withdrawn.

The symptoms of bipolar I disorder include alternating periods of depression and high or irritable mood, increased activity and restlessness, racing thoughts, talking fast, impulsive behavior, and a decreased need for sleep.

The symptoms of treatment resistant depression include decreased mood, decreased interest, increased guilty feelings, decreased energy, decreased concentration, changes in appetite, and suicidal thoughts or behavior.

Some of your symptoms may improve with treatment. If you do not think you are getting better, call your doctor.

### **What should I tell my doctor before taking olanzapine tablets?**

Olanzapine tablets may not be right for you. Before starting olanzapine tablets, tell your doctor if you have or had:

- heart problems
- seizures
- diabetes or high blood sugar levels (hyperglycemia)
- high cholesterol or triglyceride levels in your blood
- liver problems
- low or high blood pressure
- strokes or "mini-strokes" also called transient ischemic attacks (TIAs)

- Alzheimer's disease
- narrow-angle glaucoma
- enlarged prostate in men
- bowel obstruction
- breast cancer
- thoughts of suicide or hurting yourself
- any other medical condition
- are pregnant or plan to become pregnant. It is not known if olanzapine tablets will harm your unborn baby.
- If you become pregnant while receiving olanzapine tablets, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or go to <http://womensmentalhealth.org/clinical-andresearch-programs/pregnancyregistry/>.
- are breast-feeding or plan to breast-feed. Olanzapine passes into your breast milk. Talk to your doctor about the best way to feed your baby if you take olanzapine tablets.

Tell your doctor if you exercise a lot or are in hot places often.

The symptoms of bipolar I disorder, treatment resistant depression, or schizophrenia may include **thoughts of suicide** or of hurting yourself or others. If you have these thoughts at any time, tell your doctor or go to an emergency room right away.

**Tell your doctor about all the medicines that you take**, including prescription and non-prescription medicines, vitamins, and herbal supplements. Olanzapine tablets and some medicines may interact with each other and may not work as well, or cause possible serious side effects. Your doctor can tell you if it is safe to take olanzapine tablets with your other medicines. Do not start or stop any medicine while taking olanzapine tablets without talking to your doctor first.

### **How should I take olanzapine tablets?**

- Take olanzapine tablets exactly as prescribed. Your doctor may need to change (adjust) the dose of olanzapine tablets until it is right for you.
- If you miss a dose of olanzapine tablets, take the missed dose as soon as you remember. If it is almost time for the next dose, just skip the missed dose and take your next dose at the regular time. Do not take two doses of olanzapine tablets at the same time.
- **To prevent serious side effects, do not stop taking olanzapine tablets suddenly. If you need to stop taking olanzapine tablets, your doctor can tell you how to safely stop taking it.**
- **If you take too much olanzapine tablets, call your doctor or poison control center at 1-800-222-1222 right away, or get emergency treatment.**
- Olanzapine tablets can be taken with or without food.
- Olanzapine tablets are usually taken one time each day
- Call your doctor if you do not think you are getting better or have any concerns about your condition while taking olanzapine tablets, .

### **What should I avoid while taking olanzapine tablets?**

- Olanzapine tablets can cause sleepiness and may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how olanzapine tablets affects you.

- Avoid drinking alcohol while taking olanzapine tablets. Drinking alcohol while you take olanzapine tablets may make you sleepier than if you take olanzapine tablets alone.

### **What are the possible side effects of olanzapine tablets?**

**Serious side effects may happen when you take olanzapine tablets, including:**

- **See “What is the most important information I should know about olanzapine tablets?”**, which describes the increased risk of death in elderly people with dementia-related psychosis and the risks of high blood sugar, high cholesterol and triglyceride levels, and weight gain.
- **Increased incidence of stroke or “mini-strokes” called transient ischemic attacks (TIAs) in elderly people with dementia-related psychosis** (elderly people who have lost touch with reality due to confusion and memory loss). Olanzapine tablets are not approved for these patients.
- **Neuroleptic Malignant Syndrome (NMS):** NMS is a rare but very serious condition that can happen in people who take antipsychotic medicines, including olanzapine tablets. NMS can cause death and must be treated in a hospital. Call your doctor right away if you become severely ill and have any of these symptoms:
  - high fever
  - excessive sweating
  - rigid muscles
  - confusion
  - changes in your breathing, heartbeat, and blood pressure
- **Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS):** DRESS can occur with olanzapine. Features of DRESS may include rash, fever, swollen glands and other internal organ involvement such as liver, kidney, lung and heart. DRESS is sometimes fatal; therefore, tell your doctor immediately if you experience any of these signs.
- **Tardive Dyskinesia:** This condition causes body movements that keep happening and that you can not control. These movements usually affect the face and tongue. Tardive dyskinesia may not go away, even if you stop taking olanzapine tablets. It may also start after you stop taking olanzapine tablets. Tell your doctor if you get any body movements that you can not control.
- **Decreased blood pressure when you change positions, with symptoms of dizziness, fast or slow heartbeat, or fainting.**
- **Difficulty swallowing, that can cause food or liquid to get into your lungs.**
- **Seizures :** Tell your doctor if you have a seizure during treatment with olanzapine tablets.
- **Problems with control of body temperature:** You could become very hot, for instance when you exercise a lot or stay in an area that is very hot. It is important for you to drink water to avoid dehydration. Call your doctor right away if you become severely ill and have any of these symptoms of dehydration:
  - sweating too much or not at all
  - dry mouth
  - feeling very hot
  - feeling thirsty
  - not able to produce urine

**Common side effects of olanzapine tablets include:** lack of energy, dry mouth, increased appetite, sleepiness, tremor (shakes), having hard or infrequent stools, dizziness, changes in behavior, or restlessness.



**Other common side effects in teenagers (13 to 17 years old) include:** headache, stomach-area (abdominal) pain, pain in your arms or legs, or tiredness. Teenagers experienced greater increases in prolactin, liver enzymes, and sleepiness, as compared with adults.

Tell your doctor about any side effect that bothers you or that does not go away.

These are not all the possible side effects with olanzapine tablets. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store olanzapine tablets?**

- Store olanzapine tablets at room temperature, between 68°F to 77°F (20°C to 25°C).
- Keep olanzapine tablets away from light.
- Keep olanzapine tablets dry and away from moisture.

**Keep olanzapine tablets and all medicines out of the reach of children.**

#### **General information about olanzapine tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use olanzapine tablets for a condition for which it was not prescribed. Do not give olanzapine tablets to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about olanzapine tablets. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about olanzapine tablets that was written for healthcare professionals.

For more information about olanzapine tablets call 1-888-375-3784.

#### **What are the ingredients in olanzapine tablets?**

**Active ingredient:** olanzapine

**Inactive ingredients:** Tablets — crospovidone, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose (Avicel PH 101) and microcrystalline cellulose (Avicel PH 102). The color coating contains hypromellose, polyethylene glycol 400 and titanium dioxide.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Rx Only

Manufactured by:

**Dr. Reddy's Laboratories Limited**

Bachupally - 500 090 INDIA

Revised: 03/2025