


## G.1.1.1.4.2 Expert B


**Figure G-2. Characteristics of the Random Draw from the Approximated Distribution of the PM<sub>2.5</sub> Effect Specified by Expert B**  
**(1) Results for the range 4-10 µg/m<sup>3</sup>**

**Notes:**

Expert B specified a non-parametric distribution using five percentile points. We linearly interpolated the CDF between them. Panel (a) shows q-q plot of the expert percentiles and empirical percentiles for the draw. Panel (b) shows empirical CDF associated with the draw, the red "X" marks indicate corresponding expert percentiles. The distribution was conditional on causality. We created a corresponding unconditional distribution by adding extra 2 percent zeros to the draw. Panels (c) and (d) show the respective distributions.

The experts specified distributions for the percent changes in the relative risk. The distribution of the corresponding PM<sub>2.5</sub> effects was the following transformation of the percent change in relative risk  $Z$  -  $\log(1+(Z/100))$ .

**Figure G-2. Characteristics of the Random Draw from the Approximated Distribution of the PM<sub>2.5</sub> Effect Specified by Expert B (continued)**  
**(2) Results for the range >10-30 µg/m<sup>3</sup>**




### Notes:

Expert B specified a non-parametric distribution using five percentile points. We linearly interpolated the CDF between them. Panel (a) shows q-q plot of the expert percentiles and empirical percentiles for the draw. Panel (b) shows empirical CDF associated with the draw, the red "X" marks indicate corresponding expert percentiles. The distribution was conditional on causality. We created a corresponding unconditional distribution by adding extra 2 percent zeros to the draw. Panels (c) and (d) show the respective distributions.

The experts specified distributions for the percent changes in the relative risk. The distribution of the corresponding PM<sub>2.5</sub> effects was the following transformation of the percent change in relative risk  $Z - \log(1+(Z/100))$ .

#### G.1.1.1.4.3 Expert C

**Figure G-3. Histogram of the Random Draw from the Distribution of the PM<sub>2.5</sub> Effect Specified by Expert C**




#### Notes:

Expert C specified a truncated Normal Distribution. We inferred the following values for the parameters of this distribution: mean=1.20 and standard deviation=0.49.

The experts specified distributions for the percent changes in the relative risk. The distribution of the corresponding PM<sub>2.5</sub> effects was the following transformation of the percent change in relative risk  $Z - \log(1+(Z/100))$ .

## G.1.1.1.4.4 Expert D

**Figure G-4. Histogram of the Random Draw from the Distribution of the PM<sub>2.5</sub> Effect Specified by Expert D****Notes:**

Expert D specified a Triangular Distribution with minimum=0.1, maximum=1.6, and mode=0.95. The distribution was conditional on causality. We created a corresponding unconditional distribution by adding extra 5 percent zeros to the draw.

The experts specified distributions for the percent changes in the relative risk. The distribution of the corresponding PM<sub>2.5</sub> effects was the following transformation of the percent change in relative risk  $Z$  -  $\log(1+(Z/100))$ .