



National Comprehensive  
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

# **Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic**

Version 1.2023 — September 7, 2022

**NCCN.org**

**Continue**



**\*Mary B. Daly, MD, PhD/Chair** † ▢  
Fox Chase Cancer Center

**\*Tuya Pal, MD/Vice-Chair** ▢  
Vanderbilt-Ingram Cancer Center

**Zahraa AlHilli, MD** ¶  
Case Comprehensive Cancer Center/  
University Hospitals Seidman Cancer Center  
and Cleveland Clinic Taussig Cancer Institute

**Banu Arun, MD** †  
The University of Texas  
MD Anderson Cancer Center

**Saundra S. Buys, MD** ‡ † †  
Huntsman Cancer Institute  
at the University of Utah

**Heather Cheng, MD, PhD** †  
Fred Hutchinson Cancer Center

**Jane Churpek, MD, MS** ‡  
University of Wisconsin Carbone Cancer Center

**Susan M. Domchek, MD** †  
Abramson Cancer Center  
at the University of Pennsylvania

**Ahmed Elkhanany, MD** †  
O'Neal Comprehensive Cancer Center at UAB

**Susan Friedman, DVM** ¥  
FORCE: Facing Our Risk of Cancer Empowered

**Michael Goggins, MD** ▢  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**Andrea Hagemann, MD** Ω  
Siteman Cancer Center at Barnes-  
Jewish Hospital and Washington  
University School of Medicine

**Ashley Hendrix, MD** ¶  
St. Jude Children's Research Hospital/  
The University of Tennessee  
Health Science Center

**Mollie L. Hutton, MS, CGC** Δ  
Roswell Park Comprehensive Cancer Center

**Beth Y. Karlan, MD** Ω Δ  
UCLA Jonsson Comprehensive Cancer Center

**Seema Khan, MD** ¶  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**Catherine Klein, MD** † †  
University of Colorado Cancer Center

**Wendy Kohlmann, MS, CGC** Δ  
Huntsman Cancer Institute  
at the University of Utah

**Allison W. Kurian, MD, MSc** † † Δ  
Stanford Cancer Institute

**Christine Laronga, MD** ¶  
Moffitt Cancer Center

**Julie S. Mak, MS, MSc, LCGC** Δ  
UCSF Helen Diller Family  
Comprehensive Cancer Center

**John Mansour, MD** ¶  
UT Southwestern Simmons  
Comprehensive Cancer Center

**Kara Maxell, MD, PhD** †  
Abramson Cancer Center at the University of  
Pennsylvania

**Kevin McDonnell, MD, PhD** †  
City of Hope National Medical Center

**Carolyn S. Menendez, MD** ¶ Δ  
Duke Cancer Institute

**Sofia D. Merajver, MD, PhD** ‡ †  
University of Michigan  
Rogel Cancer Center

**Barbara S. Norquist, MD** Ω  
Fred Hutchinson Cancer Center

**Kenneth Offit, MD** † † Δ  
Memorial Sloan Kettering Cancer Center

**Gwen Reiser, MS, CGC** Δ  
Fred & Pamela Buffett Cancer Center

**Leigha Senter-Jamieson, MS, CGC** Δ  
The Ohio State University Comprehensive  
Cancer Center - James Cancer Hospital  
and Solove Research Institute

**Kristen Mahoney Shannon, MS, CGC** Δ  
Massachusetts General Hospital  
Cancer Center

**Rebecca Shatsky, MD** †  
UC San Diego Moores Cancer Center

**Kala Visvanathan, MD, MHS** † †  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**Jeanna Welborn, MD** †  
UC Davis Comprehensive Cancer Center

**Myra J. Wick, MD, PhD** Ω ▢  
Mayo Clinic Cancer Center

**Matthew B. Yurgelun, MD** † †  
Dana-Farber/Brigham and  
Women's Cancer Center

**NCCN**  
**Susan Darlow, PhD**  
**Mary Dwyer, MS**

¶ Breast surgical oncology	‡ Hematology/Hematology oncology
▢ Clinical genetics	‡ Internal medicine
Δ Genetic counseling	† Medical oncology
▢ Gastroenterology	§ Radiation oncology
Ω Gynecologic oncology/Gynecology	¥ Patient advocacy
	* Discussion Writing Committee Member

**Continue**



### [NCCN Genetic/Familial High-Risk Assessment Panel Members](#) [Summary of the Guidelines Updates](#)

#### Principles of Cancer Risk Assessment and Counseling

- [Pre-test Counseling \(EVAL-A 1 of 10\)](#)
- [Testing Considerations Prior to Testing \(EVAL-A 2 of 10\)](#)
- [Choice of Multi-Gene Testing \(EVAL-A 3 of 10\)](#)
- [Evaluating the Source of Genetic Testing Information \(EVAL-A 4 of 10\)](#)
- [Tumor Genomic Testing: Potential Implications for Germline Testing \(EVAL-A 5 of 10\)](#)
  - ▶ [Circulating Tumor DNA \(ctDNA\)](#)
- [Post-test Counseling \(EVAL-A 6 of 10\)](#)
  - ▶ [Positive Results](#)
  - ▶ [Negative Results](#)
  - ▶ [Variants of Uncertain Significance](#)
- [Pedigree: First-, Second-, and Third-Degree Relatives of Proband \(EVAL-B\)](#)

#### Hereditary Testing Criteria

- [General Testing Criteria \(CRIT-1\)](#)
- [Testing Criteria for High-Penetrance Breast Cancer Susceptibility Genes \(CRIT-2\)](#)
- [Testing Criteria for Ovarian Cancer Susceptibility Genes \(CRIT-4\)](#)
- [Testing Criteria for Pancreatic Cancer Susceptibility Genes \(CRIT-5\)](#)
- [Testing Criteria for High-Penetrance Prostate Cancer Susceptibility Genes \(CRIT-6\)](#)
- [Testing Criteria for Li-Fraumeni Syndrome \(CRIT-7\)](#)
- [Testing Criteria for Cowden Syndrome/PTEN Hamartoma Tumor Syndrome \(CRIT-8\)](#)

#### Gene Summary: Risks and Management

- [Testing Criteria Met \(GENE-1\)](#)
- [Cancer Risk Management Based on Genetic Test Results \(GENE-A\)](#)
- [Autosomal Recessive Risk in Cancer Genes – Multi-Gene Panel Testing \(GENE-B\)](#)

#### Management/Screening

- [BRCA Pathogenic/Likely Pathogenic Variant-Positive Management \(BRCA-A\)](#)
- [Pancreatic Cancer Screening \(PANC-A\)](#)
- [Li-Fraumeni Syndrome Management in Adults \(LIFR-A\)](#)
- [Cowden Syndrome/PHTS Management \(COWD-A\)](#)

**Clinical Trials:** NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Find an NCCN Member Institution:  
<https://www.nccn.org/home/member-institutions>.

**NCCN Categories of Evidence and Consensus:** All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

- [Abbreviations \(ABBR-1\)](#)
- [For chemoprevention options, see NCCN Guidelines for Breast Cancer Risk Reduction.](#)
- [Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines \(SUMM-1\)](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2022.



Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:

### General

- For all instances of annual mammogram, "with consideration of tomosynthesis" was removed.
- Genetic Testing Process changed to Gene Summary: Risks and Management

### New section

- Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines ([SUMM-1](#))

### Breast, Ovarian and/or Pancreatic Cancer Genetic Assessment

- Pages added for Positive results ([EVAL-A 7 of 10](#)), Negative results ([EVAL-A 8 of 10](#)), and Variants of uncertain significance (VUS) ([EVAL-A 9 of 10](#))

#### EVAL-A 1 of 10

- Pre-test counseling
  - ▶ Bullet added: Discuss that their results may be important to therapeutic decision making as directed by a qualified health care provider (eg, oncologist).
  - ▶ Footnote b added: Genetic Information Nondiscrimination Act of 2008 (GINA). Vol. Public Law No.110-233. Available at: <https://www.eeoc.gov/laws/statutes/gina.cfm>

#### EVAL-A 2 of 10

- Prior to genetic testing,
  - ▶ 1st bullet revised: ...will vary based on family structure, *which includes size of the family, age of the family members, early death, adoption, and number of male and female relatives...* The estimated likelihood of P/LP variant detection may be ~~very~~ low in families with a large number of unaffected ~~female and/or male relatives or a large number of male relatives~~.
  - ▶ 2nd bullet revised: ...molecular genetic testing via blood, *saliva*, or buccal samples... If this source of DNA is not possible, buccal samples can be considered, subject to the risk of donor DNA contamination *or malignant cells from the hematologic malignancy*.

#### EVAL-A 3 of 10

- Choice of multi-gene testing
  - ▶ 6th bullet revised: "... Not all genes included on available multi-gene tests *will change risk management compared to that based on other risk factors such as family history* ~~are necessarily clinically actionable~~..
  - ▶ 10th bullet revised: Multigene panel testing increases the likelihood of finding P/LP variants *in genes; however, without some genes do not have* clear clinical ~~significance~~ *actionability or have a clear impact on change in* medical management.
  - ▶ 11th bullet revised: When a P/LP variant with clinical implications *for patient and/or their at-risk family members* is found on tumor genomic testing, germline confirmatory testing should be done.
  - ▶ Bullet removed: In many cases the information from testing for moderate penetrance genes does not change risk management compared to that based on family history alone.

#### EVAL-A 4 of 10

- Confirmatory germline testing
  - ▶ 1st bullet revised: Confirmatory germline testing through an appropriately certified laboratory is ~~recommended~~ *clinically indicated* when a potential P/LP variant... (Also for 3rd sub-bullet)
  - ▶ 1st sub-bullet revised: ... and recent research suggests that the error rate (*40%*) is substantial.
  - ▶ 2nd sub-bullet added: Commercial laboratories utilizing consumer-initiated or direct-to-consumer (DTC) marketing of DNA sequence-based cancer predisposition tests vary substantially in providing information necessary to make informed decisions regarding results and may vary in accuracy in their variant interpretation.

[Continued](#)



**Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:**

### Breast, Ovarian and/or Pancreatic Cancer Genetic Assessment

#### EVAL-A 5 of 10

- Tumor Genomic Testing: Potential Implications for Germline Testing

- ▶ 1st bullet, 4th sub-bullet added: Tumor-only sequencing may fail to detect about 10% of clinically actionable P/LP germline variants (eg, deletion, duplication, and splicing variants).
- ▶ 2nd bullet, 4th sub-bullet added: ctDNA, detected by mutation profile, copy number changes, altered methylation patterns, fragmentation, size alterations, or other approaches, has application for disease monitoring as well as early detection. For individuals at increased hereditary risk for cancer, use of pre-symptomatic ctDNA cancer detection assays should only be offered in the setting of prospective clinical trials, because the sensitivity, false-positive rates, and positive predictive value of ctDNA tests for early-stage disease, which are needed to derive clinical utility and determine clinical validity, are not fully defined. The psychological impact of ctDNA testing remains unknown. (Corresponding references added)

#### EVAL-A 6 of 10

- Post-test counseling section extensively revised.

### Hereditary Cancer Testing Criteria

#### CRIT-1

- General testing criteria, Testing may be considered criteria

- ▶ 2nd bullet added: Personal history of serous endometrial cancer with corresponding footnote d: This is a rare subtype of uterine cancer for which there is evolving evidence of an association with BRCA1 P/LP variants.
- ▶ Statement revised: ~~The following~~ For a list of NCCN Guidelines include content focused on inherited cancer conditions, including general principles of testing and/or criteria for testing and/or cancer risk management based on a genetic test result: *Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines (SUMM-1)*.
- ▶ Footnote b revised: ...platinum therapy for prostate cancer and pancreatic cancer; *and risk-reducing surgery*

#### CRIT-2

- Testing Criteria for High-Penetrance Breast Cancer Susceptibility Genes

- ▶ Testing is clinically indicated, Personal history of breast cancer criteria was reorganized and revised by:
  - ◊ ≤50 y
  - ◊ Any age:
    - Treatment indications
    - Pathology/histology
    - Male breast cancer
    - Ancestry: Ashkenazi Jewish ancestry
    - Family history
- ▶ New criteria added under Pathology/histology: Multiple primary breast cancers (synchronous or metachronous)
- ▶ Family history, ≥1 close blood relative with ANY, prostate cancer removed: intraductal/cirbriform histology

#### CRIT-2A

- Footnote k added: Weitzel JN, et al. Breast Cancer Res Treat 2021;188:759-768.
- Footnote l revised by moving unknown or limited family structure criteria to the footnote: *Unknown or limited family structure* (Weitzel JN, et al. JAMA 2007;297:2587-2595).

#### CRIT-3

- Testing may be considered...2nd bullet added: Personal history of breast cancer diagnosed at any age with ≥ 1 close blood relative with intermediate-risk prostate cancer with intraductal/cirbriform histology



Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:

### CRIT-6

- Testing Criteria For High-Penetrance Prostate Cancer Susceptibility Genes
  - ▶ Personal history of prostate cancer with specific features:
    - ◊ By tumor characteristic, Histology
      - removed intraductal/cirriiform histology
    - ◊ By family history and ancestry,
      - two sub-bullets were added: triple-negative breast cancer at any age and male breast cancer at any age
      - 6th sub-bullet revised by removing intraductal/cirriiform histology
  - ▶ Testing may be considered in the following scenario added: Personal history of prostate cancer with intermediate-risk prostate cancer with intraductal/cirriiform histology (see Initial Risk Stratification and Staging Workup in NCCN Guidelines for Prostate Cancer) at any age

### CRIT-7

- Testing Criteria for LFS
  - ▶ 4th bullet added: Pediatric hypodiploid acute lymphoblastic leukemia
- Statement added to page: Other cancers associated with LFS but not in the testing criteria include: melanoma, colorectal, gastric, and prostate.

### CRIT-8A

- Statement added to page under major and minor criteria: Other cancers associated with PTEN but not in the testing criteria include: colorectal, kidney cancer, and melanoma.

### Cancer Risk Management Based on Genetic Test Results

#### GENE-A all pages

- Breast Cancer Risk, "First primary" added.
- Revised to "Strength of evidence of association with cancer:"
- NBN gene removed from table and added to footnote a: The following genes and others are found on some of the panels, but there is insufficient evidence to make any recommendations for breast MRI, RRSO, or RRM for: ...NBN or...
- References were updated accordingly.

#### GENE-A 1 of 10

- ATM
  - ▶ Breast cancer,
    - ◊ Absolute risk revised from "15%–40%" to "20%–40%"
    - ◊ Management revised: Annual mammogram at age 40 y and consider breast MRI with contrast starting at age 40 30–35 y...
  - ▶ Ovarian cancer, Absolute risk revised from "<3%" to "2%–3%"
  - ▶ Prostate cancer revised from "Unknown or insufficient evidence" to "Emerging evidence for association with increased risk"
  - ▶ Comments revised: "Counsel for risk of autosomal recessive condition in offspring" to "See GENE-B for reproductive implications recessive disease." Also for BRCA1, BRCA2, BRIP1, PALB2, RAD51C.
- BARD1,
  - ▶ Breast cancer, Absolute risk revised from "15%–40%" to "20%–40%"
  - ▶ Breast cancer, Strength of evidence for association with cancer revised: Strong-for triple-negative disease
  - ▶ Ovarian cancer, Evidence of increased risk changed from "None" to "No established association" Also for CDH1, CDKN2A, CHEK2, NF1, PTEN, STK11, TP53.





**Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:**  
**Cancer Risk Management Based on Genetic Test Results**

### [GENE-A 2 of 10](#)

- BRCA1
  - ▶ Breast cancer, Male breast cancer added: Absolute risk: 0.2%–1.2% by age 70 y and Strength of evidence of association with cancer: Strong
  - ▶ Prostate cancer added: Absolute risk: 7%–26% and Management: See BRCA Pathogenic Variant-Positive Management
- BRCA2
  - ▶ Breast cancer, Male breast cancer added: 1.8%–7.1% by age 70 y and Strength of evidence of association with cancer: Strong
  - ▶ Prostate cancer added: Absolute risk: 19%–61% and Management: See BRCA Pathogenic Variant-Positive Management

### [GENE-A 3 of 10](#)

- BRIP1,
  - ▶ Breast cancer, Strength of evidence of association with cancer revised: Limited; potential increase in female breast cancer (~~including triple negative~~)
  - ▶ Ovarian cancer,
    - ◊ Absolute risk revised from ">10%" to "5%–15%"
    - ◊ Management revised: Risk reduction: ~~Consider~~ **Recommend** RRSO at 45–50 y
  - ▶ Comments revised: Based on estimates from available studies, the lifetime risk of ovarian cancer in carriers of P/LP variants in BRIP1 appears to be sufficient to justify ~~consideration of~~ RRSO.
- CDH1,
  - ▶ Breast, Strength of evidence of association with cancer revised: Strong (~~with predisposition to lobular disease~~)
  - ▶ Hereditary diffuse gastric cancer (HDGC) added: Strength of evidence of association with cancer: Strong
  - ▶ Comments revised: ...However, one small study found that >50% of such individuals had gastric cancer identified at the time of risk-reducing total gastrectomy (Jacobs MF, et al. Gastroenterology 2019;157:87-96), *and penetrance for lifetime risk is increased with a positive family history of HDGC (Roberts ME, et al. JAMA Oncol 2019;5:1325-1331).*
- CDKN2A,
  - ▶ Comments revised from "General melanoma risk management is appropriate, such as annual full-body skin examination and minimizing UV exposure" to "Comprehensive skin examination by a dermatologist, supplemented with total body photography and dermoscopy is recommended biannually (Chan SH, et al. Hered Cancer Clin Pract 2021;19:21)."

### [GENE-A 4 of 10](#)

- CHEK2
  - ▶ Breast cancer,
    - ◊ Absolute risk revised from "15%–40%" to "20%–40%"
    - ◊ Management revised: Annual mammogram at age 40 y and consider breast MRI with contrast starting at age ~~40~~ 30–35 y...
    - ◊ Strength of evidence of association with cancer: Strong (~~with predisposition to ER+ disease~~)
  - ▶ Comments revised: The risks for most missense variants are unclear but for some P/LP variants, such as Ile157Thr, the risk for breast cancer appears to be lower *and does not reach the threshold for management change.*
- MSH2, MLH1, MSH6, PMS2, EPCAM
  - ▶ Ovarian cancer revised:
    - ◊ MLH1, Absolute risk revised from ">10%" to "4%–20%"
    - ◊ MSH2 /EPCAM, Absolute risk revised from: >10%" to "8%–38%"
    - ◊ MSH6, Absolute risk revised from "≤13%" to "≤1%–13%" and Strength of evidence: ~~Mixed~~ **Strong**
    - ◊ PMS2, Absolute risk revised from "<3%" to "1.3%–3%"
    - ◊ EPCAM combined with MSH2
  - ▶ Comments revised from "Counsel for risk of autosomal recessive condition in offspring" to "Counsel for biallelic risk of P/LP variants that lead to CMMRD. See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal.



Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:

### Cancer Risk Management Based on Genetic Test Results

#### [GENE-A 5 of 10](#)

- NF1
  - ▶ Breast cancer, Absolute risk revised from "15%–40%" to "20%–40%"
  - ▶ Comments revised by removing: Screening recommendations only apply to individuals with a clinical diagnosis of NF.
- PALB2,
  - ▶ Breast cancer, Strength of evidence of association with cancer revised: Strong ~~(with overrepresentation of triple-negative disease)~~
  - ▶ Ovarian cancer, Management revised from "Evidence insufficient; manage based on family history" to "Consider RRSO at age >45 y"
- PTEN,
  - ▶ Breast cancer, Strength of evidence of association with cancer revised: Strong ~~(with predisposition to luminal subtype)~~

#### [GENE-A 6 of 10](#)

- RAD51C and RAD51D:
  - ▶ Breast cancer,
    - ◊ Absolute risk revised from "15%–40%" to "20%–40%"
    - ◊ Management revised from "Insufficient data; managed based on family history" to "Annual mammogram and consider breast MRI with contrast starting at age 40 y"
    - ◊ Strength of evidence of association with cancer revised: Strong ~~for ER/PR-negative breast cancer~~
- RAD51C
  - ▶ Ovarian cancer,
    - ◊ Absolute risk revised from ">10%" to "10%–15%"
    - ◊ Management revised: Risk reduction: ~~Consider~~ *Recommend* RRSO at 45–50 y
- RAD51D:
  - ▶ Ovarian cancer
    - ◊ Absolute risk revised from ">10%" to "10%–20%"
    - ◊ Management revised: Risk reduction: ~~Consider~~ *Recommend* RRSO at 45–50 y

#### [GENE-A 7 of 10](#)

- STK11,
  - ▶ Breast cancer, Absolute risk revised from 40%–60% to 32%–54%
  - ▶ Management revised by adding: Annual mammogram and breast MRI with contrast starting at age 30 y and Risk reduction revised ~~Evidence insufficient manage based on family history~~ *Discuss option of RRM*
- TP53,
  - ▶ Breast cancer, Strength of evidence of association with cancer revised: Strong ~~(with predisposition to triple-positive disease)~~





Updates in Version 1.2023 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic from Version 2.2022 include:

### [GENE-A 8 of 10](#)

- Footnotes added:

- ▶ Footnote e: The use of MRI in these patients depends on a number of risk factors, including family history, age, breast density, and patient preference.
  - ▶ Footnote g: The higher range of risk is reflective of a prospective study of pancreatic cancer kindreds (Hsu FC, Roberts NJ, Childs E, et al. Risk of pancreatic cancer among individuals with pathogenic variants in the ATM gene. JAMA Oncol 2021;7:1664-1668).
  - ▶ Footnote h: Risks and benefits of premature surgical menopause versus risk of cancer and family history should all be carefully considered, and the panel recommends patients seek expert care.
- Footnote removed: Recent data have demonstrated no significant association between pathogenic/likely pathogenic germline PMS2 variants and risks of Lynch syndrome cancers beyond colorectal and endometrial cancer (Ten Broeke S, et al. J Clin Oncol 2018;36:2961-2968). This study did not specifically evaluate pancreatic cancer in PMS2 carriers, but we should note that it is currently unclear if individuals with germline PMS2 variants have increased risk of pancreatic cancer, even though it is a Lynch syndrome gene. There are no data to quantify the strength of association between EPCAM and pancreatic cancer, but EPCAM is generally thought to have the same cancer risks/penetrance as MSH2, given that pathogenic/likely pathogenic germline alterations in EPCAM induce constitutional silencing of MSH2

### [GENE-B](#)

- MLH1, MSH2, MSH6, PMS2, EPCAM – CMMRD revised from "CMMRD is a childhood cancer predisposition syndrome characterized by four main tumor types (hematologic malignancies, brain/central nervous system tumors, colorectal tumors and multiple intestinal polyps, and other malignancies including embryonic tumors and rhabdomyosarcoma) to "CMMRD is a childhood cancer predisposition syndrome characterized by hematologic malignancies, brain/central nervous system tumors, colorectal tumors and multiple intestinal polyps, and other malignancies including embryonic tumors and rhabdomyosarcoma."
- MSH3 and NBN removed.

### [BRCA Pathogenic/Likely Pathogenic Variant-Positive Management](#)

#### [BRCA-A 2 of 3](#)

- Ovarian and Uterine Cancer

- ▶ 3rd revised by adding: HRT recommendations should be tailored depending on each patient's personal history of breast cancer and/or breast cancer risk reduction strategies. HRT is a consideration for premenopausal patients who do not carry a diagnosis of breast cancer or have other contraindications for HRT.
- ▶ 4th bullet revising by adding: Consider preoperative menopause management consultation if patient is still premenopausal at time of RRSO.
- ▶ Bullet removed: For those patients who have not elected RRSO, transvaginal ultrasound combined with serum CA-125 for ovarian cancer screening, although of uncertain benefit, may be considered at the clinician's discretion starting at age 30–35 y.

### [Li-Fraumeni Syndrome Management in Adults](#)

#### [LIFR-A 2 of 2](#)

- Other aspects of managing LFS, 1st bullet revised: "This screening and management of LFS is complex, *and LFS is rare...*"

### [Cowden Syndrome Management](#)

#### [COWD-A 1 of 2](#)

- General

- ▶ 1st bullet added: Due to the rarity of the syndrome and complexities of diagnosing and managing individuals with Cowden syndrome, referral to a specialized team or centers with expertise is recommended.

#### [COWD-A 2 of 2](#)

- Endometrial cancer, 2nd bullet revised by adding: Risk of ovarian cancer is not elevated; therefore, ovaries can be left in situ.

### [ABBR-1](#)

- New section added: Abbreviations.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

- The decision to offer genetic testing involves three related stages:
  - 1) pre-test counseling done prior to ordering testing;
  - 2) consideration of the most appropriate tests to order; and
  - 3) post-test counseling done when results are disclosed.<sup>1-6</sup>
- It is recommended that a genetic counselor, clinical geneticist, oncologist, surgeon, oncology nurse, or other health professional with expertise and experience in cancer genetics be involved at each stage whenever possible.
- Testing should be considered in appropriate high-risk individuals where it is likely to impact the risk management and/or treatment of the tested individuals and/or their at-risk family members.

#### Pre-test counseling includes the following elements:

- Evaluate patient's needs and concerns regarding:
  - Knowledge of genetic testing for cancer risk, including benefits, risks, and limitations
  - Goals for cancer family risk assessment
- Detailed family history including:
  - Collection of a comprehensive family history
    - ◊ Assessment of family history; close blood relatives include first-, second-, and third-degree relatives on each side of the family, particularly around individuals with a diagnosis of cancer ([See EVAL-B](#))
    - ◊ Types of cancer, bilaterality, age at diagnosis, subtype, and pathology report confirmation
    - ◊ Ethnicity (specifically Ashkenazi Jewish ancestry)
- Detailed medical and surgical history including:
  - Documentation of prior genetic testing results for patients and their family members
  - Personal cancer history (eg, age, histology, laterality)
  - Pathology reports of primary cancers and/or benign lesions (eg, breast biopsies)
  - Carcinogen exposure (eg, history of radiation therapy)
  - Reproductive history
  - Hormone or oral contraceptive use
  - History of risk-reducing surgeries
- Focused physical exam (conducted by qualified clinician) when indicated:
  - CS/PHTS specific: dermatologic,<sup>a</sup> including oral mucosa, head circumference, and thyroid (enlarged or nodular on palpation)
- Generate a differential diagnosis and educate the patient on inheritance patterns, penetrance, variable expressivity, and the possibility of genetic heterogeneity
- Prepare for the possible outcomes of testing, including positive (pathogenic, likely pathogenic [P/LP]), true negative and uninformative negative, uncertain variants, and mosaic results
- Obtain written informed consent, and document the informed consent in the patient's medical record
- Discuss plan for results disclosure when appropriate, including the possibility of the patient consenting to Release of Information of test results to a close relative or spouse when results are released in case patient is deceased or incapacitated
- Discuss possible management options if a P/LP variant is identified (enhanced surveillance, risk-reducing agents, and risk-reducing surgery)
- Discuss that their results may be important to therapeutic decision making as directed by a qualified health care provider (eg, oncologist).
- Advise about possible inherited cancer risk to relatives, options for risk assessment, testing, and management
- Discuss cost of genetic testing
- Provide overview of current legislation regarding genetic discrimination and the privacy of genetic information<sup>b</sup>

<sup>a</sup> For Cowden syndrome/PTEN hamartoma tumor syndrome (CS/PHTS) dermatologic manifestations, [see CRIT-8](#) and for Peutz-Jeghers syndrome (PJS) dermatologic manifestations, [see NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#).

<sup>b</sup> Genetic Information Nondiscrimination Act of 2008 (GINA). Vol. Public Law No.110-233. Available at: <https://www.eeoc.gov/laws/statutes/gina.cfm>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References on  
EVAL-A 10 of 10](#)

[Continued](#)

**EVAL-A  
1 OF 10**



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

**Prior to genetic testing, the following should be taken into consideration:**

- The probability of P/LP variant detection associated with these criteria will vary based on family structure, which includes size of the family, age of the family members, early death, adoption, and number of male and female relatives. Individuals with unknown or limited family history/structure, such as fewer than 2 female first- or second-degree relatives having lived beyond age 45 in either lineage, may have an underestimated probability of familial P/LP variant detection. The estimated likelihood of P/LP variant detection may be low in families with a large number of unaffected and/or male relatives.
- Patients who have received an allogeneic bone marrow transplant or with active or recent hematologic malignancies should not have molecular genetic testing via blood, saliva, or buccal samples (due to unreliable test results from contamination by donor DNA) until other technologies are available. If available, DNA should be extracted from a fibroblast culture. If this source of DNA is not possible, buccal samples can be considered, subject to the risk of donor DNA contamination or malignant cells from the hematologic malignancy.
- If more than one family member is affected with cancers highly associated with a particular inherited cancer susceptibility syndrome, consider initial testing of a family member with youngest age at diagnosis, bilateral disease, multiple primary cancers, or other cancers associated with the syndrome, or most closely related to the proband/patient. If there are no available family members with cancer that is a cardinal feature of the syndrome in question, consider testing first- or second-degree family members affected with other cancers thought to be related to the gene in question (eg, prostate or pancreas with *BRCA1/2*).
- Testing for unaffected family members when no affected member is available should be considered. Significant limitations of interpreting test results should be discussed.
- In children <18 y, genetic testing is generally not recommended when results would not impact medical management.<sup>7</sup>
- LP variants are usually clinically managed similarly to pathogenic variants, while patients with variants of uncertain significance (VUS) and likely benign variants should be managed based on the cancers present in the family.
- Choice of multi-gene testing, see [EVAL-A 3 of 10](#).

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. On this page, the terms male and female refer to sex assigned at birth.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References on](#)  
[EVAL-A 10 of 10](#)

[Continued](#)

**EVAL-A**  
**2 OF 10**



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

##### Choice of multi-gene testing

- The introduction of multi-gene testing for hereditary forms of cancer has rapidly altered the clinical approach to hereditary cancer testing of at-risk patients and their families. Based on next-generation sequencing (NGS) technology, these tests simultaneously analyze a set of genes that are associated with a specific family cancer phenotype or multiple phenotypes.
- An individual's personal and/or family history may be explained by more than one inherited cancer syndrome; thus, phenotype-directed testing based on personal and family history through a tailored<sup>c</sup> multi-gene panel test is often more efficient and cost-effective and increases the yield of detecting a P/LP variant in a gene that will impact medical management for the individual or their at-risk family members.
- There may also be a role for multi-gene testing in individuals who have tested negative for a single syndrome, but whose personal or family history remains suggestive of an inherited susceptibility.
- Some individuals may carry P/LP germline variants in more than one cancer susceptibility gene; thus, consideration of a multi-gene panel for individuals already known to carry a single P/LP germline variant from phenotype-directed testing may be considered on a case-by-case basis, based on the degree of suspicion for there being additional variants.
- Because commercially available tests differ in the specific genes analyzed, variant classification, and other factors (eg, methods of DNA/RNA analysis or option to reflex from a narrow to a larger panel; provision of financial assistance for cascade testing of relatives), it is important to consider the indication for testing and expertise of the laboratory when choosing the specific laboratory and test panel.
- Multi-gene testing can include “intermediate” penetrant (moderate-risk) genes.<sup>d</sup> For many of these genes, there are limited data on the degree of cancer risk, and there may currently be no clear guidelines on risk management for carriers of P/LP variants. Not all genes included on available multi-gene tests will change risk management compared to that based on other risk factors such as family history.
- It may be possible to refine risks associated with both moderate and high-penetrance genes, taking into account the influence of gene/gene or gene/environment interactions. In addition, certain P/LP variants in a gene may pose higher or lower risk than other P/LP variants in that same gene. This information should be taken into consideration when assigning risks and management recommendations for individuals and their at-risk relatives.
- P/LP variants in many breast, ovarian, pancreatic, and prostate cancer susceptibility genes involved in DNA repair may be associated with rare autosomal recessive conditions, thus posing risks to offspring if the partner is also a carrier.
- As more genes are tested, there is an increased likelihood of finding VUS, mosaicism, and clonal hematopoiesis of indeterminate potential (CHIP).
- Multigene panel testing increases the likelihood of finding P/LP variants in genes; however, some genes do not have clear clinical actionability or have a clear impact on change in medical management.
- When a P/LP variant with clinical implications for patient and/or their at-risk family members is found on tumor genomic testing, germline confirmatory testing should be done.
- There are significant limitations in interpretation of polygenic risk scores (PRS). PRS should not be used for clinical management at this time and use is recommended in the context of a clinical trial, ideally including diverse populations. [See Discussion](#).

<sup>c</sup> Tailored is defined as a disease-focused multi-gene panel of clinically actionable cancer susceptibility genes, in contrast to large multi-gene panels of uncertain or unknown clinical relevance.

<sup>d</sup> Research is evolving, and individuals with P/LP variants in cancer susceptibility genes should be encouraged to participate in clinical trials or genetic registries. Individuals with P/LP variants are also encouraged to recontact their genetics providers every few years for updates.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References on  
EVAL-A 10 of 10](#)

[Continued](#)

**EVAL-A  
3 OF 10**



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

##### Evaluating the Source of Genetic Testing Information

- Prior to using any germline findings for medical management, it is important to establish whether the reported findings were obtained from a laboratory that is certified by the College of American Pathologists (CAP) and Clinical Laboratory Improvement Amendments (CLIA) to issue a report of germline findings directly to ordering health care providers. Some states (eg, New York) may have additional reporting requirements.
- Confirmatory germline testing through an appropriately certified laboratory is clinically indicated when a potential P/LP variant is identified through various data sources as noted below:
  - ▶ Commercial entities providing ancestry (and sometimes health) information typically do so through microarray-based single nucleotide polymorphism (SNP) testing that has not been validated for clinical use. Third-party software applications can be used by consumers to obtain an interpretation of the raw data provided by these companies. Raw data and third-party software are not able to provide information that is appropriate for medical management, as these services are not subject to quality-control processes and recent research suggests that the error rate (40%) is substantial.<sup>8</sup> In addition, the current tests only provide limited founder pathogenic variants results without the benefit of family history. More comprehensive genetic counseling and testing for pathogenic variants in other inherited cancer risk genes may be appropriate at the time of confirmation testing.
  - ▶ Commercial laboratories utilizing consumer-initiated or direct-to-consumer (DTC) marketing of DNA sequence-based cancer predisposition tests vary substantially in providing information necessary to make informed decisions regarding results and may vary in accuracy in their variant interpretation.<sup>9,10</sup>
  - ▶ Research: Patients may have participated in research studies that included germline genomic analysis.<sup>11</sup> In such cases, it is clinically indicated to review the patient's findings with a genetics professional and/or the reporting laboratory to establish whether the original report was generated by an appropriately certified laboratory, or whether confirmatory testing is clinically indicated.

[References on  
EVAL-A 10 of 10](#)  
[Continued](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

##### Tumor Genomic Testing: Potential Implications for Germline Testing

- Testing may provide information suggesting a potential germline finding. P/LP variants reported in the tumor may be of somatic or germline origin.
  - ▶ Because tumor genomic testing is designed to address treatment actionability, not germline status, a variant that may be considered as P/LP in the germline may not be reported at all, or reported as normal in the tumor if it lacks clinical implications.
  - ▶ The filtering of raw sequencing data may differ between tumor and germline testing labs so that variants reported out with one analysis may not be reported with the other.
  - ▶ Somatic P/LP variants seen in tumor specimens are common in some genes with germline implications (eg, *TP53*, *STK11*, *PTEN*) and may not indicate the need for germline testing unless the clinical/family history is consistent with a P/LP variant in the germline.
  - ▶ Tumor-only sequencing may fail to detect about 10% of clinically actionable P/LP germline variants (eg, deletion, duplication, and splicing variants).<sup>12</sup>
- Regardless of findings in the tumor, when germline testing is clinically indicated, it should be performed in a CLIA-approved lab with established experience in germline testing because:
  - ▶ The germline panel performed by some labs offering paired tumor and germline testing may have incomplete coverage and analyze only a subset of those genes of interest to the clinician.
  - ▶ The sensitivity of most tumor genomic testing is lower (particularly for intermediate-sized deletions and duplications) than germline testing.
  - ▶ Similarly, circulating tumor DNA (ctDNA) has the potential to identify both somatic and germline variants with germline treatment implications. Some ctDNA assays, but not all, will alert providers that the particular gene variant identified has a high enough variant allele frequency (VAF) that it is suspicious for germline origin. However, most commercially available assays specializing in somatic ctDNA detection are neither intended nor validated for the reporting or interpretation of germline variants. Thus, variants detected by ctDNA that are suspected to be present in the germline should be evaluated via a CLIA-approved assay specializing in detection and interpretation of germline variants.
  - ▶ ctDNA, detected by mutation profile, copy number changes, altered methylation patterns, fragmentation, size alterations, or other approaches, has application for disease monitoring as well as early detection. For individuals at increased hereditary risk for cancer, use of pre-symptomatic ctDNA cancer detection assays should only be offered in the setting of prospective clinical trials, because the sensitivity, false-positive rates, and positive predictive value of ctDNA tests for early-stage disease, which are needed to derive clinical utility and determine clinical validity, are not fully defined.<sup>13-16</sup> The psychological impact of ctDNA testing remains unknown.

[References on  
EVAL-A 10 of 10](#)

[Continued](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

##### Post-Test Counseling

- When the testing provider/facility does not include pre-test counseling or have all of the resources or expertise for facilitating follow-up testing, management, or family testing, referral to a genetics provider is recommended. In particular, referral to a genetics provider is recommended for the following test results:
  - ▶ P/LP variant identified
  - ▶ Negative results but tumor profiling, personal history, or family history remain suggestive of inherited condition
  - ▶ Any VUS result for which a provider considers using to guide management
  - ▶ A mosaic or possibly mosaic result
  - ▶ Discrepant interpretation of variants, including discordant results across laboratories
  - ▶ Interpretation of PRS, if they are being considered for use in clinical management
  - ▶ Interpretation of P/LP variants for patients tested through direct-to-consumer or consumer-initiated models
- Post-test counseling includes the following elements:
  - ▶ Discussion of results and associated medical risks
  - ▶ Interpretation of results in context of personal and family history of cancer
  - ▶ Discussion of recommended medical management options including discussion of therapeutic implications by a qualified health care provider if positive.
  - ▶ Discussion of the importance of notifying family members and offering materials/resources for informing and testing at-risk family members
  - ▶ Discussion of available resources such as high-risk clinics, disease-specific support groups, and research studies

[References on  
EVAL-A 10 of 10](#)

[Continued](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

- **Positive results:**

- ▶ Some medical centers include services that are specialized in cancer screening, risk reduction, and treatment for individuals with a P/LP variant associated with increased risk for cancer. Where available, consider referring patients to these services, either on a consultative basis or for coordination of ongoing care.
- ▶ In patients being treated for cancer, identification of a P/LP variant may affect options and recommendations for treatment of their disease. A P/LP variant in certain genes is also a component of eligibility for some clinical trials. Specific circumstances are addressed in the NCCN Treatment Guidelines for breast, ovarian, and other cancers.
- ▶ Many patients who have been diagnosed with cancer and have a P/LP variant are at increased risk for additional primary cancers in the future. Management of those risks may be appropriate after treatment of the current cancer or may be combined with treatment for a current cancer.
- ▶ Multiple sources, including these NCCN Guidelines, provide estimated lifetime risks of cancer associated with specific P/LP variants. An individual's personal risk may differ from published lifetime risks, depending on individual medical history and, importantly, based on patient age. Specifically, patients who are older will have lower remaining lifetime risks.
- ▶ Individuals with a P/LP variant should be informed of the importance of this information for their biological relatives. Knowledge of the P/LP variant may affect risk assessment and recommendations for genetic testing, early detection, and/or cancer risk reduction in those relatives. Where relationships allow, individuals should be encouraged to communicate this information to their biological relatives. A medical provider can assist by providing patients with information for relatives written in simple language and a copy of their genetic test results.
- ▶ Over time, patients with a P/LP variant benefit from re-consultation with a medical provider who is familiar with inherited risk for cancer. This re-consultation is important for:
  - ◊ Increasing compliance with screening guidelines, which is known to decrease over time
  - ◊ Re-evaluating personal choices about risk-reducing surgeries, based on changing life stage and circumstances
  - ◊ Ensuring patients are following up-to-date guidelines
  - ◊ Discussing additional genetic testing options
  - ◊ Reviewing improved risk models as appropriate
- ▶ The frequency of follow-up depends on many factors, such as age, reproductive planning, comorbidities, risk-reducing surgeries, and other risk factors.
- ▶ For patients of reproductive age, advise about options for prenatal diagnosis and assisted reproduction, including pre-implantation genetic testing. Discussion should include known risks, limitations, and benefits of these technologies. See [Discussion](#) for details.
- ▶ Biallelic P/LP variants in some genes, included on gene panels, may be associated with rare autosomal recessive conditions, such as Fanconi anemia or constitutional mismatch repair (MMR) deficiency (CMMRD) ([GENE-B](#)). Thus, for these genes, consideration should be given to carrier testing the partner for P/LP variants in the same gene if it would inform reproductive decision-making and/or risk assessment and management.<sup>17</sup>
- ▶ Some P/LP variants found in blood, saliva, or buccal samples, most notably in *TP53*, warrant consideration of testing of non-blood sample to try to distinguish between germline, constitutional mosaicism, and somatic findings.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References on](#)  
[EVAL-A 10 of 10](#)  
[Continued](#)

EVAL-A  
7 OF 10



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

- **Negative results:**

- ▶ These results reduce concern for cancer risk. However, the individual may still have increased cancer risk based on family history. Also, other family members may have a P/LP variant that the tested individual did not inherit.
- ▶ Although negative results of genetic testing are generally reassuring, other reasons that a patient can test negative include:
  - 1) A gene P/LP variant may exist in the gene that was not recognized due to limitations in technology
  - 2) P/LP variants exist in genes that were not evaluated by this testing
  - 3) Family members may harbor a genetic P/LP variant that the patient may not have inherited
- ▶ Other family members may be appropriate candidates for testing, both to assess their own cancer risk as well as to clarify the overall contribution of known P/LP variants to the family history. If another family member tests positive for a P/LP variant, this might lower concern for the individuals who tested negative. The determination of a “true negative” result depends on the specific family history of cancer, the specific P/LP variant found, and the relationship to the family member(s) who tested positive.
- ▶ When an individual has tested negative, it may still be appropriate to consider increased screening and risk reduction measures for cancer based on family history. See appropriate screening based on family history in the guidelines as outlined in [Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines \(SUMM-1\)](#). Some medical centers include specialized high-risk clinics to offer this type of family history-based screening.
- ▶ Over time an individual who tested negative may be a candidate for additional genetic testing due to additional family history or as new genes are identified to be associated with cancer risk.

[References on  
EVAL-A 10 of 10](#)

[Continued](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING

- Variants of uncertain significance (VUS)
  - ▶ VUS are alterations in the genetic code for which the impact on protein function is uncertain.
  - ▶ VUS are common, particularly with the use of large multi-gene panels. The more genes that are included on a genetic testing panel, the more likely a VUS will be identified.<sup>18</sup>
  - ▶ VUS are more commonly found during genetic testing of racial and ethnic minorities compared with non-Hispanic Whites.<sup>18</sup>
  - ▶ In VUS that are reclassified, approximately 80%–90% are reclassified to likely benign or benign and 10%–20% to P/LP.<sup>19,20</sup>
  - ▶ Variant interpretation can vary between commercial laboratories.<sup>21</sup> Resources are available to review the available data supporting pathogenic consequences of specific variants and identify discrepant results (eg, <https://www.ncbi.nlm.nih.gov/clinvar/>; <https://brcaexchange.org/about/app>; [cangene-canvaruk.org/canvig-uk](https://cangene-canvaruk.org/canvig-uk)).
  - ▶ VUS should not be used to alter medical management. Screening and risk reduction strategies should be recommended on the basis of personal and family history.
  - ▶ Testing family members for a VUS should not be done for clinical purposes, unless there are data to support discrepancy in interpretation of results. Consider a referral to research studies that aim to define the functional impact of variants such as variant reclassification programs through clinical labs or registries.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References on](#)  
[EVAL-A 10 of 10](#)

**EVAL-A**  
**9 OF 10**



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### PRINCIPLES OF CANCER RISK ASSESSMENT AND COUNSELING


#### References

- 1 Robson ME, Bradbury AR, Arun B, et al. American Society of Clinical Oncology Policy Statement Update: Genetic and Genomic Testing for Cancer Susceptibility. *J Clin Oncol* 2015;33:3660-3667.
- 2 Berliner JL, Fay AM, Cummings SA, Burnett B, Tillmanns T. NSGC practice guideline: risk assessment and genetic counseling for hereditary breast and ovarian cancer. *J Genet Couns* 2013;22:155-163.
- 3 American College of Obstetricians and Gynecologists; ACOG Committee on Practice Bulletins--Gynecology; ACOG Committee on Genetics; Society of Gynecologic Oncologists. ACOG Practice Bulletin No. 103: Hereditary breast and ovarian cancer syndrome. *Obstet Gynecol* 2009;113:957-966.
- 4 Lancaster JM, Powell CB, Chen LM, Richardson DL; SGO Clinical Practice Committee. Society of Gynecologic Oncology statement on risk assessment for inherited gynecologic cancer predispositions. *Gynecol Oncol* 2015;136:3-7.
- 5 Konstantinopoulos PA, Norquist B, Laccetti C, et al. Germline and somatic tumor testing in epithelial ovarian cancer: ASCO Guideline. *J Clin Oncol* 2020;38:1222-1245.
- 6 Weitzel JN, Blazer KR, Macdonald DJ, Culver JO, Offit K. Genetics, genomics, and cancer risk assessment: State of the art and future directions in the era of personalized medicine. *CA Cancer J Clin* 2011;61:327-359.
- 7 Committee on Bioethics; Committee on Genetics, and American College of Medical Genetics and; Genomic Social; Ethical; Legal Issues Committee. Ethical and policy issues in genetic testing and screening of children. *Pediatrics* 2013;131:620-622.
- 8 Tandy-Connor S, Gultinan J, Krempely K, et al. False-positive results released by direct-to-consumer genetic tests highlight the importance of clinical confirmation testing for appropriate patient care. *Genet Med* 2018;20:1515-1521.
- 9 Kilbride MK, Bradbury AR. Evaluating web-based direct-to-consumer genetic tests for cancer susceptibility. *JCO Precis Oncol* 2020 Mar 5;4:PO.19.00317.
- 10 Direct-to-Consumer Tests: <https://www.fda.gov/medical-devices/in-vitro-diagnostics/direct-consumer-tests>.
- 11 Green R, Berg J, Grody W, et al. ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. *Genet Med* 2013;15:565-574.
- 12 Terraf P, Pareja F, Brown DN, et al. Comprehensive assessment of germline pathogenic variant detection in tumor-only sequencing. *Ann Oncol* 2022;33:426-433.
- 13 Duffy MJ, Diamandis EP, Crown J. Circulating tumor DNA (ctDNA) as a pan-cancer screening test: is it finally on the horizon? *Clin Chem Lab Med* 2021;59:1353-1361.
- 14 Offit K, Sharkey C, Green D, et al. Regulation of laboratory-developed tests in preventive oncology: Emerging needs and opportunities. *J Clin Oncol* 2022 Aug 9;JCO2200995.
- 15 Hackshaw A, Clarke C, Hartman A-R. New genomic technologies for multi-cancer early detection: Rethinking the scope of cancer screening. *Cancer Cell* 2022;40:109-113.
- 16 Raoof S, Lee R, Jajoo K, et al. Multicancer Early Detection Technologies: A Review Informed by Past Cancer Screening Studies *Cancer Epidemiol Biomarkers Prev* 2022;31:1139-1145.
- 17 Offit K, Levrin O, Mullaney B, et al. Shared genetic susceptibility to breast cancer, brain tumors, and Fanconi anemia. *J Natl Cancer Inst* 2003;95:1548-1551.
- 18 Kurian AW, Ward KC, Abrahamse P, et al. Time trends in receipt of germline genetic testing and results for women diagnosed with breast cancer or ovarian cancer, 2012-2019. *J Clin Oncol* 2021;39:1631-1640.
- 19 Esterling L, Wijayatunge R, Brown K, et al. Impact of a cancer gene variant reclassification program over a 20-year period. *JCO Precis Oncol* 2020 Aug 27;4:PO.20.00020.
- 20 Mersch J, Brown N, Pirzadeh-Miller S, et al. Prevalence of variant reclassification following hereditary cancer genetic testing. *JAMA* 2018;320:1266-1274.
- 21 Balmaña J, Digiovanni L, Gaddam P, et al. Conflicting interpretation of genetic variants and cancer risk by commercial laboratories as assessed by the prospective registry of multiplex testing. *J Clin Oncol* 2016;34:4071-4078.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PEDIGREE: FIRST-, SECOND-, AND THIRD-DEGREE RELATIVES OF PROBAND<sup>a</sup>**



<sup>a</sup> First-degree relatives: parents, siblings, and children;  
second-degree relatives: grandparents, aunts, uncles, nieces, nephews, grandchildren, and half-siblings;  
third-degree relatives: great-grandparents, great-aunts, great-uncles, great-grandchildren, first cousins, and half aunts and uncles.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### COMMON PEDIGREE SYMBOLS<sup>b</sup>



<sup>b</sup> Bennett RL, Steinhaus KA, Uhrich SB, et al. Recommendations for standardized human pedigree nomenclature. Pedigree Standardization Task Force of the National Society of Genetic Counselors. Am J Hum Genet 1995;56:745-752.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### GENERAL TESTING CRITERIA<sup>a</sup>

#### **Testing is clinically indicated in the following scenarios:**

- Individuals with any blood relative with a known P/LP variant in a cancer susceptibility gene
- Individuals meeting the criteria below but who tested negative with previous limited testing (eg, single gene and/or absent deletion/duplication analysis) and are interested in pursuing multi-gene testing
- A P/LP variant identified on tumor genomic testing that has clinical implications if also identified in the germline
- To aid in systemic therapy and surgical decision-making<sup>b</sup>
- Individual who meets Li-Fraumeni syndrome (LFS) testing criteria ([see CRIT-7](#)) or Cowden syndrome/PTEN hamartoma tumor syndrome (PHTS) testing criteria ([see CRIT-8](#)) or Lynch syndrome [See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)
- For personal or family history of
  - ▶ Breast cancer [See Testing Criteria for High-Penetrance Breast Cancer Susceptibility Genes \(CRIT-2\)](#)
  - ▶ Ovarian cancer [See Testing Criteria for High-Penetrance Ovarian Cancer Susceptibility Genes \(CRIT-4\)](#)
  - ▶ Pancreatic cancer [See Testing Criteria for Pancreatic Cancer Susceptibility Genes \(CRIT-5\)](#)
  - ▶ Prostate cancer [See Testing Criteria for Prostate Cancer Susceptibility Genes \(CRIT-6\)](#)
  - ▶ Colorectal cancer [See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)

#### **Testing *may be* considered in the following scenario (with appropriate pre-test education and access to post-test management):**

- An individual of Ashkenazi Jewish ancestry<sup>c</sup> without additional risk factors
- Personal history of serous endometrial cancer<sup>d</sup>

For a list of NCCN Guidelines that include content focused on inherited cancer conditions, including criteria for testing and/or cancer risk management based on a genetic test result, see [Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines \(SUMM-1\)](#).

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>b</sup> Eg, PARP inhibitors for ovarian cancer, prostate cancer, pancreatic cancer, and metastatic HER2-negative breast cancer; platinum therapy for prostate cancer and pancreatic cancer; and risk-reducing surgery. See the relevant [NCCN Treatment Guidelines](#) for further details.

<sup>c</sup> Testing for three founder P/LP variants of *BRCA1/2* may be offered to individuals as early as age 18–25 years, who have one grandparent identified as of Ashkenazi Jewish ancestry, irrespective of cancer history in the family, as part of longitudinal studies. For those without access to longitudinal research studies, testing may be provided if there is access to pre-test education along with post-test counseling, additional genetic testing if indicated, and high-risk management. Testing should not be offered outside of a medical framework or clinical trial.

<sup>d</sup> This is a rare subtype of uterine cancer for which there is evolving evidence of an association with *BRCA1* P/LP variants.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR HIGH-PENETRANCE BREAST CANCER SUSCEPTIBILITY GENES (Specifically *BRCA1*, *BRCA2*, *CDH1*, *PALB2*, *PTEN*, and *TP53*. See [GENE-A](#))<sup>a,e,f,g</sup>

#### Testing is clinically indicated in the following scenarios:

- See General Testing Criteria on [CRIT-1](#).

- Personal history of breast cancer with specific features:

- ▶ ≤50 y

- ▶ Any age:

- ◊ Treatment indications

- To aid in systemic treatment decisions using PARP inhibitors for breast cancer in the metastatic setting<sup>h,i</sup> (See [NCCN Guidelines for Breast Cancer](#))
      - To aid in adjuvant treatment decisions with olaparib for high-risk,<sup>j</sup> HER2-negative breast cancer<sup>h</sup>

- ◊ Pathology/histology

- Triple-negative breast cancer
    - Multiple primary breast cancers (synchronous or metachronous)<sup>k</sup>
    - Lobular breast cancer with personal or family history of diffuse gastric cancer [See NCCN Guidelines for Gastric Cancer](#)

- ◊ Male breast cancer

- ◊ Ancestry: Ashkenazi Jewish ancestry

- ▶ Any age (continued):

- ◊ Family history<sup>l</sup>

- ≥1 close blood relative<sup>m</sup> with ANY:
        - breast cancer at age ≤50
        - male breast cancer
        - ovarian cancer
        - pancreatic cancer
        - prostate cancer with metastatic,<sup>n</sup> or high- or very-high-risk group (Initial Risk Stratification and Staging Workup in [NCCN Guidelines for Prostate Cancer](#))
      - ≥3 total diagnoses of breast cancer in patient and/or close blood relatives<sup>m</sup>
      - ≥2 close blood relatives<sup>m</sup> with either breast or prostate cancer (any grade)

Criteria met

→ [See GENE-1](#)

If testing criteria not met, consider testing criteria for other hereditary syndromes

If criteria for other hereditary syndromes not met, then cancer screening as per [NCCN Screening Guidelines](#)

- Family history of cancer only

- ▶ An affected individual (not meeting testing criteria listed above) or unaffected individual with a first- or second-degree blood relative meeting any of the criteria listed above (except unaffected individuals whose relatives meet criteria only for systemic therapy decision-making).<sup>o</sup>

- ◊ If the affected relative has pancreatic cancer or prostate cancer only first-degree relatives should be offered testing unless indicated based on additional family history.

- ▶ An affected or unaffected individual who otherwise does not meet the criteria above but has a probability >5% of a *BRCA1/2* pathogenic variant based on prior probability models (eg, Tyrer-Cuzick, BRCAPro, CanRisk)<sup>p</sup>

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued on CRIT-3](#)

[Footnotes on CRIT-2A](#)



### TESTING CRITERIA FOR HIGH-PENETRANCE BREAST CANCER SUSCEPTIBILITY GENES

- <sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).
- <sup>e</sup> Testing for pathogenic variants in other genes should take into consideration factors such as patient preferences, turnaround time, and insurance restrictions to particular labs (and thus particular panels). The prevalence of VUS increases with testing of additional genes. Individuals should have pre-test education on the challenges in managing pathogenic variants in genes associated with specific syndromes (eg, *CDH1* and *TP53* given their expanding clinical phenotypes) in the absence of a family history typical of such syndromes (does not apply for de novo pathogenic variants). Patients should also have pre-test education regarding the uncertain clinical utility of identifying certain pathogenic variants (eg, monoallelic *MUTYH*).
- <sup>f</sup> Meeting one or more of these criteria warrants further personalized risk assessment, genetic counseling, and often genetic testing and management.
- <sup>g</sup> For the purposes of these guidelines, invasive and ductal carcinoma in situ breast cancers should be included.
- <sup>h</sup> Robson M, et al. *N Engl J Med* 2017;377:523-533. Litton JK, et al. *N Engl J Med* 2018;379:753-763.
- <sup>i</sup> As indicated in the criteria, testing is recommended for all triple-negative breast cancers, and these indications are specifically for PARP inhibitor eligibility.
- <sup>j</sup> The definition of high-risk disease is that used in the phase III OlympiA trial, which compared adjuvant olaparib to placebo among *BRCA1/BRCA2* carriers with high-risk disease (Tutt ANJ, et al. *Engl J Med* 2021;384:2394-2405). The definition includes:
- Triple-negative breast cancer treated with either:
    - adjuvant chemotherapy with axillary node-positive disease or an invasive primary tumor  $\geq 2$  cm on pathology analysis
    - neoadjuvant chemotherapy with residual invasive breast cancer in the breast or resected lymph nodes.
  - Hormone receptor-positive disease treated with either:
    - adjuvant chemotherapy with  $\geq 4$  positive pathologically confirmed lymph nodes
    - neoadjuvant chemotherapy that did not have a complete pathologic response, with a CPS+EG score of 3 or higher.
  - The CPS+EG scoring system is based on a combination of clinical and pathologic stage, estrogen receptor status, and histologic grade. [See Neoadjuvant Therapy Outcomes Calculator](#) (Jeruss JS, et al. *J Clin Oncol* 2008;26:246-252; Mittendorf EA, et al. *J Clin Oncol* 2011;29:1956-1962). [See NCCN Guidelines for Breast Cancer](#) for further details.
- <sup>k</sup> Weitzel JN, et al. *Breast Cancer Res Treat* 2021;188:759-768.
- <sup>l</sup> Unknown or limited family structure (Weitzel JN, et al. *JAMA* 2007;297:2587-2595).
- <sup>m</sup> Close blood relatives include first-, second-, and third-degree relatives on the same side of the family. ([See EVAL-B](#))
- <sup>n</sup> Metastatic prostate cancer is biopsy-proven and/or with radiographic evidence and includes distant metastasis and regional bed or nodes. It is not a biochemical recurrence only. Prostate cancer-specific mortality should be a surrogate for metastatic disease for family history purposes.
- <sup>o</sup> This may be extended to an affected third-degree relative if related through two male relatives (eg, paternal grandfather's mother or sister). If the affected first-degree relative underwent genetic testing and is negative for detectable P/LP variants and there is no other family history of cancer, there is a low probability that any finding will have documented clinical utility.
- <sup>p</sup> The approximate 5% threshold for probability of carrying *BRCA1/2* pathogenic variants is utilized because of availability of prior probability models; however, it is recognized that current model estimates vary substantially, and that different thresholds may be appropriate if other genes are included in the model utilized. If genes other than *BRCA1* and *BRCA2* are to be included in models evaluating the threshold for testing, the penetrance, clinical actionability, and phenotypic features of cancers associated with P/LP variants in these genes should be considered. The panel encourages the development of validated models that include these parameters to determine eligibility and appropriateness for gene panel testing for inherited cancer risk. These models are only validated for *BRCA1/2*.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR HIGH-PENETRANCE BREAST CANCER SUSCEPTIBILITY GENES (continued)

**Testing *may be* considered in the following scenarios (with appropriate pre-test education and access to post-test management):**

- Personal history of breast cancer <60 y not meeting any of the above criteria may approach a 2.5% probability of having a PV, based on recent data.<sup>q</sup> It is cautioned that the majority of those PVs will be in moderate penetrance genes, which are over-represented in older affected individuals, and for which data on appropriate management are often lacking. Access to an experienced genetic counseling team to discuss management options is particularly important in this setting.
- Personal history of breast cancer diagnosed at any age with ≥1 close blood relative<sup>m</sup> with intermediate-risk prostate cancer with intraductal/criform histology (see Initial Risk Stratification and Staging Workup in [NCCN Guidelines for Prostate Cancer](#))
- An affected or unaffected individual who otherwise does not meet any of the above criteria but with a 2.5%–5% probability of *BRCA1/2* pathogenic variant based on prior probability models (eg, Tyrer-Cuzick, BRCAPro, CanRisk)<sup>e</sup>

**There is a low probability (<2.5%) that testing will have findings of documented high-penetrance genes in the following scenarios:**

- Female diagnosed with breast cancer at age >60 y, with no close relative<sup>m</sup> with breast, ovarian, pancreatic, or prostate cancer.
- Diagnosed with localized prostate cancer with Gleason Score <7 and no close relative<sup>m</sup> with breast, ovarian, pancreatic, or prostate cancer.

**NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.**

<sup>e</sup> Testing for pathogenic variants in other genes should take into consideration factors such as patient preferences, turnaround time, and insurance restrictions to particular labs (and thus particular panels). The prevalence of VUS increases with testing of additional genes. Individuals should have pre-test education on the challenges in managing pathogenic variants in genes associated with specific syndromes (eg, *CDH1* and *TP53* given their expanding clinical phenotypes) in the absence of a family history typical of such syndromes (does not apply for de novo pathogenic variants). Patients should also have pre-test education regarding the uncertain clinical utility of identifying certain pathogenic variants (eg, monoallelic *MUTYH*).

<sup>m</sup> Close blood relatives include first-, second-, and third-degree relatives on the same side of the family. ([See EVAL-B](#))

<sup>q</sup> Kurian A, et al. JAMA 2020;323:995-997.


**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



### TESTING CRITERIA FOR OVARIAN CANCER SUSCEPTIBILITY GENES<sup>a</sup>

(See [GENE-A](#))



NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>o</sup> This may be extended to an affected third-degree relative if related through two male relatives (eg, paternal grandfather's mother or sister). If the affected first-degree relative underwent genetic testing and is negative for detectable P/LP variants and there is no other family history of cancer, there is a low probability that any finding will have documented clinical utility.

<sup>p</sup> The approximate 5% threshold for probability of carrying *BRCA1/2* pathogenic variants is utilized because of availability of prior probability models; however, it is recognized that current model estimates vary substantially, and that different thresholds may be appropriate if other genes are included in the model utilized. If genes other than *BRCA1* and *BRCA2* are to be included in models evaluating the threshold for testing, the penetrance, clinical actionability, and phenotypic features of cancers associated with P/LP variants in these genes should be considered. The panel encourages the development of validated models that include these parameters to determine eligibility and appropriateness for gene panel testing for inherited cancer risk. These models are only validated for *BRCA1/2*.

<sup>r</sup> *BRCA*-related ovarian cancers are associated with epithelial, non-mucinous histology. Lynch syndrome can be associated with both non-mucinous and mucinous epithelial tumors. Be attentive for clinical evidence of Lynch syndrome (see [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)). Specific types of non-epithelial ovarian cancers and tumors can also be associated with other rare syndromes. Examples include an association between sex-cord tumors with annular tubules and PJS or Sertoli-Leydig tumors and *DICER1*-related disorders.

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### TESTING CRITERIA FOR PANCREATIC CANCER SUSCEPTIBILITY GENES (See [GENE-A](#))<sup>a</sup>

<b>Testing is clinically indicated in the following scenarios:</b>
• See General Testing Criteria on <a href="#">CRIT-1</a> .
• Exocrine pancreatic cancers <sup>s</sup> <ul style="list-style-type: none"><li>▶ All individuals diagnosed with exocrine pancreatic cancer<sup>t</sup></li><li>▶ First-degree relatives of individuals diagnosed with exocrine pancreatic cancer<sup>u</sup></li></ul>
• Neuroendocrine pancreatic tumors - <a href="#">See NCCN Guidelines for Neuroendocrine and Adrenal Tumors</a>



<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>s</sup> Genes that are typically tested for pancreatic cancer risk include *ATM*, *BRCA1*, *BRCA2*, *CDKN2A*, most Lynch syndrome genes (*MLH1*, *MSH2*, *MSH6*, and *EPCAM*), *PALB2*, *STK11*, and *TP53*.

<sup>t</sup> Pancreatic cancer risk is higher in individuals of Ashkenazi Jewish descent. Genetic testing of Ashkenazi Jewish patients with pancreatic cancer may have a higher yield of P/LP variants than of non-Ashkenazi Jewish patients.

<sup>u</sup> Testing of first-degree relatives should only be done if it is impossible to test the individual who has pancreatic cancer. Some second-degree relatives may meet testing criteria based on additional family history. Approximately 2%–5% of unselected cases of pancreatic adenocarcinoma will have a *BRCA1/2* P/LP variant. However, the disease is highly aggressive and the option to test the affected relative may not be available in the future. Thus, there may be significant benefit to family members in testing these patients near the time of diagnosis. In addition, increasing evidence suggests that identification of a *BRCA1/2* P/LP variant may direct use of targeted therapies for patients with pancreatic cancer ([See NCCN Guidelines for Pancreatic Adenocarcinoma](#)). (Holter S, et al. J Clin Oncol 2015;33:3124-3129. Shindo K, et al. J Clin Oncol 2017;35:3382-3390. Golan T, et al. N Engl J Med 2019;381:317-327.) Family history of pancreatic cancer of unknown histology is often assumed to be an exocrine pancreatic cancer.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR HIGH-PENETRANCE PROSTATE CANCER SUSCEPTIBILITY GENES (See [GENE-A](#))<sup>a</sup>

#### Testing is clinically indicated in the following scenarios:

- See General Tumor Criteria on [CRIT-1](#).
- Personal history of prostate cancer with specific features:
  - ▶ By tumor characteristics (any age)
    - ◊ Metastatic<sup>n</sup>
    - ◊ Histology
      - high- or very-high-risk group (see Initial Risk Stratification and Staging Workup in [NCCN Guidelines for Prostate Cancer](#))
  - ▶ By family history and ancestry
    - ◊ ≥1 close blood relative<sup>m</sup> with:
      - breast cancer at age ≤50 y
      - triple-negative breast cancer at any age
      - male breast cancer at any age
      - ovarian cancer at any age
      - pancreatic cancer at any age
      - metastatic,<sup>n</sup> high- or very-high-risk group (see Initial Risk Stratification and Staging Workup in [NCCN Guidelines for Prostate Cancer](#)) at any age
    - ◊ ≥2 close blood relatives<sup>m</sup> with either breast or prostate cancer (any grade) at any age
    - ◊ Ashkenazi Jewish ancestry<sup>c</sup>
- Family history of cancer only
  - ▶ An affected (not meeting testing criteria listed above) or unaffected individual with a first-degree blood relative meeting any of the criteria listed above (except unaffected individuals whose relatives meet criteria only for systemic therapy decision-making)<sup>o</sup>

Criteria met → [See GENE-1](#)

If testing criteria not met, consider testing criteria for other hereditary syndromes

If criteria for other hereditary syndromes not met, then cancer screening as per [NCCN Screening Guidelines](#)

#### Testing *may be considered* in the following scenario:

- Personal history of prostate cancer with intermediate-risk prostate cancer with intraductal/cribriform histology (see Initial Risk Stratification and Staging Workup in [NCCN Guidelines for Prostate Cancer](#)) at any age

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR HIGH-PENETRANCE PROSTATE CANCER SUSCEPTIBILITY GENES (See [GENE-A](#))<sup>a</sup>

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>c</sup> Testing for three founder P/LP variants of *BRCA1/2* may be offered to individuals as early as age 18–25 years, who have one grandparent identified as of Ashkenazi Jewish ancestry, irrespective of cancer history in the family, as part of longitudinal studies. For those without access to longitudinal research studies, testing may be provided if there is access to pre-test education along with post-test counseling, additional genetic testing if indicated, and high-risk management. Testing should not be offered outside of a medical framework or clinical trial.

<sup>m</sup> Close blood relatives include first-, second-, and third-degree relatives on the same side of the family ([See EVAL-B](#)).

<sup>n</sup> Metastatic prostate cancer is biopsy-proven and/or with radiographic evidence and includes distant metastasis and regional bed or nodes. It is not a biochemical recurrence only. Prostate cancer-specific mortality should be a surrogate for metastatic disease for family history purposes.

<sup>o</sup> This may be extended to an affected third-degree relative if related through two male relatives (eg, paternal grandfather's mother or sister). If the affected first-degree relative underwent genetic testing and is negative for detectable P/LP variants and there is no other family history of cancer, there is a low probability that any finding will have documented clinical utility.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR LI-FRAUMENI SYNDROME<sup>a</sup>

**Testing is clinically indicated in the following scenarios:<sup>\*</sup>**

- See General Testing Criteria on [CRIT-1](#).
- Individual from a family with a known *TP53*<sup>v</sup> P/LP variant
- Classic Li-Fraumeni syndrome (LFS) criteria:<sup>w</sup>
  - Combination of an individual diagnosed at age <45 years with a sarcoma<sup>u</sup> **AND**  
A first-degree relative diagnosed at age <45 years with cancer **AND**  
An additional first- or second-degree relative in the same lineage with cancer diagnosed at age <45 years, or a sarcoma at any age
- Chompret criteria:<sup>x</sup>
  - Individual with a tumor from LFS tumor spectrum (eg, soft tissue sarcoma, osteosarcoma, CNS tumor, breast cancer, adrenocortical carcinoma), before 46 years of age, **AND** at least one first- or second-degree relative with any of the aforementioned cancers (other than breast cancer if the proband has breast cancer) before the age of 56 years or with multiple primaries at any age **OR**
  - Individual with multiple tumors (except multiple breast tumors), two of which belong to LFS tumor spectrum with the initial cancer occurring before the age of 46 years **OR**
  - Individual with adrenocortical carcinoma, or choroid plexus carcinoma or rhabdomyosarcoma of embryonal anaplastic subtype, at any age of onset, regardless of family history **OR**
  - Breast cancer before 31 years of age
- Pediatric hypodiploid acute lymphoblastic leukemia
- Affected individual with P/LP variant identified on tumor genomic testing that may have implications if also identified on germline testing<sup>y</sup>

LFS testing criteria met → [See GENE-1](#)

If LFS testing criteria not met, consider testing criteria for other hereditary syndromes, if appropriate

Individualized recommendations according to personal and family history

<sup>\*</sup> Other cancers associated with LFS but not in the testing criteria include: melanoma, colorectal, gastric, and prostate.

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>v</sup> When this gene is included as part of a multi-gene panel, an individual does not need to meet these testing criteria if testing criteria on other testing criteria pages are met.

<sup>w</sup> Li FP, et al. Cancer Res 1988;48:5358-5362. To date, there have been no reports of Ewing sarcoma, gastrointestinal stromal tumor (GIST), desmoid tumor, or angiosarcoma in *TP53* P/LP variant carriers.

<sup>x</sup> Chompret A, et al. J Med Genet 2001;38:43-47; Bougeard G, et al. J Clin Oncol 2015;33:2345-2352.

<sup>y</sup> This should prompt a careful evaluation of personal and family history of the individual to determine the yield of germline sequencing. Somatic *TP53* P/LP variants are common in many tumor types in absence of a germline P/LP variant.

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR COWDEN SYNDROME (CS)/PTEN HAMARTOMA TUMOR SYNDROME (PHTS)<sup>a,z,aa,bb</sup>

#### Testing is clinically indicated in the following scenarios:

- See General Testing Criteria on [CRIT-1](#).
- Individual from a family with a known *PTEN*<sup>v</sup> P/LP variant
- Individual with a personal history of Bannayan-Riley-Ruvalcaba syndrome (BRRS)
- Individual meeting clinical diagnostic criteria<sup>cc</sup> for CS/PHTS
- Individual not meeting clinical diagnostic criteria<sup>cc</sup> for CS/PHTS with a personal history of:
  - ▶ Adult Lhermitte-Duclos disease (cerebellar tumors); or
  - ▶ Autism spectrum disorder and macrocephaly; or
  - ▶ Two or more biopsy-proven trichilemmomas; or
  - ▶ Two or more major criteria (one must be macrocephaly); or
  - ▶ Three major criteria, without macrocephaly; or
  - ▶ One major and ≥3 minor criteria;<sup>dd</sup> or
  - ▶ ≥4 minor criteria
- At-risk individual with a relative with a clinical diagnosis of CS/PHTS or BRRS for whom testing has not been performed
  - ▶ The at-risk individual must have the following:
    - ◇ Any one major criterion or
    - ◇ Two minor criteria
- *PTEN* P/LP variant detected by tumor genomic testing on any tumor type in the absence of germline analysis<sup>ee</sup>



[See major and minor criteria on CRIT-8A.](#)

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>v</sup> When this gene is included as part of a multi-gene panel, an individual does not need to meet these testing criteria if testing criteria on other testing criteria pages are met.

<sup>z</sup> These are testing criteria; clinical diagnostic criteria can be found on [CRIT-8A](#).

<sup>aa</sup> If two criteria involve the same structure/organ/tissue, both may be included as criteria.

<sup>bb</sup> Current evidence does not support testing for succinate dehydrogenase (SDH) gene P/LP variants in patients with PHTS. (Bayley J-P. Am J Hum Genet 2011;88:674-675).

<sup>cc</sup> Pilarski R, et al. J Natl Cancer Inst 2013;105:1607-1616. [See COWD-A](#).

<sup>dd</sup> If an individual has two or more major criteria, such as breast cancer and nonmedullary thyroid cancer, but does not have macrocephaly, one of the major criteria may be included as one of the three minor criteria to meet testing criteria.

<sup>ee</sup> This should prompt a careful evaluation of personal and family history of the individual to determine the yield of germline sequencing. Somatic *PTEN* P/LP variants are common in many tumor types in absence of germline P/LP variant.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### TESTING CRITERIA FOR COWDEN SYNDROME (CS)/PTEN HAMARTOMA TUMOR SYNDROME (PHTS)<sup>a,\*</sup>

#### Major criteria:

- Breast cancer
- Endometrial cancer
- Follicular thyroid cancer
- Multiple GI hamartomas or ganglioneuromas<sup>ff</sup>
- Macrocephaly (megalcephaly) (ie,  $\geq 97\%$ , 58 cm in adult female, 60 cm in adult male)<sup>gg</sup>
- Macular pigmentation of glans penis
- Mucocutaneous lesions<sup>hh</sup>
  - One biopsy-proven trichilemmoma
  - Multiple palmpoplantar keratoses
  - Multifocal or extensive oral mucosal papillomatosis
  - Multiple cutaneous facial papules (often verrucous)

#### Minor criteria:<sup>ii</sup>

- Autism spectrum disorder
- Colon cancer
- $\geq 3$  esophageal glycogenic acanthoses
- Lipomas
- Intellectual disability (ie, IQ  $\leq 75$ )
- Papillary or follicular variant of papillary thyroid cancer
- Thyroid structural lesions (eg, adenoma, nodule[s], goiter)
- Renal cell carcinoma
- Single GI hamartoma or ganglioneuroma
- Testicular lipomatosis
- Vascular anomalies (including multiple intracranial developmental venous anomalies)

\* Other cancers associated with *PTEN* but not in the testing criteria include: colorectal, kidney cancer, and melanoma.

### REVISED CLINICAL DIAGNOSTIC CRITERIA FOR PTEN HAMARTOMA TUMOR SYNDROME<sup>cc</sup>

#### Operational diagnosis in an individual (either of the following):

1. Three or more major criteria, but one must include macrocephaly, Lhermitte-Duclos disease, or GI hamartomas; or
2. Two major and three minor criteria.

#### Operational diagnosis in a family where one individual meets revised PTEN hamartoma tumor syndrome clinical diagnostic criteria or has a *PTEN* P/LP variant:

1. Any two major criteria with or without minor criteria; or
2. One major and two minor criteria; or
3. Three minor criteria.

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>cc</sup> Pilarski R, et al. J Natl Cancer Inst 2013;105:1607-1616. [See COWD-A](#).

<sup>ff</sup> Multiple polyp types are often seen in patients with PHTS, and less commonly may include adenomas, hyperplastic polyps, and other histologies.

<sup>gg</sup> Roche AF, et al. Pediatrics 1987;79:706-712.


<sup>hh</sup> The literature available on mucocutaneous lesions is not adequate to accurately specify the number or extent of mucocutaneous lesions required to be a major criterion for CS/PHTS. Clinical judgment should be used.

<sup>ii</sup> Insufficient evidence exists in the literature to include fibrocystic disease of the breast, fibromas, and uterine fibroids as diagnostic criteria.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





<sup>a</sup> For further details regarding the nuances of genetic counseling and testing, [see EVAL-A](#).

<sup>b</sup> If of Ashkenazi Jewish descent, in addition to the specific familial P/LP variant, test for all three founder P/LP variants.

<sup>c</sup> Additional testing may be indicated if there is also a significant family history of cancer on the side of the family without the known P/LP variant.

<sup>d</sup> If no P/LP variant is found, consider testing another family member with next highest likelihood of having a P/LP variant.

<sup>e</sup> Patients meeting CS/PHTS clinical diagnostic criteria ([see COWD-A 1 of 2](#)) should be managed as P/LP variant carriers.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
ATM	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 20%–40%<sup>3,4,5,6</sup></li><li>• <b>Management:</b><sup>b</sup><ul style="list-style-type: none"><li>▶ Screening: Annual mammogram at age 40 y and consider breast MRI with contrast starting at age 30–35 y<sup>c,d,e</sup></li><li>▶ Risk reduction: Evidence insufficient for risk-reducing mastectomy (RRM), manage based on family history</li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 2%–3%<sup>10-12</sup></li><li>• <b>Management:</b><sup>f</sup><ul style="list-style-type: none"><li>▶ Risk reduction: Evidence insufficient for risk-reducing salpingo oophorectomy (RRSO); manage based on family history</li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<b>Pancreatic cancer</b> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> ~5%–10%<sup>9,23</sup></li><li>• <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer, <a href="#">see PANC-A</a>.</li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul> <b>Prostate cancer</b> <ul style="list-style-type: none"><li>• Emerging evidence for association with increased risk<sup>24</sup></li></ul>
	Comments: Heterozygous ATM P/LP variants should not lead to a recommendation to avoid radiation therapy at this time. <a href="#">See Discussion</a> for information regarding the c.7271T>G variant. See <a href="#">GENE-B</a> for reproductive implications/ recessive disease.		
BARD1	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 20%–40%<sup>7</sup></li><li>• <b>Management:</b><ul style="list-style-type: none"><li>▶ Screening: Annual mammogram and consider breast MRI with contrast starting at age 40 y<sup>c,d,e</sup></li><li>▶ Risk reduction: Evidence insufficient for RRM, manage based on family history</li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong<sup>7-9</sup></li></ul>	<b>Evidence of increased risk:</b> No established association	<b>Other cancers</b> <ul style="list-style-type: none"><li>• Unknown or insufficient evidence</li></ul>

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.  
**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
BRCA1	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> &gt;60%<sup>5,25-29</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li><li>• <b>Strength of evidence of association with cancer:</b> Very strong (with predisposition to triple-negative disease)</li></ul> <p><b>Male breast cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 0.2%–1.2% by age 70 y<sup>30,31</sup></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 39%–58%<sup>33</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li><li>• <b>Strength of evidence of association with cancer:</b> Very strong</li></ul>	<p><b>Pancreatic cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> ≤5%<sup>31</sup></li><li>• <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer, <a href="#">see PANC-A</a>.</li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul> <p><b>Prostate cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 7%–26%<sup>34</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li></ul>
	Comment: See <a href="#">GENE-B</a> for reproductive implications/ recessive disease.		
BRCA2	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> &gt;60%<sup>5,21-25</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li><li>• <b>Strength of evidence of association with cancer:</b> Very strong</li></ul> <p><b>Male breast cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 1.8%–7.1% by age 70 y<sup>30,31,32</sup></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 13%–29%<sup>33</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li><li>• <b>Strength of evidence of association with cancer:</b> Very strong</li></ul>	<p><b>Pancreatic cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 5%–10%<sup>31</sup></li><li>• <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer, <a href="#">see PANC-A</a>.</li><li>• <b>Strength of evidence of association with cancer:</b> Very strong</li></ul> <p><b>Prostate cancer</b></p> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 19%–61%<sup>34,35</sup></li><li>• <b>Management:</b> <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li></ul> <p><b>Melanoma</b></p> <ul style="list-style-type: none"><li>• <a href="#">See BRCA Pathogenic Variant-Positive Management</a></li></ul>
	Comment: See <a href="#">GENE-B</a> for reproductive implications/ recessive disease.		

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.  
**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
<i>BRIP1</i>	<ul style="list-style-type: none"> <li><b>Absolute risk:</b> Insufficient data to define</li> <li><b>Management:</b> Insufficient data; managed based on family history</li> <li><b>Strength of evidence of association with cancer:</b> Limited; potential increase in female breast cancer<sup>8</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Absolute risk:</b> 5%–15%<sup>10-12,39</sup></li> <li><b>Management:</b> <ul style="list-style-type: none"> <li>▶ Risk reduction: Recommend RRSO at age 45–50 y<sup>h</sup></li> </ul> </li> <li><b>Strength of evidence of association with cancer:</b> Strong</li> </ul>	<b>Other cancers</b> <ul style="list-style-type: none"> <li>Unknown or insufficient evidence</li> </ul>
Comments: See <a href="#">GENE-B</a> for reproductive implications/ recessive disease. Based on estimates from available studies, the lifetime risk of ovarian cancer in carriers of P/LP variants in <i>BRIP1</i> appears to be sufficient to justify RRSO. The current evidence is insufficient to make a firm recommendation as to the optimal age for this procedure. Based on the current, limited evidence base, a discussion about surgery should be held around age 45–50 y or earlier based on a specific family history of an earlier onset of ovarian cancer.			
<i>CDH1</i>	<ul style="list-style-type: none"> <li><b>Absolute risk:</b> 41%–60%<sup>36-39</sup></li> <li><b>Management:</b><sup>b</sup> <ul style="list-style-type: none"> <li>▶ Screening: Annual mammogram and consider breast MRI with contrast starting at age 30 y<sup>c,d</sup></li> <li>▶ Risk reduction: Discuss option of RRM</li> </ul> </li> <li><b>Strength of evidence of association with cancer:</b> Strong</li> </ul>	<b>Evidence of increased risk:</b> No established association	<b>Hereditary diffuse gastric cancer (HDGC)</b> <ul style="list-style-type: none"> <li><b>Strength of evidence of association with cancer:</b> Strong</li> <li>See <a href="#">NCCN Guidelines for Gastric Cancer</a>: Principles of Genetic Risk Assessment for Gastric Cancer</li> </ul>
Comments: There is controversy over how to manage gastric cancer risk in individuals with P/LP variants in <i>CDH1</i> in the absence of a family history of gastric cancer. However, one small study found that >50% of such individuals had gastric cancer identified at the time of risk-reducing total gastrectomy (Jacobs MF, et al. Gastroenterology 2019;157:87-96), and penetrance for lifetime risk is increased with a positive family history of HDGC (Roberts ME, et al. JAMA Oncol 2019;5:1325-1331). Cleft lip with or without cleft palate has been associated with <i>CDH1</i> P/LP variants (Frebourg T, et al. J Med Genet 2006;43:138-142).			
<i>CDKN2A</i>	<b>Evidence of increased risk:</b> No established association	<b>Evidence of increased risk:</b> No established association	<b>Pancreatic cancer</b> <ul style="list-style-type: none"> <li><b>Absolute risk:</b> &gt;15%</li> <li><b>Management:</b> Screening, see <a href="#">PANC-A</a>.</li> <li><b>Strength of evidence of association with cancer:</b> Very strong</li> </ul> <b>Melanoma</b> <ul style="list-style-type: none"> <li><b>Absolute risk:</b> 28%–76% depending on other risk factors, including family history, geographic location, and other genetic modifiers<sup>40,41</sup></li> <li><b>Strength of evidence of association with cancer:</b> Strong</li> </ul>
Comments: Comprehensive skin examination by a dermatologist, supplemented with total body photography and dermoscopy is recommended biannually (Chan SH, et al. Hered Cancer Clin Pract 2021;19:21).			

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**GENE-A**  
**3 OF 10**



# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
<i>CHEK2</i>	<ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> 20%–40%<sup>5,6,42,43</sup></li> <li>• <b>Management:</b><sup>b</sup> <ul style="list-style-type: none"> <li>▶ Screening: Annual mammogram at age 40 y and consider breast MRI with contrast starting at age 30–35 y<sup>c,d,e</sup></li> <li>▶ Risk reduction: Evidence insufficient for RRM, manage based on family history</li> </ul> </li> <li>• <b>Strength of evidence of association with cancer:</b> Strong<sup>44</sup></li> </ul>	Evidence of increased risk: No established association	<b>Colorectal cancer</b> <ul style="list-style-type: none"> <li>• <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal (GENE-1)</a></li> </ul>
Comments: Risk data are based only on frameshift P/LP variants. The risks for most missense variants are unclear but for some P/LP variants, such as Ile157Thr, the risk for breast cancer appears to be lower and does not reach the threshold for management change. Management should be based on best estimates of cancer risk for the specific P/LP variant.			
<i>MSH2, MLH1, MSH6, PMS2, EPCAM</i> <sup>f</sup>	<b><u>MLH1, MSH2, MSH6, PMS2, and EPCAM</u></b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> &lt;15%<sup>45,46</sup></li> <li>• <b>Management:</b> Insufficient data; managed based on family history</li> <li>• <b>Strength of evidence of association with cancer:</b> Limited</li> </ul>	<b><u>MLH1</u></b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> 4%–20%<sup>47,48</sup></li> <li>• <b>Strength of evidence:</b> Strong</li> </ul> <b><u>MSH2 /EPCAM</u></b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> 8%–38%<sup>47,48,50,51</sup></li> <li>• <b>Strength of evidence:</b> Strong</li> </ul> <b><u>MSH6</u></b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> ≤1%–13%<sup>49,50</sup></li> <li>• <b>Strength of evidence:</b> Strong</li> </ul> <b><u>PMS2</u></b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> 1.3%–3%<sup>51</sup></li> <li>• <b>Strength of evidence:</b> Limited</li> </ul> <ul style="list-style-type: none"> <li>• <b>Management for all genes:</b> <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a></li> </ul>	<b>Pancreatic cancer</b> <ul style="list-style-type: none"> <li>• <b>Absolute risk:</b> &lt;5%–10% (excluding <i>PMS2</i>)</li> <li>• <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer (insufficient evidence for <i>PMS2</i>), <a href="#">see PANC-A</a>.</li> <li>• <b>Strength of evidence of association with cancer:</b> Strong</li> </ul> <b>Colorectal, uterine, others</b> <ul style="list-style-type: none"> <li>• <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a></li> </ul>
Comments: Counsel for biallelic risk of P/LP variants that lead to CMMRD. <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a> .			

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
<i>NF1</i>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 20%–40%<sup>52,53</sup></li><li>• <b>Management:</b><sup>b</sup><ul style="list-style-type: none"><li>▶ Screening: Annual mammogram starting at age 30 y and consider breast MRI with contrast from ages 30–50 y<sup>c,d</sup></li><li>▶ Risk reduction: Evidence insufficient for RRM, manage based on family history</li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<b>Evidence of increased risk:</b> No established association	<b>Malignant peripheral nerve sheath tumors, GIST, others</b> <ul style="list-style-type: none"><li>• Recommend referral to <i>NF1</i> specialist for evaluation and management</li></ul>
Comments: At this time, there are no data to suggest an increased breast cancer risk after age 50 y. Consider possibility of false-positive MRI results due to presence of breast neurofibromas.			
<i>PALB2</i>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 41%–60%<sup>5,8,22,54</sup></li><li>• <b>Management:</b><sup>b</sup><ul style="list-style-type: none"><li>▶ Screening: Annual mammogram and breast MRI with contrast at 30 y<sup>c,d</sup></li><li>▶ Risk reduction: Discuss option of RRM</li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul> <b>Male breast cancer</b> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 0.9% by age 70 y<sup>22</sup></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 3%–5%<sup>10-12,22,61,62</sup></li><li>• <b>Management:</b><sup>f</sup><ul style="list-style-type: none"><li>▶ Risk reduction: Consider RRSO at age &gt;45 y<sup>h</sup></li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<b>Pancreatic cancer</b> <ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 5%–10%</li><li>• <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer, <a href="#">see PANC-A</a></li><li>• <b>Strength of evidence of association with cancer:</b> Limited</li></ul> <b>Other cancers</b> <ul style="list-style-type: none"><li>• Unknown or insufficient evidence</li></ul>
Comments: See <a href="#">GENE-B</a> for reproductive implications/ recessive disease.			
<i>PTEN</i>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 40%–60% (historical cohort data), &gt;60% (projected estimates)<sup>55-58</sup></li><li>• <b>Management:</b><sup>b</sup> <a href="#">See Cowden Syndrome Management</a></li><li>• <b>Strength of evidence of association with cancer:</b> Strong<sup>59,60</sup></li></ul>	<b>Evidence of increased risk:</b> No established association	<b>Thyroid, colorectal, endometrial, renal cancers</b> <ul style="list-style-type: none"><li>• <a href="#">See Cowden Syndrome Management</a></li></ul>

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.  
**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**GENE-A**  
**5 OF 10**





# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
RAD51C	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 20%–40%<sup>5,7,44</sup></li><li>• <b>Management:</b> Annual mammogram and consider breast MRI with contrast starting at age 40 y</li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 10%–15%<sup>10-12,63,64</sup></li><li>• <b>Management:</b><ul style="list-style-type: none"><li>▶ Risk reduction: Recommend RRSO at 45–50 y<sup>h</sup></li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<b>Other cancers</b> <ul style="list-style-type: none"><li>• Unknown or insufficient evidence</li></ul>
	<b>Comments:</b> See GENE-B for reproductive implications/ recessive disease. Based on estimates from available studies, the lifetime risk of ovarian cancer in carriers of P/LP variants in RAD51C appears to be sufficient to justify RRSO. The current evidence is insufficient to make a firm recommendation as to the optimal age for this procedure. Based on the current, limited evidence base, a discussion about surgery should be held around age 45–50 y or earlier based on a specific family history of an earlier onset ovarian cancer.		
RAD51D	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 20%–40%<sup>5,7,44</sup></li><li>• <b>Management:</b> Annual mammogram and consider breast MRI with contrast starting at age 40 y</li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<ul style="list-style-type: none"><li>• <b>Absolute risk:</b> 10%–20%<sup>10-12,63,64</sup></li><li>• <b>Management:</b><ul style="list-style-type: none"><li>▶ Risk reduction: Recommend RRSO at 45–50 y<sup>h</sup></li></ul></li><li>• <b>Strength of evidence of association with cancer:</b> Strong</li></ul>	<b>Other cancers</b> <ul style="list-style-type: none"><li>• Unknown or insufficient evidence</li></ul>
	<b>Comments:</b> Based on estimates from available studies, the lifetime risk of ovarian cancer in carriers of P/LP variants in RAD51D appears to be sufficient to justify RRSO. The current evidence is insufficient to make a firm recommendation as to the optimal age for this procedure. Based on the current, limited evidence base, a discussion about surgery should be held around age 45–50 y or earlier based on a specific family history of an earlier onset ovarian cancer.		

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

GENE-A  
6 OF 10



# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### CANCER RISK MANAGEMENT BASED ON GENETIC TEST RESULTS<sup>a,1,2</sup>

The inclusion of a gene in this table below does not imply the endorsement either for or against multi-gene testing for moderate-penetrance genes.

Gene	Breast Cancer Risk and Management (First primary)	Epithelial Ovarian Cancer Risk and Management	Pancreatic Cancer Risk and Management <sup>13-22</sup> and Other Cancer Risks
STK11	<b>Absolute risk:</b> 32%–54% <sup>65,66</sup> • <b>Management:</b> ▶ Screening: Annual mammogram and breast MRI with contrast starting at age 30 y <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a> - Peutz-Jeghers syndrome ▶ Risk reduction: Discuss option of RRM • <b>Strength of evidence of association with cancer:</b> Strong	<b>Evidence of increased risk:</b> No established association	<b>Pancreatic cancer</b> • <b>Absolute risk:</b> >15% • <b>Management:</b> Screening, <a href="#">see PANC-A</a> • <b>Strength of evidence of association with cancer:</b> Strong  <b>Non-Epithelial Ovarian Cancer (Sex cord with annular tubules)</b> • <b>Absolute risk:</b> >10% <sup>65</sup> • <b>Management:</b> <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a> - Peutz-Jeghers syndrome • <b>Strength of evidence of association with cancer:</b> Strong  <b>Other cancers</b> • <a href="#">See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal</a> - Peutz-Jeghers syndrome
	Comments: Case-control studies have consistently demonstrated germline <i>STK11</i> pathogenic variants to be associated with high lifetime risks of pancreatic cancer. However, these variants are rare, and the risk estimates have wide confidence intervals.		
TP53	• <b>Absolute risk:</b> >60% <sup>5,67</sup> • <b>Management:</b> <a href="#">See Li-Fraumeni Syndrome Management</a> • <b>Strength of evidence of association with cancer:</b> Strong <sup>68</sup>	<b>Evidence of increased risk:</b> No established association	<b>Pancreatic cancer</b> • <b>Absolute risk:</b> 5%–10% • <b>Management:</b> Screen P/LP variant carriers with a family history of pancreatic cancer, <a href="#">see PANC-A</a> • <b>Strength of evidence of association with cancer:</b> Limited  <b>Other cancers</b> • <a href="#">See Li-Fraumeni Syndrome Management</a>

Footnotes on [GENE-A 8 of 10](#)

References on [GENE-A 9 of 10](#) and [GENE-A 10 of 10](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### FOOTNOTES FOR TABLES

- <sup>a</sup> The following genes and others are found on some of the panels, but there is insufficient evidence to make any recommendations for breast MRI, RRSO, or RRM for: *FANCC*, *MRE11*, *MUTYH* heterozygotes, *NBN*, *RAD50*, *RECQL*, *RINT1*, *SLX4*, *SMARCA4*, or *XRCC2*; or for prostate cancer management for *HOXB13*.
- <sup>b</sup> Screening and risk-reduction management is extrapolated from *BRCA1/2* data based on risk levels.
- <sup>c</sup> May be modified based on family history (typically beginning screening 5–10 years earlier than the youngest diagnosis in the family but not later than stated in the table) or specific gene P/LP variant.
- <sup>d</sup> For patients with P/LP variants who are treated for breast cancer and have not had bilateral mastectomy, screening should continue as described.
- <sup>e</sup> The use of MRI in these patients depends on a number of risk factors, including family history, age, breast density, and patient preference.
- <sup>f</sup> Transvaginal ultrasound combined with serum CA-125 for ovarian cancer screening, although of uncertain benefit, may be considered at the clinician's discretion.
- <sup>g</sup> The higher range of risk is reflective of a prospective study of pancreatic cancer kindreds (Hsu FC, Roberts NJ, Childs E, et al. Risk of pancreatic cancer among individuals with pathogenic variants in the ATM gene. JAMA Oncol 2021;7:1664-1668).
- <sup>h</sup> Risks and benefits of premature surgical menopause versus risk of cancer and family history should all be carefully considered, and the panel recommends patients seek expert care.

<b><u>Strength of Evidence of Association with Cancer</u></b> <ul style="list-style-type: none"><li>• <b><u>Very strong</u></b>: Prospective cohort studies in a population-based setting have demonstrated risk.</li><li>• <b><u>Strong</u></b>: Traditional case-control studies or more than three case-control studies including those with cases ascertained by commercial laboratories or those without controls from the same population. Traditional case-control study: A retrospective study that compares patients with a disease or specific outcome (cases) with patients without the disease or outcome (controls).</li><li>• <b><u>Limited</u></b>: Small sample size or case series</li><li>• <b><u>None</u></b></li></ul>	<ul style="list-style-type: none"><li>• <b>Population risk (per SEER registry data)</b><ul style="list-style-type: none"><li>▶ Breast cancer: 12%–13%</li><li>▶ Ovarian cancer: 1%–2%</li><li>▶ Pancreatic cancer: 1%–2%</li></ul></li></ul>
--	--

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

### REFERENCES FOR TABLES

- 1 Tung N, Domchek SM, Stadler Z, et al. Counselling framework for moderate-penetrance cancer-susceptibility mutations. *Nat Rev Clin Oncol* 2017;13:581-588.
- 2 Domchek SM, Robson ME. Update on genetic testing in gynecologic cancer. *J Clin Oncol* 2019;37:2501-2509.
- 3 Marabelli M, Cheng SC, Parmigiani G. Penetrance of ATM gene mutations in breast cancer: a meta-analysis of different measures of risk. *Genet Epidemiol* 2016;40:425-431.
- 4 van Os NJ, Roeleveld N, Weemaes CM, et al. Health risks for ataxia-telangiectasia mutated heterozygotes: a systematic review, meta-analysis and evidence-based guideline. *Clin Genet* 2016;90:105-117.
- 5 Breast Cancer Association Consortium. Pathology of tumors associated with pathogenic germline variants in 9 breast cancer susceptibility genes. *JAMA Oncol* 2022;8:e216744.
- 6 Lowry KP, Geuzinge HA, Stout NK, et al. Breast cancer screening strategies for women with ATM, CHEK2, and PALB2 pathogenic variants: A comparative modeling analysis. *JAMA Oncol* 2022;8:587-596.
- 7 Breast Cancer Association Consortium. Breast cancer risk genes – Association analysis in more than 113,000 women. *N Engl J Med* 2021;384:428-439.
- 8 Hu C, Polley EC, Yadav S, et al. The contribution of germline predisposition gene mutations to clinical subtypes of invasive breast cancer from a clinical genetic testing cohort. *J Natl Cancer Inst* 2020;112:1231-1241.
- 9 Shimelis H, LaDuca H, Hu C, et al. Triple-negative breast cancer risk genes identified by multigene hereditary cancer panel testing. *J Natl Cancer Inst* 2018;110:855-862.
- 10 Lilyquist J, LaDuca H, Polley E, et al. Frequency of mutations in a large series of clinically ascertained ovarian cancer cases tested on multi-gene panels compared to reference controls. *Gynecol Oncol* 2017;147:375-380.
- 11 Kurian A, Hughes E, Handorf E, et al. Breast and ovarian cancer penetrance estimates derived from germline multiple-gene sequencing results in women. *Precis Oncol* 2017;1:1-12.
- 12 Norquist BM, Harrell MI, Brady MF, et al. Inherited mutations in women with ovarian carcinoma. *JAMA Oncol* 2016;2:482-490.
- 13 Grant RC, Selander I, Connor AA, et al. Prevalence of germline mutations in cancer predisposition genes in patients with pancreatic cancer. *Gastroenterology* 2015;148:556-564.
- 14 Holter S, Borgida A, Dodd A, et al. Germline BRCA mutations in a large clinic-based cohort of patients with pancreatic adenocarcinoma. *J Clin Oncol* 2015;33:3124-3129.
- 15 Hu C, Hart SN, Polley EC, et al. Association between inherited germline mutations in cancer predisposition genes and risk of pancreatic cancer. *JAMA* 2018;319:2401-2409.
- 16 Jones S, Hruban RH, Kamiyama M, et al. Exomic sequencing identifies PALB2 as a pancreatic cancer susceptibility gene. *Science* 2009;324:217.
- 17 Lowery MA, Wong W, Jordan EJ, et al. Prospective evaluation of germline alterations in patients with exocrine pancreatic neoplasms. *J Natl Cancer Inst* 2018;110:1067-1074.
- 18 Rainone M, Singh I, Salo-Mullen EE, et al. An emerging paradigm for germline testing in pancreatic ductal adenocarcinoma and immediate implications for clinical practice: a review. *JAMA Oncol* 2020;6:764-771.
- 19 Roberts NJ, Jiao Y, Yu J, et al. ATM mutations in patients with hereditary pancreatic cancer. *Cancer Discov* 2012;2:41-46.
- 20 Salo-Mullen EE, O'Reilly EM, Kelsen DP, et al. Identification of germline genetic mutations in patients with pancreatic cancer. *Cancer* 2015;121:4382-4388.
- 21 Shindo K, Yu J, Suenaga M, et al. Deleterious germline mutations in patients with apparently sporadic pancreatic adenocarcinoma. *J Clin Oncol* 2017;35:3382-3390.
- 22 Yang X, Leslie G, Doroszk A, et al. Cancer risks associated with germline PALB2 pathogenic variants: an international study of 524 families. *J Clin Oncol* 2020;38:674-685.
- 23 Hsu FC, Roberts NJ, Childs E, et al. Risk of pancreatic cancer among individuals with pathogenic variants in the ATM gene. *JAMA Oncol* 2021;7:1664-1668.
- 24 Karlsson Q, Brook MN, Dadaev S, et al. Rare germline variants in ATM predispose to prostate cancer: a PRACTICAL consortium study. *Eur Urol Oncol* 2021;4:570-579.
- 25 Chen S, Parmigiani G. Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol* 2007;25:1329-1333.
- 26 Ford D, Easton DF, Bishop DT, et al. Risks of cancer in BRCA1-mutation carriers. Breast cancer linkage consortium. *Lancet* 1994;343:692-695.
- 27 King MC, Marks JH, Mandell JB, New York Breast Cancer Study G. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* 2003;302:643-646.
- 28 Mavaddat N, Peock S, Frost D, et al. Cancer risks for BRCA1 and BRCA2 mutation carriers: results from prospective analysis of embrace. *J Natl Cancer Inst* 2013;105:812-822.
- 29 van den Broek AJ, van 't Veer LJ, Hoening MJ, et al. Impact of age at primary breast cancer on contralateral breast cancer risk in BRCA1/2 mutation carriers. *J Clin Oncol* 2016;34:409-418.
- 30 Tai YC, Domchek S, Parmigiani G, Chen S. Breast cancer risk among male BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst* 2007;99:1811-1814.
- 31 Li S, Silvestri V, Leslie G, et al. Cancer risks associated with BRCA1 and BRCA2 pathogenic variants. *J Clin Oncol* 2022;40:1529-1541.
- 32 Evans D, Susnerwala I, Dawson J, et al. Risk of breast cancer in male BRCA2 carriers. *J Med Genet* 2010;47:710-711.
- 33 Chen J, Bae E, Zhang L, et al. Penetrance of breast and ovarian cancer in women who carry a BRCA1/2 mutation and do not use risk-reducing salpingo-oophorectomy: an updated meta-analysis. *JNCI Cancer Spectr* 2020;4:pkaa029.
- 34 Lecarpentier J, Silvestri V, Kuchenbaecker KB, et al. Prediction of breast and prostate cancer risks in male BRCA1 and BRCA2 mutation carriers using polygenic risk scores. *J Clin Oncol* 2017;35:2240-2250.
- 35 Nyberg T, Frost D, Barrowdale D, et al. Prostate cancer risks for male BRCA1 and BRCA2 mutation carriers: a prospective cohort study. *Eur Urol* 2020;77:24-35.
- 36 Hansford S, Kaurah P, Li-Chang H, et al. Hereditary diffuse gastric cancer syndrome: CDH1 mutations and beyond. *JAMA Oncol* 2015;1:23-32.
- 37 Kaurah P, MacMillan A, Boyd N, et al. Founder and recurrent CDH1 mutations in families with hereditary diffuse gastric cancer. *JAMA* 2007;297:2360-2372.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**GENE-A**  
**9 OF 10**



### REFERENCES FOR TABLES

- 38 Xicola RM, Li S, Rodriguez N, et al. Clinical features and cancer risk in families with pathogenic CDH1 variants irrespective of clinical criteria. *J Med Genet* 2019;56:838-843.
- 39 Weber-Lasselle N, Hauke J, Ramser J, et al. BRIP1 loss-of-function mutations confer high risk for familial ovarian cancer, but not familial breast cancer. *Breast Cancer Res* 2018;20:7.
- 40 Bishop DT, Demenais F, Goldstein AM, et al. Geographical variation in the penetrance of CDKN2A 36 mutations for melanoma. *J Natl Cancer Inst* 2002;94:894-903.
- 41 Begg CB, Orlow I, Hummer AJ, et al. Lifetime risk of melanoma in CDKN2A mutation carriers in a population-based sample. *J Natl Cancer Inst* 2005;97:1507-1515.
- 42 Cybulski C, Wokolorczyk D, Jakubowska A, et al. Risk of breast cancer in women with a CHEK2 mutation with and without a family history of breast cancer. *J Clin Oncol* 2011;29:3747-3752.
- 43 Weischer M, Bojesen SE, Ellervik C, et al. CHEK2\*1100delc genotyping for clinical assessment of breast cancer risk: meta-analyses of 26,000 patient cases and 27,000 controls. *J Clin Oncol* 2008;26:542-548.
- 44 Hu C, Hart S, Gnanaolivu R, et al. A population-based study of genes previously implicated in breast cancer. *N Engl J Med* 2021;384:440-451.
- 45 Goldberg M, Bell K, Aronson M, et al. Association between the Lynch syndrome gene MSH2 and breast cancer susceptibility in a Canadian familial cancer registry. *J Med Genet* 2017;54:742-746.
- 46 Harkness EF, Barrow E, Newton K, et al. Lynch syndrome caused by mlh1 mutations is associated with an increased risk of breast cancer: a cohort study. *J Med Genet* 2015;52:553-556.
- 47 Bonadona V, Bonaiti B, Olschwang S, et al. Cancer risks associated with germline mutations in MLH1, MSH2, and MSH6 genes in Lynch syndrome. *JAMA* 2011;305:2304-2310.
- 48 Engel C, Loeffler M, Steinke V, et al. Risks of less common cancers in proven mutation carriers with lynch syndrome. *J Clin Oncol* 2012;30:4409-4415.
- 49 Bonadona V, Bonaiti B, Olschwang S, et al. Cancer risks associated with germline mutations in MLH1, MSH2, and MSH6 genes in Lynch syndrome. *JAMA* 2011;305:2304-2310.
- 50 Moller P, Seppala TT, Bernstein I, et al. Cancer risk and survival in path\_MMR carriers by gene and gender up to 75 years of age: a report from the Prospective Lynch Syndrome Database. *Gut* 2018;67:1306-1316.
- 51 Dominguez-Valentin M, Sampson JR, Seppala TT, et al. Cancer risks by gene, age, and gender in 6350 carriers of pathogenic mismatch repair variants: findings from the prospective Lynch syndrome database. *Genet Med* 2020;22:15-25.
- 52 Uusitalo E, Rantanen M, Kallionpaa RA, et al. Distinctive cancer associations in patients with neurofibromatosis type 1. *J Clin Oncol* 2016;34:1978-1986.
- 53 Seminog OO, Goldacre MJ. Age-specific risk of breast cancer in women with neurofibromatosis type 1. *Br J Cancer* 2015;112:1546-1548.
- 54 Antoniou AC, Casadei S, Heikkinen T, et al. Breast-cancer risk in families with mutations in PALB2. *N Engl J Med* 2014;371:497-506.
- 55 Pilarski R. Cowden syndrome: A critical review of the clinical literature. *J Genet Couns* 2009;18:13-27.
- 56 Pilarski R, Burt R, Kohlman W, et al. Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria. *J Natl Cancer Inst* 2013;105:1607-1616.
- 57 Buben V, Bonnet F, Brouste V, et al. High cumulative risks of cancer in patients with PTEN hamartoma tumour syndrome. *J Med Genet* 2013;50:255-263.
- 58 Tan MH, Mester JL, Ngeow J, et al. Lifetime cancer risks in individuals with germline PTEN mutations. *Clin Cancer Res* 2012;18:400-407.
- 59 Walsh S, Carter M, Tubridy N, McDermott EW. Lhermitte-Duclos and Cowden diseases: breast cancer as an unusual initial presentation of these overlapping conditions. *BMJ Case Rep* 2011;2011:bcr0820114730.
- 60 Schrager CA, Schneider D, Gruener AC, et al. Clinical and pathological features of breast disease in Cowden's syndrome: an underrecognized syndrome with an increased risk of breast cancer. *Hum Pathol* 1998;29:47-53.
- 61 Lukomska A, Menkiszak J, Gronwald J, et al. Recurrent mutations in *BRCA1*, *BRCA2*, *RAD51C*, *PALB2* and *CHEK2* in Polish patients with ovarian cancer. *Cancers* 2021;13:849.
- 62 Song H, Dicks E, Tyrer J, et al. Population-based targeted sequencing of 54 candidate genes identifies PALB2 as a susceptibility gene for high-grade serous ovarian cancer. *J Med Genet* 2021;58:305-313.
- 63 Song H, Dicks E, Ramus SJ, et al. Contribution of germline mutations in the RAD51B, RAD51C, and RAD51D genes to ovarian cancer in the population. *J Clin Oncol* 2015;33:2901-2907.
- 64 Yang X, Song H, Leslie G, et al. Ovarian and breast cancer risks associated with pathogenic variants in RAD51C and RAD51D. *J Natl Cancer Inst* 2020;112:1242-1250.
- 65 Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res* 2006;12:3209-3215.
- 66 Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial 60Peutz-Jeghers syndrome. *Gastroenterology* 2000;119:1447-1453.
- 67 Mai PL, Best AF, Peters JA, et al. Risks of first and subsequent cancers among TP53 mutation carriers in the national cancer institute Li-Fraumeni syndrome cohort. *Cancer* 2016;122:3673-3681.
- 68 Packwood K, Martland G, Sommerlad M, et al. Breast cancer in patients with germline TP53 pathogenic variants have typical tumour characteristics: the Cohort study of TP53 carrier early onset breast cancer (COPE study). *J Pathol Clin Res* 2019;5:189-198.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 1.2023

## Gene Summary: Risks and Management

### AUTOSOMAL RECESSIVE RISK IN CANCER GENES – MULTI-GENE PANEL TESTING

<b>GENE and CONDITION</b>	<b>DESCRIPTION</b>
<b>ATM –</b> <a href="#">Ataxia-Telangiectasia (AT)</a>	AT is characterized by progressive cerebellar ataxia, telangiectasias, immune defects, and a predisposition to malignancy. Cells of individuals with AT are abnormally sensitive to ionizing radiation and resistant to inhibition of DNA synthesis by ionizing radiation.
<b>BRCA1 –</b> <a href="#">Fanconi anemia complementation group S (FANCS)</a>	There are rare reports of compound heterozygous or homozygous <i>BRCA1</i> P/LP variants causing FANCS. FANCS is characterized by developmental delay apparent from infancy, short stature, microcephaly, and coarse dysmorphic features. It is associated with defective DNA repair and increased chromosomal breakage.
<b>BRCA2 –</b> <a href="#">Fanconi anemia complementation group D1</a>	Fanconi anemia is characterized by developmental abnormalities in major organ systems, early-onset bone marrow failure, and a high predisposition to cancer. Bone marrow failure with pancytopenia often presents in the first decade of life. Biallelic pathogenic variants in <i>BRCA2</i> are associated with early-onset acute leukemia and solid tumors with a cumulative probability of any malignancy of 97% by age 6 years.
<b>BRIP1 –</b> <a href="#">Fanconi anemia complementation group J (FANCJ)</a>	Fanconi anemia is characterized by developmental abnormalities in major organ systems, early-onset bone marrow failure, and a high predisposition to cancer. Bone marrow failure with pancytopenia often presents in the first decade of life.
<b>MLH1, MSH2, MSH6, PMS2, EPCAM –</b> <a href="#">CMMRD</a>	CMMRD is a childhood cancer predisposition syndrome characterized by hematologic malignancies, brain/central nervous system tumors, colorectal tumors and multiple intestinal polyps, and other malignancies including embryonic tumors and rhabdomyosarcoma.
<b>PALB2 –</b> <a href="#">Fanconi anemia complementation group N (FANCN)</a>	Fanconi anemia is characterized by developmental abnormalities in major organ systems, early-onset bone marrow failure, and an increased lifetime risk of cancer. Bone marrow failure with pancytopenia often presents in the first decade of life. Pathogenic variants in <i>PALB2</i> are associated with solid tumors, such as medulloblastomas and Wilms tumors.
<b>RAD51C –</b> <a href="#">Fanconi anemia complementation group O</a>	Fanconi anemia is characterized by developmental abnormalities in major organ systems, early-onset bone marrow failure, and a high predisposition to cancer. Bone marrow failure with pancytopenia often presents in the first decade of life.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### BRCA PATHOGENIC/LIKELY PATHOGENIC VARIANT-POSITIVE MANAGEMENT

#### GENERAL

- Education regarding signs and symptoms of cancer(s), especially those associated with *BRCA* gene P/LP variants.

#### BREAST CANCER

- Female
  - ▶ Breast awareness<sup>a</sup> starting at age 18 years.
  - ▶ Clinical breast exam, every 6–12 months,<sup>b</sup> starting at age 25 years.
  - ▶ Breast screening<sup>c,d</sup>
    - ◊ Age 25–29 years, annual breast MRI<sup>e</sup> screening with contrast<sup>f</sup> (or mammogram, only if MRI is unavailable) or individualized based on family history if a breast cancer diagnosis before age 30 is present.
    - ◊ Age 30–75 years, annual mammogram and breast MRI<sup>e</sup> screening with contrast.
    - ◊ Age >75 years, management should be considered on an individual basis.
    - ◊ For individuals with a *BRCA* P/LP variant who are treated for breast cancer and have not had a bilateral mastectomy, screening with annual mammogram and breast MRI should continue as described above.
  - ▶ Discuss option of RRM
    - ◊ Counseling should include a discussion regarding degree of protection, reconstruction options, and risks. In addition, the family history and residual breast cancer risk with age and life expectancy should be considered during counseling.
  - ▶ Address psychosocial and quality-of-life aspects of undergoing RRM.
- Consider risk reduction agents as options for breast cancer, including discussion of risks and benefits ([See Discussion](#) for details). ([See NCCN Guidelines for Breast Cancer Risk Reduction](#)).
- Male
  - ▶ Breast self-exam training and education starting at age 35 years.
  - ▶ Clinical breast exam, every 12 months, starting at age 35 years.
  - ▶ Consider annual mammogram screening in men with gynecomastia starting at age 50 or 10 years before the earliest known male breast cancer in the family (whichever comes first).<sup>g</sup>

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

<sup>a</sup> Females should be familiar with their breasts and promptly report changes to their health care provider. Periodic, consistent breast self exam (BSE) may facilitate breast self awareness. Premenopausal individuals may find BSE most informative when performed at the end of menses.

<sup>b</sup> Randomized trials comparing clinical breast exam versus no screening have not been performed. Rationale for recommending clinical breast exam every 6–12 mo is the concern for interval breast cancers.

<sup>c</sup> The appropriateness of imaging modalities and scheduling is still under study. Lowry KP, Lee JM, Kong CY, et al. Cancer 2012;118:2021-2030.

<sup>d</sup> Lehman CD, et al. J Natl Cancer Inst 2016;108.

<sup>e</sup> The criteria for high-quality breast MRI include a dedicated breast coil, the ability to perform biopsy under MRI guidance, radiologists experienced in breast MRI, and regional availability. Breast MRI is preferably performed on days 7–15 of a menstrual cycle for premenopausal patients. [FDA Drug Safety Communication](#): FDA identifies no harmful effects to date with brain retention of gadolinium-based contrast agents for MRIs; review to continue.

<sup>f</sup> Breast MRI is preferred due to the theoretical risk of radiation exposure in P/LP variant carriers.

<sup>g</sup> There are only limited data to support screening for male breast cancer. Gao Y, et al Radiology 2019;293:282-291.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



# NCCN Guidelines Version 1.2023

## BRCA-Pathogenic/Likely Pathogenic Variant - Positive Management

### BRCA PATHOGENIC/LIKELY PATHOGENIC VARIANT-POSITIVE MANAGEMENT

#### OVARIAN/UTERINE CANCER

- Recommend RRSO,<sup>h</sup> typically between 35 and 40 years, and upon completion of childbearing. Because ovarian cancer onset in patients with *BRCA2* P/LP variants is an average of 8–10 years later than in patients with *BRCA1* P/LP variants, it is reasonable to delay RRSO for management of ovarian cancer risk until age 40–45 years in patients with *BRCA2* P/LP variants unless age at diagnosis in the family warrants earlier age for consideration of prophylactic surgery. See Risk-Reducing Salpingo-Oophorectomy (RRSO) Protocol in [NCCN Guidelines for Ovarian Cancer](#) - Principles of Surgery.
  - ▶ Counseling includes a discussion of reproductive desires, extent of cancer risk, degree of protection for breast and ovarian cancer, management of menopausal symptoms, hormone replacement therapy (HRT), and related medical issues.
  - ▶ Salpingectomy alone is not the standard of care for risk reduction, although clinical trials of interval salpingectomy and delayed oophorectomy are ongoing. The concern for risk-reducing salpingectomy alone is that individuals are still at risk for developing ovarian cancer. In addition, in premenopausal individuals, oophorectomy likely reduces the risk of developing breast cancer but the magnitude is uncertain and may be gene-specific.
- Limited data suggest that there may be a slightly increased risk of serous uterine cancer among individuals with a *BRCA1* P/LP variant. The clinical significance of these findings is unclear. Further evaluation of the risk of serous uterine cancer in the *BRCA* population needs to be undertaken. The provider and patient should discuss the risks and benefits of concurrent hysterectomy at the time of RRSO for individuals with a *BRCA1* P/LP variant prior to surgery.
- Individuals who undergo hysterectomy at the time of RRSO are candidates for estrogen-alone HRT, which is associated with a decreased risk of breast cancer compared to combined estrogen and progesterone, which is required when the uterus is left in situ (Chlebowski R, et al. JAMA Oncol 2015;1:296-305). HRT recommendations should be tailored depending on each patient's personal history of breast cancer and/or breast cancer risk reduction strategies. HRT is a consideration for premenopausal patients who do not carry a diagnosis of breast cancer or have other contraindications for HRT.
- Address psychosocial and quality-of-life aspects of undergoing RRSO. Consider preoperative menopause management consultation if patient is still premenopausal at time of RRSO.
- Consider risk reduction agents as options for ovarian cancer, including discussion of risks and benefits ([See Discussion](#) for details).

<sup>h</sup> Given the high rate of occult neoplasms, special attention should be given to sampling and pathologic review of the ovaries and fallopian tubes. ([See Discussion](#) for details.) [See the College of American Pathologists, Protocol for the Examination of Specimens from Patients with Carcinoma of the Ovary.](#) [See NCCN Guidelines for Ovarian Cancer](#) for treatment of findings.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



### BRCA PATHOGENIC/LIKELY PATHOGENIC VARIANT-POSITIVE MANAGEMENT

#### **PANCREATIC CANCER**

- For pancreatic cancer screening recommendations, [see PANC-A](#).

#### **PROSTATE CANCER**

- Starting at age 40 years: ([See Guidelines for Prostate Cancer Early Detection](#))
  - Recommend prostate cancer screening for *BRCA2* carriers.
  - Consider prostate cancer screening for *BRCA1* carriers.

#### **MELANOMA**

- No specific screening guidelines exist for melanoma, but general melanoma risk management is appropriate, such as annual full-body skin examination and minimizing UV exposure.

#### **RISK TO RELATIVES**

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\)](#)

#### **REPRODUCTIVE OPTIONS**

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\)](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

### PANCREATIC CANCER SCREENING

- Emerging data have examined the efficacy of pancreatic cancer screening in select individuals at increased risk for exocrine pancreatic cancer. To date, most such studies have restricted pancreatic cancer screening to individuals with:
  1. A known P/LP germline variant in a pancreatic cancer susceptibility gene (*ATM*, *BRCA1*, *BRCA2*, *CDKN2A*, *MLH1*, *MSH2*, *MSH6*, *EPCAM*, *PALB2*, *STK11*, *TP53*; [see GENE-A](#)) and a family history of pancreatic cancer (first-degree or second-degree relative) from the same side of the family as the germline P/LP variant; or
  2. A family history of exocrine pancreatic cancer in  $\geq 2$  first-degree relatives from the same side of the family, even in the absence of a known P/LP germline variant (many centers would enroll individuals with one affected first-degree relative and one second-degree relative); or
  3. A family history of exocrine pancreatic cancer in  $\geq 3$  first- and/or second-degree relatives from the same side of the family, even in the absence of a known P/LP germline variant.
- These studies have typically started screening with contrast-enhanced MRI/magnetic resonance cholangiopancreatography (MRCP) and/or endoscopic ultrasound (EUS) in such high-risk individuals.
- Potential benefits of pancreatic cancer screening include a suggestion of downstaging, compared to historical data, in that 75%–90% of screen-detected pancreatic cancer has been surgically resectable at diagnosis (which is markedly higher than historical rates of resectability with pancreatic cancers detected due to symptoms).<sup>a,b</sup> There has also been a suggestion of improved mortality compared to historical data, with one study demonstrating an 85% 3-year overall survival rate after screen-detected pancreatic cancer in high-risk individuals,<sup>a</sup> and another study demonstrating a 24% 5-year overall survival rate following screen-detected pancreatic cancer in individuals with germline c.67G>C *CDKN2A* variants.<sup>b</sup> One study<sup>a</sup> also demonstrated 100% overall survival among 10 individuals with screen-detected precursor lesions (intraductal papillary mucinous neoplasms [IPMN] with high-grade dysplasia and/or high-grade pancreatic intraepithelial neoplasia [PanIN]) treated with surgical resection.
- Although evidence for downstaging has emerged in recent studies, longer-term studies are needed to determine if this downstaging translates to improved survival. Evidence from patients with sporadic forms of pancreatic ductal adenocarcinoma suggest that long-term survival is common for patients who present with stage I disease. Since many patients who undergo pancreatic surveillance have pancreatic abnormalities, mostly subcentimeter pancreatic cysts (42% of high-risk individuals in one study<sup>c</sup> had at least one pancreatic mass/cyst and/or duct abnormality), there is potential for unnecessary interventions (such as fine-needle aspiration [FNA] and in some cases surgery). Although there is much more experience with evaluating and managing pancreatic cysts and other pancreatic imaging abnormalities, determination of the overall risk/benefits of pancreatic surveillance requires further study. Results of surveillance of high-risk individuals performed in tertiary care/high-volume centers under clinical trial settings may not be the same as those performed in routine clinical practice. Data are beginning to better define which screen-detected lesions in high-risk individuals should be considered to be at particularly high risk for neoplastic progression (eg, those with a solid pancreatic mass, those with pancreatic duct abnormalities, those with growing pancreatic cysts<sup>a</sup>), but further data are needed to better define the threshold for surgical intervention in high-risk individuals undergoing pancreatic cancer screening.

<sup>a</sup> Canto MI, et al. *Gastroenterology* 2018;155:740-751.

<sup>b</sup> Vasen H, et al. *J Clin Oncol* 2016;34:2010-2019.

<sup>c</sup> Canto MI, et al. *Gastroenterology* 2012;142:796-804.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



### PANCREATIC CANCER SCREENING

- For individuals considering pancreatic cancer screening, the panel recommends that screening be performed in experienced high-volume centers. The panel recommends that such screening only take place after an in-depth discussion about the potential limitations to screening, including cost, the high incidence of benign or indeterminate pancreatic abnormalities, and uncertainties about the potential benefits of pancreatic cancer screening.
- Consider screening using annual contrast-enhanced MRI/MRCP and/or EUS, with consideration of shorter screening intervals, based on clinical judgment, for individuals found to have potentially concerning abnormalities on screening. The panel emphasizes that most small cystic lesions found on screening will not warrant biopsy, surgical resection, or any other intervention.
- For all individuals with P/LP germline variants in *STK11*
  - ▶ Consider pancreatic cancer screening beginning at age 30–35 years (or 10 years younger than the earliest exocrine pancreatic cancer diagnosis in the family, whichever is earlier).
- For all individuals with P/LP germline variants in *CDKN2A*
  - ▶ Consider pancreatic cancer screening beginning at age 40 years (or 10 years younger than the earliest exocrine pancreatic cancer diagnosis in the family, whichever is earlier).
- For individuals with P/LP germline variants in one of the other pancreatic cancer susceptibility genes (*ATM*, *BRCA1*, *BRCA2*, *MLH1*, *MSH2*, *MSH6*, *EPCAM*, *PALB2*, *TP53*), [see GENE-A](#).
  - ▶ Consider pancreatic cancer screening beginning at age 50 years (or 10 years younger than the earliest exocrine pancreatic cancer diagnosis in the family, whichever is earlier) for individuals with exocrine pancreatic cancer in ≥1 first- or second-degree relatives from the same side of (or presumed to be from the same side of) the family as the identified P/LP germline variant.<sup>d</sup>
  - ▶ The panel does not currently recommend pancreatic cancer screening for carriers of P/LP variants in genes other than *STK11* and *CDKN2A* in the absence of a close family history of exocrine pancreatic cancer.

### Hereditary Pancreatitis Genes

- For individuals with P/LP variants in *PRSS1* or other hereditary pancreatitis genes AND a clinical phenotype consistent with hereditary pancreatitis<sup>e</sup>
  - ▶ Consider pancreatic cancer screening 20 years after onset of pancreatitis, or at age 40 years, whichever is earlier.

<sup>d</sup> Abe T, et al. J Clin Oncol 2019;37:1070-1080.

<sup>e</sup> The panel recognizes that patients with hereditary pancreatitis (sometimes caused by pathogenic germline variants in *PRSS1*, *SPINK1*, and other genes) have increased lifetime risks of pancreatic cancer. The clinical significance of pathogenic germline variants in these genes is unclear, when such variants are identified in individuals lacking a clinical history of pancreatitis. As such, the panel recommends germline testing for *PRSS1*, *SPINK1*, and other pancreatitis genes in individuals with a personal and/or family history of exocrine pancreatic cancer only if there is a personal and/or family history suggestive of hereditary pancreatitis.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### LI-FRAUMENI SYNDROME MANAGEMENT IN ADULTS

#### **BREAST CANCER (female)**

- Breast awareness<sup>a</sup> starting at age 18 y.
- Clinical breast exam, every 6–12 mo, starting at age 20 y.<sup>b</sup>
- Breast screening
  - ▶ Age 20–29<sup>b</sup> y, annual breast MRI<sup>c</sup> screening with contrast.<sup>d</sup>
  - ▶ Age 30–75 y, annual breast MRI<sup>c</sup> screening with contrast and mammogram.
  - ▶ Age >75 y, management should be considered on an individual basis.
  - ▶ For individuals with a *TP53* P/LP variant who are treated for breast cancer, and who have not had a bilateral mastectomy, screening with annual breast MRI and mammogram should continue as described above.
- Discuss option of RRM
  - ▶ Counseling should include a discussion regarding degree of protection, reconstruction options, and risks. In addition, the family history and residual breast cancer risk with age and life expectancy should be considered during counseling.
- Address psychosocial and quality-of-life aspects of undergoing RRM.

#### **OTHER CANCER RISKS**

- Comprehensive physical exam including neurologic examination with high index of suspicion for rare cancers and second malignancies in cancer survivors every 6–12 mo.
- Colonoscopy and upper endoscopy every 2–5 y starting at 25 y or 5 y before the earliest known colon or gastric cancer in the family, respectively.
- Annual dermatologic examination starting at 18 y.
- Annual whole body MRI<sup>e,f,g</sup> (category 2B).
- Annual brain MRI (category 2B) may be performed as part of the whole body MRI or as a separate exam.

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

<sup>a</sup> Females should be familiar with their breasts and promptly report changes to their health care provider. Periodic, consistent BSE may facilitate breast self awareness. Premenopausal individuals may find BSE most informative when performed at the end of menses.

<sup>b</sup> Or at the age of the earliest diagnosed breast cancer in the family, if younger than age 20 y.

<sup>c</sup> High-quality breast MRI limitations include having: a need for a dedicated breast coil, the ability to perform biopsy under MRI guidance by experienced radiologists in breast MRI, and regional availability. Breast MRI is preferably performed on days 7–15 of a menstrual cycle for premenopausal individuals.

<sup>d</sup> Or mammogram with consideration of tomosynthesis, if MRI is unavailable. Breast MRI is preferred because of concerns regarding the risk of radiation exposure in P/LP variant carriers.

<sup>e</sup> Whole body MRI is not uniformly available. If whole body MRI is not available, then individuals with LFS are encouraged to participate in clinical trials or consider alternate comprehensive imaging methods. Other components of screening are being evaluated in protocols, including biochemical screening and regular blood screening for hematologic malignancies. [FDA Drug Safety Communication](#): FDA identifies no harmful effects to date with brain retention of gadolinium-based contrast agents for MRIs; review to continue.

<sup>f</sup> Ballinger M, Best A, Mai P, et al. Baseline surveillance in Li-Fraumeni syndrome using whole-body magnetic resonance imaging: a meta-analysis. *JAMA Oncol* 2017;3:1634-1639.

<sup>g</sup> Screening through whole body MRI has been broadly demonstrated to be feasible and of potential utility in the early detection of cancer among classic LFS families, though it also results in the detection of false-positive findings and possible cancer overdiagnosis. Furthermore, screening utility has not been evaluated among those with a germline *TP53* P/LP variant without a classic family history of LFS, who are increasingly identified through multi-gene panel tests.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**LIFR-A**  
**1 OF 2**





### LI-FRAUMENI SYNDROME MANAGEMENT IN ADULTS

#### **OTHER ASPECTS OF MANAGING LFS**

- This screening and management of LFS is complex, and LFS is rare; it is preferred that individuals with LFS be followed at centers with expertise in the management of this syndrome.
- Because of the remarkable risk of additional primary neoplasms, screening may be considered for cancer survivors with LFS and a good prognosis from their prior tumor(s).
- Address limitations of screening for many cancers associated with LFS.
- Pediatricians should be apprised of the risk of childhood cancers in affected families and review screening recommendations for children with LFS.<sup>h</sup>
- Therapeutic RT for cancer should be avoided when possible; diagnostic radiation should be minimized to the extent feasible without sacrificing accuracy.
- Provide additional surveillance based on family history of cancer.
- Provide education regarding signs and symptoms of cancer.
- Address psychosocial and quality-of-life aspects of the complex management of LFS.
- There is controversy over how to manage cancer risk in incidental *TP53* carriers who do not meet classic LFS criteria; some data suggest lower cancer risks in *TP53* P/LP carriers who do not have a family history consistent with LFS.

#### **REPRODUCTIVE OPTIONS**

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\)](#).

#### **RISK TO RELATIVES**

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\)](#).

#### **TESTING CONSIDERATIONS**

- Somatic *TP53* variants frequently confound germline testing results. Late post-zygotic aberrant clonal expansions containing a pathogenic *TP53* variant, limited to the hematologic compartment or to a tumor, may be detected in the blood or saliva through germline testing, particularly using NGS technology. The phenomenon of aberrant clonal expansion is well described and is most often due to CHIP, which can be demonstrated in healthy populations at increasing frequency with increasing age.<sup>i</sup> This finding has important clinical implications regarding potential application of unwarranted clinical interventions. Further, the finding of clonal hematopoiesis itself may portend adverse clinical outcomes, such as the development of hematologic neoplasia and increased non-hematologic mortality.
- Blood and/or saliva is an unsuitable source of DNA for germline testing for cases with a history of hematologic abnormalities. Careful examination of the patient's complete blood count (CBC) and peripheral blood smear may be warranted in all cases reporting the discovery of a *TP53* P/LP variant, and testing of non-lymphoid ancillary tissues may help to delineate bona fide mosaic involvement of different germ layers.<sup>j</sup>

<sup>h</sup> For additional information on the management of children with LFS, see Kratz C, et al. Clin Cancer Res 2017;23:e38-e45.

<sup>i</sup> Jaiswal S, et al. N Engl J Med 2014;371:2488-2498; Genovese G, et al. N Engl J Med 2014;371:2477-2487.

<sup>j</sup> Weitzel J, et al. Genet Med 2018;20:809-816.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Cowden Syndrome/PTEN Hamartoma Tumor Syndrome Management

### COWDEN SYNDROME/PHTS MANAGEMENT

#### **GENERAL**

- Due to the rarity of the syndrome and complexities of diagnosing and managing individuals with Cowden syndrome, referral to a specialized team or centers with expertise is recommended.
- Annual comprehensive physical exam starting at age 18 y or 5 y before the youngest age of diagnosis of a component cancer in the family (whichever comes first), with particular attention to thyroid exam.
- Education regarding the signs and symptoms of cancer.

#### **BREAST CANCER** (female)

- Breast awareness<sup>a</sup> starting at age 18 years.
- Clinical breast exam, every 6–12 months, starting at age 25 years or 5–10 years before the earliest known breast cancer in the family (whichever comes first).
- Breast screening
  - ▶ Annual mammography and breast MRI screening with contrast starting at age 35 years or 10 years before the earliest known breast cancer in the family (whichever comes first).<sup>b,c</sup>
  - ▶ Age >75 years, management should be considered on an individual basis.
  - ▶ For individuals with a *PTEN* P/LP variant who are treated for breast cancer, and have not had a bilateral mastectomy, screening with annual mammogram and breast MRI should continue as described above.
- Discuss option of RRM in individuals with P/LP variants identified. For those with clinical CS/PHTS syndrome, consideration of risk-reducing surgery should be based on family history.
  - ▶ Counseling should include a discussion regarding degree of protection, reconstruction options, and risks. In addition, the family history and residual breast cancer risk with age and life expectancy should be considered during counseling.
- Address psychosocial and quality-of-life aspects of undergoing RRM.

#### **COLORECTAL**

- Colonoscopy, starting at age 35 y unless symptomatic or if close relative with colon cancer before age 40 y, then start 5–10 y before the earliest known colon cancer in the family. Colonoscopy should be done every 5 y or more frequently if patient is symptomatic or polyps are found.

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. In this guideline, the terms male and female refer to sex assigned at birth.

<sup>a</sup> Females should be familiar with their breasts and promptly report changes to their health care provider. Periodic, consistent BSE may facilitate breast self awareness. Premenopausal individuals may find BSE most informative when performed at the end of menses.

<sup>b</sup> The appropriateness of imaging modalities and scheduling is still under study.

<sup>c</sup> High-quality breast MRI limitations include having: a need for a dedicated breast coil, the ability to perform biopsy under MRI guidance by experienced radiologists in breast MRI, and regional availability. Breast MRI is preferably performed on days 7–15 of a menstrual cycle for premenopausal females. [FDA Drug Safety Communication](#): FDA identifies no harmful effects to date with brain retention of gadolinium-based contrast agents for MRIs; review to continue.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**Continued**

**COWD-A**  
**1 OF 2**



# NCCN Guidelines Version 1.2023

## Cowden Syndrome/PTEN Hamartoma Tumor Syndrome Management

### COWDEN SYNDROME/PHTS MANAGEMENT

#### ENDOMETRIAL CANCER

- For endometrial cancer screening,<sup>d</sup> consider starting by age 35 years.
  - ▶ Encourage patient education and prompt response to symptoms (eg, abnormal bleeding). Patients are encouraged to keep a calendar in order to identify irregularities in their menstrual cycle.
  - ▶ Because endometrial cancer can often be detected early based on symptoms, individuals should be educated regarding the importance of prompt reporting and evaluation of any abnormal uterine bleeding or postmenopausal bleeding. The evaluation of these symptoms should include endometrial biopsy.
  - ▶ Endometrial cancer screening does not have proven benefit in individuals with CS/PHTS. However, endometrial biopsy is both highly sensitive and highly specific as a diagnostic procedure. Screening via endometrial biopsy every 1 to 2 years can be considered.
  - ▶ Transvaginal ultrasound to screen for endometrial cancer in postmenopausal individuals has not been shown to be sufficiently sensitive or specific as to support a positive recommendation, but may be considered at the clinician's discretion. Transvaginal ultrasound is not recommended as a screening tool in premenopausal individuals due to the wide range of endometrial stripe thickness throughout the normal menstrual cycle.
- Discuss option of hysterectomy<sup>e</sup> upon completion of childbearing and counsel regarding degree of protection, extent of cancer risk, and reproductive desires. Risk of ovarian cancer is not elevated; therefore, ovaries can be left in situ.
- Address psychosocial and quality-of-life aspects of undergoing risk-reducing hysterectomy.

#### KIDNEY

- Consider renal ultrasound starting at age 40 y, then every 1–2 y.

#### NEUROLOGIC

- Consider psychomotor assessment in children at diagnosis and brain MRI if there are symptoms.

#### SKIN

- There may be an increased risk of melanoma, and the prevalence of other skin characteristics with CS/PHTS may independently make routine dermatology evaluations of value. Annual dermatology exams are recommended.

#### THYROID

- Annual thyroid ultrasound starting at age 7 y. This may also be considered for children at 50% risk of inheriting a known P/LP variant whose parents wish to delay genetic testing until age 18 y.

#### RISK TO RELATIVES

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\).](#)

#### REPRODUCTIVE OPTIONS

- [Principles of Cancer Risk Assessment and Counseling \(EVAL-A\).](#)

<sup>d</sup> There are limited data regarding the lifetime risk of endometrial cancer in CS/PHTS. Surveillance screening and surgical intervention should be on an individual basis.

<sup>e</sup> Oophorectomy is not indicated for CS/PHTS alone.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines

NCCN Guideline	Specific Sections Included in Table of Contents	Genes and/or Syndromes Included/Mentioned in Guideline
<b>Treatment Guidelines</b>		
<a href="#">Acute Lymphocytic Leukemia (ALL)</a>	<a href="#">Familial/Genetic Alterations in ALL</a>	<i>RUNX1, ETV6, PAX5, IKZF1, TP53</i>
<a href="#">Acute Myeloid Leukemia (AML)</a>	<a href="#">Risk Stratification by Genetics in Non-APL AML</a>	<i>RUNX1, ANKRD26, CEBPA, DDX41, ETV6, GATA2, MBD4, MECOM/EVI1 complex, SAMD9/SAMD9L, TERC/TERT, ATG2B/GSKIP</i>
<a href="#">Basal Cell Skin Cancer</a>		Gorlin syndrome ( <i>PTCH1</i> ), xeroderma pigmentosa
<a href="#">Bladder Cancer/ Urothelial Cancer (including renal pelvis and ureter)</a>		Lynch syndrome (LS)
<a href="#">Breast Cancer</a>		Refers to Genetic/Familial BOP (Breast, Ovarian, Pancreatic) Guidelines
<a href="#">Colon Cancer</a>		Refers to Genetic/Familial Colorectal Cancer (CRC) Guidelines
<a href="#">Esophageal and Esophagogastric Junction (EGJ) Cancers</a>	<a href="#">Principles of Genetic Risk Assessment for Esophageal and Esophagogastric Junction (EGJ) Cancers</a>	<i>RHBDF2</i> ; Bloom syndrome (BS)/ <i>BLM, RECQL3</i> ; Fanconi anemia (FA)/ <i>FANCD1, BRCA2, PALB2</i>
<a href="#">Gastric Cancer</a>	<a href="#">Principles of Genetic Risk Assessment for Gastric Cancer</a>	<i>CDH1</i> ; Lynch syndrome (LS)/ <i>MLH1, MSH2, MSH6, PMS2, EPCAM</i> ; juvenile polyposis syndrome (JPS)/ <i>SMAD4, BMPR1A</i> ; Peutz-Jeghers syndrome (PJS)/ <i>STK11</i> ; familial adenomatous polyposis (FAP)/ <i>APC</i> ; ataxia-telangiectasia ( <i>ATM</i> ); BS ( <i>BLM, RECQL3</i> ); <i>BRCA1, BRCA2, TP53</i> ; xeroderma pigmentosum; Cowden syndrome (CS)/ <i>PTEN</i>
<a href="#">Gastrointestinal Stromal Tumors (GISTs)</a>	<a href="#">Principles of Mutation Testing</a>	<i>KIT, PDGFRA, SDHB, NF1</i>
<a href="#">Hepatobiliary Cancers</a>		Hemochromatosis, LS, <i>BRCA1/2</i>
<a href="#">Kidney Cancer</a>	<a href="#">Hereditary Renal Cell Carcinoma section</a>	Von-Hippel Lindau (VHL) syndrome; hereditary papillary renal carcinoma (HPRC)/ <i>MET</i> ; Birt-Hogg-Dube syndrome (BHDS)/ <i>FLCN</i> ; tuberous sclerosis complex (TSC)/ <i>TSC1, TSC2</i> ; hereditary leiomyomatosis and renal cell carcinoma (HLRCC)/ <i>FH</i> ; <i>BAP1</i> tumor predisposition syndrome (TPDS)/ <i>BAP1</i> ; hereditary paraganglioma/pheochromocytoma (PGL/PCC) syndrome/ <i>SDHA/SDHB/SDHC/SDHD</i>

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines

NCCN Guideline	Specific Sections Included in Table of Contents	Genes and/or Syndromes Included/Mentioned
<b>Treatment Guidelines (Continued)</b>		
<a href="#">Malignant Pleural Mesothelioma</a>	<a href="#">Principles of Pathologic Review</a>	<i>BAP1</i> tumor predisposition syndrome
<a href="#">Melanoma: Cutaneous</a>	<a href="#">Risk Factors for Development of Single or Multiple Primary Melanomas</a>	<i>CDKN2a</i> , <i>CDK4</i> , <i>MC1R</i> , <i>BRCA2</i> , <i>BAP1</i> (including uveal), <i>TERT</i> , <i>MITF</i> , <i>PTEN</i> , xeroderma pigmentosum
<a href="#">Myelodysplastic Syndromes</a>	<ul style="list-style-type: none"> <li><a href="#">Genetic Familial High-Risk Assessment: Hereditary Myeloid Malignancy Predisposition Syndromes</a></li> <li><a href="#">Gene Mutations Associated With Hereditary Myeloid Malignancies</a></li> </ul>	<i>CEBPA</i> , <i>DDX41</i> , <i>ATG2B/GSKIP</i> , Xeroderma Pigmentosum C/ <i>XPC</i> , <i>ERCC6L2</i> , <i>ANKRD26</i> , <i>ETV6</i> , <i>GATA2</i> , <i>RUNX1</i> , <i>LIG-4</i> , MIRAGE syndrome/ <i>SAMD9</i> , Ataxia-pancytopenia syndrome/ <i>SAMD9L</i> , <i>SRP72</i> , Diamond-Blackfan anemia, Fanconi anemia, Shwachman-Diamond syndrome, Short telomere syndromes, Congenital neutropenia, Myeloid neoplasms associated with Down syndrome, Constitutional mismatch repair deficiency, <i>BRCA1/BRCA2</i> , Li-Fraumeni syndrome/ <i>TP53</i> , RASopathies, Other rare DNA repair syndromes/ <i>BLM</i> , <i>MBD4</i> , <i>XPC</i>
<a href="#">Neuroendocrine and Adrenal Tumors</a>	<a href="#">Principles of Hereditary Cancer Risk Assessment and Genetic Counseling</a>	Hereditary PGL/PCC syndrome/ <i>MAX</i> , <i>SDHA</i> , <i>SDHAF2</i> , <i>SDHB</i> , <i>SDHC</i> , <i>SDHD</i> , <i>TMEM127</i> ; multiple endocrine neoplasia type 1 (MEN1); MEN type 2 (MEN2)/ <i>RET</i> ; MEN type 4 (MEN4)/ <i>CDKN1B</i> ; neurofibromatosis type 1 ( <i>NF1</i> ); tuberous sclerosis complex ( <i>TSC1</i> , <i>TSC2</i> ); VHL syndrome; Li-Fraumeni syndrome/ <i>TP53</i> ; LS ( <i>MLH1</i> , <i>EPCAM/MSH2</i> , <i>MSH6</i> , <i>PMS2</i> ); familial adenomatous polyposis (FAP)/ <i>APC</i>
<a href="#">Non-Small Cell Lung Cancer</a>	<a href="#">Principles of Molecular and Biomarker Analysis</a>	<i>EGFR</i> p.T790M
<a href="#">Ovarian Cancer</a>	<a href="#">Principles of Cancer Risks Assessment and Counseling</a>	<i>BRCA1</i> , <i>BRCA2</i>
<a href="#">Pancreatic Cancer</a>		<i>ATM</i> , <i>BRCA1</i> , <i>BRCA2</i> , <i>CDKN2A</i> , <i>MLH1</i> , <i>MSH2</i> , <i>MSH6</i> , <i>PALB2</i> , <i>PMS2</i> , <i>STK11</i> , and <i>TP53</i>
<a href="#">Pediatric Acute Lymphoblastic Leukemia</a>	<a href="#">Genetic Risk Groups for B-ALL</a>	Li-Fraumeni Syndrome/ <i>TP53</i> association with hypodiploid ALL
<a href="#">Prostate Cancer</a>	<a href="#">Principles of Genetics and Molecular/Biomarker Analysis</a>	<i>BRCA1</i> , <i>BRCA2</i> , <i>ATM</i> , <i>PALB2</i> , <i>CHEK2</i> , LS/ <i>MLH1</i> , <i>MSH2</i> , <i>MSH6</i> , <i>PMS2</i>
<a href="#">Rectal Cancer</a>		LS, FAP, attenuated FAP (AFAP)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 1.2023

## Breast, Ovarian, and/or Pancreatic Cancer

### Genetic Assessment

#### Summary of Genes and/or Syndromes Included/Mentioned in Other NCCN Guidelines

NCCN Guideline	Specific Sections Included in Table of Contents	Genes and/or Syndromes Included/Mentioned
<b>Treatment Guidelines (Continued)</b>		
<a href="#">Soft Tissue Sarcoma</a>	<a href="#">Principles of Cancer Risks Assessment and Counseling</a>	Neurofibromatosis/ <i>NF1</i> ; Li-Fraumeni syndrome (LFS)/ <i>TP53</i> ; LS; Familial Adenomatous Polyposis (FAP)
<a href="#">Thyroid Carcinoma</a>	<ul style="list-style-type: none"><li>• <a href="#">Germline Mutation of <i>RET</i> Proto-oncogene</a></li><li>• <a href="#">Principles of Cancer Risk Assessment and Counseling</a></li></ul>	MEN2/ <i>RET</i>
<a href="#">Uterine Neoplasms</a>	<ul style="list-style-type: none"><li>• <a href="#">Principles of Pathology and Molecular Analysis</a> (Endometrial Cancer)</li><li>• <a href="#">Principles of Pathology and Molecular Analysis</a> (Uterine Sarcomas)</li></ul>	LS, <i>SMARCA4</i>
<a href="#">Wilms Tumor (Nephroblastoma)</a>	<ul style="list-style-type: none"><li>• <a href="#">Syndromes and Congenital Anomalies Associated with Wilms Tumor</a></li></ul>	Denys-Drash Syndrome, Frasier Syndrome, Beckwith-Wiedemann Syndrome
<b>Detection, Prevention, and Risk Reduction Guidelines</b>		
<a href="#">Colorectal Cancer Screening</a>	<ul style="list-style-type: none"><li>• <a href="#">Risk Assessment for Colorectal Cancer</a></li><li>• <a href="#">Increased Risk Based on Personal History of Childhood, Adolescent, and Young Adult Cancer</a></li><li>• <a href="#">Increased Risk Based on Positive Family History</a></li></ul>	LS
<a href="#">Prostate Cancer Early Detection</a>	<a href="#">Baseline Evaluation, Risk Assessment, and Early Detection Evaluation</a>	<i>BRCA2</i> , <i>HOXB13</i> , (and indicate that germline mutation timing and testing less clear for <i>BRCA1</i> , <i>ATM</i> and LS genes)
<b>Supportive Care Guidelines</b>		
<a href="#">Survivorship</a>	<a href="#">Principles of Cancer Risk Assessment and Counseling</a>	Refers to some of the other guidelines containing inherited cancer content
<b>Special Populations</b>		
<a href="#">Adolescent and Young Adult (AYA)</a>	<a href="#">Comprehensive Initial Assessment</a>	

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





### ABBREVIATIONS

<b>AFAB</b>	<b>assigned female at birth</b>	<b>GI</b>	<b>gastrointestinal</b>	<b>RRM</b>	<b>risk-reducing mastectomy</b>
<b>AMAB</b>	<b>assigned male at birth</b>	<b>GIST</b>	<b>gastrointestinal stromal tumor</b>	<b>RRSO</b>	<b>risk-reducing salpingo-oophorectomy</b>
<b>AT</b>	<b>ataxia-telangiectasia</b>			<b>RT</b>	<b>radiation therapy</b>
<b>BRRS</b>	<b>Bannayan-Riley-Ruvalcaba syndrome</b>	<b>HDGC</b>	<b>hereditary diffuse gastric cancer</b>	<b>SEER</b>	<b>Surveillance, Epidemiology, and End Results</b>
<b>BSE</b>	<b>breast self-examination</b>	<b>HRT</b>	<b>hormone replacement therapy</b>	<b>SNP</b>	<b>single nucleotide polymorphism</b>
<b>CBC</b>	<b>complete blood count</b>	<b>HER2</b>	<b>human epidermal growth factor receptor 2</b>		
<b>CHIP</b>	<b>clonal hematopoiesis of indeterminant potential</b>	<b>LS</b>	<b>Lynch syndrome</b>	<b>UV</b>	<b>ultraviolet</b>
<b>CLIA</b>	<b>Clinical Laboratory Improvement Amendments</b>	<b>LFS</b>	<b>Li-Fraumeni syndrome</b>	<b>TSC</b>	<b>tuberous sclerosis complex</b>
<b>CMMRD</b>	<b>constitutional mismatch repair deficiency</b>	<b>MRCP</b>	<b>magnetic resonance cholangiopancreatography</b>	<b>VHL</b>	<b>Von-Hippel Lindau</b>
<b>CNS</b>	<b>central nervous system</b>	<b>MRI</b>	<b>magnetic resonance imaging</b>	<b>VUS</b>	<b>variant of uncertain significance</b>
<b>CPS + EG</b>	<b>clinical-pathologic stage + estrogen receptor status and histologic grade</b>	<b>NF</b>	<b>neurofibromatosis</b>		
<b>CS</b>	<b>Cowden syndrome</b>	<b>NGS</b>	<b>next-generation sequencing</b>		
<b>ctDNA</b>	<b>circulating tumor DNA</b>	<b>PARP</b>	<b>poly-ADP ribose polymerase</b>		
<b>DTC</b>	<b>direct to consumer</b>	<b>P/LP</b>	<b>pathogenic/likely pathogenic</b>		
<b>EUS</b>	<b>endoscopic ultrasound</b>	<b>PHTS</b>	<b>PTEN hamartoma tumor syndrome</b>		
<b>FA</b>	<b>Fanconi anemia</b>	<b>PJS</b>	<b>Peutz-Jeghers syndrome</b>		
<b>FNA</b>	<b>fine-needle aspiration</b>	<b>PRS</b>	<b>polygenic risk scores</b>		
		<b>PTEN</b>	<b>phosphatase and tensin homolog</b>		
		<b>PV</b>	<b>pathogenic variant</b>		



# NCCN Guidelines Version 1.2023

## Genetic Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic Cancer

NCCN Categories of Evidence and Consensus	
<b>Category 1</b>	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2A</b>	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2B</b>	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
<b>Category 3</b>	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

**Note:** All recommendations are category 2A unless otherwise indicated.  
**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

## Discussion

This discussion corresponds to the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic. Last updated: March 9, 2022.

## Table of Contents

<b>Overview</b>	<b>MS-2</b>	MLH1, MSH2, MSH6, PMS2, EPCAM	MS-26
<b>Literature Search Criteria and Guidelines Update Methodology</b>	<b>MS-3</b>	NBN	MS-27
<b>Genetic Risk Assessment and Counseling</b>	<b>MS-3</b>	NF1	MS-27
Evaluating the Source of Genetic Testing Information	MS-5	PALB2	MS-28
Tumor Testing	MS-5	RAD51C and RAD51D	MS-28
Multi-Gene Testing	MS-5	STK11	MS-29
Pre- and Post-Test Counseling	MS-7	NCCN Genetic Testing Criteria	MS-29
<b>High-Penetrance Breast and/or Ovarian Cancer Susceptibility</b>		Testing Criteria Related to Prostate Cancer	MS-30
<b>Genes</b>	<b>MS-9</b>	Systemic Therapy Decision-Making	MS-30
BRCA-Related Breast/Ovarian Cancer Syndrome	MS-10	Ashkenazi Jewish Ancestry	MS-31
Breast Cancer Risk	MS-10	Breast Cancer Population Testing	MS-31
Ovarian Cancer Risk	MS-11	Probability Models	MS-32
Prostate Cancer Risk	MS-12	Li-Fraumeni Syndrome	MS-32
Pancreatic Cancer Risk	MS-13	Risk Assessment, Counseling, and Management	MS-35
Other Cancer and Health Risks	MS-13	Cowden Syndrome/ <i>PTEN</i> Hamartoma Tumor Syndrome	MS-37
Risk Management	MS-14	Risk Assessment, Counseling, and Management	MS-40
Other Pathogenic/Likely Pathogenic Variants Associated with		<b>Hereditary Pancreatic Cancer</b>	<b>MS-42</b>
Breast/Ovarian Cancer	MS-22	Pancreas Screening	MS-43
ATM	MS-24	<b>Table 1. Glossary of Relevant Genetic Terms (from the National</b>	
BARD1	MS-24	<b>Cancer Institute [NCI])</b>	<b>MS-45</b>
BRIP1	MS-25	<b>Table 2. Genetic Test Results to Determine the Presence of a</b>	
CDH1	MS-25	<b>Cancer-Predisposing Gene</b>	<b>MS-46</b>
CHEK2	MS-25	<b>References</b>	<b>MS-47</b>

### Overview

All cancers develop as a result of pathogenic or likely pathogenic (P/LP) variants in certain genes, such as those involved in the regulation of cell growth and/or DNA repair,<sup>1,2</sup> although not all of these P/LP variants are inherited from a parent. For example, sporadic P/LP variants can occur in somatic/tumor cells only, and de novo P/LP variants can occur for the first time in a germ cell (ie, egg or sperm) or in the fertilized egg itself during early embryogenesis. However, family studies have long documented an increased risk for several forms of cancer among first-degree relatives (ie, parents, siblings, children) and second-degree relatives (ie, grandparents, aunts or uncles, grandchildren, nieces or nephews) of affected individuals. These individuals may have an increased susceptibility to cancer as the result of one or more P/LP variants present in parental germline cells; cancers developing in these individuals may be classified as hereditary or familial cancers.

Hereditary cancers are often characterized by P/LP variants associated with increased risk for certain cancers (ie, a high-penetrance phenotype) and transmission to offspring through the mother and/or father.<sup>3,4</sup> They often have an early age of onset and exhibit an autosomal dominant inheritance pattern (ie, occur when the individual has a P/LP variant in only one copy of a gene). Familial cancers share some but not all features of hereditary cancers. For example, although familial breast cancers occur in a given family more frequently than in the general population, they generally do not exhibit the inheritance patterns or onset age consistent with hereditary cancers. Familial cancers may be associated with chance clustering of sporadic cancer cases within families, genetic variation in lower penetrance genes, a shared environment, or combinations of these factors.<sup>5-8</sup>

An individual suspected of being at risk for hereditary cancer should be offered genetic counseling.<sup>9,10</sup> This is consistent with recommendations

from the U.S. Preventive Services Task Force (USPSTF).<sup>11</sup> Assessment of an individual's risk for familial or hereditary cancer is based on a thorough evaluation of the personal and family history. With respect to hereditary cancers, advances in molecular genetics have identified a number of genes associated with inherited susceptibility to breast, ovarian, and pancreatic cancers (eg, *BRCA1/2*, *TP53*, *CDH1*) and have provided a means of characterizing the specific P/LP variant present in certain individuals and families exhibiting an increased risk for cancer. The field of cancer genetics has implications for all aspects of cancer management of individuals with hereditary or familial cancers, including prevention, screening, and treatment.<sup>12</sup>

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic were developed with an acute awareness of the preliminary nature of much of our knowledge regarding the clinical application of the rapidly emerging field of molecular genetics, and with an appreciation for the need for flexibility when applying these guidelines to individual families. Furthermore, it should be emphasized that these Guidelines were not developed as a substitute for professional genetic counseling. Rather, they are intended to: 1) serve as a resource for health care providers to identify individuals who may benefit from cancer risk assessment and genetic counseling; 2) provide genetics health care professionals with an updated tool for the assessment of individual breast cancer, ovarian cancer, and pancreatic cancer risk and to guide decisions related to genetic testing; and 3) facilitate a multidisciplinary approach in the management of individuals at increased risk for hereditary breast, ovarian, and pancreatic cancer. The current NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic focus primarily on assessment of P/LP variants associated with increased risk of breast, ovarian, pancreatic, and prostate cancer, including *BRCA1*, *BRCA2*, *CDH1*, *PALB2*, *PTEN*, and *TP53*, and recommended approaches to

genetic counseling/testing and management strategies in individuals with these P/LP variants. Where possible, P/LP variants in more recently identified genes have been addressed to the extent possible given the limited information available. Recommendations regarding P/LP variants associated with pancreatic cancer, and pancreas screening for individuals harboring such variants, were added to the Guidelines in the 2020 update. Additionally, testing criteria for those with or at risk for prostate cancer have also been included in the Guidelines.

A glossary of genetic terms is included in [Table 1](#) for reference.

### Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic, an electronic search of the PubMed database was performed to obtain key literature using the following search terms: (hereditary breast cancer) or (familial breast cancer) or (hereditary ovarian cancer) or (familial ovarian cancer) or (Li-Fraumeni syndrome) or (Cowden syndrome) or (pten hamartoma tumor syndrome) or (brca breast cancer) or (brca ovarian cancer) or (hereditary pancreas cancer) OR (hereditary pancreatic cancer) OR (familial pancreas cancer) OR (familial pancreatic cancer) OR (brca pancreas cancer) OR (brca pancreatic cancer) OR (cancer genetic testing) or (cancer genetic counseling). The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.<sup>13</sup>

NCCN recommendations have been developed to be inclusive of individuals of all sexual and gender identities to the greatest extent possible. When citing data and recommendations from other organizations, the terms *men*, *male*, *women*, and *female* will be used to be consistent with the cited sources.

### Genetic Risk Assessment and Counseling

Cancer genetic risk assessment and genetic counseling is a multi-step process involving the identification and counseling of individuals at risk for familial or hereditary cancer. The purpose of cancer genetic counseling is to educate individuals about the genetic, biological, and environmental factors related to a cancer diagnosis and/or risk for disease to help derive personal meaning from cancer genetic information, and to empower them to make educated, informed decisions about genetic testing, cancer screening, and cancer prevention. Many patients undergoing genetic testing do not receive proper counseling.<sup>14</sup> Further, testing rates are inadequate among some high-risk populations, including members of racial/ethnic minority groups.<sup>15,16</sup> A genetic counselor, clinical geneticist, oncologist, surgeon, oncology nurse, or other health professional with expertise and experience in cancer genetics should be involved in every stage of the process.<sup>9</sup>

Testing is clinically indicated in individuals for whom there is a personal or family history suggesting genetic cancer susceptibility and for whom results will aid in risk management and treatment. The selection of genes for which testing is indicated is based on the personal and familial characteristics that determine the individual's prior probability of being a carrier of a P/LP variant, and on the psychosocial degree of readiness of the person to receive genetic test results. Genetic risk assessment is a dynamic process and can change if additional relatives are diagnosed with cancer. The genetic testing strategy is greatly facilitated when a P/LP variant has already been identified in another family member. In that case, the genetic testing laboratory can limit the search for P/LP variants in additional family members to the same location in the gene. However, if there is reason to suspect more than one P/LP variant in the family, then broader testing may be considered.



For the majority of families in whom presence of a P/LP variant is unknown, it is best to consider testing an affected family member first, especially a family member with early-onset disease, bilateral disease, or multiple primaries, because that individual has the highest likelihood for a positive test result. The testing of the unaffected individual (or of unaffected family members) should only be considered when no affected family member is available for testing. In such cases, the unaffected individual or unaffected close relative with the highest likelihood of testing positive for the P/LP variant should be tested. This may include the relative closest to the family member with the youngest age at diagnosis, bilateral disease, multiple primary tumors, or other cancers associated with a suspected hereditary syndrome. A negative test result in such cases, however, is considered indeterminate and does not provide the same level of information as when there is a known P/LP variant in the family. Thus, one should be mindful that, when testing unaffected individuals (in the absence of having tested affected family members), significant limitations may exist in interpreting the test results, and testing multiple family members may be indicated since absence of a P/LP variant in one unaffected relative does not rule out a P/LP variant in others. The maternal and paternal sides of the family should be considered independently for familial patterns of cancer. “Limited” family structure is defined as two or fewer first- or second-degree female relatives who survive past age 45 (on either side of the family) and/or possessing no or inadequate information about one’s birth parents.<sup>17</sup>

Individuals who have received allogeneic hematopoietic cell transplantation (HCT) should not have molecular genetic testing performed on blood samples, as these blood cells would represent donor-derived DNA. In such cases, DNA of the individual being tested should be extracted from a fibroblast culture, if available. If this is not possible, buccal cells may be considered as an alternative source for DNA; however, a study has reported that over time, buccal epithelial cells are

replaced by donor-derived cells in allogeneic HCT recipients.<sup>18,19</sup> Therefore, genetic testing using buccal swab samples may be limited given this known risk of donor DNA contamination. Fibroblasts are also indicated when testing individuals with active or recent hematologic malignancies.<sup>20</sup>

A counseling dilemma is posed by the finding of a variant of uncertain significance (VUS), a genetic alteration that may actually represent a benign polymorphism unrelated to an increased cancer risk or may indicate an increased cancer risk. More than 90% of the time, if a VUS is reclassified, it is downgraded to benign or likely benign.<sup>21</sup> These patients should be considered for referral to research studies that aim to define the functional impact of the gene variant, such as variant reclassification programs through clinical labs or registries. Some examples of these programs and registries include ClinVar (the archival database at the National Center for Biotechnology Information [NCBI]); the NIH-funded Clinical Genome Resource (ClinGen; <https://www.clinicalgenome.org/>); the international Evidence-based Network for the Interpretation of Germline Mutant Alleles (ENIGMA; <https://enigmaconsortium.org/>); and the International Society for Gastrointestinal Hereditary Tumors (InSIGHT; <http://insight-group.org/>). It is important to note that there may be inconsistencies among how some programs and registries interpret the clinical actionability of some VUS, which may lead to confusion regarding medical management.<sup>22-24</sup> Family members should not be tested for a VUS for the purposes of clinical management unless there are conflicting data between laboratories regarding the classification of a variant. In the event where there are discrepancies in classification, careful consideration must be taken regarding family history, testing family members, and if other functional studies could aid in variant classification. Clinicians and scientists should work together to develop a VUS classification system as more information is discovered in research studies.<sup>25</sup> Carriers of a VUS or likely benign variant should be managed based on family history of cancer.



Carriers of a P/LP variant should be encouraged to participate in clinical trials or genetic registries. Carriers should be encouraged to recontact their genetics providers every few years for updates, as laboratories may issue amended reports as the knowledge base surrounding hereditary cancer risk expands.

### Evaluating the Source of Genetic Testing Information

Reports regarding germline findings that may impact medical management should come from laboratories that are certified by the College of American Pathologists (CAP) and Clinical Laboratory Improvement Amendments (CLIA), with some U.S. states (eg, New York) having additional reporting requirements. Recently, there has been an increase in genetic test results through direct-to-consumer (DTC) services or through tumor profiling. The testing typically used by companies providing ancestry information directly to consumers is microarray-based single nucleotide polymorphism (SNP) testing that has not been validated for clinical use. These companies do not provide comprehensive genetic analysis that includes gross deletion or duplication analysis. Third-party services are available to assist patients with interpreting their raw data, but these services are not government-regulated. In addition to the errors inherent in working with raw uncurated data from DTC labs, other limitations of these services include inadequate informed consent process, uncertain clinical validity and utility, and lack of medical oversight.<sup>26</sup> Currently available tests also only provide limited founder P/LP variant results without the benefit of family history. An analysis of concordance between DTC testing results and results from confirmatory testing for 49 patients showed a false-positive rate of 40%, as well as variant classification errors in 8 patients.<sup>27</sup> Given the limitations of the information obtained from DTC services, confirmatory germline testing by a certified laboratory is recommended, and changes to a patient's medical management based solely on DTC testing results are not recommended.<sup>27</sup>

Incidental germline findings discovered through other sources (eg, participation in a research study) should be reviewed by a genetics professional.<sup>28</sup> Confirmatory testing in these cases may be recommended, especially if the reporting laboratory is not appropriately certified.

### Tumor Testing

Tumor profiling can be considered complementary to germline testing. However, the absence of a P/LP variant for a given gene from tumor profiling does not rule out the possibility of a germline P/LP variant in that gene. Tumor testing tends to be designed to address treatment actionability and prognosis.<sup>29</sup> Therefore, a variant interpreted as P/LP in the germline may be interpreted as normal or as a VUS in the tumor, if that variant has no clear clinical implications. In addition, the sensitivity of most tumor testing is lower (particularly for intermediate-sized deletions and duplications) than that for most dedicated germline tests, sometimes due to filtering out of germline findings reported in tumor sequencing results. If a patient meets testing criteria for germline testing for a given gene, then confirmatory germline testing should be considered through a CLIA-approved lab despite tumor profiling results. Circulating tumor DNA (ctDNA) assays may be used by some labs. ctDNA has the potential to identify both somatic and germline variants.<sup>30</sup> However, since the primary intent of tumor testing is to inform treatment decision-making, ctDNA assays are not validated for reporting or interpretation of germline variants. If a germline variant that could impact medical management is detected with a ctDNA assay, then confirmatory testing with a CLIA-approved assay intended for detection and interpretation of germline results is recommended.

### Multi-Gene Testing

Next-generation sequencing allows for the sequencing of multiple genes simultaneously. This is referred to as multi-gene testing. Multi-gene testing can detect P/LP variants not found in single-gene testing.<sup>31-33</sup> Since more

than one gene can explain an inherited cancer syndrome, phenotype-directed testing based on personal and family history through a multi-gene panel test is often more efficient and/or cost-effective.<sup>34-36</sup> Multi-gene testing may also be considered for those who tested negative for one particular syndrome, but whose personal and family history is suggestive of an inherited susceptibility.<sup>34,37</sup> It has become routine practice at many institutions to now order phenotypically directed multi-gene panel tests to assess for P/LP changes in multiple relevant genes simultaneously.<sup>38</sup>

There are several issues to consider regarding multi-gene testing. First, commercially available tests may differ significantly on a number of factors, such as number of genes analyzed, turnaround time, insurance coverage, laboratory expertise, variant reclassification protocol, methods of DNA/RNA analysis, and availability of financial assistance for cascade testing of relatives, among others. Therefore, the specific laboratory and multi-gene test should be chosen carefully.<sup>34</sup> In addition, P/LP variants identified for more than one gene add complexity that may lead to difficulty in making risk management recommendations.<sup>37</sup> A management plan based on genetic test results should only be developed for identified P/LP variants that are clinically actionable.

A major dilemma regarding multi-gene testing is that there are limited data and a lack of clear guidelines regarding degree of cancer risk associated with some of the genes assessed, and how to communicate and manage risk for carriers of these genes.<sup>39-43</sup> This issue is compounded by the low incidence rates of hereditary disease, leading to a difficulty in conducting adequately powered studies.<sup>39</sup> Multi-gene tests include moderate-penetrance genes, and they often also include low-penetrance genes for which there are little available data regarding degree of cancer risk and guidelines for risk management.<sup>34,44</sup> Analysis from a prospective, multicenter cohort study including 2,984 patients with cancer unselected based on cancer type, disease stage, family history of cancer, age of

diagnosis, and ethnicity showed that, with use of an 80-gene panel test, a P/LP variant was found in 13.3%, with a highly penetrant variant found in 5%.<sup>45</sup> About half of the identified variants were of moderate or low penetrance, and a VUS was also found in about half the sample. The use of tailored panels that are disease-focused and include clinically actionable cancer susceptibility genes is preferred over large panels that include genes of uncertain clinical relevance. Also, certain variants in a gene may be associated with a different degree of risk than other variants in that gene. For example, the presence of the c.7271T>G missense P/LP variant in *ATM* is associated with an increased risk for early-onset breast cancer,<sup>46-48</sup> but the association between other *ATM* variants and breast cancer susceptibility is less clear.<sup>49-52</sup>

Multi-gene tests also increase the likelihood of detecting a VUS.<sup>32-34,40,53-55</sup> However, as multi-gene testing is increasingly used, the frequency of a variant being interpreted as a VUS is expected to decrease. There is also an increase in the chance of finding genotypically distinct cell lines (ie, genetic mosaicism) with next-generation sequencing.<sup>56</sup> Clones of non-cancerous cells (ie, aberrant clonal expansion) containing a P/LP *TP53* variant have been found in healthy adults undergoing multi-gene testing. This phenomenon can often be attributed to clonal hematopoiesis, a condition in which a hematopoietic stem cell begins making blood cells with the same acquired P/LP variant.<sup>20</sup> When there is no evidence of a hematologic malignancy, then it is referred to as clonal hematopoiesis of indeterminate potential (CHIP). Age-related CHIP is associated with increased risk of hematologic malignancies,<sup>57,58</sup> but may also lead to unnecessary clinical intervention. Ancillary testing of non-lymphoid non-cancerous tissue can be used to help determine the true presence of a germline variant.<sup>20</sup>

Polygenic risk scores (PRS) are now sometimes included in some genetic test reports. PRS are groups of SNPs associated with a specific disorder



or disease, such as hereditary cancer. Some studies evaluating the validity of PRS for identifying those at risk of hereditary cancer to date have been conducted, primarily with breast and prostate cancers. Two studies identified PRS that were strongly associated with ER-negative breast cancer in carriers of a *BRCA1* P/LP variant, overall breast cancer in carriers of a *BRCA2* P/LP variant, and high-grade serous ovarian cancer in carriers of both *BRCA1* and *BRCA2* P/LP variants.<sup>59,60</sup> Another study of male carriers of a *BRCA1/2* P/LP variant identified PRS associated with breast cancer risk and prostate cancer risk.<sup>61</sup> Studies have also evaluated the potential clinical utility of incorporating PRS into a risk-stratified approach for screening for prostate cancer<sup>62</sup> and for identifying age of onset of aggressive prostate cancer.<sup>63</sup> Studies of PRS have largely been done with those of European ancestry.<sup>64</sup> Studies with larger samples from more diverse populations are needed. Given the lack of validation of PRS, these should not be used to inform medical management at this time.

### Pre- and Post-Test Counseling

For individuals potentially meeting established criteria for one or more of the hereditary cancer syndromes, genetic testing should be considered along with appropriate pre- and post-test counseling. Pre-test counseling should include a discussion of why the test is being offered and how test results may impact medical management, cancer risks associated with the P/LP variant in question, the significance of possible test results (see Table 2), the likelihood of a positive result, technical aspects and accuracy of the test, cost considerations, risks of genetic discrimination, psychosocial aspects, confidentiality issues, the potential significance of the test results for family members, and other topics.<sup>7</sup> A discussion of confidentiality issues should include an explanation of the federal Genetic Information Nondiscrimination Act (GINA) enacted in 2008, which prohibits most health insurers and employers from discrimination on the basis of genetic test results.<sup>65</sup> Since some patients with cancer who have a poor prognosis may be unable to receive results directly, a plan for results

disclosure should be discussed, such as the patient consenting to Release of Information of test results to a spouse or other close relative. A detailed family history should be collected, which involves development of an expanded pedigree, beginning with the health of the individual diagnosed with cancer and proceeding outward to include first-, second-, and third-degree relatives on both the maternal and paternal sides. Factors that limit the informativeness of the pedigree are small family size, a small number of individuals of the susceptible gender for sex-limited cancers, reduced penetrance, early deaths in family members (which precludes the possibility that they will develop adult diseases), prophylactic surgeries that remove an organ from subsequent risk for cancer (eg, hysterectomy for uterine fibroids in which the ovaries are also removed), adoptions, and inaccurate or incomplete information on family members (eg, in the case of adoption or divorce).<sup>5,66</sup> It is also important to know the ancestry/ethnicity of the individual, since members of certain groups (eg, Ashkenazi Jewish) have increased risks of carrying P/LP variants for specific diseases. Any family members who received genetic testing should also be noted, as well as testing results. Finally, a detailed medical and surgical history from the proband should be collected, and a physical examination should be performed by a qualified clinician when appropriate.

The presentation of testing information is most effective when tailored to the age and education of the person undergoing counseling, and that individual's personal exposure to the disease, level of risk, and social environment.<sup>7</sup> Information could be delivered in person or over the phone.<sup>67,68</sup> Telehealth (ie, real-time two-way videoconference) is also increasingly utilized as a feasible alternative for in-person genetic counseling.<sup>68</sup> Remote options (telephone, telehealth) have the potential to help improve genetic testing rates in areas with inadequate access.<sup>68</sup>



Post-test counseling includes disclosure of results, a discussion of the associated medical risks, an assessment of the impact of the results on the emotional state of the individual, a discussion of the impact of the results on the medical management of the individual, and how and where the patient will be screened for cancer risk.<sup>9</sup> Counseling should include making the individual aware of any available resources, such as disease-specific support groups, high-risk clinics, advocacy groups, and research studies.<sup>69</sup> The counselor should discuss the importance of genetic counseling and testing for at-risk relatives.

Since some P/LP variants are associated with rare autosomal recessive conditions (eg, Fanconi anemia is associated with *BRCA2*, *BRIP1*, and *PALB2* variants), the proband should be advised regarding possible inherited cancer risk to relatives and his/her options for risk assessment and management. Testing of a partner of a carrier of a P/LP variant may also be considered to inform reproductive decision-making.<sup>70</sup> See *Autosomal Recessive Risk in Cancer Genes – Multi-Gene Panel Testing* in the algorithm for a full list of the P/LP variants covered in these Guidelines that are associated with autosomal recessive conditions.

Pre- and post-test genetic counseling with involvement of an expert in cancer genetics is recognized as the gold standard. However, during the meeting for the 2020 update, the panel acknowledged that most genetic testing is conducted by providers with limited expertise in genetics and often without pre-test genetic counseling.<sup>71-73</sup> Shortages in genetics health providers,<sup>74</sup> expansion of testing indications, aggressive marketing, and increased accessibility of testing due to plummeting costs inclusive of DTC models for testing provide the impetus for the panel to identify scenarios in which referral to a genetics health provider should be considered. These scenarios are as follows: identification of a P/LP variant; negative results despite family history suggestive of inherited disease; VUS result for which provider considers altering clinical management; mosaic or possibly

mosaic result; discrepant interpretation of variants (eg, discordant results across laboratories); interpretation of PRS; and detection of P/LP variants from DTC testing.

Counseling may be warranted for those with negative or indeterminate results, as reasons for a negative result include the following scenarios: P/LP variant exists in a gene variant that was not recognized due to limitations in technology; P/LP variant in a gene variant that was not evaluated; and potential presence of a P/LP variant in a family member that was not detected in the individual.

The full list of elements that should be included in pre- and post-test genetic counseling can be found in the *Principles of Cancer Risk Assessment and Counseling* in the algorithm.

### *Reproductive Options*

The outcomes of genetic testing can have a profound impact on family planning decisions for individuals of reproductive age who are found to be carriers of a P/LP variant. Counseling for reproductive options such as prenatal diagnosis and assisted reproduction using preimplantation genetic testing (PGT) may therefore be warranted for couples expressing concern over their future offspring's carrier status of a P/LP variant. Such counseling should include a comprehensive discussion of the potential risks, benefits, and limitations of reproductive options, including cost.

Prenatal diagnosis involves postimplantation genetic analysis of an early embryo, utilizing chorionic villi or amniotic fluid cell samples; genetic testing is typically conducted between week 12 and week 16 of gestation, and testing results may potentially lead to a couple's decision to terminate pregnancy.<sup>75,76</sup> PGT has emerged as an alternative method of genetic testing in early embryos. PGT involves the testing of 1 or 2 cells from embryos in very early stages of development (ie, 6–8 cells) after in vitro fertilization (IVF). This procedure allows for the selection of unaffected

embryos to be transferred to the uterus,<sup>75,76</sup> and may therefore offer the advantage of avoiding potential termination of pregnancy. The PGT process requires the use of IVF regardless of the fertility status of the couple (ie, also applies to couples without infertility issues), and IVF may not always lead to a successful pregnancy. Lastly, the technology or expertise may not be readily available in a couple's geographic location.

Various factors, both medical and personal, must be weighed in the decision to utilize prenatal diagnosis or PGT. Medical considerations may include factors such as the age of onset of the hereditary cancer, penetrance, severity or associated morbidity and mortality of the cancer, and availability of effective cancer risk reduction methods or effective treatments.<sup>75,76</sup> Although the use of prenatal diagnosis or PGT is relatively well established for severe hereditary disorders with very high penetrance and/or early onset (eg, Fanconi anemia), its use in conditions associated with lower penetrance and/or later onset (eg, hereditary breast or ovarian cancer syndrome) remains somewhat controversial from both an ethical and regulatory standpoint. Personal considerations for the decision to utilize prenatal diagnosis or PGT may include individual ethical beliefs, value systems, cultural and religious beliefs, and social and economic factors. Successful births have been reported with the use of PGT and IVF in carriers of a *BRCA1/2* P/LP variant,<sup>77,78</sup> but data in the published literature are still very limited. In addition, data pertaining to long-term safety or outcomes of PGT and assisted reproduction in carriers of a P/LP variant are not yet available.

### High-Penetrance Breast and/or Ovarian Cancer Susceptibility Genes

Specific patterns of hereditary breast and ovarian cancers have been found to be linked to P/LP variants in the *BRCA1/2* genes.<sup>79,80</sup> In addition, two very rare hereditary cancer syndromes exhibiting an increased risk for breast cancer are Li-Fraumeni syndrome (LFS) and Cowden syndrome,

which are related to germline P/LP variants in the *TP53* and *PTEN* genes, respectively.<sup>81,82</sup> *PALB2* is also considered a high penetrance breast cancer susceptibility gene.<sup>83-85</sup> These hereditary syndromes share several features beyond elevation of breast cancer risk. These syndromes arise from germline P/LP variants that are not within sex-linked genes; hence, the variants can be inherited from either parent. The syndromes are associated with breast cancer onset at an early age and development of other types of cancer, and exhibit an autosomal dominant inheritance pattern (see [Table 1](#)). Offspring of an individual with one of these hereditary syndromes have a 50% chance of inheriting the P/LP variant. In addition, individuals with these hereditary syndromes share increased risks for multiple cases of early-onset disease as well as bilateral disease. The P/LP variants associated with these hereditary syndromes are considered to be highly penetrant. In addition, the manifestations (ie, expression) of these hereditary syndromes are often variable in individuals within a single family (eg, age of onset, tumor site, number of primary tumors). The risk of developing cancer in individuals with one of these hereditary syndromes depends on numerous variables including the gender and age of the individual.

Prior to 2020, the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian (Breast, Ovarian, and Pancreatic as of 2020) focused largely on testing criteria for *BRCA1/2* and appropriate risk management for carriers of a *BRCA1* or *BRCA2* P/LP variant. Sections on LFS and Cowden syndrome/*PTEN* hamartoma tumor syndrome (PHTS) were also included. Based on strong evidence that genes beyond *BRCA1/2*, *TP53*, and *PTEN* confer markedly increased risk of breast and/or ovarian cancers, these Guidelines have been expanded; see the sections below on other P/LP variants associated with breast/ovarian cancer.

### BRCA-Related Breast/Ovarian Cancer Syndrome

Both the *BRCA1* and *BRCA2* genes encode for proteins involved in tumor suppression. *BRCA1/2* P/LP variants can be highly penetrant (for definition, see [Table 1](#)), although the probability of cancer development in carriers of *BRCA1/2* P/LP variants is variable, even within families with the same variant.<sup>86-88</sup> At present, it is unclear whether penetrance is related only to the specific P/LP variant identified in a family or whether additional factors, either genetic or environmental, affect disease expression. Epigenetic modification can also influence disease penetrance for a P/LP variant.<sup>89</sup> It is generally accepted, however, that carriers of *BRCA1/2* P/LP variants have an excessive risk for both breast and ovarian cancer that warrants consideration of more intensive screening and preventive strategies.

### Breast Cancer Risk

Estimates of penetrance range from a 41% to 90% lifetime risk for breast cancer, with an increased risk for contralateral breast cancer.<sup>85,90-97</sup> A prospective cohort study including 9856 unaffected *BRCA1/2* carriers showed that a cumulative risk of breast cancer by 80 years of age was 72% for carriers of a *BRCA1* P/LP variant and 69% for carriers of a *BRCA2* P/LP variant.<sup>98</sup> Estimates of cumulative risk for contralateral breast cancer 20 years after breast cancer diagnosis are 40% for carriers of a *BRCA1* P/LP variant and 26% for carriers of a *BRCA2* P/LP variant, though this risk is age-dependent and greatest in those diagnosed with breast cancer at an early age (ie, <40 years).<sup>98</sup>

While the evidence is mixed, we do not currently have evidence to support that *BRCA*-associated breast cancers are more aggressive and/or have poorer outcomes. A meta-analysis including 13 studies showed that carriers of a *BRCA1* P/LP variant with breast cancer had worse overall survival (OS) compared to those without a *BRCA1* or *BRCA2* P/LP variant (hazard ratio [HR], 1.50; 95% CI, 1.11–2.04), while OS did not significantly

differ between those harboring a *BRCA2* P/LP variant and those without a *BRCA1* or *BRCA2* P/LP variant (HR, 0.97; 95% CI, 0.78–1.22).<sup>99</sup> Another meta-analysis including 60 studies and 105,220 patients with breast cancer also found that carriers of a *BRCA1* P/LP variant had worse OS compared to non-carriers (HR, 1.30; 95% CI, 1.11–1.52; *P* = .001).<sup>100</sup> Carriers of a *BRCA2* P/LP variant had worse breast cancer-specific survival compared to non-carriers (HR, 1.29; 95% CI, 1.03–1.62; *P* = .03), though OS was not significantly different. This meta-analysis also showed that, among patients with triple-negative breast cancer, *BRCA1/2* P/LP variants are associated with better OS (HR, 0.49; 95% CI, 0.26–0.92; *P* = .03). However, this subgroup analysis only included two studies. A third meta-analysis including 66 studies also showed that a *BRCA2* P/LP variant was associated with worse breast cancer-specific survival (HR, 1.57; 95% CI, 1.29–1.86), but study results were too heterogeneous for the analysis to be conclusive.<sup>101</sup> Results of the prospective cohort Prospective Outcomes in Sporadic versus Hereditary breast cancer (POSH) study including 2,733 women with breast cancer showed no significant differences in OS between carriers of a *BRCA1/2* P/LP variant and non-carriers 2, 5, and 10 years after diagnosis.<sup>102</sup>

*BRCA1/2* P/LP variants are associated with early-onset breast cancer. In a sample of 21,401 families who met German Consortium for Hereditary Breast and Ovarian Cancer testing criteria for *BRCA1/2* P/LP variants, a P/LP variant was detected in 13.7% of families with a single case of breast cancer diagnosed at younger than 36 years of age.<sup>103</sup> An analysis of 6,478 patients who were diagnosed with breast cancer before 50 years of age showed that carriers of a *BRCA1* P/LP variant had worse OS compared to patients who were not carriers of a P/LP *BRCA1/2* variant (HR, 1.28; 95% CI, 1.05–1.57; *P* = .01), but this association was no longer statistically significant when taking into account disease and treatment characteristics (HR, 1.20; 95% CI, 0.97–1.47; *P* = .09).<sup>104</sup> *BRCA2* P/LP variants were not



significantly associated with decreased OS in these analyses, except for the first 5 years of follow-up (HR, 1.56; 95% CI, 1.06–2.28;  $P = .02$ ).

Some histopathologic features have been reported to occur more frequently in breast cancers of individuals with a germline *BRCA1/2* P/LP variant. For example, several studies have shown that *BRCA1*-related breast cancer is more likely to be characterized as ER-/PR-negative and HER2-negative (ie, “triple negative”).<sup>85,105–110</sup> Studies have reported *BRCA1* P/LP variants in 4.4% to 16% of patients with triple-negative breast cancer.<sup>85,110–119</sup> One cohort study showed an absolute lifetime risk of 40% for hormone receptor-positive (ER+ and/or PR+) breast cancer in carriers of a P/LP *BRCA2* variant.<sup>85</sup> The Breast Cancer Association Consortium and the CARRIERS case-control studies showed associations between a *BRCA2* P/LP variant and increased risk of ER-positive breast cancer (1.46%; OR, 5.68; 95% CI, 4.65–6.96 and 1.09%; OR, 4.66; 95% CI, 3.52–6.23, respectively).<sup>118,119</sup> Another case-control study showed that the 20-year survival rate in carriers of a *BRCA2* P/LP variant with ER-positive tumors was 62.2%, compared to 83.7% in those with ER-negative tumors, though this difference was only statistically significant in those younger than age 50 ( $n = 199$ ; 68.3% vs. 91.3%, respectively;  $P = .03$ ).<sup>120</sup> A case-control study of carriers of the Icelandic founder *BRCA2* variant 999del5 showed that ER-positive disease was associated with increased mortality risk, compared to those with ER-negative disease (HR, 1.94; 95% CI, 1.22–3.07;  $P = .005$ ).<sup>121</sup> However, prevalence of ER-negative disease was not significantly greater in carriers of a P/LP *BRCA2* variant than in non-carriers (75.6% vs. 70.2%, respectively;  $P = .11$ ). The explanation for the association between *BRCA2* P/LP variant with ER-positive tumors and poor survival outcomes is currently unknown and warrants investigation, though one hypothesized explanation includes difference in estrogen signaling pathways and increased sensitivity to ovarian hormones for these tumors.<sup>121,122</sup>

Among patients with triple-negative disease, carriers of a P/LP *BRCA* variant were diagnosed at a younger age compared with non-carriers.<sup>112,123</sup> In a study of a large cohort of patients with triple-negative breast cancer ( $N = 403$ ), the median age of diagnosis among carriers of a P/LP *BRCA1* variant ( $n = 65$ ) was 39 years.<sup>111</sup> Patients in this population-based study were unselected for family history or age. Among the group of patients with early-onset (age at diagnosis <40 years) triple-negative breast cancer ( $n = 106$ ), the incidence of *BRCA1* P/LP variants was 36%; the incidence was 27% among those diagnosed before 50 years of age ( $n = 208$ ). Results from the prospective cohort POSH study showed that, among 558 patients with triple-negative breast cancer, 2-year OS was greater in carriers of a *BRCA1/2* P/LP variant than in non-carriers (95% vs. 91%, respectively; HR, 0.59; 95% CI, 0.35–0.99;  $P = .047$ ), but 5- and 10-year OS did not differ significantly between these groups.<sup>102</sup>

Carriers of a P/LP *BRCA1/2* variant who were assigned male at birth also have a greater risk for cancer susceptibility.<sup>124</sup> Among male patients with breast cancer unselected for family history, 4% to 14% tested positive for a germline *BRCA2* P/LP variant.<sup>125–128</sup> For males carrying a P/LP *BRCA2* variant, the cumulative lifetime risk for breast cancer has been estimated at 7% to 8%.<sup>129,130</sup> The cumulative lifetime risk for male carriers of a P/LP *BRCA1* variant is 1.2%.<sup>130</sup> In contrast, for males who are not carriers of a P/LP *BRCA1/2* variant, the lifetime risk for breast cancer has been estimated at approximately 0.1% (1 in 1000).<sup>127,131</sup>

### Ovarian Cancer Risk

Increased risks for cancers of the ovary, fallopian tube, and peritoneum are observed in carriers of a P/LP *BRCA1/2* variant.<sup>132–134</sup> In the setting of an invasive ovarian cancer diagnosis, a P/LP *BRCA1* variant has been found in 3.8% to 14.5% of patients, and a P/LP *BRCA2* variant has been found in 4.2% to 5.7% of patients.<sup>96,135–138</sup> *BRCA1* variants have an estimated 48.3% (95% CI, 38.8%–57.9%) cumulative risk of ovarian

cancer by age 70, while the cumulative risk by age 70 is 20.0% (95% CI, 13.3%–29.0%) for carriers of a P/LP *BRCA2* variant.<sup>139</sup>

Several studies have reported more favorable survival outcomes among carriers of a P/LP *BRCA1/2* variant in patients with ovarian cancer compared with non-carrier patients.<sup>140–146</sup> Survival outcomes appear to be most favorable for carriers of a P/LP *BRCA2* variant.<sup>140,145,147</sup> Additionally, *BRCA2* P/LP variants were associated with significantly higher response rates (compared with non-carriers or with carriers of a *BRCA1* P/LP variant) to primary chemotherapy. In contrast, *BRCA1* P/LP variants were not associated with prognosis or improved chemotherapy response.<sup>145</sup>

The histology of ovarian cancers in carriers of a P/LP *BRCA1/2* variant is more likely to be characterized as serous adenocarcinoma and high grade compared with ovarian cancers in non-carriers, although endometrioid and clear cell ovarian cancers also have been reported in the former population.<sup>134,136,148–151</sup> P/LP variants are also associated with non-mucinous ovarian carcinoma as opposed to mucinous.<sup>135,137</sup> Mucinous epithelial ovarian carcinomas may be associated with other P/LP variants, such as *TP53*,<sup>152</sup> which are implicated in LFS (see below). Non-epithelial ovarian carcinomas (eg, germ cell and sex cord-stromal tumors) are not significantly associated with a *BRCA1/2* P/LP variant,<sup>153</sup> though ovarian sex cord tumor with annular tubules is associated with *STK11* P/LP variants.<sup>154,155</sup> Current data show that ovarian low malignant potential tumors (ie, borderline epithelial ovarian tumors) are also not associated with a *BRCA1/2* P/LP variant.<sup>135</sup>

In studies of carriers of a P/LP *BRCA1/2* variant who underwent risk-reducing salpingo-oophorectomy (RRSO), occult gynecologic neoplasia, both invasive carcinoma and intraepithelial lesions, were identified in 4.5% to 9% of cases based on rigorous pathologic examinations of the ovaries and fallopian tubes.<sup>156–158</sup> Tubal intraepithelial carcinoma (TIC) is thought to represent an early precursor lesion for serous ovarian cancers, and TIC

(with or without other lesions) was detected in 5% to 8% of cases from patients carrying a P/LP *BRCA1/2* variant who underwent RRSO.<sup>156,159,160</sup> The fimbriae or distal tube was reported to be the predominant site of origin for these early malignancies found in carriers of a P/LP *BRCA1/2* variant.<sup>156,160,161</sup> Although TIC appeared to present more frequently among carriers of a P/LP *BRCA1/2* variant compared with non-carriers undergoing RRSO,<sup>160,161</sup> TIC has also been documented among patients with serous carcinomas unselected for family history or *BRCA* P/LP variant status.<sup>162</sup> Because TIC was identified in individuals who underwent surgery for risk reduction (for carriers of a P/LP *BRCA1/2* variant) or other gynecologic indications, the incidence and significance of these early lesions within the general population is unclear.

### Prostate Cancer Risk

Germline *BRCA1/2* P/LP variants are associated with increased risk for prostate cancer.<sup>163–166</sup> Carriers of a P/LP *BRCA1* variant have an estimated 29% (95% CI, 17%–45%) cumulative lifetime risk of prostate cancer, while the cumulative lifetime risk is 60% (95% CI, 43%–78%) for carriers of a P/LP *BRCA2* variant.<sup>167</sup> There is evidence that advanced or metastatic prostate cancer is associated with carrying a *BRCA2* P/LP variant; it is not yet known if an aggressive phenotype is also associated with *BRCA1* P/LP variant.<sup>168–170</sup> An international study including 5,545 patients with prostate cancer (with European ancestry) showed that the frequency of a *BRCA2* P/LP variant was significantly higher in patients with aggressive disease (ie, died from prostate cancer, metastatic disease, T4 disease, or T3 with Gleason score  $\geq 8$ ) than in patients with non-aggressive disease (OR, 3.19; 95% CI, 1.94–5.25).<sup>170</sup> A study of a large cohort of patients from Spain with prostate cancer (N = 2019) showed that carriers of a P/LP *BRCA1/2* variant had significantly higher rates of aggressive prostate cancer (Gleason score  $\geq 8$ ), nodal involvement, and distant metastasis compared with non-carriers.<sup>171</sup> In a sample of 692 patients with metastatic prostate cancer, unselected for family history or

age at diagnosis, 5.3% carried a *BRCA2* P/LP variant, and 0.9% carried a *BRCA1* P/LP variant.<sup>169</sup> In addition, analyses from a treatment center database showed that *BRCA1/2* and *ATM* (see below under *NCCN Genetic Testing Criteria: Testing Criteria Related to Prostate Cancer*) P/LP variant rates were highest in patients with metastatic disease (8.2%). This study also showed that carriers with prostate cancer had significantly decreased survival, compared with patients who were non-carriers (5 years vs. 16 years, respectively;  $P < .001$ ).<sup>172</sup> This association remained statistically significant when controlling for race, age, PSA, and Gleason score. Ashkenazi Jewish ancestry is also associated with *BRCA1/2* P/LP variants in patients with prostate cancer, with rates for *BRCA1* being 0% to 2% and rates for *BRCA2* being 1% to 3%.<sup>163,173-176</sup>

### Pancreatic Cancer Risk

Prior to more widespread testing of individuals with pancreatic cancer for germline variants in cancer predisposition genes, studies showed that *BRCA1/2* P/LP variant rates in pancreatic cancer cases ranged from 1% to 11% for *BRCA1* and 0% to 17% for *BRCA2*.<sup>177-185</sup> However, some of these studies included only patients with familial pancreatic cancer<sup>180,181,184</sup> or those of Ashkenazi Jewish ancestry,<sup>182</sup> both of whom may have a greater likelihood of testing positive for a *BRCA1/2* P/LP variant. More recent studies that used panel testing confirm that some pancreatic cancers harbor actionable *BRCA1/2* P/LP variants (0%–3% for *BRCA1* and 1%–6% for *BRCA2*).<sup>186-190</sup> Patients with pancreatic cancer and Ashkenazi Jewish ancestry may have a greater likelihood of testing positive for a *BRCA1/2* P/LP variant, with prevalence of detected P/LP variants in this group ranging from 5.5% to 19%, with P/LP variants being more common for *BRCA2*.<sup>182,183,185,191</sup>

More information on genes associated with pancreatic cancer can be found below, under *Hereditary Pancreatic Cancer*.

### Other Cancer and Health Risks

Some studies have suggested an increased risk specifically of serous uterine cancer in carriers of a P/LP *BRCA1/2* variant.<sup>192-196</sup> Analyses from a multicenter prospective cohort study including 1,083 carriers of a P/LP *BRCA1* variant who underwent RRSO without hysterectomy showed an increased risk for serous and/or serous-like endometrial cancer.<sup>197</sup> However, it has been suggested that the increased risk for endometrial cancer observed in some carriers of *BRCA1/2* P/LP variants may be due to the use of tamoxifen therapy by these patients rather than the presence of a P/LP variant.<sup>198,199</sup> A meta-analysis including five studies of patients with uterine serous cancer and Ashkenazi Jewish ancestry showed that *BRCA1/2* P/LP variant prevalence was greater in those with uterine serous cancer than in controls (also of Ashkenazi Jewish ancestry) (OR, 5.4; 95% CI, 2.2–13.1).<sup>192</sup> In a retrospective case control study including 2627 Jewish Israeli carriers of a P/LP *BRCA1/2* variant (88% Ashkenazi Jewish), risk of developing uterine cancer was increased, with an observed-to-expected ratio of 3.98 (95% CI, 2.17–6.67;  $P < .001$ ).<sup>195</sup> This association persisted regardless of uterine cancer histology. Despite some evidence of increased risk of uterine cancer in carriers of a P/LP *BRCA1/2* variant, the absolute risk is low.

Studies that investigated associations between *BRCA2* P/LP variant and cutaneous melanoma have drawn inconsistent conclusions, though there is some evidence of an association.<sup>200</sup> One study showed that women carrying a P/LP *BRCA2* variant have an elevated risk for leukemia (standardized incidence ratio [SIR], 4.76; 95% CI, 1.21–12.96;  $P = .03$ ), particularly women who have received chemotherapy (SIR, 8.11; 95% CI, 2.06–22.07;  $P = .007$ ).<sup>201</sup> Analyses of data from the Swedish Family-Cancer Database showed that carriers of a P/LP *BRCA1/2* variant who also have family history of breast and ovarian cancer are at increased risk of gastric cancer by age 70 (SIR, 1.88; 95% CI, 1.05–3.12).<sup>202</sup> A 1999 analysis from the Breast Cancer Linkage Consortium suggested that this



risk might be particularly elevated in carriers of a *BRCA2* P/LP variant (RR, 2.59; 95% CI, 1.46–4.61).<sup>203</sup> Finally, an analysis of 490 families with a known *BRCA1/2* P/LP variant showed an increased risk for ocular melanoma in carriers of a P/LP *BRCA2* variant (RR, 99.4; 95% CI, 11.1–359.8), though absolute risk is low.<sup>204</sup>

In cases where both partners carry a P/LP *BRCA2* variant, there may be a high risk for the offspring to develop Fanconi anemia, a rare autosomal recessive condition.<sup>70</sup> A review of 27 cases of Fanconi anemia with biallelic P/LP variants in *BRCA2* (FA-D1) showed a 97% cumulative risk of malignancy by age 5.2 years (79% risk of leukemia by age 10 years, 83% risk of any solid tumor by age 6.7 years, 85% risk of a brain tumor by age 9 years, and 63% risk of a Wilms tumor by age 6.7 years).<sup>205</sup> Some case reports have also identified biallelic *BRCA1* P/LP variants causing Fanconi anemia-like disorder,<sup>206,207</sup> particularly FANCS, a severe form of Fanconi anemia characterized by developmental delay, short stature, and microcephaly.<sup>208,209</sup>

### Risk Management

Recommendations for the medical management of *BRCA*-related cancers are based on an appreciation of the early onset and increased risk for associated cancers. An individual from a family with a known *BRCA1/2* P/LP variant who tests negative for the familial variant should be followed according to the recommendations for the general population (eg, the NCCN Guidelines for Breast Cancer Screening and Diagnosis [available at [www.NCCN.org](http://www.NCCN.org)] for breast cancer).

### Breast Cancer Risk Management

#### Screening

Mammography has served as the standard screening modality for detection of breast cancer during the last few decades. There are currently no data indicating that mammography on its own reduces mortality in

women with genetically increased risk for breast cancer.<sup>210</sup> Also, false-negative mammography results are common and have been correlated with factors such as presence of a *BRCA1/2* P/LP variant and high breast tissue density,<sup>211–214</sup> both of which may occur more frequently among younger women. Rapidly growing or aggressive breast tumors—also more common among younger women—have also been associated with decreased sensitivity of mammographic screening methods.<sup>211,215</sup> Prospective studies on comparative surveillance modalities in women at high risk for familial breast cancer (ie, confirmed or suspected *BRCA1/2* P/LP variant based on family history) have consistently reported higher sensitivity of MRI screening (77%–94%) compared with mammography (33%–59%) in detecting breast cancers. False-positive rates were higher with MRI in some reports, resulting in a slightly lower or similar specificity with MRI screening (81%–98%) compared with mammography (92%–100%).<sup>216–221</sup> The sensitivity with ultrasound screening (33%–65%) appeared similar to that of mammography in this high-risk population.<sup>217,219–221</sup> In a prospective screening trial (conducted from 1997–2009) that evaluated the performance of annual MRI and mammography in women (aged 25–65 years; N = 496) with confirmed P/LP *BRCA1/2* variant, sensitivity with MRI was significantly higher compared with mammography during the entire study period (86% vs. 19%;  $P < .0001$ ).<sup>222</sup> Factors such as age, P/LP variant type, or invasiveness of the tumor did not significantly influence the relative sensitivity of the two screening modalities. Importantly, the large majority (97%) of cancers detected by MRI screening were early-stage tumors.<sup>222</sup> At a median follow-up of 8 years from diagnosis, none of the surviving patients (n = 24) had developed distant recurrence. In an analysis of 606 women with either a family history of breast cancer or who harbor a P/LP variant associated with increased risk for breast cancer, sensitivity of breast MRI screening was reported to be 79%, while specificity was reported to be 86%.<sup>223</sup>

All of these studies discussed above evaluated a screening strategy that was conducted on an annual basis, and many of the studies included individuals without known *BRCA1/2* P/LP variant status. A study of 1219 carriers of a P/LP *BRCA1* variant and 732 carriers of a P/LP *BRCA2* variant showed that the increased sensitivity of mammography in addition to MRI was greater for carriers of a P/LP *BRCA2* variant (12.6%) than for carriers of a P/LP *BRCA1* variant (3.9%).<sup>224</sup> In a retrospective study, a different screening interval was evaluated, using alternating mammography and MRI screening every 6 months in women with a confirmed P/LP *BRCA1/2* variant (N = 73).<sup>225</sup> After a median follow-up of 2 years, 13 breast cancers were detected among 11 women; 12 of the tumors were detected by MRI screening but not by mammography obtained 6 months earlier. The sensitivity and specificity with MRI screening was 92% and 87%, respectively.<sup>225</sup>

The optimal surveillance approach in individuals assigned female at birth who are at high risk for familial breast cancer remains uncertain, especially for those between the ages of 25 and 30 years. Some studies have reported an unlikely association between radiation exposure from mammography and increased risk for breast cancer in carriers of a P/LP *BRCA1/2* variant.<sup>226,227</sup> Thus, one of the potential benefits of incorporating MRI modalities into surveillance strategies may include minimizing the radiation risks associated with mammography, in addition to the higher sensitivity of MRI screening in detecting tumors. The use of MRI, however, may potentially be associated with higher false-positive results and higher costs relative to mammography. The combined use of digital mammography (two-dimensional, 2D) in conjunction with digital breast tomosynthesis (DBT) appears to improve cancer detection and reduce false-positive call back rates.<sup>228-237</sup> Tomosynthesis allows acquisition of three-dimensional (3D) data using a moving x-ray and digital detector. These data are reconstructed using computer algorithms to generate thin sections of images. The combined use of 2D and DBT results in double

the radiation exposure compared with mammography alone. However, this increase in radiation dose falls below dose limits of radiation set by the U.S. Food and Drug Administration (FDA) for standard mammography. The radiation dose can be minimized by newer tomosynthesis techniques that create a synthetic 2D image, which may obviate the need for a conventional digital image.<sup>229,238,239</sup> When mammography is performed, the panel recommends that tomosynthesis be considered. In carriers of a *BRCA1/2* P/LP variant who are younger than 30 years of age, breast MRI screening is preferred over mammography due to the potential radiation exposure risk and less sensitivity for detection of tumors associated with mammography. Studies have reported that deposits of gadolinium, a component of MRI contrast agents, remain in the brain of some patients who undergo 4 or more contrast MRI scans, long after the last administration.<sup>240,241</sup> Retention of gadolinium has also been seen in the bone.<sup>242</sup> In 2017, the FDA issued an update stating that its review of available data had not identified adverse health effects from gadolinium retained in the brain and that patients should read a medication guide prior to receiving gadolinium. However, review of the evidence will continue.

The appropriate imaging modalities and surveillance intervals are still under investigation. In a report based on a computer simulation model that evaluated different annual screening strategies in carriers of a P/LP *BRCA1/2* variant, a screening approach that included annual MRI starting at 25 years of age combined with alternating digital mammography/MRI starting at 30 years of age was shown to be the most effective strategy when radiation risks, life expectancy, and false-positive rates were considered.<sup>243</sup> Future prospective trials are needed to evaluate the different surveillance strategies in individuals at high risk for familial breast cancer. For a woman who is a carrier of a *BRCA1/2* P/LP variant, training in breast awareness with regular monthly practice should begin at 18 years of age, and clinical breast examinations should be conducted every 6 to 12 months, beginning at 25 years of age. Between the ages of 25 and

29 years, the woman should have annual breast MRI screening with contrast (to be performed on days 7–15 of menstrual cycle for premenopausal individuals) or annual mammograms only if MRI is not available. The age to begin screening can be individualized if the family history includes a breast diagnosis prior to 30 years of age.<sup>216,218,221,244,245</sup> Breast MRI screening is preferred over mammogram in the 25- to 29-year age group. High-quality breast MRI screening should consist of the following: dedicated breast coil, ability to perform biopsy under MRI guidance, experienced radiologists in breast MRI, and regional availability. Between 30 and 75 years of age, annual mammogram and breast MRI with contrast should both be done. After 75 years of age, management should be considered on an individual basis. In females treated for breast cancer who have not had bilateral mastectomy, mammography and breast MRI screening with contrast should continue as recommended based on age. Emerging evidence suggests that abbreviated-protocol breast MRI is a screening strategy that warrants further investigation in carriers of a *BRCA1/2* P/LP variant.<sup>246,247</sup>

Carriers of a *BRCA1/2* P/LP variant who were assigned male at birth should have an annual clinical breast examination and undergo training in breast self-examination with regular monthly practice starting at 35 years of age. Data to support breast screening in these individuals are limited. A 12-year longitudinal observational study evaluated the outcomes of mammography screening in 1869 men who were at increased risk of developing breast cancer (ie, personal or family history of breast cancer and/or germline P/LP variant associated with breast cancer, mostly *BRCA1* and *BRCA2*).<sup>248</sup> Node-negative breast cancer was identified in five men (18 per 1000 examinations), which is greater than the cancer detection rates in both average-risk and high-risk women who undergo breast screening. Harboring a P/LP variant ( $n = 47$ ) was associated with breast cancer (OR, 7; 95% CI, 2–29;  $P = .006$ ). Annual mammogram screening in men with gynecomastia may be considered, beginning at age

50 or 10 years before the earliest known breast cancer in the family (whichever comes first).

### Bilateral Total Mastectomy

Two meta-analyses show that prophylactic bilateral mastectomy reduces the risk for breast cancer.<sup>249,250</sup> Only one of these analyses showed that risk-reducing surgery is significantly associated with reduced mortality.<sup>250</sup> Retrospective studies and small prospective studies provide support for concluding that RRM provides a high degree of protection against breast cancer in women with carrying a P/LP *BRCA1/2* variant.<sup>251–254</sup>

It is important that the potential psychosocial effects of RRM are addressed. A 2018 Cochrane review including 20 studies that evaluated psychosocial effects of RRM showed that patients are generally satisfied with their decision, with reported decreases in worry about breast cancer, but negative impacts on body image and sexuality have also been reported. Additional research is needed to further evaluate the psychosocial impact of RRM.<sup>255</sup> RRM is also associated with long-term physical symptoms, such as lower sensitivity to touch, pain, tingling, infection, and edema.<sup>250</sup> Multidisciplinary consultations are recommended prior to surgery and should include the discussions of the risks and benefits of surgery, and surgical breast reconstruction options. Immediate breast reconstruction is an option following RRM, and early consultation with a reconstructive surgeon is recommended for those considering either immediate or delayed breast reconstruction.<sup>256</sup> Nipple-sparing mastectomy has been suggested to be a safe and effective risk reduction strategy for patients carrying a *BRCA1/2* P/LP variant,<sup>257</sup> although more data and longer follow-up are needed.

The NCCN Guidelines Panel supports discussion of the option of RRM for individuals assigned female at birth on a case-by-case basis. Counseling for this risk-reducing surgery should include discussion of extent of cancer risk reduction/protection, risks associated with surgeries, breast



reconstructive options, and management of menopausal symptoms. Since risk of breast cancer remains increased with age in carriers of a *BRCA1/2* P/LP variant,<sup>92</sup> age and life expectancy should also be considered during this counseling, as well as family history. It is important to address the psychosocial and quality-of-life aspects of undergoing risk-reducing surgical procedures.<sup>258</sup>

### Chemoprevention

The use of selective estrogen receptor modulators (ie, tamoxifen, raloxifene) has been shown to reduce the risk for invasive breast cancer in postmenopausal individuals considered at high risk for developing breast cancer, especially ER-positive disease.<sup>259-266</sup> However, only limited data are available on the specific use of these agents in patients with *BRCA1/2* P/LP variants. As previously discussed, patients with *BRCA1/2* P/LP variants who are diagnosed with breast cancer have elevated risks for developing contralateral breast tumors. In one of the largest prospective series of carriers of a P/LP *BRCA1/2* variant evaluated, the mean cumulative lifetime risks for contralateral breast cancer were estimated to be 83% for carriers of a P/LP *BRCA1* variant and 62% for carriers of a P/LP *BRCA2* variant.<sup>95</sup> Patients carrying a P/LP *BRCA1/2* variant who have intact contralateral breast tissue (and who do not undergo oophorectomy or receive chemoprevention) have an estimated 40% risk for contralateral breast cancer at 10 years, though risk is dependent on age of first breast cancer diagnosis.<sup>267</sup> Case-control studies from the Hereditary Breast Cancer Clinical Study Group reported that the use of tamoxifen protected against contralateral breast cancer with an odds ratio (OR) of 0.38 (95% CI, 0.19–0.74) to 0.50 (95% CI, 0.30–0.85) among carriers of a P/LP *BRCA1* variant and 0.42 (95% CI, 0.17–1.02) to 0.63 (95% CI, 0.20–1.50) among carriers of a P/LP *BRCA2* variant.<sup>268,269</sup> This translates to an approximately 45% to 60% risk reduction for contralateral tumors among carriers of a P/LP *BRCA1/2* variant with breast cancer. The data were not consistent in regard to the protective effects of tamoxifen in

the subset of carriers of a P/LP *BRCA1/2* variant who also underwent oophorectomy. In addition, no data were available on the estrogen receptor status of the tumors. An evaluation of the subset of healthy carriers of a P/LP *BRCA1/2* variant in the Breast Cancer Prevention Trial revealed that breast cancer risk was reduced by 62% in carriers of a P/LP *BRCA2* variant receiving tamoxifen relative to placebo (risk ratio, 0.38; 95% CI, 0.06–1.56).<sup>270</sup> However, an analysis of 288 women who developed breast cancer during their participation in this trial showed that tamoxifen use was not associated with a reduction in breast cancer risk in carriers of a P/LP *BRCA1* variant.<sup>270</sup> These findings may be related to the greater likelihood for development of estrogen receptor-negative tumors in carriers of a P/LP *BRCA1* variant relative to carriers of a P/LP *BRCA2* variant. However, this analysis was limited by the very small number of individuals with a P/LP *BRCA1/2* variant (n = 19; 7% of participants diagnosed with breast cancer). Common single-nucleotide polymorphisms have been identified in genes (*ZNF423* and *CTSO*) that are involved in estrogen-dependent regulation of *BRCA1* expression.<sup>271</sup> These gene variants were associated with alterations in breast cancer risk during treatment with selective estrogen receptor modulators, and may eventually pave the way for predicting the likelihood of benefit with these chemopreventive approaches in individual patients.

The aromatase inhibitors (AIs) exemestane and anastrozole have been demonstrated to be effective in preventing breast cancer in postmenopausal women considered to be at high risk of developing breast cancer.<sup>272,273</sup> However, to date, there is little evidence supporting the use of AIs as an effective chemopreventive approach for individuals with a *BRCA1/2* P/LP variant. A retrospective study showed that AIs may reduce the risk of contralateral breast cancer in women with a *BRCA1/2* P/LP variant and ER-positive breast cancer who take AIs as adjuvant therapy, but these data are currently published in abstract form only.<sup>274</sup>

Studies on the effect of oral contraceptive use on breast cancer risk among carriers of a P/LP *BRCA1/2* variant have reported conflicting data. In one case-control study, use of oral contraceptives was associated with a modest but statistically significant increase in breast cancer risk among carriers of a P/LP *BRCA1* variant (OR, 1.20; 95% CI, 1.02–1.40), with breast cancer risk in these carriers being associated with 5 or more years of oral contraceptive use (OR, 1.33; 95% CI, 1.11–1.60), breast cancer diagnosed before 40 years of age (OR, 1.38; 95% CI, 1.11–1.72), and use of oral contraceptives before 1975 (OR, 1.42; 95% CI, 1.17–1.75).<sup>275</sup> Oral contraceptive use was not significantly associated with breast cancer in carriers of a *BRCA2* P/LP variant in this study. In another case-control study, use of oral contraceptives for at least 5 years was associated with a significantly increased risk for breast cancer in carriers of a P/LP *BRCA2* variant (OR, 2.06; 95% CI, 1.08–3.94); results were similar when only the cases with oral contraceptive use on or after 1975 were considered.<sup>276</sup> Oral contraceptive use for at least 1 year was not significantly associated with breast cancer risk in carriers of a P/LP *BRCA1* or *BRCA2* variant in this study. In a third case-control study, the use of low-dose oral contraceptives for at least 1 year was associated with significantly decreased risks for breast cancer among carriers of a P/LP *BRCA1* variant (OR, 0.22; 95% CI, 0.10–0.49;  $P < .001$ ), though not for carriers of a P/LP *BRCA2* variant.<sup>277</sup> Two meta-analyses<sup>278,279</sup> and another case-control study<sup>280</sup> showed that oral contraceptive use is not significantly associated with breast cancer risk in carriers of a P/LP *BRCA1/2* variant.

Differences in the study design employed by these case-control studies make it difficult to compare outcomes between studies, and likely account for the conflicting results. The design of these studies might have differed with regard to factors such as the criteria for defining the “control” population for the study (eg, non-*BRCA1/2* carriers vs. P/LP variant carriers without a cancer diagnosis), consideration of family history of breast or ovarian cancer, baseline demographics of the population studied

(eg, nationality, ethnicity, geographic region, age groups), age of onset of breast cancer, and formulations or duration of oral contraceptives used. Larger prospective trials are needed to elucidate the impact of oral contraceptives on breast cancer risk in carriers of a *BRCA1/2* P/LP variant.

### *Ovarian/Uterine Cancer Risk Management*

#### **Screening**

Studies assessing whether ovarian cancer screening procedures are sufficiently sensitive or specific have yielded mixed results. The UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS), which assessed multimodality screening with transvaginal ultrasound (TVUS) and CA-125 versus either TVUS alone or no screening, showed that multimodality screening is more effective at detecting early-stage cancer; however, after a median of 11 years of follow-up, a significant mortality reduction was not observed.<sup>281,282</sup> In phase II of the UK Familial Ovarian Cancer Screening Study (UK FOCSS), 4348 women with an estimated lifetime ovarian cancer risk no less than 10% underwent ovarian cancer screening via serum CA-125 tests every 4 months (with the risk of ovarian cancer algorithm [ROCA] used to interpret results) and TVUS (annually or within 2 months if abnormal ROCA score).<sup>283</sup> Thirteen patients were diagnosed with ovarian cancer as a result of the screening protocol, with 5 of the 13 patients being diagnosed with early-stage cancer. Sensitivity, positive predictive value, and negative predictive value of the screening protocol for detecting ovarian cancer within 1 year were 94.7%, 10.8%, and 100%, respectively. A third study including 3692 women who were at increased familial/genetic risk of ovarian cancer (ie, known P/LP *BRCA1/2* variant in the family and/or family history of multiple breast and/or ovarian cancers) showed that a ROCA-based screening protocol (ie, serum CA-125 testing every 3 months with annual TVUS annually or sooner depending on CA-125 test results) identified 6 incidental ovarian cancers,

of which 50% were early stage.<sup>284</sup> The results of these studies suggest a potential stage shift when a ROCA-based ovarian cancer screening protocol is followed in high-risk women, though it remains unknown whether this screening protocol impacts survival. For those who have not elected RRSO, which is the recommended risk management option for ovarian cancer in carriers of a P/LP *BRCA1/2* variant (see Discussion below on *Bilateral Salpingo-Oophorectomy*), TVUS and serum CA-125 may be considered at the clinician's discretion starting at 30 to 35 years of age.

### **Bilateral Salpingo-Oophorectomy**

Carriers of a confirmed *BRCA1/2* P/LP variant are at increased risk for both breast and ovarian cancers (including fallopian tube cancer and primary peritoneal cancer).<sup>132,133</sup> Although the risk for ovarian cancer is generally considered to be lower than the risk for breast cancer in carriers of a P/LP *BRCA1/2* variant,<sup>90,91,285</sup> the absence of reliable methods of early detection and the poor prognosis associated with advanced ovarian cancer have lent support for the performance of bilateral RRSO after completion of childbearing.

An observational prospective study of 5,783 women carrying a P/LP *BRCA1/2* variant showed that ovarian cancer is more prevalent in individuals carrying a *BRCA1* (4.2%) P/LP variant than those carrying a *BRCA2* (0.6%) P/LP variant.<sup>286</sup> In carriers of a P/LP *BRCA1* variant, prevalence of ovarian, fallopian tube, and peritoneal cancers found during risk-reducing surgery was 1.5% for those younger than 40 years of age and 3.8% in those between the ages of 40 and 49 years.<sup>286</sup> The highest incidence rate for carriers of a P/LP *BRCA1* variant was observed between the ages of 50 and 59 years (annual risk, 1.7%); for carriers of a P/LP *BRCA2* variant, the highest incidence rate was observed between the ages of 60 and 69 years (annual risk, 0.6%). Therefore, the

recommended age for RRSO should be younger for carriers of a P/LP *BRCA1* variant than for carriers of a P/LP *BRCA2* variant.

The effectiveness of RRSO in reducing the risk for ovarian cancer in carriers of a *BRCA1/2* P/LP variant has been demonstrated in a number of studies. For example, results of a meta-analysis involving 10 studies of carriers of a *BRCA1/2* P/LP variant showed an approximately 80% reduction in the risk for ovarian or fallopian cancer following RRSO.<sup>287</sup> In a large prospective study of women who carried deleterious *BRCA1/2* variants (N = 1079), RRSO significantly reduced the risk for *BRCA1*-associated gynecologic tumors (including ovarian, fallopian tube, or primary peritoneal cancers) by 85% compared with observation during a 3-year follow-up period (HR, 0.15; 95% CI, 0.04–0.56; *P* = .005).<sup>288</sup> An observational study of 5,783 women carrying a P/LP *BRCA1/2* variant showed that risk-reducing oophorectomy reduces risk for ovarian, fallopian, or peritoneal cancer by 80% (HR, 0.20; 95% CI, 0.13–0.30) and all-cause mortality by 77% (HR, 0.23; 95% CI, 0.13–0.39).<sup>286</sup> RRSO reduces mortality at all ages in carriers of a P/LP *BRCA1* variant, but among carriers of a P/LP *BRCA2* variant, RRSO is only associated with reduced mortality in those between the ages of 41 and 60 years.<sup>286</sup>

A 1% to 4.3% residual risk for a primary peritoneal carcinoma has been reported in some studies.<sup>157,287,289-292</sup> An analysis of 36 carriers of a *BRCA1/2* P/LP variant who developed peritoneal carcinomatosis following RRSO showed that 86% were carriers of a *BRCA1* P/LP variant specifically.<sup>293</sup> When comparing to 113 carriers of a P/LP *BRCA1/2* variant who did not develop peritoneal carcinomatosis following RRSO, women who eventually developed peritoneal carcinomatosis were older at time of RRSO (*P* = .025) and had a greater percentage of serous tubal intraepithelial carcinoma (STIC) in their RRSO specimen (*P* < .001), supporting the removal of the fallopian tubes as part of the risk-reducing procedure. Further, an analysis from a multicenter prospective cohort



study (N = 1,083) showed an increased risk for serous and/or serous-like endometrial cancer in women carrying a P/LP *BRCA1* variant who underwent RRSO without hysterectomy.<sup>197</sup>

RRSO may provide an opportunity for gynecologic cancer detection in carriers of a P/LP *BRCA1/2* variant. An analysis of 966 RRSO procedures showed that invasive or intraepithelial ovarian, tubal, or peritoneal neoplasms were detected in 4.6% of carriers of a P/LP *BRCA1* variant and 3.5% of carriers of a P/LP *BRCA2* variant.<sup>294</sup> Carrying a *BRCA1/2* P/LP variant was associated with detection of clinically occult neoplasms during RRSO ( $P = .006$ ).

In early studies, RRSO was reported to reduce the risk for breast cancer in carriers of a P/LP *BRCA1/2* variant.<sup>249,287,291,292,295-298</sup> In the case-control international study by Eisen et al, a 56% (OR, 0.44; 95% CI, 0.29–0.66;  $P < .001$ ) and a 43% (OR, 0.57; 95% CI, 0.28–1.15;  $P = .11$ ) breast cancer risk reduction (adjusted for oral contraceptive use and parity) were reported following RRSO in carriers of a *BRCA1* and a *BRCA2* P/LP variant, respectively.<sup>295</sup> A study comparing breast cancer risk in women carrying a P/LP *BRCA1/2* variant who had undergone RRSO with carriers of these P/LP variants who opted for surveillance only also showed reduced breast cancer risk in women who underwent RRSO (HR, 0.47; 95% CI, 0.29–0.77).<sup>292</sup> These studies were further supported by a meta-analysis that found similar reductions in breast cancer risk of approximately 50% for carriers of a P/LP *BRCA1/2* variant following RRSO.<sup>287</sup>

Results of a prospective cohort study suggested that RRSO may be associated with a greater reduction in breast cancer risk for carriers of a P/LP *BRCA2* variant compared with carriers of a *BRCA1* P/LP variant.<sup>288</sup> Another retrospective analysis including 676 women with stage I or II breast cancer and a P/LP *BRCA1/2* variant showed that oophorectomy was associated with decreased risk of mortality from breast cancer in

carriers of a P/LP *BRCA1* variant (HR, 0.38; 95% CI, 0.19–0.77;  $P = .007$ ), but not in carriers of a P/LP *BRCA2* variant ( $P = .23$ ).<sup>299</sup>

The reduction in breast cancer risk following RRSO was questioned in a prospective cohort study from the Netherlands (N = 822), which did not find a statistically significant difference in breast cancer incidence between carriers of a *BRCA1/2* P/LP variant who opted for an RRSO and women who did not, regardless of whether the P/LP variant was *BRCA1* or *BRCA2*.<sup>300</sup> Study investigators argued that previous study findings showing a 50% decrease in breast cancer risk may have been influenced by bias, specifically inclusion of patients with a history of breast or ovarian cancer in the comparison group and immortal person-time bias. One study that corrected for immortal person-time bias as a result of this analysis continued to find a protective effect of RRSO on breast cancer incidence in carriers of a P/LP *BRCA1/2* variant (HR, 0.59; 95% CI, 0.42–0.82;  $P < .001$ ).<sup>301</sup> Another prospective cohort analysis including 1,289 carriers of a P/LP *BRCA1/2* variant unaffected with breast cancer (196 eventually being diagnosed) also showed that, when RRSO was treated as a time-dependent variable, it was no longer associated with breast cancer risk.<sup>302</sup> A meta-analysis including 19 studies of the association between RRSO and breast cancer risk and mortality showed a protective effect in studies published earlier than 2016, but not in studies published in 2016 or later ( $n = 3$ ).<sup>296</sup>

Results from one of the earlier studies showed that greater reductions in breast cancer risk were observed in women carrying a P/LP *BRCA1* variant who had an RRSO at 40 years of age or younger (OR, 0.36; 95% CI, 0.20–0.64) relative to carriers of a P/LP *BRCA1* variant aged 41 to 50 years who had this procedure (OR, 0.50; 95% CI, 0.27–0.92).<sup>295</sup> A nonsignificant reduction in breast cancer risk was found for women aged 51 years or older, although only a small number of women were included in this group.<sup>295</sup> However, results from another early study also suggested

that RRSO after 50 years of age is not associated with a substantial decrease in breast cancer risk.<sup>291</sup> A 2017 study showed that oophorectomy was not significantly associated with decreased risk of breast cancer in carriers of a P/LP *BRCA1/2* variant ( $N = 3,722$ ).<sup>303</sup> However, stratified analyses in carriers of a P/LP *BRCA2* variant who were diagnosed with breast cancer before 50 years of age showed that oophorectomy was associated with an 82% reduction in breast cancer (HR, 0.18; 95% CI, 0.05–0.63;  $P = .007$ ). The risk reduction in carriers of a P/LP *BRCA1* variant was not statistically significant ( $P = .51$ ). A 2020 study including 853 premenopausal carriers of a P/LP *BRCA1/2* variant showed the opposite; that premenopausal RRSO decreased breast cancer risk in carriers of a *BRCA1* P/LP variant (HR, 0.45; 95% CI, 0.22–0.92), but not in carriers of a *BRCA2* P/LP variant (HR, 0.77; 95% CI, 0.35–1.67).<sup>304</sup> Analysis for this study began observation 6 months after genetic testing to avoid event-free time bias.

A large case series published in 2021 addressed the permanent exposure hypothesis that has potentially dampened the strength of the conclusions drawn from previous studies on the association between RRSO and breast cancer risk reduction.<sup>305</sup> Specifically, some of these earlier studies assumed that this association remains constant each year following RRSO. This study, which included 876 families with a known *BRCA1* or *BRCA2* P/LP variant, showed that RRSO reduced risk of breast cancer within 5 years following the surgery (HR, 0.28; 95% CI, 0.10–0.63 and HR, 0.19; 95% CI, 0.06–0.71, respectively). More than 5 years after RRSO, breast cancer risk reduction diminished but continued to be significant for carriers of a *BRCA1* P/LP variant (HR, 0.64; 95% CI, 0.38–0.97), while the reduction was no longer statistically significant for carriers of a *BRCA2* P/LP variant (HR, 0.99; 95% CI, 0.84–1.00).

Studies suggest a benefit of RRSO on breast cancer risk, but the magnitude of the effect is not well-understood, and evidence is mixed

regarding age at which RRSO should be undertaken, and the specific P/LP variant (*BRCA1* vs. *BRCA2*) carried.

Two systematic reviews showed that hormone replacement therapy (HRT) does not negate the reduction in breast cancer risk associated with the surgery.<sup>306,307</sup> One of these reviews showed that breast cancer risk tended to be lower in women who received estrogen only, compared to estrogen plus progesterone (OR, 0.62; 95% CI, 0.29–1.31).<sup>306</sup> It is important to have a discussion about the potential risks and benefits of HRT in carriers of a P/LP variant following RRSO, given the limitations inherent in nonrandomized studies.<sup>308,309</sup>

Salpingectomy (surgical removal of the fallopian tube with retention of the ovaries) rates are increasing, especially in women younger than 50 years of age.<sup>310</sup> Despite some evidence regarding the safety and feasibility of this procedure,<sup>310,311</sup> more data are needed regarding its efficacy in reducing the risk for ovarian cancer.<sup>258,312</sup> Further, carriers of a P/LP *BRCA1/2* variant who undergo salpingectomy without oophorectomy may not get the reduction in breast cancer risk that research suggests carriers of a P/LP *BRCA1/2* variant who undergo oophorectomy may receive. Therefore, at this time, the panel does not recommend risk-reducing salpingectomy alone as the standard of care in carriers of a P/LP *BRCA1/2* variant. Clinical trials of interval salpingectomy with delayed oophorectomy are ongoing (eg, NCT02321228, NCT01907789).

Some studies suggest a link between *BRCA* P/LP variants and development of serous uterine cancer (primarily with *BRCA1*), although the overall risk for uterine cancer was not increased when controlling for tamoxifen use.<sup>192,193,197</sup> Individuals who undergo hysterectomy at the time of RRSO are candidates for estrogen-alone HRT, which is associated with a decreased risk of breast cancer, compared to combined estrogen and progesterone, which is required when the uterus is left in situ.<sup>313</sup> For patients who choose to undergo RRSO, the provider may discuss the risks

and benefits of concurrent hysterectomy, but more data are needed to determine the magnitude of the association between *BRCA* variants and development of serous uterine cancer.

The NCCN Guidelines Panel recommends RRSO for carriers of a known *BRCA1/2* P/LP variant, typically between 35 and 40 years of age for carriers of a *BRCA1* P/LP variant. Since ovarian cancer onset tends to be later in carriers of a *BRCA2* P/LP variant, it is reasonable to delay RRSO for management of ovarian cancer risk until between 40 and 45 years of age, unless age at diagnosis in the family warrants earlier age for consideration of this prophylactic surgery.<sup>286</sup> Peritoneal washings should be performed at surgery, and pathologic assessment should include fine sectioning of the ovaries and fallopian tubes.<sup>158,159</sup> The protocol published by CAP (2009) can be consulted for details on specimen evaluation.<sup>314</sup> See the NCCN Guidelines for Ovarian Cancer for treatment of findings (available at [www.NCCN.org](http://www.NCCN.org)).

The decision to undergo RRSO is a complex one and should be made ideally in consultation with a gynecologic oncologist, especially when the patient wishes to undergo RRSO before the age at which it is typically recommended. Topics that should be addressed include impact on reproduction, impact on breast and ovarian cancer risk, risks associated with premature menopause (eg, osteoporosis, cardiovascular disease, cognitive changes, changes to vasomotor symptoms, sexual concerns), and other medical issues. The panel recommends that a gynecologic oncologist help patients considering RRSO understand how it may impact quality of life.

### Chemoprevention

With respect to the evidence regarding the effect of oral contraceptives on cancer risks in carriers of a known *BRCA1/2* P/LP variant, case-control studies have demonstrated that oral contraceptives reduced the risk for ovarian cancer by 45% to 50% in carriers of a P/LP *BRCA1* variant and by

60% in carriers of a P/LP *BRCA2* variant.<sup>315,316</sup> Moreover, risks appeared to decrease with longer duration of oral contraceptive use.<sup>316</sup> In a meta-analysis conducted in a large number of carriers of a P/LP *BRCA1/2* variant with (n = 1503) and without (n = 6315) ovarian cancer, use of oral contraceptives significantly reduced the risk for ovarian cancer by approximately 50% for both the carriers of a P/LP *BRCA1* variant (summary relative risk [SRR], 0.51; 95% CI, 0.40–0.65) and carriers of a P/LP *BRCA2* variant (SRR, 0.52; 95% CI, 0.31–0.87).<sup>278</sup> Another meta-analysis including one cohort study (N = 3,181) and three case-control studies (1,096 cases and 2,878 controls) also showed an inverse association between ovarian cancer and having ever used oral contraceptives (OR, 0.58; 95% CI, 0.46–0.73).<sup>279</sup>

### Risk Management for Other Cancers

Screening for prostate cancer starting at 40 years of age is recommended for carriers of a P/LP *BRCA2* variant and should be considered for carriers of a P/LP *BRCA1* variant.<sup>166</sup> See the NCCN Guidelines for Prostate Cancer Early Detection (available at [www.NCCN.org](http://www.NCCN.org)). General melanoma risk management is also indicated, such as annual full body skin exam and minimizing UV exposure. There are no specific screening guidelines for melanoma, though more information can be found at the website for the Skin Cancer Foundation ([www.skincancer.org](http://www.skincancer.org)). Information on pancreas screening can be found below under *Hereditary Pancreatic Cancer*.

### Other P/LP Variants Associated with Breast/Ovarian Cancer

Prior to 2020, the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic focused largely on testing criteria for *BRCA1/2*, *PTEN*, and *TP53* and appropriate risk management for carriers of these P/LP variants. There is now strong evidence that genes beyond *BRCA1/2* confer markedly increased risk of breast and/or ovarian cancers. These genes include *ATM*, *BARD1*, *BRIP1*, *CDH1*,





*CHEK2*, *MSH2*, *MSH6*, *MLH1*, *PMS2*, *EPCAM*, *NBN*, *NF1*, *PALB2*, *RAD51C*, *RAD51D*, and *STK11*. The panel's recommendations for cancer risk management intervention for carriers of P/LP variants associated with breast and/or ovarian cancer risk are based on absolute lifetime risk estimates. Cancer risk management intervention may be recommended when a carrier's absolute risk exceeds that of the average-risk population (ie, 12%–13% for breast cancer and 1%–2% for ovarian cancer, based on SEER registry data<sup>317,318</sup>).<sup>42,319</sup> Strength of the evidence supporting risk estimates should also be evaluated when determining appropriate risk management for carriers of a P/LP variant. For example, prospective cohort studies in a population-based setting can be considered very strong evidence, while limited conclusions can be drawn from case series or studies with small samples.<sup>319</sup>

The investigators of an analysis of breast cancer risk in carriers of moderately penetrant P/LP variants posited that, based on an absolute risk approach, screening with mammography in these carriers should begin when the estimated 5-year risk of developing breast cancer exceeds 1%, consistent with recommendations for the average-risk population.<sup>42</sup> Likewise, breast MRI screening in these carriers should begin when the estimated 5-year risk of developing breast cancer exceeds 2.2%. However, for practical reasons, beginning MRI and mammographic screening at the same time is a reasonable approach. The age at which breast screening is recommended may be impacted by the presence of risk factors such as family history of breast cancer, especially early-onset breast cancer.<sup>42</sup> In those with a family history of early-onset breast cancer, breast screening may begin 5 to 10 years earlier than the youngest breast cancer diagnosis in the family. In individuals assigned female at birth treated for breast cancer who have not had bilateral mastectomy, breast screening should continue as recommended based on age. When mammography is performed, the panel recommends that tomosynthesis be considered. Currently there is insufficient evidence to recommend risk-

reducing mastectomy in carriers of moderately penetrant P/LP variants,<sup>42</sup> though this option may be considered and discussed in the presence of a family history of breast cancer.

Discussion of RRSO may be considered when risk of developing ovarian cancer exceeds that of the average-risk population. The panel uses a threshold of 10% for a recommendation to discuss RRSO. For P/LP variants for which lifetime risk estimates are 5% (eg, *PALB2*), RRSO may be considered based on family history. The decision to carry out RRSO should not be made lightly, given the impact of premature menopause.<sup>42</sup> RRSO is the standard of care for ovarian cancer risk management in carriers of a P/LP variant in an ovarian cancer susceptibility gene. However, some may choose to not receive an RRSO. TVUS and serum CA-125 may be considered at the clinician's discretion, though there is no known benefit (see *BRCA-Related Breast/Ovarian Cancer Syndrome: Risk Management*, above).

The P/LP variants described below may be included concurrently in panel testing (see *Multi-Gene Testing* above). Lower penetrance genes that may be included as part of multi-gene testing but for which there is currently insufficient evidence of an association with breast and/or ovarian cancer include: *FANCC*, *MRE11A*, *MUTYH* heterozygotes, *RECQL4*, *RAD50*, *RINT1*, *SLX4*, *SMARCA4*, and *XRCC2*. Risk management recommendations for these genes should take into account family history and other clinical factors. A more comprehensive review of these lower-penetrance genes is described in another publication.<sup>320</sup>

Information regarding testing criteria and risk management for LFS (associated with germline *TP53* P/LP variant) and Cowden syndrome/PHTS (associated with germline *PTEN* P/LP variant) can be found in their respective sections, below.

### ATM

P/LP variants in the *ATM* (ataxia-telangiectasia mutated) gene may increase risk for breast cancer. A meta-analysis including 19 studies showed that the cumulative lifetime risk for breast cancer in individuals with an *ATM* P/LP variant is 6% by age 50 years and 33% by age 80 years.<sup>321</sup> A meta-analysis of three cohort studies of relatives with ataxia-telangiectasia showed an estimated RR of 2.8 (90% CI, 2.2–3.7;  $P < .001$ ).<sup>322</sup> Other analyses of patients with breast cancer showed that about 1% had an *ATM* P/LP variant.<sup>85,115,118,119,323-326</sup>

The association between specific types of *ATM* genetic variants and breast cancer susceptibility is less clear,<sup>49-52</sup> with some evidence showing that certain missense P/LP variants may act in a dominant-negative fashion to increase cancer risk, relative to truncating P/LP variants.<sup>49,50</sup> A meta-analysis including five studies showed that carriers of an *ATM* P/LP variant have a 38% lifetime risk of developing breast cancer, with carriers of the c.7271T>G missense P/LP variant having a 69% risk of developing breast cancer by 70 years of age.<sup>46</sup> An analysis from a case-control study (42,671 breast cancer cases and 42,164 controls) showed a significant association between the c.7271T>G variant and breast cancer risk (OR, 11.60; 95% CI, 1.50–89.90;  $P = .001$ ).<sup>47</sup> An analysis of 27 families in which P/LP *ATM* variants were identified showed an association between the c.7271T>G variant and increased risk for breast cancer (HR, 8.0; 95% CI, 2.3–27.4;  $P < .001$ ).<sup>48</sup>

The panel recommends annual mammogram for carriers with a P/LP *ATM* variant beginning at 40 years of age, with consideration of annual breast MRI. There are no data on the benefit of risk-reducing mastectomy for carriers of a P/LP *ATM* variant,<sup>42</sup> but this procedure may be considered based on family history. Results of the case-control WECARE study suggested that radiation exposure may be associated with increased risk for contralateral breast cancer in women who are carriers of very rare *ATM*

missense variants.<sup>327</sup> However, these variants are not P/LP, and a meta-analysis including five studies showed that radiation therapy (with conventional dosing) is not contraindicated in patients with a heterozygous *ATM* P/LP variant.<sup>46</sup> Therefore, radiation therapy does not need to be avoided in these carriers who are diagnosed with cancer.

Large studies of patients with ovarian cancer have shown that there may be a slightly increased risk for ovarian cancer in carriers of an *ATM* P/LP variant,<sup>147,324,328,329</sup> but there is currently insufficient evidence to recommend RRSO in these carriers.<sup>319</sup> Given the association between *ATM* and development of the autosomal recessive condition ataxia telangiectasia, counseling for carriers of *ATM* P/LP variants should include a discussion of reproductive options. *ATM* P/LP variants have been found in patients with pancreatic cancer (see *Hereditary Pancreatic Cancer*, below).<sup>187,330</sup>

### BARD1

A modest association between breast cancer and P/LP variants in the *BRCA1*-associated RING domain 1 (*BARD1*) gene has been found in case-control studies with a prevalence rate of 0.1% to 0.51% in patients with breast cancer.<sup>85,323,324,331-333</sup> Studies show that *BARD1* is prevalent in 0.41% to 0.90% of patients with triple-negative breast cancer.<sup>85,117-119</sup> The Breast Cancer Association Consortium and the CARRIERS case-control studies also found associations between a *BARD1* P/LP variant and increased risk of triple-negative breast cancer (0.42%; OR, 9.29; 95% CI, 4.58–18.85 and 0.41%; OR, 3.18; 95% CI, 1.16–7.42, respectively).<sup>118,119</sup> The panel recommends annual mammogram for carriers of a P/LP *BARD1* variant beginning at 40 years of age, with consideration of annual breast MRI. Risk-reducing mastectomy is not recommended in carriers of a *BARD1* P/LP variant, but this procedure may be considered based on family history.

### **BRIP1**

Panel testing of germline DNA in patients with ovarian cancer has shown that the prevalence rate of P/LP variants in the *BRCA1* interaction protein C-terminal helicase 1 gene (*BRIP1*), a Fanconi anemia gene, is about 1%.<sup>147,324,328,329,334</sup> An analysis of 3236 women with epithelial ovarian cancer, 3431 controls, and 2000 unaffected high-risk women from an ovarian cancer screening trial (UKFOCSS) showed that *BRIP1* is associated with an increased risk for ovarian cancer ( $P < .001$ ), with the RR for invasive epithelial ovarian cancer being 11.22 (95% CI, 3.22–34.10;  $P < .001$ ) and 14.09 for high-grade serous disease (95% CI, 4.04–45.02;  $P < .001$ ).<sup>335</sup> An analysis of an Icelandic population (656 ovarian cancer cases, 3913 controls) also showed an association between *BRIP1* and increased risk for ovarian cancer (OR, 8.13; 95% CI, 4.74–13.95;  $P < .001$ ).<sup>336</sup> The cumulative lifetime risk of developing ovarian cancer by 80 years of age in carriers of a *BRIP1* P/LP variant is estimated to be 5.8% (95% CI, 3.6–9.1),<sup>335</sup> though lifetime risk of developing ovarian cancer may also be as high as 12%.<sup>319</sup> The panel recommends that RRSO in carriers of a *BRIP1* P/LP variant be considered beginning at 45 to 50 years of age. A discussion about risk-reducing surgery may be initiated earlier if there is a family history of early-onset ovarian cancer. Ultimately, large prospective trials are needed to make a firm age recommendation regarding when a discussion about RRSO should begin in these variant carriers.

Regarding breast cancer, a case-control study including 10,901 patients with triple-negative breast cancer showed that *BRIP1* was prevalent in 0.43% of cases.<sup>117</sup> The panel has determined that more evidence is needed to provide breast screening recommendations in these carriers. *BRIP1* is associated with Fanconi anemia (group FANCI), inherited in an autosomal recessive manner. Therefore, counseling for carriers of *BRIP1* P/LP variants should include a discussion of reproductive options.

### **CDH1**

Germline P/LP variants in *CDH1* are associated with hereditary diffuse gastric cancer and lobular breast cancer, and studies have reported a cumulative lifetime risk for breast cancer of 39% to 52%.<sup>337-340</sup> Given the considerable risk for lobular breast cancer in carriers of a *CDH1* P/LP variant, the panel recommends screening with annual mammogram (or consideration of breast MRI) beginning at 30 years of age. Alternatively, screening may begin 5 to 10 years earlier than the youngest breast cancer diagnosis in the family. Risk-reducing mastectomy may be discussed with these carriers.

There is controversy over how best to manage gastric cancer risk in individuals harboring a *CDH1* P/LP variant in the absence of a family history of gastric cancer. A small study found that more than half of the individuals with a *CDH1* P/LP variant who lacked a family history of gastric cancer had early-stage signet ring cell adenocarcinoma identified at the time of risk-reducing gastrectomy.<sup>341</sup> See the NCCN Guidelines for Gastric Cancer (available at [www.NCCN.org](http://www.NCCN.org)) for screening recommendations for gastric cancer for individuals with a *CDH1* P/LP variant. A report of two cases showed that *CDH1* P/LP variant may also be associated with cleft lip with or without cleft palate.<sup>342</sup>

### **CHEK2**

Another breast cancer susceptibility gene that has been identified is *CHEK2* (cell cycle checkpoint kinase 2). Panel testing of germline DNA in large samples of patients with breast cancer has shown that the prevalence rate of a *CHEK2* P/LP variant is about 1% to 2%.<sup>323-326,329</sup> Deleterious *CHEK2* P/LP variants have been reported to occur with a higher frequency in Northern and Eastern European countries compared with North America.<sup>320,343-345</sup> The cumulative lifetime risk for breast cancer in women with *CHEK2* P/LP variants and familial breast cancer has been estimated to range from approximately 28% to 37%, and is higher in



women with stronger family histories of breast cancer than in those without.<sup>346,347</sup> The estimated RR for breast cancer, based on data from two large case-control studies, was 3.0 (90% CI, 2.6–3.5).<sup>322</sup> The Breast Cancer Association Consortium and the CARRIERS case-control studies showed associations between a *CHEK2* P/LP variant and increased risk of ER-positive breast cancer (1.58%; OR, 2.67; 95% CI, 2.30–3.11 and 1.11%; OR, 2.60; 95% CI, 2.05–3.31, respectively).<sup>118,119</sup>

Studies investigating the association between breast cancer risk and specific *CHEK2* variants have primarily been based on the truncating variant 1100delC. An analysis from the Copenhagen General Population Study (N = 86,975) showed that *CHEK2* 1100delC heterozygotes had an increased risk for breast cancer when analyses were stratified by age and sex (HR, 2.08; 95% CI, 1.51–2.85).<sup>348</sup> A case-control study (10,860 cases and 9,065 controls) carried out by the CHEK2 Breast Cancer Case-Control Consortium of Europe and Australia showed that the 1100delC variant is associated with increased risk for breast cancer, even in women unselected for family history (OR, 2.34; 95% CI, 1.72–3.20;  $P < .001$ ).<sup>349</sup> Another case-control study (44,777 cases and 42,997 controls) showed that heterozygous 1100delC carriers have a significantly increased risk of developing ER-positive breast cancer (OR, 2.55; 95% CI, 2.10–3.10;  $P < .001$ ), but not ER-negative breast cancer (OR, 1.32; 95% CI, 0.93–1.88;  $P = 0.12$ ).<sup>350</sup> Results from a meta-analysis including 18 case-control studies (26,336 cases and 44,219 controls) showed that the missense variant I157T is associated with a modestly increased risk for breast cancer (OR, 1.58; 95% CI, 1.42–1.75;  $P < .001$ ).<sup>351</sup> There is a lack of data regarding the association between other missense *CHEK2* P/LP variants and risk for breast cancer.

The panel recommends annual mammogram for carriers of a P/LP *CHEK2* variant beginning at 40 years of age, with consideration of annual breast MRI. There are no data on the benefit of risk-reducing mastectomy for

carriers of a P/LP *CHEK2* variant, but this procedure may be considered based on family history.

### ***MLH1, MSH2, MSH6, PMS2, EPCAM***

Lynch syndrome results from a germline P/LP variant in 1 of 4 DNA MMR genes (*MLH1*, *MSH2*, *MSH6*, or *PMS2*).<sup>352</sup> Additionally, deletions in the *EPCAM* gene, which lead to hypermethylation of the *MSH2* promoter and subsequent *MSH2* silencing, cause Lynch syndrome.<sup>353,354</sup> Individuals with Lynch syndrome are at increased risk for endometrial and ovarian cancers (up to 60% and 24%, respectively).<sup>355–359</sup> However, estimates may vary depending on the specific gene. For example, there is less evidence of increased risk for ovarian cancer in carriers of a P/LP *PMS2* variant.<sup>360–362</sup> Estimates also vary for *MSH6*, in which cumulative lifetime risk for ovarian cancer ranges from 1% to 13%.<sup>355,359</sup> Total abdominal hysterectomy and/or bilateral salpingo-oophorectomy are options that may be considered for risk reduction in those who have completed childbearing and carry a mismatch repair gene linked to Lynch syndrome.<sup>363–367</sup> There is no clear evidence to support routine screening for gynecologic cancers in these carriers of these P/LP variants. Annual endometrial sampling may be considered, but the benefit is uncertain.<sup>363,368–371</sup> Routine TVUS and serum CA-125 testing are not endorsed because they have not been shown to be sufficiently sensitive or specific<sup>363,368–372</sup>; however, there may be circumstances where these tests may be helpful.

Some studies have suggested that some of the mismatch repair genes linked to Lynch syndrome (*MLH1* and *MSH2*) may be associated with increased risk for breast cancer.<sup>373,374</sup> However, there is currently not enough evidence for the panel to recommend breast screening for individuals with Lynch syndrome beyond that which is recommended for the average-risk population.

Patients of reproductive age should be advised regarding their options for prenatal diagnosis and assisted reproduction, including PGT. This

discussion should include known risks, limitations, and benefits of these technologies. If both partners are a carrier of a P/LP variant in the same MMR gene (eg, if both partners carry a P/LP variant in the *PMS2* gene), then they should also be advised about the risk for constitutional MMR deficiency (CMMRD) syndrome, a rare recessive syndrome.<sup>375</sup> More information regarding Lynch syndrome can be found in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal (available at [www.NCCN.org](http://www.NCCN.org)).

### NBN

The *NBN* gene is responsible for producing the protein nibrin. An association between the *NBN* gene and breast cancer risk has previously been suggested.<sup>323,324,329</sup> Two meta-analyses showed a significant association between the variant 657del5 and breast cancer risk, with ORs ranging from 2.42 to 2.66.<sup>376,377</sup> However, two more recent and much larger case-control studies have suggested no established association between *NBN* and breast cancer risk.<sup>118,119</sup> Given the limited evidence, the panel does not recommend breast cancer risk management for carriers of an *NBN* P/LP variant beyond what is recommended for the general population, including carriers of a 657del5 P/LP variant.

Some studies have shown a potential increase in ovarian cancer risk in carriers of an *NBN* P/LP variant, with ORs ranging from 1.85 to 2.30.<sup>147,324,378</sup> A recent study including 6001 patients with ovarian cancer found an *NBN* P/LP variant in 0.35%.<sup>329</sup> There is currently insufficient evidence to recommend RRSO in these carriers at this time. The *NBN* gene is associated with development of the autosomal recessive condition Nijmegen breakage syndrome, which is characterized by hypergonadotropic hypogonadism in those assigned female at birth, microcephaly, growth failure early in life, respiratory infections, and childhood lymphomas. Therefore, counseling for carriers of *NBN* P/LP variants should include a discussion of reproductive options.

### NF1

Neurofibromatosis type 1 (NF1) is an autosomal dominant hereditary cancer syndrome that is caused by an *NF1* P/LP variant. NF1 is a neurocutaneous syndrome characterized by café-au-lait spots and axillary/inguinal freckling, associated with non-cancerous tumors of the nerve tissues. Individuals with NF1 have an increased risk for malignant peripheral nerve sheath tumors, other CNS tumors, and gastrointestinal stromal tumors.<sup>379-383</sup> A population-based study in Finland of 1404 patients with NF1 showed an estimated lifetime cancer risk of 59.6%.<sup>379</sup> This study showed a significant association between NF1 and increased risk for breast cancer (SIR, 3.04; 95% CI, 2.06–4.31;  $P < .001$ ). Among patients with breast cancer, NF1 was associated with poorer survival, with 5-year survival rates for patients with NF1 being 67.9%, compared to 87.8% in patients without NF1. Excess incidence was highest in women younger than 40 years of age (SIR, 11.10; 95% CI, 5.56–19.50;  $P < .001$ ). A population-based study in England of 848 patients with NF1 also showed an increased risk for breast cancer (SIR, 3.5; 95% CI, 1.9–5.9), especially among women younger than 50 years (SIR, 4.9; 95% CI, 2.4–8.8).<sup>384</sup>

Given the increased risk for early-onset breast cancer in carriers of these P/LP variants, annual breast screening with mammography should begin at 30 years of age.<sup>383,385</sup> Screening with breast MRI could also be considered. These screening recommendations apply only to individuals with a clinical diagnosis of NF1. The presence of neurofibromas in the breast may lead to false-positive MRI results, but more data are needed to determine the sensitivity and specificity of breast MRI in individuals with NF1. A prospective study of patients with NF1 from the United Kingdom ( $N = 448$ ) showed that breast cancer risk in carriers of these P/LP variants is not significantly increased at 50 years of age and beyond.<sup>382</sup> Case-control analyses of women with NF1 from England showed that RR estimates for women aged 30 to 39 years was 6.5 (95% CI, 2.6–13.5) and 4.4 for women aged 40 to 49 years (95% CI, 2.5–7.0).<sup>386</sup> RR estimates then drop



for women aged 50 to 59 years (RR, 2.6; 95% CI, 1.5–4.2) and continue to drop as age increases (RR, 1.9; 95% CI, 1.0–3.3 for women aged 60–69 years and RR, 0.8; 95% CI, 0.2–2.2 for women aged 70–79 years). These studies show that, beginning at age 50, breast cancer risk in women with NF1 may not significantly differ from that of women in the general population. Therefore, breast MRI screening in patients with NF1 may be discontinued at 50 years of age. There are no data regarding the benefit of risk-reducing mastectomy for carriers of *NF1* P/LP variants. Therefore, risk-reducing mastectomy is not recommended in these patients, but this procedure may be considered based on family history. Complications related to NF1 (eg, neurologic complications) may appear early in life, and these have the potential to be severe.<sup>387</sup> Therefore, referral to a neurofibromatosis specialist for management is recommended.<sup>383</sup>

### **PALB2**

*PALB2* (partner and localizer of *BRCA2*) is a Fanconi anemia gene. *PALB2* P/LP variants are associated with increased risk for breast cancer, with studies of patients with breast cancer showing that 0.4% to 3% harbor a *PALB2* P/LP variant.<sup>85,114,323–326,329,331,388,389</sup> A meta-analysis of three studies estimated an RR of 5.3 (90% CI, 3.0–9.4),<sup>322</sup> while an analysis including 524 families with a known P/LP *PALB2* variant estimated an RR of 7.18 (95% CI, 5.82–8.85) for female breast cancer.<sup>84</sup> The Breast Cancer Association Consortium study and the CARRIERS study showed associations between a *PALB2* P/LP variant and increased risk of triple-negative breast cancer (1.02%; OR, 10.36; 95% CI, 6.42–16.71 and 1.64%; OR, 13.03; 95% CI, 7.08–23.75, respectively).<sup>118,119</sup> *PALB2* P/LP variant is associated with a 33% to 53% lifetime risk of breast cancer.<sup>83–85</sup> The risk increases with increasing number of relatives affected with breast cancer. The analysis, which included 524 families with a known P/LP *PALB2* variant, showed that lifetime risk of breast cancer is as high as 76% when there is a family history of two first-degree relatives with breast cancer.<sup>84</sup> In a study of patients with breast cancer from Poland who

underwent genetic testing, contralateral breast cancer was reported in 10% of *PALB2* carriers.<sup>389</sup> This study also showed that 10-year survival among *PALB2* carriers with breast cancer was 48%, compared to 72% in carriers of a *BRCA1* P/LP variant and 75% in non-carriers ( $P < .001$ ).

The panel recommends annual mammogram for carriers of a *PALB2* P/LP variant assigned female at birth beginning at 30 years of age. Breast MRI screening may also be considered. Risk-reducing mastectomy for carriers of a *PALB2* P/LP variant may be considered based on family history. Some studies suggest an association between *PALB2* and increased ovarian cancer risk.<sup>147,378,390</sup> The most robust data to date showing an association between *PALB2* and increased ovarian cancer risk comes from the international study, which included 524 families with a known P/LP *PALB2* variant.<sup>84</sup> This study showed a 5% lifetime risk of ovarian cancer in carriers of a *PALB2* P/LP variant. Evidence is currently insufficient to recommend RRSO in all carriers of these P/LP variants, but it may be considered if there is a family history of ovarian cancer. *PALB2* is associated with Fanconi anemia, inherited in an autosomal recessive manner.<sup>391</sup> Therefore, counseling for carriers of *PALB2* P/LP variants should include a discussion of reproductive options.

### **RAD51C and RAD51D**

Genes in the *RAD51* protein family are involved in homologous recombination and DNA repair. *RAD51C* and *RAD51D* have been shown to be associated with increased risk for ovarian cancer. Panel testing of germline DNA in women with ovarian cancer has shown that the prevalence rate of the *RAD51C* or *RAD51D* P/LP variant is about 1%.<sup>147,324,328,334</sup> In a comparison of 1132 probands with a family history of ovarian cancer and 1156 controls, *RAD51C* was associated with an increased risk for ovarian cancer (RR, 5.88; 95% CI, 2.91–11.88;  $P < .001$ ).<sup>392</sup> Analyses from the same trial (911 probands and 1060 controls) also showed an association between *RAD51D* and increased risk for

ovarian cancer (RR, 6.30; 95% CI, 2.86–13.85;  $P < .011$ ).<sup>393</sup> In a case-control analysis of 3429 women with epithelial ovarian cancer and 2772 controls, both *RAD51C* (OR, 5.2; 95% CI, 1.1–24;  $P = .035$ ) and *RAD51D* (OR, 12.0; 95% CI, 1.5–90;  $P = .019$ ) were associated with an increased risk for ovarian cancer.<sup>394</sup> A study including 6178 and 6690 families with a known P/LP *RAD51C* and *RAD51D* variant, respectively, showed that the cumulative risk of developing ovarian cancer by age 80 was 11% for carriers of a *RAD51C* P/LP variant and 13% for carriers of a *RAD51D* P/LP variant.<sup>395</sup>

The panel recommends that RRSO in carriers of *RAD51C* and *RAD51D* P/LP variants be considered beginning at 45 to 50 years of age. A discussion about risk-reducing surgery may be initiated earlier if there is a family history of early-onset ovarian cancer. As with *BRIP1* P/LP variants, and large prospective trials are needed to make a firm age recommendation regarding when a discussion about RRSO should begin in carriers of *RAD51C* and *RAD51D* P/LP variants.<sup>319</sup>

Regarding breast cancer, studies have shown prevalence rates of 0.23% to 0.45% for *RAD51C* and 0.29% to 0.38% for *RAD51D* in patients with triple-negative breast cancer.<sup>114,117,396</sup> Case-control analyses from a large study including 56,480 breast tumors showed that both *RAD51C* and *RAD51D* P/LP variants ( $n = 68$  and  $n = 29$ , respectively) were significantly associated with triple-negative disease (OR, 4.5; 95% CI, 2.61–7.50 for *RAD51C* and OR, 4.14; 95% CI, 1.80–7.04 for *RAD51D*).<sup>85</sup> The Breast Cancer Association Consortium study and the CARRIERS study showed associations between increased risk of ER-negative breast cancer and both *RAD51C* P/LP variant (OR, 3.99; 95% CI, 2.20–7.26 and OR, 2.19; 95% CI, 0.97–4.49, respectively) and *RAD51D* P/LP variant (OR, 2.92; 95% CI, 1.47–5.78 and OR, 3.93; 95% CI, 1.40–10.29, respectively), with prevalence rates of 0.26% and 0.24% for *RAD51C*, respectively, and 0.17% and 0.18% for *RAD51D*, respectively.<sup>118,119</sup> The panel asserts that

there is currently insufficient evidence to recommend breast cancer screening in carriers of these variants. *RAD51C* is associated with Fanconi anemia, inherited in an autosomal recessive manner. Therefore, counseling for carriers of a *RAD51C* P/LP variant should include a discussion of reproductive options.

### STK11

Germline *STK11* P/LP variants are associated with PJS, an autosomal dominant disorder characterized by gastrointestinal polyps, mucocutaneous pigmentation, and elevated risk for gastrointestinal cancers as well as breast or non-epithelial ovarian cancers, such as Sertoli-Leydig tumors. Breast cancer risk in women with PJS is 8% at 40 years of age, 13% at 50 years of age, 31% at 60 years of age, and 45% at 70 years of age.<sup>155</sup> There are no data on the benefit of risk-reducing mastectomy for carriers of *STK11* P/LP variants. Therefore, risk-reducing mastectomy is not recommended in these patients, but this procedure may be considered based on family history. Absolute risk of developing non-epithelial ovarian cancer (sex cord with annular tubules) is 18% to 21%.<sup>154,155</sup> Information regarding screening for patients with PJS can be found in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal (available at [www.NCCN.org](http://www.NCCN.org)).

### NCCN Genetic Testing Criteria

The NCCN genetic testing criteria for high-penetrance breast, ovarian, pancreatic, and prostate cancer are organized into three sections: 1) testing is clinically indicated; 2) testing may be considered; and 3) there is a low probability of testing results having documented clinical utility (ie, finding of high-penetrance genes). The testing criteria listed are for cancer susceptibility genes with strong or moderate evidence of actionability for breast, ovarian, pancreatic, and prostate cancer (eg, *BRCA1/2*, *CDH1*, *PALB2*, *PTEN*, and *TP53* for breast cancer; additionally, testing criteria for Li-Fraumeni syndrome and Cowden syndrome continue to be contained in

their own dedicated sections; see below). Included genes may change with emerging clinical data. Further, the personal and/or family history criteria included may suggest the possibility of additional syndromes and would necessitate additional unlisted genes to be evaluated.

The NCCN Panel recommends that individuals from a family with a known P/LP variant in a breast, ovarian, pancreatic, and/or prostate cancer susceptibility gene be tested for the known variant. In individuals from a family without a known P/LP variant, germline multigene testing is recommended for those individuals who meet the testing criteria described in the *Hereditary Cancer Testing Criteria* section in the algorithm. Multi-gene testing may be considered for individuals who meet testing criteria and who previously underwent single-gene and/or absent deletion/duplication analysis but tested negative. Both first- and second-degree relatives of individuals who meet these testing criteria are also eligible for testing, except for second-degree relatives of individuals with pancreatic cancer or prostate cancer, for whom prior probability of a high-penetrance cancer susceptibility gene is low in the absence of additional family history of cancer; only first-degree relatives of these affected individuals should be offered testing, unless indicated based on additional family history.

### Testing Criteria Related to Prostate Cancer

Approximately 11% of patients with prostate cancer and at least 1 additional primary cancer carry germline P/LP variants associated with increased cancer risk.<sup>397</sup> As described above, germline *BRCA1/2* P/LP variants are associated with increased risk for prostate cancer (see *BRCA-Related Breast/Ovarian Cancer Syndrome*, above).<sup>163-166</sup> *ATM* P/LP variants have been found in patients with prostate cancer,<sup>165,166,172,397,398</sup> but there is currently insufficient evidence to recommend prostate screening for these cancers in carriers of an *ATM* P/LP variant.<sup>166</sup> *HOXB13* P/LP variants have also been found in 1.4% to 4.5% of men with prostate cancer.<sup>165,397,399</sup> Prostate tumors with intraductal or cribriform

histology may be more likely to harbor somatic and/or germline MMR gene alterations than those with adenocarcinoma histology.<sup>400,401</sup> Intraductal histology specifically is common in patients with prostate cancer who carry a germline *BRCA2* P/LP variant.<sup>402</sup> By definition, intraductal carcinoma includes cribriform proliferation of malignant cells, as long as they remain confined to a preexisting gland that is surrounded by basal cells. These features are seen frequently with an adjacent invasive cribriform component and would be missed without the use of basal cell markers.

For the 2021 Guidelines update, the panel expanded testing criteria related to prostate cancer. Specifically, cribriform histology is now a testing criterion, and any patient in the high- or very high-risk stratification group as defined in the NCCN Guidelines for Prostate Cancer (available at [www.NCCN.org](http://www.NCCN.org)) is eligible for testing without any additional testing criteria. References to high-grade (Gleason score  $\geq 7$ ) prostate cancer have been removed, and patients with any prostate cancer diagnosis (including very-low- and low-risk disease as defined in the NCCN Guidelines for Prostate Cancer, available at [www.NCCN.org](http://www.NCCN.org)) meet testing criteria if additional family history criteria are met or if the patient is of Ashkenazi Jewish descent. A diagnosis of metastatic prostate cancer without additional personal or family history of cancer continues to be a testing criterion.

### Systemic Therapy Decision-Making

Some of the NCCN treatment guidelines for *BRCA*-related cancers (Breast, Ovarian, Pancreatic Adenocarcinoma, Prostate; available at [www.NCCN.org](http://www.NCCN.org)) now recommend treatment with PARP (poly ADP-ribose polymerase) inhibitors for patients with germline or somatic *BRCA1/2* P/LP variants, as PARP inhibitors have been demonstrated to be active in these patients. These agents include olaparib<sup>403,404</sup> and talazoparib<sup>405</sup> for HER2-negative metastatic and HER2-negative high-risk early-stage (olaparib only) breast cancer; niraparib,<sup>406</sup> olaparib,<sup>407,408</sup> and rucaparib<sup>409</sup> for



chemotherapy-refractory ovarian cancer; olaparib<sup>410</sup> and rucaparib<sup>411</sup> for metastatic castration-resistant prostate cancer that has progressed following previous treatment; and olaparib as a maintenance therapy option for metastatic pancreatic cancer.<sup>412</sup> Even though the focus of these Guidelines continues to be on management of breast, ovarian, and/or pancreatic cancer risk in individuals with associated hereditary syndromes, the Guidelines now identify intent to aid in systemic therapy and surgical decision-making as a scenario in which germline testing is clinically indicated. If a P/LP variant is detected through tumor profiling that has clinical implications if identified in the germline, then germline testing for this variant is indicated.

### **Ashkenazi Jewish Ancestry**

The rate of the three founder P/LP variants in those of Ashkenazi Jewish ancestry is 2.2% to 2.5%.<sup>413-415</sup> Studies have shown that genetic testing based on clinical guidelines emphasizing family history of breast, ovarian, pancreatic, prostate, or other cancers missed about 38% to 56% of P/LP variant carriers in those of Ashkenazi ancestry.<sup>413,414,416,417</sup> Therefore, there is some evidence to support population-based genetic testing for individuals with Ashkenazi Jewish ancestry. However, there are concerns about the demand on genetic counseling resources, the preparedness of health care professionals to provide cancer genetic counseling and management, and participants' fears and concerns about testing, including those regarding privacy, stigmatization, and the need for appropriate medical and or surgical management in patients and family members found to have a founder P/LP variant. Thus, universal testing for founder *BRCA1/2* P/LP variants in individuals of Ashkenazi Jewish ancestry, regardless of personal or family history, should be offered primarily in the setting of longitudinal research studies. If there is no access to longitudinal studies, then testing may be offered when pre- and post-test genetic counseling are available (see above). There remains a vital need for longitudinal data from research studies exploring various methods of

providing population-based genetic testing of individuals with Ashkenazi Jewish ancestry in the United States.

### **Breast Cancer Population Testing**

In 2019, the American Society of Breast Surgeons published a consensus statement recommending genetic testing for all patients with breast cancer.<sup>418</sup> This recommendation was based on studies showing that criteria in testing guidelines miss some patients with breast cancer who harbor a P/LP variant<sup>419,420</sup> and that population-based multi-gene testing is more cost-effective than testing based on personal and family history criteria.<sup>36,421</sup> However, only 4.4% of patients with a high-penetrance mutation (*BRCA1/2*, *PALB2*, *TP53*, *PTEN*) were missed in the Beitsch et al study.<sup>419,16</sup> Analyses from studies of postmenopausal patients with breast cancer showed rates of 3.6% to 5.6% harboring a P/LP variant, supporting possible population testing for this group of patients.<sup>422,423</sup> Further analyses of this population have suggested universal testing for those age 65 or younger, as two studies showed that about 7% of these patients harbor a P/LP variant associated with breast cancer.<sup>422,424</sup> A follow-up analysis of one of these studies examined age 60 as a cut-off for universal testing of patients with breast cancer and found that 8.2% of these patients harbor a P/LP variant associated with breast cancer.<sup>425</sup> In this analysis, about 2% of patients diagnosed with breast cancer at age 60 or younger who did not meet other testing criteria harbored a highly penetrant P/LP variant associated with breast cancer. This percentage increased to about 5% when expanding the genes to include *ATM*, *CHEK2*, and *NF1*. It is not likely that patients diagnosed with breast cancer older than age 60 who do not meet other testing criteria will harbor a highly penetrant P/LP variant associated with breast cancer.

Additional tailoring of testing criteria in patients with breast cancer could be done based on histopathology or the presence of multiple primary breast cancers. An analysis of women older than age 65 (N = 26,707)

from population-based case-control studies showed that 3.42% of women with ER-negative breast cancer and 3.01% of women with triple-negative breast cancer harbored a P/LP variant in a high-penetrance breast cancer susceptibility gene (*BRCA1*, *BRCA2*, and *PALB2*).<sup>426</sup> Multiple studies also show that individuals with multiple primary breast cancers may be more likely than individuals with a single breast cancer to harbor a P/LP variant associated with breast cancer (7.1%–13.2% vs. 4.2%–9.4%).<sup>326,427,428</sup>

The panel continues to endorse a risk-stratified approach and does not endorse universal testing of all patients with breast cancer due to limitations of this approach, such as low specificity, shortages in trained genetics health professionals to provide appropriate pre- and post-test genetic counseling, and lack of evidence to support risk management for genes included in many multi-gene panels. Though all patients with breast cancer should be evaluated to determine the appropriateness of germline genetic testing, testing should ultimately be based on patient characteristics, such as those specified in the *Testing Criteria for High-Penetrance Breast Cancer Susceptibility Genes* in the algorithm.<sup>16</sup>

### Probability Models

Decision models developed to estimate the likelihood that a *BRCA1/2* P/LP variant is present include BRCAPRO,<sup>429,430</sup> Penn II,<sup>431</sup> and the Breast and Ovarian Analysis of Disease Incidence and Carrier Estimation Algorithm (BOADICEA).<sup>429</sup> A lifetime risk for breast cancer of 20% to 25% or greater as assessed by models based largely on family history has been used in some guidelines to identify a woman as being at high risk for breast cancer. For example, this risk threshold was used in updates to the American Cancer Society (ACS) guidelines on breast screening, which incorporate MRI.<sup>244,432</sup> Penn II has been validated in families with two or more cases of breast and/or ovarian cancer.<sup>431,433</sup> Therefore, caution should be taken in applying this model to individuals with only one case of breast or ovarian cancer. In addition, this model was developed

specifically to evaluate the likelihood of a *BRCA1/2* P/LP variant, and not the appropriateness of multi-gene testing.

If an individual does not meet the criteria for testing for high-penetrance breast and/or ovarian cancer susceptibility genes that are described above, then testing may be considered in those who are determined to have a 2.5% to 5% probability of harboring a *BRCA1/2* P/LP variant, based on probability models validated for *BRCA1/2* (eg, Tyrer-Cuzick, BRCAPRO, BOADICEA). However, the panel cautions that model estimates vary substantially, and different thresholds may be applied if other genes are utilized in a specific model. If genes other than *BRCA1/2* are to be included in models that evaluate the threshold for testing, then penetrance, clinical actionability, and phenotypic features of cancers associated with these genes should be taken into account. Models that take these parameters into account to determine eligibility and appropriateness of multi-gene testing should be developed and validated. Subgroup analyses of 1075 carriers of a *BRCA1/2* P/LP variant from the Breast Cancer Prospective Family Study Cohort showed that BRCAPRO underpredicted breast cancer risk, but BOADICEA was well-validated.<sup>434</sup> In 2020, the web-based CanRisk tool was developed to apply BOADICEA for clinical use and is now available, though further development and testing is needed to increase acceptability of the tool by clinicians.<sup>435</sup>

### Li-Fraumeni Syndrome

LFS is a rare hereditary cancer syndrome associated with germline *TP53* P/LP variants.<sup>82</sup> It has been estimated to be involved in only about 1% of hereditary breast cancer cases,<sup>436</sup> although results from other studies suggest that germline *TP53* P/LP variants may be more common than previously believed, with estimates of 1 in 5000 to 1 in 20,000.<sup>437,438</sup> The tumor suppressor gene, *TP53*, is located on chromosome 17,<sup>439,440</sup> and the protein product of the *TP53* gene (ie, p53) is located in the cell nucleus and binds directly to DNA. It has been called the “guardian of the genome”



and plays important roles in controlling the cell cycle and apoptosis.<sup>439-441</sup> Germline P/LP variants in the *TP53* gene have been observed in greater than 50% (and in >70% in some studies) of families meeting the classic definition of LFS (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm).<sup>82,437,442</sup> Additional studies are needed to investigate the possibility of other P/LP variants in families meeting these criteria not carrying germline *TP53* P/LP variants.<sup>443</sup>

LFS is a highly penetrant cancer syndrome associated with a high lifetime risk for cancer. An analysis from the NCI Li-Fraumeni Syndrome Study (N = 286) showed a cumulative lifetime cancer incidence of nearly 100%.<sup>444</sup> LFS is characterized by a wide spectrum of neoplasms occurring at a young age. It is associated with soft tissue sarcomas, osteosarcomas (although Ewing sarcoma is less likely to be associated with LFS), premenopausal breast cancer, colon cancer, gastric cancer, adrenocortical carcinoma, bronchoalveolar carcinoma, and brain tumors.<sup>82,437,441,445-451</sup> Sarcoma, breast cancer, adrenocortical tumors, and certain brain tumors have been referred to as the “core” cancers of LFS since they account for the majority of cancers observed in individuals with germline *TP53* P/LP variants, and, in one study, at least one of these cancers was found in one or more members of all families with a germline *TP53* P/LP variant.<sup>437</sup> Hypodiploid acute lymphoblastic leukemia is also strongly associated with LFS.<sup>452,453</sup> While case reports have suggested an association between melanoma and LFS, risk estimates are currently not available.<sup>454,455</sup>

The NCI Li-Fraumeni Syndrome Study (N = 286) showed that the cumulative incidence rates by 70 years of age in women are 54%, 15%, 6%, and 5% for breast cancer, soft tissue sarcoma, brain cancer, and osteosarcoma, respectively.<sup>444</sup> The cumulative incidence rates by age 70 years in men are 22%, 19%, and 11% for soft tissue sarcoma, brain cancer, and osteosarcoma, respectively. Case-control analyses from a

large study including 56,480 breast tumors showed that *TP53* P/LP variants ( $n = 82$ ) were significantly associated with HER2-positive disease, regardless of whether disease was ER-positive (OR, 11.95; 95% CI, 5.84–23.0) or negative (OR, 22.71; 95% CI, 10.45–45.49).<sup>85</sup> These results are supported by two earlier retrospective studies that reported a very high frequency of HER2-positive breast tumors (67%–83% of evaluated breast tumors) among patients with germline *TP53* P/LP variants.<sup>456,457</sup> A cohort study including 45 patients diagnosed with breast cancer and harboring a germline *TP53* P/LP variant showed that 36.1% had triple-positive (HER2+/ER+/PR+) breast cancer.<sup>458</sup> Taken together, results suggest that amplification of HER2 may arise in conjunction with germline *TP53* P/LP variants. This association warrants further investigation, as such patients may potentially benefit from chemoprevention therapies that incorporate HER2-targeted agents.

Individuals with LFS often present with certain cancers (eg, soft tissue sarcomas, brain tumors, adrenocortical carcinomas) in early childhood,<sup>447</sup> and have an increased risk of developing multiple primary cancers during their lifetimes.<sup>459</sup> Results of a segregation analysis of data collected on the family histories of 159 patients with childhood soft tissue sarcoma showed carriers of germline *TP53* P/LP variants to have estimated cancer risks of approximately 60% and 95% by 45 and 70 years, respectively.<sup>460</sup> Although similar cancer risks are observed in males and females with LFS when gender-specific cancers are not considered, breast cancer in those assigned female at birth is commonly associated with the syndrome.<sup>437</sup> It is important to mention that estimations of cancer risks associated with LFS are limited to at least some degree by selection bias since dramatically affected kindreds are more likely to be identified and become the subject of further study.

A number of different sets of criteria have been used to help identify individuals with LFS. For the purposes of the NCCN Guidelines, two sets

of these criteria are used to facilitate the identification of individuals who are candidates for testing for *TP53* P/LP variants.

Classic LFS criteria, based on a study by Li and Fraumeni involving 24 LFS kindreds, include the following:<sup>448</sup> a member of a kindred with a known *TP53* P/LP variant; a combination of an individual diagnosed at 45 years of age or younger with a sarcoma and a first-degree relative diagnosed with cancer at 45 years of age or younger; and an additional first- or second-degree relative in the same lineage with cancer diagnosed at younger than 45 years of age or a sarcoma diagnosed at any age (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm). Classic LFS criteria have been estimated to have a high positive predictive value (estimated at 56%) as well as a high specificity, although the sensitivity is relatively low (estimated at 40%).<sup>437</sup> Thus, it is not uncommon for individuals with patterns of cancer outside of these criteria to be carriers of germline *TP53* P/LP variants.<sup>450,461</sup> Classic LFS criteria make up one set of criteria included in the guidelines to guide selection of individuals for *TP53* P/LP variant testing (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm).

Other groups have broadened the classic LFS criteria to facilitate identification of individuals with LFS.<sup>462-464</sup> For example, criteria for *TP53* testing proposed by Chompret and colleagues recommend testing for patients with multiple primary tumors of at least two “core” tumor types (ie, sarcoma, breast cancer, adrenocortical carcinoma, brain tumors) diagnosed at younger than 36 years of age or patients with adrenocortical carcinoma diagnosed at any age, regardless of family history (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm).<sup>463</sup> The Chompret criteria have an estimated positive predictive value of 20% to 35%,<sup>437,463</sup> and, when incorporated as part of *TP53* testing criteria in conjunction with classic LFS criteria, have been shown to improve the sensitivity to 95% (ie, the Chompret criteria added to classic LFS criteria detected 95% of

patients with *TP53* P/LP variants).<sup>437</sup> The Chompret criteria are the second set of criteria included in the NCCN Guidelines. Although not part of the original published criteria set forth by Chompret et al, the panel recommends adopting the 2015 Revised Chompret Criteria and testing individuals with choroid plexus carcinoma or rhabdomyosarcoma of embryonal anaplastic subtype diagnosed at any age and regardless of family history (for inclusion in criterion 3), based on reports of considerable incidence of *TP53* P/LP variants found in patients with these rare forms of cancer.<sup>437,446,465-467</sup> The panel supports the broader age cut-offs proposed by Tinat et al, based on a study in a large number of families, which detected germline *TP53* P/LP variants in affected individuals with later tumor onsets.<sup>465,467</sup>

Patients with early-onset breast cancer (age of diagnosis ≤30 years) who were assigned female at birth, with or without family history of core tumor types, are another group for whom *TP53* gene P/LP variant testing may be considered.<sup>466</sup> Several studies have investigated the likelihood of a germline *TP53* P/LP variant in this population.<sup>437,465,468-471</sup> Among women younger than 30 years of age with breast cancer and without a family history, the incidence of *TP53* P/LP variants has been reported at 3% to 8%.<sup>437,469,471,472</sup> Other studies have found an even lower incidence of germline *TP53* P/LP variants in this population. For example, Bougeard et al reported that only 0.7% of unselected women with breast cancer before 33 years of age were carriers of a germline *TP53* P/LP variant.<sup>465</sup> Furthermore, Ginsburg and colleagues found no germline *TP53* P/LP variants in 95 unselected women with early-onset breast cancer who previously tested negative for *BRCA1/2* P/LP variants.<sup>468</sup> When taking into account family history of LFS-associated tumors, the *TP53* germline P/LP variant prevalence increases. For example, in a study including 83 patients with *BRCA1/2* P/LP variant-negative early-onset breast cancer (age of diagnosis ≤35 years), deleterious *TP53* P/LP variants were identified in 3 of 4 patients (75%) with a family history of at least 2 LFS-

associated tumors (breast cancer, bone or soft tissue sarcoma, brain tumors, or adrenocortical carcinoma) and in 1 of 17 patients (6%) with a family history of breast cancer only.<sup>470</sup> In another study, all women younger than 30 years of age with breast cancer who had a first- or second-degree relative with at least one of the core cancer types (n = 5) had germline *TP53* P/LP variants.<sup>437</sup>

A member of a family with a known *TP53* P/LP variant is considered to be at sufficient risk to warrant variant testing, even in the absence of any other risk factors. Individuals not meeting testing criteria should be followed according to recommendations tailored to his/her personal cancer history and family history, and testing for other hereditary syndromes may be considered. When *TP53* is included on multigene panels, testing criteria for LFS do not need to be met. If a *TP53* P/LP variant is detected through tumor profiling, and there are clinical implications if a *TP53* P/LP variant is identified in the germline, then germline testing for a *TP53* variant may be considered, depending on a careful examination of the individual's personal and family history. *TP53* P/LP variants are common in tumors.<sup>473,474</sup> Therefore, if a *TP53* somatic P/LP variant is found in the absence of paired germline analysis, then germline testing may not be warranted unless there is clinical suspicion of a germline P/LP variant.

### **Risk Assessment, Counseling, and Management**

The approach to families with other hereditary breast cancer syndromes, such as LFS, reflects that of hereditary breast/ovarian cancer in many ways. However, there are some syndrome-specific differences with regard to assessment and management. In the case of LFS, there are multiple associated cancers, both pediatric and adult, that should be reflected in the expanded pedigree (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm). Cancers associated with LFS include but are not limited to premenopausal breast cancer, bone and soft tissue sarcomas, CNS tumor, adrenocortical carcinoma, hypodiploid acute lymphoblastic

leukemia, unusually early onset of other adenocarcinomas, or other childhood cancers.<sup>437,453,459,466</sup> Verification of these sometimes very rare cancers is particularly important.

Employment of a screening protocol that includes MRI may improve early cancer detection in individuals with LFS.<sup>475</sup> In 2017, the panel made revisions to the LFS management recommendations following revisions to the "Toronto protocol," screening recommendations developed by a multi-institutional group of experts.<sup>476</sup> NCCN recommendations for management of LFS apply specifically to adults with LFS, and discussions with patients should address the limitations of screening for the many cancers associated with this syndrome. Pediatricians should be made aware of the risk for childhood cancers in affected families and review with these families the screening recommendations for children with LFS.<sup>476</sup> It is also important to address the psychosocial and quality-of-life aspects of this syndrome. Given the complexity of LFS management, individuals with LFS should be followed at centers with expertise in management of this syndrome.

For those at risk for breast cancer, training and education in breast self-examination should start at 18 years of age, with the patient performing regular self-examination on a monthly basis. For members of families with LFS, breast cancer surveillance by clinical breast examination is recommended every 6 to 12 months, beginning at 20 years of age (or at the age of the earliest known breast cancer in the family, if younger than 20 years of age) because of the very early age of breast cancer onset seen in these families. Recommendations for breast screening in LFS are similar to those for *BRCA*-related breast and ovarian cancer syndrome management, although screening is begun at an earlier age. They include annual breast MRI screening with contrast (preferred) or mammogram if MRI is not available for individuals assigned female at birth aged 20 to 29 years; annual mammogram and breast MRI screening with contrast in



individuals assigned female at birth aged 30 to 75 years; and management on an individual basis for individuals assigned female at birth older than 75 years. For individuals assigned female at birth who have a family history of breast cancer diagnosed earlier than 20 years of age, breast MRI screening with contrast may begin at the earliest age of diagnosis. In patients assigned female at birth who were treated for breast cancer and who have not had bilateral mastectomy, mammography and breast MRI screening with contrast should continue as recommended based on age. When mammography is performed, the panel recommends that tomosynthesis be considered. As with carriers of a *BRCA1/2* P/LP variant, breast MRI screening in individuals assigned female at birth who are younger than 30 years of age is preferred over mammography due to the potential radiation exposure risk and less sensitivity for detection of tumors.

Although there are no data regarding risk reduction surgery in individuals with LFS who were assigned female at birth, options for risk-reducing mastectomy should be discussed on a case-by-case basis. Counseling for risk-reducing surgeries may include discussion of extent of cancer risk reduction/protection, risks associated with surgeries, degree of age-specific cancer risk, reconstructive options, and competing risks from other cancers. Family history and life expectancy should also be considered during this counseling.

Many of the other cancers associated with germline *TP53* P/LP variants do not lend themselves to early detection. Thus, additional recommendations are general and include comprehensive physical examinations (including neurologic examination) every 6 to 12 months, especially when there is a high index of suspicion for second malignancies in cancer survivors and rare cancers (see *Testing Criteria for Li-Fraumeni Syndrome* in the algorithm). Clinicians should address screening limitations for other cancers associated with LFS. Colonoscopy and upper

endoscopy should be done every 2 to 5 years, starting at 25 years of age, or 5 years before the earliest known colon or gastric cancer diagnosis in family history (whichever comes first). Education regarding signs and symptoms of cancer is important. Patients should be advised about the risk to relatives, and genetic counseling for relatives is recommended. Annual dermatologic examination should be done beginning at 18 years of age.

Whole-body MRI for screening of cancers associated with LFS is being evaluated in multiple international trials. Use of whole-body MRI is appealing due to its wide anatomic coverage and the potential to cut down on the number of imaging studies that a patient undergoes.<sup>477</sup> A meta-analysis including 578 individuals with *TP53* P/LP variants across 13 prospective cohorts showed that baseline whole-body MRI identified cancer in 7% of the sample, with 83% of the cancers being localized and able to treat with curative intent.<sup>478</sup> In a prospective observational study, a clinical surveillance protocol for carriers of a *TP53* P/LP variant from families affected by LFS was incorporated.<sup>479</sup> The surveillance protocol included biochemical methods (ie, bloodwork to evaluate 17-OH-progesterone, total testosterone, dehydroepiandrosterone sulfate, androstenedione, complete blood count [CBC], erythrocyte sedimentation rate, and lactate dehydrogenase; and 24-hour urine cortisol) and imaging techniques, such as annual brain MRI, annual rapid whole-body MRI, ultrasound of the abdomen and pelvis, and colonoscopy.<sup>480</sup> For surveillance of breast cancers, the protocol was similar to the NCCN Guidelines for LFS Management in Adults.<sup>479</sup> Eleven-year follow-up of this study, which included 89 carriers of a *TP53* P/LP variant, showed that this surveillance protocol may be beneficial, with 84% (16 out of 19) of patients who were diagnosed with cancer and had chosen to undergo surveillance being alive at final follow-up, compared to 49% (21 out of 43) of patients who were diagnosed with cancer and had chosen to not undergo surveillance ( $P = .012$ ).<sup>480</sup> Five-year OS was greater for patients

undergoing surveillance (88.8%) compared to patients not undergoing surveillance (59.6%;  $P = .013$ ). The clinical surveillance protocol employed was shown to be feasible, though further evaluation is warranted.<sup>479</sup> Based on these study results the panel recommends annual whole-body MRI as a category 2B recommendation. This is consistent with recommendations described in the Toronto protocol.<sup>476</sup> The panel acknowledges that this surveillance method may not be uniformly available. Patients who do not have access to whole-body MRI should be encouraged to enroll in clinical trials, or alternative comprehensive imaging methods may be used. The panel also acknowledges that whole-body MRI screening of all individuals with LFS may result in false positives and overdiagnosis.<sup>478,481</sup> Further, the utility of whole-body MRI has not been evaluated in individuals with a *TP53* P/LP variant who don't have a classic family history of LFS, a group that is increasingly being identified through multi-gene testing. The brain may be examined as part of whole-body MRI or as a separate exam.

Only very limited data exist on the use of prenatal diagnostics/genetic testing for *TP53* P/LP variants in families with LFS.<sup>482,483</sup> For a general discussion on the topic of reproductive options and counseling considerations, see the section above on *Reproductive Options* under *Genetic Risk Assessment and Counseling*.

### Cowden Syndrome/*PTEN* Hamartoma Tumor Syndrome

The spectrum of disorders resulting from germline P/LP variants in *PTEN*<sup>484</sup> are referred to as PHTS. The spectrum of PHTS includes Cowden syndrome, Bannayan-Riley-Ruvalcaba syndrome (BRRS), adult Lhermitte-Duclos disease (LDD), Proteus-like syndrome,<sup>81,485,486</sup> and autism spectrum disorders with macrocephaly.<sup>81,486,487</sup> Cowden syndrome is rare, with an incidence of 1 in 200,000, although it is likely to be underestimated due to difficulties associated with making a clinical diagnosis of the disease.<sup>488,489</sup> Cowden syndrome is an autosomal dominant disorder, and most cases are associated with germline *PTEN*

P/LP variants, though one study found that germline *KILLIN* methylation may also be associated with this syndrome.<sup>490</sup> The frequency of germline *PTEN* P/LP variant in Cowden syndrome cases is high, at approximately 80%.<sup>491</sup>

Hamartomas (benign tumors resulting from an overgrowth of normal tissue) are a common manifestation of the PHTS syndromes. Cowden syndrome is associated with multiple hamartomatous and/or cancerous lesions in various organs and tissues, including the skin, mucous membranes, breast, thyroid, endometrium, and brain.<sup>81,492</sup> However, it has been suggested that patients with other PHTS diagnoses associated with *PTEN* P/LP variants should be assumed to have Cowden syndrome-associated cancer risks.

The lifetime risk for breast cancer for women diagnosed with Cowden syndrome/PHTS has been estimated at 40% to 60%, with an average age of 38 to 50 years at diagnosis.<sup>81,493</sup> Some studies (as discussed above) have reported a higher cumulative lifetime risk for breast cancer (77%–85%) in individuals with Cowden syndrome/PHTS or *PTEN* P/LP variants.<sup>494-496</sup> There have been only two cases of breast cancer reported in men with Cowden syndrome/PHTS.<sup>493</sup> Although many women with Cowden syndrome/PHTS experience benign breast disease,<sup>81</sup> there is no evidence that the rate is higher than in the general population.<sup>493</sup>

Thyroid disease, including benign multinodular goiter, adenomatous nodules, and follicular adenomas, has been reported to occur in approximately 30% to 68% of adults with *PTEN* P/LP variants,<sup>486,497</sup> and the lifetime risk for thyroid cancer (follicular or papillary) has been estimated at 3% to 10%.<sup>81,498</sup> However, data tend to be aggregated, so it is difficult to calculate rates for multinodular goiter versus solitary nodules.<sup>493</sup> A retrospective chart review of 47 children with *PTEN* P/LP variants showed that 26% had abnormal thyroid imaging.<sup>499</sup> The youngest reported



case of thyroid cancer in a child with Cowden syndrome/PHTS was at age 7.<sup>500</sup>

Macrocephaly (defined as head circumference greater than the 97<sup>th</sup> percentile)<sup>501</sup> is a common finding in patients with Cowden syndrome/PHTS. It has been estimated that approximately 80% to 100% of individuals with this syndrome will exhibit this clinical finding.<sup>493</sup> Adult LDD and autism spectrum disorder characterized by macrocephaly are strongly associated with Cowden syndrome/PHTS.<sup>485,491,495,502</sup> A rare, slow-growing, benign hamartomatous lesion of the brain, LDD, is a dysplastic gangliocytoma of the cerebellum.<sup>81,495</sup> In a multicenter prospective study examining 3042 probands who met clinical criteria for Cowden syndrome/PHTS, 6% met criteria for LDD.<sup>497</sup> In a study of individuals meeting the diagnostic criteria for Cowden syndrome/PHTS, the cumulative lifetime risk for LDD was reported to be 32%.<sup>495</sup> The preponderance of evidence supports a strong association between adult-onset LDD and the presence of a *PTEN* P/LP variant,<sup>491,503</sup> although exceptions have been reported.<sup>504</sup> In addition, there is a relatively large body of evidence to support that 10% to 20% of individuals with autism spectrum disorder and macrocephaly carry germline *PTEN* P/LP variants.<sup>487,505-508</sup>

As in many other hereditary cancer syndromes, affected individuals are more likely to develop bilateral and multifocal cancer in paired organs.<sup>491</sup> Although not well defined, women with Cowden syndrome/PHTS may have a 5% to 10% risk for endometrial cancer.<sup>81,509</sup> <sup>493</sup> While many women with Cowden syndrome/PHTS may also have uterine fibroids, this risk is not likely to be much greater than in women without Cowden syndrome/PHTS or *PTEN* P/LP variant.<sup>493</sup>

In addition, brain tumors and vascular malformations affecting any organ are occasionally seen in individuals with Cowden syndrome/PHTS, although the risks for developing these conditions are not well

defined.<sup>81,493</sup> It is important to note, however, that most of the data on the frequencies of the clinical features of Cowden syndrome/PHTS are from compilations of case reports of relatively young individuals who may have subsequently developed additional signs of the disease (ie, new cancerous lesions), and these data are also likely to be confounded by selection bias.<sup>81</sup> Furthermore, a considerable number of these studies were published prior to the establishment in 1996 of the International Cowden Consortium operational diagnostic criteria for the syndrome, which were based on published data and the expert opinion of individuals representing a group of centers mainly in North America and Europe.<sup>81,510</sup>

Benign skin lesions are experienced by most to all Cowden syndrome/PHTS patients.<sup>486,492,499</sup> Skin lesions associated with Cowden syndrome/PHTS include trichilemmomas (ie, benign tumors derived from the outer root sheath epithelium of a hair follicle), oral papillomas, mucocutaneous neuromas (hamartoma of the peripheral nerve sheath), palmoplantar keratoses, penile pigmentation, lipomas and vascular anomalies, and fibromas.<sup>493,499,511</sup> Trichilemmomas associated with Cowden syndrome/PHTS tend to appear on the face, particularly the eyes, mouth, nose, and forehead.<sup>493</sup> Most individuals with Cowden syndrome/PHTS exhibit characteristic mucocutaneous lesions by their twenties, and such lesions have been reported to occur in 99% of individuals with Cowden syndrome/PHTS, showing nearly complete penetrance, although this may be a reflection of selection bias in the cases reported.<sup>149,485</sup> The presence of three or more mucocutaneous neuromas is considered a major diagnostic criterion of Cowden syndrome/PHTS,<sup>493</sup> while the presence of two or more trichilemmomas has been reported to be pathognomonic for Cowden syndrome/PHTS.<sup>512,513</sup> However, since most of the evidence regarding trichilemmomas is from the older literature, it is possible that the association with Cowden syndrome/PHTS is somewhat overestimated.<sup>81</sup> There are reports of individuals with a solitary trichilemmoma who do not have Cowden syndrome/PHTS.<sup>512,513</sup>

Nevertheless, due to the strong association between these lesions and Cowden syndrome/PHTS and the difficulty in clinically distinguishing between a trichilemmoma and another mucocutaneous lesion, it is important that a diagnosis of trichilemmoma is histologically confirmed.

It was previously estimated that about half of individuals with Cowden syndrome/PHTS have gastrointestinal polyps.<sup>514</sup> However, this was almost certainly an underestimate.<sup>514,515</sup> In an analysis of 67 *PTEN* P/LP variant carriers undergoing colonoscopy, colorectal polyps were found in 92.5% of patients.<sup>514</sup> About half of the patients undergoing colonoscopy had hyperplastic polyps, and about 25% had polyps that were hamartomatous, ganglioneuromatous, or adenomatous.<sup>514</sup> Adenomatous or hyperplastic polyps were associated with development of colorectal cancer in this sample. Out of 39 carriers of a *PTEN* P/LP variant undergoing esophagogastroduodenoscopy (EGD), upper gastrointestinal polyps were found in 67% of patients.<sup>514</sup> A systematic review of published case series (N = 102) regarding gastrointestinal manifestations in Cowden syndrome/PHTS and component syndromes showed that 92.5% of these patients had polyps, with 64% having 50 or more.<sup>516</sup> Histologies were described as: hyperplastic (44%), adenomatous (40%), hamartomatous (38%), ganglioneuroma (33%), and inflammatory (24.5%). Other studies have also reported ganglioneuromatous polyps (ie, rare, benign peripheral nervous system tumors) in this population.<sup>493,517</sup> A retrospective chart review of 47 children with *PTEN* P/LP variants showed that only 13% had gastrointestinal polyps, but 34% had other gastrointestinal symptoms such as abdominal pain, rectal bleeding, and/or constipation.<sup>499</sup> Early-onset (<50 years of age) colorectal cancer has been reported in 13% of patients with *PTEN* P/LP variant-associated Cowden syndrome/PHTS, suggesting that routine colonoscopy may be warranted in this population.<sup>514</sup> The lifetime risk for colorectal cancer has been estimated as 9% to 16%.<sup>495,496</sup>

Several studies have projected lifetime estimates of cancer risk that are significantly higher than previously estimated. In a study of patients meeting diagnostic criteria for Cowden syndrome/PHTS (N = 211; identified from published literature and records from a single institution), the cumulative lifetime risk for any cancer was 89%.<sup>495</sup> *PTEN* P/LP variants had been identified in 97 of 105 patients (92%) who underwent testing. The cumulative lifetime cancer risks for all evaluable patients (n = 210) were 81% for female breast cancer, 21% for thyroid cancer, 19% for endometrial cancer, 15% for renal cancer, and 16% for colorectal cancer.<sup>495</sup> In a prospective study that evaluated genotype-phenotype associations between *PTEN* P/LP variants and cancer risks,<sup>496</sup> deleterious germline P/LP variants in *PTEN* were identified in 368 patients. Calculation of age-adjusted SIRs using cancer incidence data from the SEER database showed elevated SIRs among individuals with *PTEN* P/LP variants for breast cancer (25), thyroid cancer (51), endometrial cancer (43), colorectal cancer (10), renal cancer (31), and melanoma (8.5). The estimated cumulative lifetime cancer risks were 85% for breast, 35% for thyroid, 28% for endometrial, 9% for colorectal, 34% for renal, and 6% for melanoma.<sup>496</sup> In another study in individuals with PHTS found to have deleterious germline *PTEN* P/LP variants (N = 154; detailed information available in n = 146), age- and gender-adjusted SIRs were elevated for female breast cancer (39), endometrial cancer (49), female thyroid cancer (43), male thyroid cancer (199.5), female melanoma (28), and male melanoma (39).<sup>494</sup> The cumulative lifetime risks in these individuals were 77% for female breast cancer and 38% for thyroid cancer. The cumulative lifetime risk for any cancer was 85% overall, and women with Cowden syndrome/PHTS were found to have a 2-fold greater cancer risk compared with men with Cowden syndrome/PHTS.<sup>494</sup> It is important to note, however, that all three of these studies suffer from significant ascertainment biases, in that patients were usually selected for *PTEN* testing based on the presence of these malignancies, which would inflate the projected lifetime cancer estimates. An observational study of 180

patients with *PTEN* P/LP variants used Kaplan-Meier methods to estimate that female carriers (n = 99) have an 87% cumulative risk of developing any cancer and/or LDD by 60 years of age, while male carriers have a cumulative risk of 56%.<sup>518</sup>

The BRRS variant of Cowden syndrome/PHTS has been characterized by the presence of multiple lipomas, gastrointestinal hamartomatous polyps, macrocephaly, hemangiomas, developmental delay, and pigmented macules on the glans penis,<sup>519</sup> although formal diagnostic criteria have not been established for this syndrome. *PTEN* gene P/LP variants testing in individuals characterized with BRRS have been reported in approximately 60% of these patients.<sup>520</sup> Further, in another study, 10% of patients with BRRS for whom a *PTEN* P/LP variant test was negative were shown to be carriers of large *PTEN* gene deletions.<sup>502</sup>

### **Risk Assessment, Counseling, and Management**

The assessment of individuals suspected of having Cowden syndrome/PHTS incorporates both a history of the benign and malignant conditions associated with the syndrome and a targeted physical examination, including the skin and oral mucosa, breast, and thyroid gland and head circumference (see *Testing Criteria for Cowden Syndrome/PHTS* in the algorithm). The NCCN Guidelines Panel has established a list of criteria to help indicate which individuals are candidates for testing for *PTEN* P/LP variants (see *Testing Criteria for Cowden Syndrome/PHTS* in the algorithm). These criteria are used to assess the need for further risk assessment and genetic testing. When *PTEN* is included on multi-gene panels, these testing criteria do not need to be met. Clinical diagnostic criteria have also been developed to help identify clinical features associated with Cowden syndrome/PHTS (see *Revised Clinical Diagnostic Criteria for PTEN Hamartoma Tumor Syndrome* in the algorithm, and discussed below under *Clinical Diagnostic Criteria*). Patients who meet clinical diagnostic criteria for Cowden

syndrome/PHTS as described in this section are candidates for testing for *PTEN* P/LP variants.

### *Testing Criteria*

Testing criteria for Cowden syndrome/PHTS are grouped into three general categories. A patient is considered for testing for *PTEN* P/LP variants based on whether he/she meets certain criteria or combinations of criteria from these three categories. The first criteria category includes individuals meeting diagnostic criteria for Cowden syndrome<sup>521</sup>: a personal history of BRRS, adult LDD, autism spectrum disorder with macrocephaly, or two or more biopsy-proven trichilemmomas. Any individual presenting with one or more of these diagnoses warrants *PTEN* testing. Previously, some of the criteria from this group have been referred to as “pathognomonic,” although it is unlikely that any of these conditions can stand alone as a definitive diagnostic criterion for Cowden syndrome/PHTS. Another criterion that can be considered to be sufficient to warrant testing for *PTEN* P/LP variants is a family history that includes the presence of a known *PTEN* P/LP variant.

The next category of criteria represents “major” features associated with Cowden syndrome/PHTS and are described in the Guidelines (see *Testing Criteria for Cowden Syndrome/PHTS* in the algorithm).<sup>486,489,497,501,521</sup> With respect to decisions related to the presence of mucocutaneous lesions, the panel did not consider the available literature to be adequate to accurately specify the number or extent of these lesions required for the condition to be defined as a major criterion for Cowden syndrome/PHTS, and clinical judgment is needed when evaluating such lesions. An individual exhibiting two or more major criteria where one criterion is macrocephaly meets the testing threshold. An individual with three or more major criteria (without macrocephaly) is also considered to meet the threshold for testing. In addition, individuals exhibiting one major criterion with three or more minor criteria (see *Testing*



Criteria for Cowden Syndrome/PHTS in the algorithm) also meet the testing threshold; if an individual exhibits two or more major criteria but does not have macrocephaly, then one of the major criteria may be included as one of the three minor criteria to meet the testing threshold.

The final category of criteria represents features with a “minor” association with Cowden syndrome/PHTS.<sup>486,489,497,521</sup> These criteria are described in the Guidelines (see *Testing Criteria for Cowden Syndrome/PHTS* in the algorithm). An individual would need to exhibit four or more minor criteria or, as discussed above, three or more minor criteria and one major criterion to meet testing.

Lastly, an individual who has a first-degree relative diagnosed with Cowden syndrome/PHTS or BRRS for whom testing has not been performed would also meet the threshold for *PTEN* testing if the individual meets at least one major criterion or two or more minor criteria. *PTEN* P/LP variants are commonly found in tumor tissue.<sup>522-524</sup> If a *PTEN* variant is detected through tumor profiling and would be classified as P/LP if present in the germline, then germline testing for *PTEN* should be considered.

### Clinical Diagnostic Criteria

The frequency of *PTEN* P/LP variant in individuals meeting International Cowden Consortium diagnostic criteria for Cowden syndrome has previously been estimated at about 80%.<sup>493,520</sup> However, evaluation of data based on samples analyzed at a single academic pathology laboratory (N = 802 evaluable) reported a much lower frequency (34%) of *PTEN* P/LP variants among individuals meeting diagnostic criteria<sup>489</sup> for Cowden syndrome.<sup>486</sup> The authors concluded that the current Consortium diagnostic criteria are not as sensitive in identifying individuals with *PTEN* P/LP variants as previously estimated. Since *PTEN* P/LP variants are relatively rare, recommendations regarding Cowden syndrome diagnostic criteria may be based on studies with a small number of patients. Studies

with larger samples have their flaws as well, as patients are selected for testing based on the number and magnitude of clinical features, which may lead to overestimation of the features of Cowden syndrome.<sup>493</sup> A review was conducted examining each consortium diagnostic criterion, and revised criteria were proposed that are more stringent and take into account clinical features that are often seen in PHTS.<sup>493</sup> The criteria were designed by focusing on clinical features associated with *PTEN* P/LP variants. The panel recommends using these criteria for clinical diagnosis of PHTS (see *Revised Clinical Diagnostic Criteria for PTEN Hamartoma Tumor Syndrome* in the algorithm).

### Screening Recommendations

Cancer is the major health risk associated with Cowden syndrome/PHTS. Therefore, the NCCN Panel has outlined guidelines for prevention and early detection screening of commonly associated cancers with Cowden syndrome/PHTS. Current medical management recommendations for individuals with Cowden syndrome/PHTS include annual physical examinations, starting at 18 years of age (or 5 years before the youngest age of diagnosis of a component cancer in the family).

The recommendations for individuals with Cowden syndrome/PHTS who were assigned female at birth focus on primary and secondary prevention options for breast cancer since this is the most commonly associated cancer in individuals with Cowden syndrome/PHTS based on the available literature. Individuals assigned female at birth should begin regular monthly breast self-examinations at 18 years of age and have a semiannual clinical breast examination beginning at 25 years of age or 5 to 10 years earlier than the earliest known breast cancer in the family (whichever comes first). Individuals assigned female at birth should also have an annual mammogram and breast MRI screening with contrast starting at 35 years of age, or 10 years earlier than the earliest known breast cancer in the family (whichever comes first). After 75 years of age,

management should be considered on an individual basis. In patients treated for breast cancer who were assigned female at birth and who have not had bilateral mastectomy, mammography and breast MRI screening with contrast should continue as recommended based on age. When mammography is performed, the panel recommends that tomosynthesis be considered.

Although there are no data regarding risk reduction surgery in individuals with Cowden syndrome who were assigned female at birth, the option of RRM and hysterectomy should be discussed. Oophorectomy is not indicated for Cowden syndrome alone. Counseling for risk-reducing surgeries may include discussion of extent of cancer risk reduction/protection, risks associated with surgeries, reconstructive options, and reproductive desires. It is also important to address the psychosocial and quality-of-life aspects of undergoing risk-reducing surgical procedures.

Given that Cowden syndrome is rare, there are no data on screening for endometrial cancer in these patients, though consideration of screening can begin as early as age 35. The panel recommends patient education regarding the symptoms of endometrial cancer including the necessity of a prompt response to symptoms such as abnormal bleeding. Prompt reporting promotes early detection of endometrial cancer. The evaluation of these symptoms should include an endometrial biopsy. Though endometrial cancer screening does not have proven benefit in individuals with Cowden syndrome, endometrial biopsy is highly sensitive and specific as a diagnostic procedure. Therefore, screening through endometrial biopsy every 1 to 2 years may be considered.

Routine TVUS to screen for endometrial cancer in postmenopausal individuals has not been shown to be sufficiently sensitive or specific to warrant a positive recommendation but may be considered at the clinician's discretion. However, TVUS is not recommended as a screening

tool in premenopausal individuals due to the wide range of endometrial strip thickness throughout the normal menstrual cycle.

Individuals with Cowden syndrome/PHTS have approximately at least a 3% to 10% lifetime risk of developing thyroid cancer,<sup>81</sup> compared to about 1% in the general population.<sup>525</sup> An annual thyroid ultrasound should be performed, starting at age 7.<sup>526</sup> Children at risk of a *PTEN* P/LP variant (based on a parent's carrier status) whose parents wish to delay genetic testing may also undergo annual thyroid ultrasound, since this is a noninvasive procedure. Colonoscopy is recommended starting at 35 years of age, or earlier if symptomatic. If a close relative was diagnosed with colon cancer before 40 years of age, then colonoscopy screening should begin 5 to 10 years before the age of the earliest known diagnosis. Colonoscopy should be performed every 5 years or more frequently in cases where the patient is symptomatic or polyps are found. To screen for renal cell carcinoma, renal ultrasound should be considered every 1 to 2 years beginning at 40 years of age. Annual dermatologic examination is recommended. If there are symptoms in children, then assessment of psychomotor abilities should be considered, as well as a brain MRI. Education regarding the signs and symptoms of cancer is important; patients should also be advised about the risk to relatives, and genetic counseling is recommended for at-risk relatives.

No published data exist on the use of prenatal diagnostics/genetic testing for *PTEN* P/LP variants in families with Cowden syndrome. For a general discussion on the topic of reproductive options and counseling considerations, see the Discussion section above on *Reproductive Options* under *Genetic Risk Assessment and Counseling*.

### Hereditary Pancreatic Cancer

Pancreatic cancer is thought to have a familial or hereditary component in approximately 10% of cases.<sup>189,190,527-529</sup> Harboring a P/LP variant has



been found to be associated with a greater incidence of pancreatic cancer than family history alone (without the presence of an associated germline variant).<sup>530</sup> Germline P/LP variants commonly found in pancreatic adenocarcinoma include *BRCA1*, *BRCA2*, *CDKN2A*, mismatch repair genes associated with Lynch syndrome (ie, *MSH2*, *MLH1*, *MSH6*, *PMS2*, *EPCAM*), *ATM*, *PALB2*, *STK11*, and *TP53*.<sup>84,183,185,187,190,330,528,530-537</sup> *BRCA2* and *CDKN2A* are generally the most prevalent, with rates in moderate- to high-risk families ranging from 2% to 6% for *BRCA2* and 1.5% to 2.5% for *CDKN2A*.<sup>180,184,189,190</sup> In addition, hereditary pancreatitis, which is associated with a significantly increased risk for pancreatic cancer, is associated with the genes *PRSS1* and *SPINK1*.<sup>528</sup> Patients with pancreatic cancer and Ashkenazi Jewish ancestry may have a greater likelihood of testing positive for a *BRCA1/2* P/LP variant, with prevalence of detected P/LP variants in this group ranging from 5.5% to 19%, with P/LP variants being more common for *BRCA2*.<sup>182,183,185,191</sup>

Given the considerable rate of predisposing P/LP variants in patients with pancreatic cancer, as well as the fact that typical clinical factors (eg, young age of onset, family history of cancer) are poorly predictive for identifying carriers of a P/LP variant, universal genetic testing for these individuals is warranted. Given the elevated rates of P/LP variants in pancreatic cancer and that pancreatic cancer risk increases when there is a family history,<sup>538</sup> testing of first-degree relatives of patients may be beneficial. However, testing the patient is preferred. Testing of second-degree relatives is generally not recommended but may be considered in select cases. Given that mortality rates for this cancer are high,<sup>539,540</sup> it may be beneficial to family members to test patients near the time of diagnosis, since the option to test the patient may not be available for very long. Family history of pancreatic cancer with unknown histology is often presumed to be exocrine. Detecting a germline P/LP variant can potentially aid in treatment decision-making, particularly regarding systemic therapy options (see *Systemic Therapy Decision-Making* above).

## Pancreas Screening

Evidence to support screening for pancreatic cancer comes from studies including those who harbor an associated germline P/LP variant and/or those who have a particularly strong family history of pancreatic cancer (two or more first-degree relatives on the same side of the family, or three or more first- or second-degree relatives on the same side of the family). An analysis of outcomes from three European centers including 411 asymptomatic individuals showed that pancreatic cancer was detected in 7% of carriers of a *CDKN2A* P/LP variant and less than 1% of those with familial pancreatic cancer.<sup>541</sup> For the carriers of a *CDKN2A* P/LP variant for whom a lesion was detected, 75% were resectable, with a 5-year OS rate of 24%. Another analysis from six high-volume centers in Italy including 187 high-risk individuals, abnormalities were detected in about 28%.<sup>542</sup> Out of the cysts detected, 62.2% were branch-duct intraductal papillary mucinous neoplasms. Pancreatic adenocarcinomas made up 2.6% of the findings (n = 5). A third analysis including screening of 354 asymptomatic high-risk individuals showed suspicious pancreas lesions in 19%.<sup>543</sup> Out of the lesions detected from screening, 90% were resectable, and the 3-year OS rate was 85% in those with resectable lesions. The considerable rate of resectable asymptomatic lesions found from routine screening of high-risk individuals demonstrates the potential for downstaging (ie, identification of lesions at an earlier stage). There is also the potential for impact on mortality rates, though long-term studies are needed in this area. Lesions detected through routine screening may not always require resection (eg, sporadic branch-duct intraductal papillary mucinous neoplasms). Therefore, larger long-term studies are needed to further determine the risks and benefits of routine pancreas screening in high-risk individuals, as well as the threshold for surgical intervention.<sup>543</sup>

With the exception of *CDKN2A* and *STK11*, pancreas cancer screening in individuals who have a P/LP variant associated with increased risk of exocrine pancreatic cancer (ie, *ATM*, *BRCA1*, *BRCA2*, *MSH2*, *MLH1*,

*MSH6, EPCAM, PALB2, TP53*) is not recommended unless there is additional family history of pancreatic cancer (at least 1 first- or second-degree relative).<sup>544</sup> If family history criteria are met, then pancreas screening may be considered at age 50, or 10 years younger than the earliest pancreatic cancer diagnosis in the family, whichever is earlier.<sup>544</sup>

The International Cancer of the Pancreas Screening Consortium recommendations for pancreas screening in individuals with increased risk for hereditary pancreatic cancer do not include carriers of a *TP53* P/LP variant in this group,<sup>544</sup> as there are very limited data on pancreatic cancer screening in these carriers. However, the NCCN Guidelines panel recommends that pancreatic cancer screening be considered in carriers of a *TP53* P/LP variant, if there is additional family history of pancreatic cancer (at least 1 first- or second-degree relative), as there is some evidence of a modestly increased risk of pancreatic cancer in these carriers.<sup>187,190</sup>

For carriers of a *CDKN2A* or *STK11* P/LP variant, no additional family history is needed to warrant screening. For carriers of a *CDKN2A* P/LP variant, screening may be considered at age 40, or 10 years younger than the earliest pancreatic cancer diagnosis in the family, whichever is earlier.<sup>544</sup> For carriers of a *STK11* P/LP variant, screening may be considered beginning at ages 30 to 35, or 10 years younger than the earliest pancreatic cancer diagnosis in the family, whichever is earlier.<sup>366,544</sup>

Hereditary pancreatitis is associated with increased lifetime risk of exocrine pancreatic cancer and is sometimes caused by P/LP variants such as *PRSS1* and *SPINK1*.<sup>545-548</sup> However, the clinical significance of these variants is unclear without a clinical history of pancreatitis. Therefore, germline testing for *PRSS1*, *SPINK1*, and other genes associated with pancreatitis is generally not recommended unless one's personal or family history is suggestive of hereditary pancreatitis.<sup>547</sup>

Pancreas cancer screening is recommended in individuals harboring one of these variants only in the presence of a clinical phenotype consistent with hereditary pancreatitis. For individuals meeting these criteria, screening may begin at age 40, or 20 years after onset of pancreatitis, whichever is earlier.<sup>544</sup>

When screening is recommended, it may be done with contrast-enhanced MRI/MRCP and/or endoscopic ultrasound (EUS).<sup>543,544,549</sup> MRI and EUS have been shown to be superior in detection of subcentimeter pancreatic cysts, compared to CT.<sup>549</sup> Screening at a high-volume center of expertise is recommended, preferably in the context of a research study. In those for whom screening shows potentially concerning features that suggest progression, shorter screening intervals may be indicated.

**Table 1. Glossary of Relevant Genetic Terms (from the National Cancer Institute [NCI])**

***Autosomal dominant***

Autosomal dominant inheritance refers to genetic conditions that occur when a P/LP variant is present in one copy of a given gene (ie, the person is heterozygous).

***Autosomal recessive***

Autosomal recessive inheritance refers to genetic conditions that occur only when P/LP variants are present in both copies of a given gene (ie, the person is homozygous for a P/LP variant, or carries two different variants of the same gene, a state referred to as compound heterozygosity).

***de novo mutation***

An alteration in a gene that is present for the first time in one family member as a result of a P/LP variant in a germ cell (egg or sperm) of one of the parents, or a P/LP variant that arises in the fertilized egg itself during early embryogenesis. Also called new P/LP variant.

***Familial***

A phenotype or trait that occurs with greater frequency in a given family than in the general population; familial traits may have a genetic and/or nongenetic etiology.

***Family history***

The genetic relationships within a family combined with the medical history of individual family members. When represented in diagram form using standardized symbols and terminology, it is usually referred to as a pedigree or family tree.

***Founder effect***

A P/LP variant observed with high frequency in a population founded by a small ancestral group that was once geographically or culturally isolated, in which one or more of the founders was a carrier of the mutant gene.

***Germline***

The cells from which eggs or sperm (ie, gametes) are derived.

***Kindred***

An extended family.

***Pedigree***

A graphic illustration of family history.

***Penetrance***

A characteristic of a genotype; it refers to the likelihood that a clinical condition will occur when a particular genotype is present.

***Proband***

The individual through whom a family with a genetic disorder is ascertained. In males this is called a propositus, and in females it is called a probanda.

***Sporadic cancer***

This term has two meanings. It is sometimes used to differentiate cancers occurring in people who do not have a germline P/LP variant that confers increased susceptibility to cancer from cancers occurring in people who are known to carry a variant. Cancer developing in people who do not carry a high-risk P/LP variant is referred to as sporadic cancer. The distinction is not absolute, because genetic background may influence the likelihood of cancer even in the absence of a specific predisposing variant. Alternatively, sporadic is also sometimes used to describe cancer occurring in individuals without a family history of cancer.

**Table 2. Genetic Test Results to Determine the Presence of a Cancer-Predisposing Gene**

<i>Result</i>	<i>Description</i>
<i>True-positive</i>	The person is a carrier of an alteration in a known cancer-predisposing gene.
<i>True-negative</i>	The person is not a carrier of a known cancer-predisposing gene that has been positively identified in another family member.
<i>Indeterminate (uninformative)</i>	The person is not a carrier of a known cancer-predisposing gene, and the carrier status of other family members is either also negative or unknown.
<i>Inconclusive (variants of uncertain significance)</i>	The person is a carrier of an alteration in a gene that currently has no known significance.





### References

1. Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759-767. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2188735>.
2. Vogelstein B, Kinzler KW. The multistep nature of cancer. *Trends Genet* 1993;9:138-141. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8516849>.
3. Lynch HT, Watson P, Conway TA, Lynch JF. Clinical/genetic features in hereditary breast cancer. *Breast Cancer Res Treat* 1990;15:63-71. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2322650>.
4. Pharoah PD, Day NE, Duffy S, et al. Family history and the risk of breast cancer: a systematic review and meta-analysis. *Int J Cancer* 1997;71:800-809. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9180149>.
5. Berliner JL, Fay AM. Risk assessment and genetic counseling for hereditary breast and ovarian cancer: recommendations of the National Society of Genetic Counselors. *J Genet Couns* 2007;16:241-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17508274>.
6. Foulkes WD. Inherited susceptibility to common cancers. *N Engl J Med* 2008;359:2143-2153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19005198>.
7. Trepanier A, Ahrens M, McKinnon W, et al. Genetic cancer risk assessment and counseling: recommendations of the national society of genetic counselors. *J Genet Couns* 2004;13:83-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15604628>.
8. Pharoah PD, Antoniou A, Bobrow M, et al. Polygenic susceptibility to breast cancer and implications for prevention. *Nat Genet* 2002;31:33-36. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11984562>.
9. Lancaster JM, Powell CB, Chen LM, Richardson DL. Society of Gynecologic Oncology statement on risk assessment for inherited gynecologic cancer predispositions. *Gynecol Oncol* 2015;136:3-7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25238946>.
10. Shiovitz S, Korde LA. Genetics of breast cancer: a topic in evolution. *Ann Oncol* 2015;26:1291-1299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25605744>.
11. Moyer VA. Risk assessment, genetic counseling, and genetic testing for BRCA-related cancer in women: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2014;160:271-281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24366376>.
12. Weitzel JN, Blazer KR, MacDonald DJ, et al. Genetics, genomics, and cancer risk assessment: state of the art and future directions in the era of personalized medicine. *CA Cancer J Clin* 2011;61:327-359. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21858794>.
13. U.S. National Library of Medicine-Key MEDLINE® Indicators. Available at: [http://www.nlm.nih.gov/bsd/bsd\\_key.html](http://www.nlm.nih.gov/bsd/bsd_key.html). Accessed July 24, 2014.
14. Kurian AW, Li Y, Hamilton AS, et al. Gaps in incorporating germline genetic testing into treatment decision-making for early-stage breast cancer. *J Clin Oncol* 2017;35:2232-2239. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28402748>.
15. Ademuyiwa FO, Salyer P, Ma Y, et al. Assessing the effectiveness of the National Comprehensive Cancer Network genetic testing guidelines in identifying African American breast cancer patients with deleterious genetic mutations. *Breast Cancer Res Treat* 2019;178:151-159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31325073>.
16. Pal T, Agnese D, Daly M, et al. Points to consider: is there evidence to support BRCA1/2 and other inherited breast cancer genetic testing for all breast cancer patients? A statement of the American College of Medical Genetics and Genomics (ACMG). *Genet Med* 2020;22:681-685. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31831881>.
17. Weitzel JN, Lagos VI, Cullinane CA, et al. Limited family structure and BRCA gene mutation status in single cases of breast cancer. *JAMA*



2007;297:2587-2595. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17579227>.

18. Hong YC, Liu HM, Chen PS, et al. Hair follicle: a reliable source of recipient origin after allogeneic hematopoietic stem cell transplantation. Bone Marrow Transplant 2007;40:871-874. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17704789>.

19. Tran SD, Pillemer SR, Dutra A, et al. Differentiation of human bone marrow-derived cells into buccal epithelial cells in vivo: a molecular analytical study. Lancet 2003;361:1084-1088. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12672312>.

20. Weitzel JN, Chao EC, Nehoray B, et al. Somatic TP53 variants frequently confound germ-line testing results. Genet Med 2018;20:809-816. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29189820>.

21. Mersch J, Brown N, Pirzadeh-Miller S, et al. Prevalence of variant reclassification following hereditary cancer genetic testing. JAMA 2018;320:1266-1274. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30264118>.

22. Balmana J, Digiovanni L, Gaddam P, et al. Conflicting interpretation of genetic variants and cancer risk by commercial laboratories as assessed by the prospective registry of multiplex testing. J Clin Oncol 2016;34:4071-4078. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27621404>.

23. Vail PJ, Morris B, van Kan A, et al. Comparison of locus-specific databases for BRCA1 and BRCA2 variants reveals disparity in variant classification within and among databases. J Community Genet 2015;6:351-359. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25782689>.

24. Lincoln SE, Yang S, Cline MS, et al. Consistency of BRCA1 and BRCA2 variant classifications among clinical diagnostic laboratories. JCO Precis Oncol 2017;1. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28782058>.

25. Eccles DM, Mitchell G, Monteiro AN, et al. BRCA1 and BRCA2 genetic testing-pitfalls and recommendations for managing variants of uncertain clinical significance. Ann Oncol 2015;26:2057-2065. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26153499>.

26. Badalato L, Kalokairinou L, Borry P. Third party interpretation of raw genetic data: an ethical exploration. Eur J Hum Genet 2017;25:1189-1194. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28832567>.

27. Tandy-Connor S, Guiltinan J, Krempely K, et al. False-positive results released by direct-to-consumer genetic tests highlight the importance of clinical confirmation testing for appropriate patient care. Genet Med 2018;20:1515-1521. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29565420>.

28. Green RC, Berg JS, Grody WW, et al. ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. Genet Med 2013;15:565-574. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23788249>.

29. Robson ME, Bradbury AR, Arun B, et al. American Society of Clinical Oncology Policy statement update: genetic and genomic testing for cancer susceptibility. J Clin Oncol 2015;33:3660-3667. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26324357>.

30. Slavin TP, Banks KC, Chudova D, et al. Identification of incidental germline mutations in patients with advanced solid tumors who underwent cell-free circulating tumor DNA sequencing. J Clin Oncol 2018;36:JCO1800328. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30339520>.

31. Walsh T, Casadei S, Coats KH, et al. Spectrum of mutations in BRCA1, BRCA2, CHEK2, and TP53 in families at high risk of breast cancer. JAMA 2006;295:1379-1388. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16551709>.

32. Kurian AW, Hare EE, Mills MA, et al. Clinical evaluation of a multiple-gene sequencing panel for hereditary cancer risk assessment. J Clin



Oncol 2014;32:2001-2009. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24733792>.

33. Kurian AW, Ward KC, Hamilton AS, et al. Uptake, results, and outcomes of germline multiple-gene sequencing after diagnosis of breast cancer. JAMA Oncol 2018;4:1066-1072. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29801090>.

34. Hall MJ, Forman AD, Pilarski R, et al. Gene panel testing for inherited cancer risk. J Natl Compr Canc Netw 2014;12:1339-1346. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25190699>.

35. Hall MJ, Obeid E, Daly MB. Multigene panels to evaluate hereditary cancer risk: reckless or relevant? J Clin Oncol 2016;34:4186-4187.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27551136>.

36. Manchanda R, Patel S, Gordeev VS, et al. Cost-effectiveness of population-based BRCA1, BRCA2, RAD51C, RAD51D, BRIP1, PALB2 mutation testing in unselected general population women. J Natl Cancer Inst 2018;110:714-725. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29361001>.

37. Walsh T, Lee MK, Casadei S, et al. Detection of inherited mutations for breast and ovarian cancer using genomic capture and massively parallel sequencing. Proc Natl Acad Sci U S A 2010;107:12629-12633. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20616022>.

38. LaDuca H, Polley EC, Yussuf A, et al. A clinical guide to hereditary cancer panel testing: evaluation of gene-specific cancer associations and sensitivity of genetic testing criteria in a cohort of 165,000 high-risk patients. Genet Med 2020;22:407-415. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31406321>.

39. Bombard Y, Bach PB, Offit K. Translating genomics in cancer care. J Natl Compr Canc Netw 2013;11:1343-1353. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24225968>.

40. Rainville IR, Rana HQ. Next-generation sequencing for inherited breast cancer risk: counseling through the complexity. Curr Oncol Rep 2014;16:371. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24488544>.

41. Blazer KR, Slavin T, Weitzel JN. Increased reach of genetic cancer risk assessment as a tool for precision management of hereditary breast cancer. JAMA Oncol 2016;2:723-724. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26869327>.

42. Tung N, Domchek SM, Stadler Z, et al. Counselling framework for moderate-penetrance cancer-susceptibility mutations. Nat Rev Clin Oncol 2016;13:581-588. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/27296296>.

43. van Marcke C, De Leener A, Berliere M, et al. Routine use of gene panel testing in hereditary breast cancer should be performed with caution. Crit Rev Oncol Hematol 2016;108:33-39. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27931838>.

44. LaDuca H, Stuenkel AJ, Dolinsky JS, et al. Utilization of multigene panels in hereditary cancer predisposition testing: analysis of more than 2,000 patients. Genet Med 2014;16:830-837. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24763289>.

45. Samadder NJ, Riegert-Johnson D, Boardman L, et al. Comparison of universal genetic testing vs guideline-directed targeted testing for patients with hereditary cancer syndrome. JAMA Oncol 2021;7:230-237. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33126242>.

46. van Os NJ, Roeleveld N, Weemaes CM, et al. Health risks for ataxia-telangiectasia mutated heterozygotes: a systematic review, meta-analysis and evidence-based guideline. Clin Genet 2016;90:105-117. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26662178>.

47. Southey MC, Goldgar DE, Winqvist R, et al. PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS. J Med Genet 2016;53:800-811. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27595995>.



48. Goldgar DE, Healey S, Dowty JG, et al. Rare variants in the ATM gene and risk of breast cancer. *Breast Cancer Res* 2011;13:R73. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21787400>.
49. Brunet J, Gutierrez-Enriquez S, Torres A, et al. ATM germline mutations in Spanish early-onset breast cancer patients negative for BRCA1/BRCA2 mutations. *Clin Genet* 2008;73:465-473. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18384426>.
50. Heikkinen K, Rapakko K, Karppinen SM, et al. Association of common ATM polymorphism with bilateral breast cancer. *Int J Cancer* 2005;116:69-72. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15756685>.
51. Thompson D, Antoniou AC, Jenkins M, et al. Two ATM variants and breast cancer risk. *Hum Mutat* 2005;25:594-595. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15880680>.
52. Tommiska J, Jansen L, Kilpivaara O, et al. ATM variants and cancer risk in breast cancer patients from Southern Finland. *BMC Cancer* 2006;6:209. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16914028>.
53. Mauer CB, Pirzadeh-Miller SM, Robinson LD, Euhus DM. The integration of next-generation sequencing panels in the clinical cancer genetics practice: an institutional experience. *Genet Med* 2014;16:407-412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24113346>.
54. Tung N, Battelli C, Allen B, et al. Frequency of mutations in individuals with breast cancer referred for BRCA1 and BRCA2 testing using next-generation sequencing with a 25-gene panel. *Cancer* 2015;121:25-33. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25186627>.
55. Kapoor NS, Curcio LD, Blakemore CA, et al. Multigene panel testing detects equal rates of pathogenic BRCA1/2 mutations and has a higher diagnostic yield compared to limited BRCA1/2 analysis alone in patients at risk for hereditary breast cancer. *Ann Surg Oncol* 2015;22:3282-3288. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26219241>.
56. Renaux-Petel M, Charbonnier F, Thery JC, et al. Contribution of de novo and mosaic TP53 mutations to Li-Fraumeni syndrome. *J Med Genet* 2018;55:173-180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29070607>.
57. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;371:2488-2498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25426837>.
58. Genovese G, Kahler AK, Handsaker RE, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med* 2014;371:2477-2487. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25426838>.
59. Kuchenbaecker KB, McGuffog L, Barrowdale D, et al. Evaluation of polygenic risk scores for breast and ovarian cancer risk prediction in BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst* 2017;109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28376175>.
60. Barnes DR, Rookus MA, McGuffog L, et al. Polygenic risk scores and breast and epithelial ovarian cancer risks for carriers of BRCA1 and BRCA2 pathogenic variants. *Genet Med* 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32665703>.
61. Lecarpentier J, Silvestri V, Kuchenbaecker KB, et al. Prediction of breast and prostate cancer risks in male BRCA1 and BRCA2 mutation carriers using polygenic risk scores. *J Clin Oncol* 2017;35:2240-2250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28448241>.
62. Pashayan N, Pharoah PD, Schleutker J, et al. Reducing overdiagnosis by polygenic risk-stratified screening: findings from the Finnish section of the ERSPC. *Br J Cancer* 2015;113:1086-1093. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26291059>.
63. Seibert TM, Fan CC, Wang Y, et al. Polygenic hazard score to guide screening for aggressive prostate cancer: development and validation in large scale cohorts. *BMJ* 2018;360:j5757. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29321194>.



64. Lewis CM, Vassos E. Polygenic risk scores: from research tools to clinical instruments. *Genome Med* 2020;12:44. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32423490>.
65. Genetic Information Non-Discrimination Act of 2008 (GINA). Vol. Public Law No. 110-233. Available at: <https://www.eeoc.gov/laws/statutes/gina.cfm>.
66. Calzone KA, Soballe PW. Genetic testing for cancer susceptibility. *Surg Clin North Am* 2008;88:705-721. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18672137>.
67. Forrest LE, Young MA. Clinically significant germline mutations in cancer-causing genes identified through research studies should be offered to research participants by genetic counselors. *J Clin Oncol* 2016;34:898-901. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26786918>.
68. Cohen SA, Bradbury A, Henderson V, et al. Genetic counseling and testing in a community setting: quality, access, and efficiency. *Am Soc Clin Oncol Educ Book* 2019;39:e34-e44. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31099680>.
69. Berliner JL, Fay AM, Cummings SA, et al. NSGC practice guideline: risk assessment and genetic counseling for hereditary breast and ovarian cancer. *J Genet Couns* 2013;22:155-163. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23188549>.
70. Offit K, Levran O, Mullaney B, et al. Shared genetic susceptibility to breast cancer, brain tumors, and Fanconi anemia. *J Natl Cancer Inst* 2003;95:1548-1551. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14559878>.
71. Cragun D, Camperlengo L, Robinson E, et al. Differences in BRCA counseling and testing practices based on ordering provider type. *Genet Med* 2015;17:51-57. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24922460>.
72. Katz SJ, Ward KC, Hamilton AS, et al. Gaps in receipt of clinically indicated genetic counseling after diagnosis of breast cancer. *J Clin Oncol* 2018;36:1218-1224. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29528794>.
73. Vadaparampil ST, Scherr CL, Cragun D, et al. Pre-test genetic counseling services for hereditary breast and ovarian cancer delivered by non-genetics professionals in the state of Florida. *Clin Genet* 2015;87:473-477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24735105>.
74. Hoskovec JM, Bennett RL, Carey ME, et al. Projecting the supply and demand for certified genetic counselors: a workforce study. *J Genet Couns* 2018;27:16-20. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29052810>.
75. Offit K, Kohut K, Clagett B, et al. Cancer genetic testing and assisted reproduction. *J Clin Oncol* 2006;24:4775-4782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16840542>.
76. Offit K, Sagi M, Hurley K. Preimplantation genetic diagnosis for cancer syndromes: a new challenge for preventive medicine. *JAMA* 2006;296:2727-2730. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17164459>.
77. Jasper MJ, Liebelt J, Hussey ND. Preimplantation genetic diagnosis for BRCA1 exon 13 duplication mutation using linked polymorphic markers resulting in a live birth. *Prenat Diagn* 2008;28:292-298. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18302307>.
78. Sagi M, Weinberg N, Eilat A, et al. Preimplantation genetic diagnosis for BRCA1/2--a novel clinical experience. *Prenat Diagn* 2009;29:508-513. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19248143>.
79. Blackwood MA, Weber BL. BRCA1 and BRCA2: from molecular genetics to clinical medicine. *J Clin Oncol* 1998;16:1969-1977. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9586917>.



80. Venkitaraman AR. Cancer susceptibility and the functions of BRCA1 and BRCA2. *Cell* 2002;108:171-182. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11832208>.
81. Pilarski R. Cowden syndrome: a critical review of the clinical literature. *J Genet Couns* 2009;18:13-27. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18972196>.
82. Schneider KA, Garber J. Li-Fraumeni syndrome. *GeneReviews*; 2013. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK1311/>.
83. Antoniou AC, Casadei S, Heikkinen T, et al. Breast-cancer risk in families with mutations in PALB2. *N Engl J Med* 2014;371:497-506. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25099575>.
84. Yang X, Leslie G, Doroszk A, et al. Cancer risks associated with germline PALB2 pathogenic variants: an international study of 524 families. *J Clin Oncol* 2020;38:674-685. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31841383>.
85. Hu C, Polley EC, Yadav S, et al. The contribution of germline predisposition gene mutations to clinical subtypes of invasive breast cancer from a clinical genetic testing cohort. *J Natl Cancer Inst* 2020;112:1231-1241. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32091585>.
86. Abeliovich D, Kaduri L, Lerer I, et al. The founder mutations 185delAG and 5382insC in BRCA1 and 6174delT in BRCA2 appear in 60% of ovarian cancer and 30% of early-onset breast cancer patients among Ashkenazi women. *Am J Hum Genet* 1997;60:505-514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9042909>.
87. Levy-Lahad E, Catane R, Eisenberg S, et al. Founder BRCA1 and BRCA2 mutations in Ashkenazi Jews in Israel: frequency and differential penetrance in ovarian cancer and in breast-ovarian cancer families. *Am J Hum Genet* 1997;60:1059-1067. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9150153>.
88. Petrucelli N, Daly MB, Bars Culver JO, Feldman GL. BRCA1 and BRCA2 hereditary breast/ovarian cancer. *GeneReviews*; 2011. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK1247/>.
89. Avraham A, Feldman S, Cho SS, et al. Breast-specific epigenetic regulation of deltaNp73 and its role in DNA-damage-response of BRCA1-mutated human mammary epithelial cells. *Cancers (Basel)* 2020;12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32825620>.
90. Prevalence and penetrance of BRCA1 and BRCA2 mutations in a population-based series of breast cancer cases. Anglian Breast Cancer Study Group. *Br J Cancer* 2000;83:1301-1308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11044354>.
91. Antoniou A, Pharoah PD, Narod S, et al. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet* 2003;72:1117-1130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12677558>.
92. Chen S, Parmigiani G. Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol* 2007;25:1329-1333. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17416853>.
93. Ford D, Easton DF, Bishop DT, et al. Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Lancet* 1994;343:692-695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7907678>.
94. King MC, Marks JH, Mandell JB, New York Breast Cancer Study G. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* 2003;302:643-646. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14576434>.
95. Mavaddat N, Peock S, Frost D, et al. Cancer risks for BRCA1 and BRCA2 mutation carriers: results from prospective analysis of EMBRACE. *J Natl Cancer Inst* 2013;105:812-822. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23628597>.



96. Risch HA, McLaughlin JR, Cole DE, et al. Population BRCA1 and BRCA2 mutation frequencies and cancer penetrances: a kin-cohort study in Ontario, Canada. *J Natl Cancer Inst* 2006;98:1694-1706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17148771>.

97. van den Broek AJ, van 't Veer LJ, Hoening MJ, et al. Impact of age at primary breast cancer on contralateral breast cancer risk in BRCA1/2 mutation carriers. *J Clin Oncol* 2016;34:409-418. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26700119>.

98. Kuchenbaecker KB, Hopper JL, Barnes DR, et al. Risks of breast, ovarian, and contralateral breast cancer for BRCA1 and BRCA2 mutation carriers. *JAMA* 2017;317:2402-2416. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28632866>.

99. Zhong Q, Peng HL, Zhao X, et al. Effects of BRCA1- and BRCA2-related mutations on ovarian and breast cancer survival: a meta-analysis. *Clin Cancer Res* 2015;21:211-220. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25348513>.

100. Baretta Z, Mocellin S, Goldin E, et al. Effect of BRCA germline mutations on breast cancer prognosis: A systematic review and meta-analysis. *Medicine (Baltimore)* 2016;95:e4975. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27749552>.

101. van den Broek AJ, Schmidt MK, van 't Veer LJ, et al. Worse breast cancer prognosis of BRCA1/BRCA2 mutation carriers: what's the evidence? A systematic review with meta-analysis. *PLoS One* 2015;10:e0120189. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25816289>.

102. Copson ER, Maishman TC, Tapper WJ, et al. Germline BRCA mutation and outcome in young-onset breast cancer (POSH): a prospective cohort study. *Lancet Oncol* 2018;19:169-180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29337092>.

103. Kast K, Rhiem K, Wappenschmidt B, et al. Prevalence of BRCA1/2 germline mutations in 21 401 families with breast and ovarian cancer. *J*

*Med Genet* 2016;53:465-471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26928436>.

104. Schmidt MK, van den Broek AJ, Tollenaar RA, et al. Breast cancer survival of BRCA1/BRCA2 mutation carriers in a hospital-based cohort of young women. *J Natl Cancer Inst* 2017;109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28376189>.

105. Atchley DP, Albarracin CT, Lopez A, et al. Clinical and pathologic characteristics of patients with BRCA-positive and BRCA-negative breast cancer. *J Clin Oncol* 2008;26:4282-4288. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18779615>.

106. Eerola H, Heikkilä P, Tamminen A, et al. Relationship of patients' age to histopathological features of breast tumours in BRCA1 and BRCA2 and mutation-negative breast cancer families. *Breast Cancer Res* 2005;7:R465-469. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15987451>.

107. Lakhani SR, Reis-Filho JS, Fulford L, et al. Prediction of BRCA1 status in patients with breast cancer using estrogen receptor and basal phenotype. *Clin Cancer Res* 2005;11:5175-5180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16033833>.

108. Lakhani SR, Van De Vijver MJ, Jacquemier J, et al. The pathology of familial breast cancer: predictive value of immunohistochemical markers estrogen receptor, progesterone receptor, HER-2, and p53 in patients with mutations in BRCA1 and BRCA2. *J Clin Oncol* 2002;20:2310-2318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11981002>.

109. Lee E, McKean-Cowdin R, Ma H, et al. Characteristics of triple-negative breast cancer in patients with a BRCA1 mutation: results from a population-based study of young women. *J Clin Oncol* 2011;29:4373-4380. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22010008>.

110. Young SR, Pilarski RT, Donenberg T, et al. The prevalence of BRCA1 mutations among young women with triple-negative breast cancer. *BMC Cancer* 2009;9:86. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19298662>.



111. Fostira F, Tsiitlaidou M, Papadimitriou C, et al. Prevalence of BRCA1 mutations among 403 women with triple-negative breast cancer: implications for genetic screening selection criteria: a Hellenic Cooperative Oncology Group study. *Breast Cancer Res Treat* 2012;134:353-362. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22434525>.

112. Gonzalez-Angulo AM, Timms KM, Liu S, et al. Incidence and outcome of BRCA mutations in unselected patients with triple receptor-negative breast cancer. *Clin Cancer Res* 2011;17:1082-1089. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21233401>.

113. Rummel S, Varner E, Shriver CD, Ellsworth RE. Evaluation of BRCA1 mutations in an unselected patient population with triple-negative breast cancer. *Breast Cancer Res Treat* 2013;137:119-125. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23192404>.

114. Couch FJ, Hart SN, Sharma P, et al. Inherited mutations in 17 breast cancer susceptibility genes among a large triple-negative breast cancer cohort unselected for family history of breast cancer. *J Clin Oncol* 2015;33:304-311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25452441>.

115. Tung N, Lin NU, Kidd J, et al. Frequency of germline mutations in 25 cancer susceptibility genes in a sequential series of patients with breast cancer. *J Clin Oncol* 2016;34:1460-1468. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26976419>.

116. Buys SS, Sandbach JF, Gammon A, et al. A study of over 35,000 women with breast cancer tested with a 25-gene panel of hereditary cancer genes. *Cancer* 2017;123:1721-1730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28085182>.

117. Shimelis H, LaDuca H, Hu C, et al. Triple-negative breast cancer risk genes identified by multigene hereditary cancer panel testing. *J Natl Cancer Inst* 2018;110:855-862. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30099541>.

118. Breast Cancer Association Consortium, Dorling L, Carvalho S, et al. Breast Cancer Risk Genes - Association Analysis in More than 113,000

Women. *N Engl J Med* 2021;384:428-439. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33471991>.

119. Hu C, Hart SN, Gnanaolivu R, et al. A population-based study of genes previously implicated in breast cancer. *N Engl J Med* 2021;384:440-451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33471974>.

120. Metcalfe K, Lynch HT, Foulkes WD, et al. Oestrogen receptor status and survival in women with BRCA2-associated breast cancer. *Br J Cancer* 2019;120:398-403. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30723304>.

121. Jonasson JG, Stefansson OA, Johannsson OT, et al. Oestrogen receptor status, treatment and breast cancer prognosis in Icelandic BRCA2 mutation carriers. *Br J Cancer* 2016;115:776-783. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27537391>.

122. Olafsdottir EJ, Borg A, Jensen MB, et al. Breast cancer survival in Nordic BRCA2 mutation carriers-unconventional association with oestrogen receptor status. *Br J Cancer* 2020;123:1608-1615. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32939053>.

123. Lee LJ, Alexander B, Schnitt SJ, et al. Clinical outcome of triple negative breast cancer in BRCA1 mutation carriers and noncarriers. *Cancer* 2011;117:3093-3100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21264845>.

124. Liede A, Karlan BY, Narod SA. Cancer risks for male carriers of germline mutations in BRCA1 or BRCA2: a review of the literature. *J Clin Oncol* 2004;22:735-742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14966099>.

125. Basham VM, Lipscombe JM, Ward JM, et al. BRCA1 and BRCA2 mutations in a population-based study of male breast cancer. *Breast Cancer Res* 2002;4:R2. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11879560>.

126. Couch FJ, Farid LM, DeShano ML, et al. BRCA2 germline mutations in male breast cancer cases and breast cancer families. *Nat Genet*





1996;13:123-125. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/8673091>.

127. Ding YC, Steele L, Kuan CJ, et al. Mutations in BRCA2 and PALB2 in male breast cancer cases from the United States. *Breast Cancer Res Treat* 2011;126:771-778. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20927582>.

128. Friedman LS, Gayther SA, Kurosaki T, et al. Mutation analysis of BRCA1 and BRCA2 in a male breast cancer population. *Am J Hum Genet* 1997;60:313-319. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/9012404>.

129. Evans DG, Susnerwala I, Dawson J, et al. Risk of breast cancer in male BRCA2 carriers. *J Med Genet* 2010;47:710-711. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20587410>.

130. Tai YC, Domchek S, Parmigiani G, Chen S. Breast cancer risk among male BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst* 2007;99:1811-1814. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18042939>.

131. What are the key statistics about breast cancer in men? 2015. Available at:  
<http://www.cancer.org/cancer/breastcancerinmen/detailedguide/breast-cancer-in-men-key-statistics>. Accessed May 28, 2015.

132. Levine DA, Argenta PA, Yee CJ, et al. Fallopian tube and primary peritoneal carcinomas associated with BRCA mutations. *J Clin Oncol* 2003;21:4222-4227. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/14615451>.

133. Piver MS, Jishi MF, Tsukada Y, Nava G. Primary peritoneal carcinoma after prophylactic oophorectomy in women with a family history of ovarian cancer. A report of the Gilda Radner Familial Ovarian Cancer Registry. *Cancer* 1993;71:2751-2755. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/8467455>.

134. Arts-de Jong M, de Bock GH, van Asperen CJ, et al. Germline BRCA1/2 mutation testing is indicated in every patient with epithelial ovarian cancer: a systematic review. *Eur J Cancer* 2016;61:137-145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27209246>.

135. Pal T, Permuth-Wey J, Betts JA, et al. BRCA1 and BRCA2 mutations account for a large proportion of ovarian carcinoma cases. *Cancer* 2005;104:2807-2816. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/16284991>.

136. Schrader KA, Hurlburt J, Kalloger SE, et al. Germline BRCA1 and BRCA2 mutations in ovarian cancer: utility of a histology-based referral strategy. *Obstet Gynecol* 2012;120:235-240. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22776961>.

137. Zhang S, Royer R, Li S, et al. Frequencies of BRCA1 and BRCA2 mutations among 1,342 unselected patients with invasive ovarian cancer. *Gynecol Oncol* 2011;121:353-357. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/21324516>.

138. Song H, Cicek MS, Dicks E, et al. The contribution of deleterious germline mutations in BRCA1, BRCA2 and the mismatch repair genes to ovarian cancer in the population. *Hum Mol Genet* 2014;23:4703-4709. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24728189>.

139. Chen J, Bae E, Zhang L, et al. Penetrance of breast and ovarian cancer in women who carry a BRCA1/2 mutation and do not use risk-reducing salpingo-oophorectomy: an updated meta-analysis. *JNCI Cancer Spectr* 2020;4:pkaa029. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/32676552>.

140. Alsop K, Fereday S, Meldrum C, et al. BRCA mutation frequency and patterns of treatment response in BRCA mutation-positive women with ovarian cancer: a report from the Australian Ovarian Cancer Study Group. *J Clin Oncol* 2012;30:2654-2663. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22711857>.

141. Bolton KL, Chenevix-Trench G, Goh C, et al. Association between BRCA1 and BRCA2 mutations and survival in women with invasive



epithelial ovarian cancer. JAMA 2012;307:382-390. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22274685>.

142. Cass I, Baldwin RL, Varkey T, et al. Improved survival in women with BRCA-associated ovarian carcinoma. Cancer 2003;97:2187-2195. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12712470>.

143. Chetrit A, Hirsh-Yechezkel G, Ben-David Y, et al. Effect of BRCA1/2 mutations on long-term survival of patients with invasive ovarian cancer: the national Israeli study of ovarian cancer. J Clin Oncol 2008;26:20-25. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18165636>.

144. Tan DS, Rothermundt C, Thomas K, et al. "BRCAness" syndrome in ovarian cancer: a case-control study describing the clinical features and outcome of patients with epithelial ovarian cancer associated with BRCA1 and BRCA2 mutations. J Clin Oncol 2008;26:5530-5536. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18955455>.

145. Yang D, Khan S, Sun Y, et al. Association of BRCA1 and BRCA2 mutations with survival, chemotherapy sensitivity, and gene mutator phenotype in patients with ovarian cancer. JAMA 2011;306:1557-1565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21990299>.

146. Dong F, Davineni PK, Howitt BE, Beck AH. A BRCA1/2 mutational signature and survival in ovarian high-grade serous carcinoma. Cancer Epidemiol Biomarkers Prev 2016;25:1511-1516. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27496093>.

147. Norquist BM, Harrell MI, Brady MF, et al. Inherited mutations in women with ovarian carcinoma. JAMA Oncol 2016;2:482-490. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26720728>.

148. Bjorge T, Lie AK, Hovig E, et al. BRCA1 mutations in ovarian cancer and borderline tumours in Norway: a nested case-control study. Br J Cancer 2004;91:1829-1834. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15477862>.

149. Jazaeri AA, Lu K, Schmandt R, et al. Molecular determinants of tumor differentiation in papillary serous ovarian carcinoma. Mol Carcinog

2003;36:53-59. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/12557260>.

150. Lakhani SR, Manek S, Penault-Llorca F, et al. Pathology of ovarian cancers in BRCA1 and BRCA2 carriers. Clin Cancer Res 2004;10:2473-2481. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15073127>.

151. Press JZ, De Luca A, Boyd N, et al. Ovarian carcinomas with genetic and epigenetic BRCA1 loss have distinct molecular abnormalities. BMC Cancer 2008;8:17. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18208621>.

152. Rechsteiner M, Zimmermann AK, Wild PJ, et al. TP53 mutations are common in all subtypes of epithelial ovarian cancer and occur concomitantly with KRAS mutations in the mucinous type. Exp Mol Pathol 2013;95:235-241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23965232>.

153. Werness BA, Ramus SJ, DiCioccio RA, et al. Histopathology, FIGO stage, and BRCA mutation status of ovarian cancers from the Gilda Radner Familial Ovarian Cancer Registry. Int J Gynecol Pathol 2004;23:29-34. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14668547>.

154. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. Gastroenterology 2000;119:1447-1453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11113065>.

155. Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. Clin Cancer Res 2006;12:3209-3215. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16707622>.

156. Callahan MJ, Crum CP, Medeiros F, et al. Primary fallopian tube malignancies in BRCA-positive women undergoing surgery for ovarian cancer risk reduction. J Clin Oncol 2007;25:3985-3990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17761984>.



157. Finch A, Shaw P, Rosen B, et al. Clinical and pathologic findings of prophylactic salpingo-oophorectomies in 159 BRCA1 and BRCA2 carriers. *Gynecol Oncol* 2006;100:58-64. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16137750>.

158. Powell CB, Chen LM, McLennan J, et al. Risk-reducing salpingo-oophorectomy (RRSO) in BRCA mutation carriers: experience with a consecutive series of 111 patients using a standardized surgical-pathological protocol. *Int J Gynecol Cancer* 2011;21:846-851. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21670699>.

159. Powell CB, Kenley E, Chen LM, et al. Risk-reducing salpingo-oophorectomy in BRCA mutation carriers: role of serial sectioning in the detection of occult malignancy. *J Clin Oncol* 2005;23:127-132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15625367>.

160. Shaw PA, Rouzbahman M, Pizer ES, et al. Candidate serous cancer precursors in fallopian tube epithelium of BRCA1/2 mutation carriers. *Mod Pathol* 2009;22:1133-1138. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19543244>.

161. Medeiros F, Muto MG, Lee Y, et al. The tubal fimbria is a preferred site for early adenocarcinoma in women with familial ovarian cancer syndrome. *Am J Surg Pathol* 2006;30:230-236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16434898>.

162. Kindelberger DW, Lee Y, Miron A, et al. Intraepithelial carcinoma of the fimbria and pelvic serous carcinoma: Evidence for a causal relationship. *Am J Surg Pathol* 2007;31:161-169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17255760>.

163. Agalliu I, Gern R, Leanza S, Burk RD. Associations of high-grade prostate cancer with BRCA1 and BRCA2 founder mutations. *Clin Cancer Res* 2009;15:1112-1120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19188187>.

164. Leongamornlert D, Mahmud N, Tymrakiewicz M, et al. Germline BRCA1 mutations increase prostate cancer risk. *Br J Cancer*

2012;106:1697-1701. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22516946>.

165. Nicolosi P, Ledet E, Yang S, et al. Prevalence of germline variants in prostate cancer and implications for current genetic testing guidelines. *JAMA Oncol* 2019;5:523-528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30730552>.

166. Giri VN, Knudsen KE, Kelly WK, et al. Implementation of germline testing for prostate cancer: Philadelphia Prostate Cancer Consensus Conference 2019. *J Clin Oncol* 2020;38:2798-2811. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32516092>.

167. Nyberg T, Frost D, Barrowdale D, et al. Prostate cancer risks for male BRCA1 and BRCA2 mutation carriers: a prospective cohort study. *Eur Urol* 2020;77:24-35. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31495749>.

168. Abida W, Armenia J, Gopalan A, et al. Prospective genomic profiling of prostate cancer across disease states reveals germline and somatic alterations that may affect clinical decision making. *JCO Precis Oncol* 2017;2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28825054>.

169. Pritchard CC, Mateo J, Walsh MF, et al. Inherited DNA-repair gene mutations in men with metastatic prostate cancer. *N Engl J Med* 2016;375:443-453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27433846>.

170. Darst BF, Dadaev T, Saunders E, et al. Germline sequencing DNA repair genes in 5545 men with aggressive and nonaggressive prostate cancer. *J Natl Cancer Inst* 2021;113:616-625. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32853339>.

171. Castro E, Goh C, Olmos D, et al. Germline BRCA mutations are associated with higher risk of nodal involvement, distant metastasis, and poor survival outcomes in prostate cancer. *J Clin Oncol* 2013;31:1748-1757. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23569316>.



172. Na R, Zheng SL, Han M, et al. Germline mutations in ATM and BRCA1/2 distinguish risk for lethal and indolent prostate cancer and are associated with early age at death. *Eur Urol* 2017;71:740-747. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27989354>.

173. Kirchhoff T, Kauff ND, Mitra N, et al. BRCA mutations and risk of prostate cancer in Ashkenazi Jews. *Clin Cancer Res* 2004;10:2918-2921. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15131025>.

174. Gallagher DJ, Gaudet MM, Pal P, et al. Germline BRCA mutations denote a clinicopathologic subset of prostate cancer. *Clin Cancer Res* 2010;16:2115-2121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20215531>.

175. Hamel N, Kotar K, Foulkes WD. Founder mutations in BRCA1/2 are not frequent in Canadian Ashkenazi Jewish men with prostate cancer. *BMC Med Genet* 2003;4:7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12911837>.

176. Nastiuk KL, Mansukhani M, Terry MB, et al. Common mutations in BRCA1 and BRCA2 do not contribute to early prostate cancer in Jewish men. *Prostate* 1999;40:172-177. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10398279>.

177. Goggins M, Schutte M, Lu J, et al. Germline BRCA2 gene mutations in patients with apparently sporadic pancreatic carcinomas. *Cancer Res* 1996;56:5360-5364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8968085>.

178. Lal G, Liu G, Schmocker B, et al. Inherited predisposition to pancreatic adenocarcinoma: role of family history and germ-line p16, BRCA1, and BRCA2 mutations. *Cancer Res* 2000;60:409-416. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10667595>.

179. Murphy KM, Brune KA, Griffin C, et al. Evaluation of candidate genes MAP2K4, MADH4, ACVR1B, and BRCA2 in familial pancreatic cancer: deleterious BRCA2 mutations in 17%. *Cancer Res* 2002;62:3789-3793. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12097290>.

180. Couch FJ, Johnson MR, Rabe KG, et al. The prevalence of BRCA2 mutations in familial pancreatic cancer. *Cancer Epidemiol Biomarkers Prev* 2007;16:342-346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17301269>.

181. Ghiorzo P, Fornarini G, Sciallero S, et al. CDKN2A is the main susceptibility gene in Italian pancreatic cancer families. *J Med Genet* 2012;49:164-170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22368299>.

182. Lucas AL, Shakya R, Lipsyc MD, et al. High prevalence of BRCA1 and BRCA2 germline mutations with loss of heterozygosity in a series of resected pancreatic adenocarcinoma and other neoplastic lesions. *Clin Cancer Res* 2013;19:3396-3403. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959126/>.

183. Holter S, Borgida A, Dodd A, et al. Germline BRCA mutations in a large clinic-based cohort of patients with pancreatic adenocarcinoma. *J Clin Oncol* 2015;33:3124-3129. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25940717>.

184. Zhen DB, Rabe KG, Gallinger S, et al. BRCA1, BRCA2, PALB2, and CDKN2A mutations in familial pancreatic cancer: a PACGENE study. *Genet Med* 2015;17:569-577. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25356972>.

185. Salo-Mullen EE, O'Reilly EM, Kelsen DP, et al. Identification of germline genetic mutations in patients with pancreatic cancer. *Cancer* 2015;121:4382-4388. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26440929>.

186. Mandelker D, Zhang L, Kemel Y, et al. Mutation detection in patients with advanced cancer by universal sequencing of cancer-related genes in tumor and normal DNA vs guideline-based germline testing. *JAMA* 2017;318:825-835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28873162>.

187. Shindo K, Yu J, Suenaga M, et al. Deleterious germline mutations in patients with apparently sporadic pancreatic adenocarcinoma. *J Clin*





Oncol 2017;35:3382-3390. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28767289>.

188. Huang KL, Mashl RJ, Wu Y, et al. Pathogenic germline variants in 10,389 adult cancers. Cell 2018;173:355-370 e314. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29625052>.

189. Chaffee KG, Oberg AL, McWilliams RR, et al. Prevalence of germline mutations in cancer genes among pancreatic cancer patients with a positive family history. Genet Med 2018;20:119-127. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28726808>.

190. Hu C, Hart SN, Polley EC, et al. Association between inherited germline mutations in cancer predisposition genes and risk of pancreatic cancer. JAMA 2018;319:2401-2409. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29922827>.

191. Ferrone CR, Levine DA, Tang LH, et al. BRCA germline mutations in Jewish patients with pancreatic adenocarcinoma. J Clin Oncol 2009;27:433-438. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19064968>.

192. de Jonge MM, Mooyaart AL, Vreeswijk MP, et al. Linking uterine serous carcinoma to BRCA1/2-associated cancer syndrome: A meta-analysis and case report. Eur J Cancer 2017;72:215-225. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28049106>.

193. Lavie O, Ben-Arie A, Segev Y, et al. BRCA germline mutations in women with uterine serous carcinoma--still a debate. Int J Gynecol Cancer 2010;20:1531-1534. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21119368>.

194. Saule C, Mouret-Fourme E, Briaux A, et al. Risk of serous endometrial carcinoma in women with pathogenic BRCA1/2 variant after risk-reducing salpingo-oophorectomy. J Natl Cancer Inst 2018;110.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28954295>.

195. Laitman Y, Michaelson-Cohen R, Levi E, et al. Uterine cancer in Jewish Israeli BRCA1/2 mutation carriers. Cancer 2019;125:698-703.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30489631>.

196. de Jonge MM, de Kroon CD, Jenner DJ, et al. Endometrial cancer risk in women with germline BRCA1 or BRCA2 mutations: multicenter cohort study. J Natl Cancer Inst 2021;113:1203-1211. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33710348>.

197. Shu CA, Pike MC, Jotwani AR, et al. Uterine cancer after risk-reducing salpingo-oophorectomy without hysterectomy in women with BRCA mutations. JAMA Oncol 2016;2:1434-1440. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27367496>.

198. Beiner ME, Finch A, Rosen B, et al. The risk of endometrial cancer in women with BRCA1 and BRCA2 mutations. A prospective study. Gynecol Oncol 2007;104:7-10. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16962648>.

199. Lee YC, Milne RL, Lheureux S, et al. Risk of uterine cancer for BRCA1 and BRCA2 mutation carriers. Eur J Cancer 2017;84:114-120.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28802188>.

200. Gumaste PV, Penn LA, Cymerman RM, et al. Skin cancer risk in BRCA1/2 mutation carriers. Br J Dermatol 2015;172:1498-1506. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25524463>.

201. Iqbal J, Nussenzweig A, Lubinski J, et al. The incidence of leukaemia in women with BRCA1 and BRCA2 mutations: an International Prospective Cohort Study. Br J Cancer 2016;114:1160-1164. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26986251>.

202. Lorenzo Bermejo J, Hemminki K. Risk of cancer at sites other than the breast in Swedish families eligible for BRCA1 or BRCA2 mutation testing. Ann Oncol 2004;15:1834-1841. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15550590>.





203. Cancer risks in BRCA2 mutation carriers. The Breast Cancer Linkage Consortium. J Natl Cancer Inst 1999;91:1310-1316. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10433620>.

204. Moran A, O'Hara C, Khan S, et al. Risk of cancer other than breast or ovarian in individuals with BRCA1 and BRCA2 mutations. Fam Cancer 2012;11:235-242. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22187320>.

205. Alter BP, Rosenberg PS, Brody LC. Clinical and molecular features associated with biallelic mutations in FANCD1/BRCA2. J Med Genet 2007;44:1-9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16825431>.

206. Keupp K, Hampp S, Hubbel A, et al. Biallelic germline BRCA1 mutations in a patient with early onset breast cancer, mild Fanconi anemia-like phenotype, and no chromosome fragility. Mol Genet Genomic Med 2019;7:e863. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31347298>.

207. Chirita-Emandi A, Andreescu N, Popa C, et al. Biallelic variants in BRCA1 gene cause a recognisable phenotype within chromosomal instability syndromes reframed as BRCA1 deficiency. J Med Genet 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32843487>.

208. Sawyer SL, Tian L, Kahkonen M, et al. Biallelic mutations in BRCA1 cause a new Fanconi anemia subtype. Cancer Discov 2015;5:135-142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25472942>.

209. Freire BL, Homma TK, Funari MFA, et al. Homozygous loss of function BRCA1 variant causing a Fanconi-anemia-like phenotype, a clinical report and review of previous patients. Eur J Med Genet 2018;61:130-133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29133208>.

210. Berg WA. How well does supplemental screening magnetic resonance imaging work in high-risk women? J Clin Oncol 2014;32:2193-2196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24934782>.

211. Buist DS, Porter PL, Lehman C, et al. Factors contributing to mammography failure in women aged 40-49 years. J Natl Cancer Inst 2004;96:1432-1440. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15467032>.

212. Mandelson MT, Oestreicher N, Porter PL, et al. Breast density as a predictor of mammographic detection: comparison of interval- and screen-detected cancers. J Natl Cancer Inst 2000;92:1081-1087. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10880551>.

213. Tilanus-Linthorst M, Verhoog L, Obdeijn IM, et al. A BRCA1/2 mutation, high breast density and prominent pushing margins of a tumor independently contribute to a frequent false-negative mammography. Int J Cancer 2002;102:91-95. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12353239>.

214. van Gils CH, Otten JD, Verbeek AL, et al. Effect of mammographic breast density on breast cancer screening performance: a study in Nijmegen, The Netherlands. J Epidemiol Community Health 1998;52:267-271. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9616416>.

215. Gilliland FD, Joste N, Stauber PM, et al. Biologic characteristics of interval and screen-detected breast cancers. J Natl Cancer Inst 2000;92:743-749. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10793111>.

216. Kriege M, Brekelmans CT, Boetes C, et al. Efficacy of MRI and mammography for breast-cancer screening in women with a familial or genetic predisposition. N Engl J Med 2004;351:427-437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15282350>.

217. Kuhl CK, Schrading S, Leutner CC, et al. Mammography, breast ultrasound, and magnetic resonance imaging for surveillance of women at high familial risk for breast cancer. J Clin Oncol 2005;23:8469-8476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16293877>.

218. Leach MO, Boggis CR, Dixon AK, et al. Screening with magnetic resonance imaging and mammography of a UK population at high familial risk of breast cancer: a prospective multicentre cohort study (MARIBS).



Lancet 2005;365:1769-1778. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/15910949>.

219. Riedl CC, Ponhold L, Flory D, et al. Magnetic resonance imaging of the breast improves detection of invasive cancer, preinvasive cancer, and premalignant lesions during surveillance of women at high risk for breast cancer. Clin Cancer Res 2007;13:6144-6152. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17947480>.

220. Sardanelli F, Podo F, D'Agnolo G, et al. Multicenter comparative multimodality surveillance of women at genetic-familial high risk for breast cancer (HIBCRIT study): interim results. Radiology 2007;242:698-715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17244718>.

221. Warner E, Plewes DB, Hill KA, et al. Surveillance of BRCA1 and BRCA2 mutation carriers with magnetic resonance imaging, ultrasound, mammography, and clinical breast examination. JAMA 2004;292:1317-1325. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15367553>.

222. Passaperuma K, Warner E, Causer PA, et al. Long-term results of screening with magnetic resonance imaging in women with BRCA mutations. Br J Cancer 2012;107:24-30. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22588560>.

223. Lehman CD, Lee JM, DeMartini WB, et al. Screening MRI in women with a personal history of breast cancer. J Natl Cancer Inst 2016;108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26744477>.

224. Phi XA, Saadatmand S, De Bock GH, et al. Contribution of mammography to MRI screening in BRCA mutation carriers by BRCA status and age: individual patient data meta-analysis. Br J Cancer 2016;114:631-637. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/26908327>.

225. Le-Petross HT, Whitman GJ, Atchley DP, et al. Effectiveness of alternating mammography and magnetic resonance imaging for screening women with deleterious BRCA mutations at high risk of breast cancer. Cancer 2011;117:3900-3907. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/21365619>.

226. Goldfrank D, Chuai S, Bernstein JL, et al. Effect of mammography on breast cancer risk in women with mutations in BRCA1 or BRCA2. Cancer Epidemiol Biomarkers Prev 2006;15:2311-2313. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17119064>.

227. Narod SA, Lubinski J, Ghadirian P, et al. Screening mammography and risk of breast cancer in BRCA1 and BRCA2 mutation carriers: a case-control study. Lancet Oncol 2006;7:402-406. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/16648044>.

228. Ciatto S, Houssami N, Bernardi D, et al. Integration of 3D digital mammography with tomosynthesis for population breast-cancer screening (STORM): a prospective comparison study. Lancet Oncol 2013;14:583-589. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23623721>.

229. Skaane P, Bandos AI, Gullien R, et al. Comparison of digital mammography alone and digital mammography plus tomosynthesis in a population-based screening program. Radiology 2013;267:47-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23297332>.

230. Rafferty EA, Park JM, Philpotts LE, et al. Assessing radiologist performance using combined digital mammography and breast tomosynthesis compared with digital mammography alone: results of a multicenter, multireader trial. Radiology 2013;266:104-113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23169790>.

231. Friedewald SM, Rafferty EA, Conant EF. Breast cancer screening with tomosynthesis and digital mammography-reply. JAMA 2014;312:1695-1696. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/25335157>.

232. Lourenco AP, Barry-Brooks M, Baird GL, et al. Changes in recall type and patient treatment following implementation of screening digital breast tomosynthesis. Radiology 2015;274:337-342. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/25247407>.

233. Rose SL, Tidwell AL, Ice MF, et al. A reader study comparing prospective tomosynthesis interpretations with retrospective readings of

the corresponding FFDM examinations. Acad Radiol 2014;21:1204-1210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25107868>.

234. Destounis S, Arieno A, Morgan R. Initial experience with combination digital breast tomosynthesis plus full field digital mammography or full field digital mammography alone in the screening environment. J Clin Imaging Sci 2014;4:9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24744966>.

235. Margolies L, Cohen A, Sonnenblick E, et al. Digital breast tomosynthesis changes management in patients seen at a tertiary care breast center. ISRN Radiol 2014;2014:658929. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24967297>.

236. Lang K, Andersson I, Rosso A, et al. Performance of one-view breast tomosynthesis as a stand-alone breast cancer screening modality: results from the Malmö Breast Tomosynthesis Screening Trial, a population-based study. Eur Radiol 2016;26:184-190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25929946>.

237. Gilbert FJ, Tucker L, Gillan MG, et al. Accuracy of digital breast tomosynthesis for depicting breast cancer subgroups in a UK retrospective reading study (TOMMY Trial). Radiology 2015;277:697-706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26176654>.

238. Zuckerman SP, Conant EF, Keller BM, et al. Implementation of synthesized two-dimensional mammography in a population-based digital breast tomosynthesis screening program. Radiology 2016;160366. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27467468>.

239. Skaane P, Bandos AI, Eben EB, et al. Two-view digital breast tomosynthesis screening with synthetically reconstructed projection images: comparison with digital breast tomosynthesis with full-field digital mammographic images. Radiology 2014;271:655-663. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24484063>.

240. McDonald RJ, McDonald JS, Kallmes DF, et al. Gadolinium deposition in human brain tissues after contrast-enhanced MR imaging in

adult patients without intracranial abnormalities. Radiology 2017;285:546-554. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28653860>.

241. Stojanov D, Aracki-Trenkic A, Benedeto-Stojanov D. Gadolinium deposition within the dentate nucleus and globus pallidus after repeated administrations of gadolinium-based contrast agents-current status. Neuroradiology 2016;58:433-441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26873830>.

242. Darrah TH, Prutsman-Pfeiffer JJ, Poreda RJ, et al. Incorporation of excess gadolinium into human bone from medical contrast agents. Metallomics 2009;1:479-488. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21305156>.

243. Lowry KP, Lee JM, Kong CY, et al. Annual screening strategies in BRCA1 and BRCA2 gene mutation carriers: a comparative effectiveness analysis. Cancer 2012;118:2021-2030. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21935911>.

244. Saslow D, Boetes C, Burke W, et al. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. CA Cancer J Clin 2007;57:75-89. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17392385>.

245. Stoutjesdijk MJ, Boetes C, Jager GJ, et al. Magnetic resonance imaging and mammography in women with a hereditary risk of breast cancer. J Natl Cancer Inst 2001;93:1095-1102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11459871>.

246. Naranjo ID, Sogani J, Saccarelli C, et al. MRI screening of BRCA mutation carriers: comparison of standard protocol and abbreviated protocols with and without T2-weighted images. AJR Am J Roentgenol 2021. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34935399>.

247. Thomas M. Abbreviated breast MRI-good, but not all the same. AJR Am J Roentgenol 2022. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34985317>.





248. Gao Y, Goldberg JE, Young TK, et al. Breast cancer screening in high-risk men: a 12-year longitudinal observational study of male breast imaging utilization and outcomes. *Radiology* 2019;293:282-291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31526252>.

249. Li X, You R, Wang X, et al. Effectiveness of prophylactic surgeries in BRCA1 or BRCA2 mutation carriers: a meta-analysis and systematic review. *Clin Cancer Res* 2016;22:3971-3981. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26979395>.

250. Honold F, Camus M. Prophylactic mastectomy versus surveillance for the prevention of breast cancer in women's BRCA carriers. *Medwave* 2018;18:e7161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30052622>.

251. Hartmann LC, Schaid DJ, Woods JE, et al. Efficacy of bilateral prophylactic mastectomy in women with a family history of breast cancer. *N Engl J Med* 1999;340:77-84. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9887158>.

252. Hartmann LC, Sellers TA, Schaid DJ, et al. Efficacy of bilateral prophylactic mastectomy in BRCA1 and BRCA2 gene mutation carriers. *J Natl Cancer Inst* 2001;93:1633-1637. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11698567>.

253. Meijers-Heijboer H, van Geel B, van Putten WL, et al. Breast cancer after prophylactic bilateral mastectomy in women with a BRCA1 or BRCA2 mutation. *N Engl J Med* 2001;345:159-164. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11463009>.

254. Rebbeck TR, Friebel T, Lynch HT, et al. Bilateral prophylactic mastectomy reduces breast cancer risk in BRCA1 and BRCA2 mutation carriers: the PROSE Study Group. *J Clin Oncol* 2004;22:1055-1062. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14981104>.

255. Carbine NE, Lostumbo L, Wallace J, Ko H. Risk-reducing mastectomy for the prevention of primary breast cancer. *Cochrane Database Syst Rev* 2018;4:CD002748. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29620792>.

256. Morrow M, Mehrara B. Prophylactic mastectomy and the timing of breast reconstruction. *Br J Surg* 2009;96:1-2. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19109821>.

257. Jakub JW, Peled AW, Gray RJ, et al. Oncologic safety of prophylactic nipple-sparing mastectomy in a population with BRCA mutations: a multi-institutional study. *JAMA Surg* 2018;153:123-129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28903167>.

258. Hartmann LC, Lindor NM. The role of risk-reducing surgery in hereditary breast and ovarian cancer. *N Engl J Med* 2016;374:454-468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26840135>.

259. Cummings SR, Eckert S, Krueger KA, et al. The effect of raloxifene on risk of breast cancer in postmenopausal women: results from the MORE randomized trial. Multiple outcomes of raloxifene evaluation. *JAMA* 1999;281:2189-2197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10376571>.

260. Cuzick J, Sestak I, Bonanni B, et al. Selective oestrogen receptor modulators in prevention of breast cancer: an updated meta-analysis of individual participant data. *Lancet* 2013;381:1827-1834. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23639488>.

261. Lippman ME, Cummings SR, Disch DP, et al. Effect of raloxifene on the incidence of invasive breast cancer in postmenopausal women with osteoporosis categorized by breast cancer risk. *Clin Cancer Res* 2006;12:5242-5247. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16951244>.

262. Martino S, Cauley JA, Barrett-Connor E, et al. Continuing outcomes relevant to Evista: breast cancer incidence in postmenopausal osteoporotic women in a randomized trial of raloxifene. *J Natl Cancer Inst* 2004;96:1751-1761. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15572757>.

263. Vogel VG, Costantino JP, Wickerham DL, et al. Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene





(STAR) P-2 trial. JAMA 2006;295:2727-2741. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16754727>.

264. Vogel VG, Costantino JP, Wickerham DL, et al. Update of the National Surgical Adjuvant Breast and Bowel Project Study of Tamoxifen and Raloxifene (STAR) P-2 Trial: preventing breast cancer. Cancer Prev Res (Phila) 2010;3:696-706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20404000>.

265. Powles TJ, Ashley S, Tidy A, et al. Twenty-year follow-up of the Royal Marsden randomized, double-blinded tamoxifen breast cancer prevention trial. J Natl Cancer Inst 2007;99:283-290. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17312305>.

266. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for the prevention of breast cancer: current status of the National Surgical Adjuvant Breast and Bowel Project P-1 study. J Natl Cancer Inst 2005;97:1652-1662. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16288118>.

267. Metcalfe K, Lynch HT, Ghadirian P, et al. Contralateral breast cancer in BRCA1 and BRCA2 mutation carriers. J Clin Oncol 2004;22:2328-2335. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15197194>.

268. Gronwald J, Tung N, Foulkes WD, et al. Tamoxifen and contralateral breast cancer in BRCA1 and BRCA2 carriers: an update. Int J Cancer 2006;118:2281-2284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16331614>.

269. Narod SA, Brunet JS, Ghadirian P, et al. Tamoxifen and risk of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers: a case-control study. Hereditary Breast Cancer Clinical Study Group. Lancet 2000;356:1876-1881. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11130383>.

270. King MC, Wieand S, Hale K, et al. Tamoxifen and breast cancer incidence among women with inherited mutations in BRCA1 and BRCA2: National Surgical Adjuvant Breast and Bowel Project (NSABP-P1) Breast

Cancer Prevention Trial. JAMA 2001;286:2251-2256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11710890>.

271. Ingle JN, Liu M, Wickerham DL, et al. Selective estrogen receptor modulators and pharmacogenomic variation in ZNF423 regulation of BRCA1 expression: individualized breast cancer prevention. Cancer Discov 2013;3:812-825. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23764426>.

272. Goss PE, Ingle JN, Ales-Martinez JE, et al. Exemestane for breast-cancer prevention in postmenopausal women. N Engl J Med 2011;364:2381-2391. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21639806>.

273. Cuzick J, Sestak I, Forbes JF, et al. Anastrozole for prevention of breast cancer in high-risk postmenopausal women (IBIS-II): an international, double-blind, randomised placebo-controlled trial. Lancet 2014;383:1041-1048. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24333009>.

274. Nemati Shafaei M, Gutierrez-Barrera AM, Lin HY, Arun B. Aromatase inhibitors and the risk of contralateral breast cancer in BRCA mutation carriers. J Clin Oncol 2015;33:3-3. Available at: [http://ascopubs.org/doi/abs/10.1200/jco.2015.33.28\\_suppl.3](http://ascopubs.org/doi/abs/10.1200/jco.2015.33.28_suppl.3).

275. Narod SA, Dube MP, Klijn J, et al. Oral contraceptives and the risk of breast cancer in BRCA1 and BRCA2 mutation carriers. J Natl Cancer Inst 2002;94:1773-1779. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12464649>.

276. Haile RW, Thomas DC, McGuire V, et al. BRCA1 and BRCA2 mutation carriers, oral contraceptive use, and breast cancer before age 50. Cancer Epidemiol Biomarkers Prev 2006;15:1863-1870. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17021353>.

277. Milne RL, Knight JA, John EM, et al. Oral contraceptive use and risk of early-onset breast cancer in carriers and noncarriers of BRCA1 and BRCA2 mutations. Cancer Epidemiol Biomarkers Prev 2005;14:350-356. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15734957>.



278. Iodice S, Barile M, Rotmensz N, et al. Oral contraceptive use and breast or ovarian cancer risk in BRCA1/2 carriers: a meta-analysis. *Eur J Cancer* 2010;46:2275-2284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20537530>.

279. Moorman PG, Havrilesky LJ, Gierisch JM, et al. Oral contraceptives and risk of ovarian cancer and breast cancer among high-risk women: a systematic review and meta-analysis. *J Clin Oncol* 2013;31:4188-4198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24145348>.

280. Lee E, Ma H, McKean-Cowdin R, et al. Effect of reproductive factors and oral contraceptives on breast cancer risk in BRCA1/2 mutation carriers and noncarriers: results from a population-based study. *Cancer Epidemiol Biomarkers Prev* 2008;17:3170-3178. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18990759>.

281. Jacobs IJ, Menon U, Ryan A, et al. Ovarian cancer screening and mortality in the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS): a randomised controlled trial. *Lancet* 2016;387:945-956. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26707054>.

282. Menon U, Gentry-Maharaj A, Hallett R, et al. Sensitivity and specificity of multimodal and ultrasound screening for ovarian cancer, and stage distribution of detected cancers: results of the prevalence screen of the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS). *Lancet Oncol* 2009;10:327-340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19282241>.

283. Rosenthal AN, Fraser LSM, Philpott S, et al. Evidence of stage shift in women diagnosed with ovarian cancer during phase II of the United Kingdom Familial Ovarian Cancer Screening Study. *J Clin Oncol* 2017;35:1411-1420. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28240969>.

284. Skates SJ, Greene MH, Buys SS, et al. Early detection of ovarian cancer using the risk of ovarian cancer algorithm with frequent CA125 testing in women at increased familial risk - combined results from two screening trials. *Clin Cancer Res* 2017;23:3628-3637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28143870>.

285. Satagopan JM, Boyd J, Kauff ND, et al. Ovarian cancer risk in Ashkenazi Jewish carriers of BRCA1 and BRCA2 mutations. *Clin Cancer Res* 2002;8:3776-3781. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12473589>.

286. Finch AP, Lubinski J, Moller P, et al. Impact of oophorectomy on cancer incidence and mortality in women with a BRCA1 or BRCA2 mutation. *J Clin Oncol* 2014;32:1547-1553. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24567435>.

287. Rebbeck TR, Kauff ND, Domchek SM. Meta-analysis of risk reduction estimates associated with risk-reducing salpingo-oophorectomy in BRCA1 or BRCA2 mutation carriers. *J Natl Cancer Inst* 2009;101:80-87. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19141781>.

288. Kauff ND, Domchek SM, Friebel TM, et al. Risk-reducing salpingo-oophorectomy for the prevention of BRCA1- and BRCA2-associated breast and gynecologic cancer: a multicenter, prospective study. *J Clin Oncol* 2008;26:1331-1337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18268356>.

289. Kauff ND, Satagopan JM, Robson ME, et al. Risk-reducing salpingo-oophorectomy in women with a BRCA1 or BRCA2 mutation. *N Engl J Med* 2002;346:1609-1615. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12023992>.

290. Kemei Y, Kauff ND, Robson ME, et al. Four-year follow-up of outcomes following risk-reducing salpingo-oophorectomy in BRCA mutation carriers [abstract]. *J Clin Oncol* (Meeting Abstracts) 2005;23(Supple 16):Abstract 1013. Available at: [http://meeting.ascopubs.org/cgi/content/abstract/23/16\\_suppl/1013](http://meeting.ascopubs.org/cgi/content/abstract/23/16_suppl/1013).

291. Rebbeck TR, Levin AM, Eisen A, et al. Breast cancer risk after bilateral prophylactic oophorectomy in BRCA1 mutation carriers. *J Natl Cancer Inst* 1999;91:1475-1479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10469748>.

292. Rebbeck TR, Lynch HT, Neuhausen SL, et al. Prophylactic oophorectomy in carriers of BRCA1 or BRCA2 mutations. *N Engl J Med*

- 2002;346:1616-1622. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12023993>.
293. Harmsen MG, Piek JMJ, Bulten J, et al. Peritoneal carcinomatosis after risk-reducing surgery in BRCA1/2 mutation carriers. *Cancer* 2018;124:952-959. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29315498>.
294. Sherman ME, Piedmonte M, Mai PL, et al. Pathologic findings at risk-reducing salpingo-oophorectomy: primary results from Gynecologic Oncology Group Trial GOG-0199. *J Clin Oncol* 2014;32:3275-3283. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25199754>.
295. Eisen A, Lubinski J, Klijn J, et al. Breast cancer risk following bilateral oophorectomy in BRCA1 and BRCA2 mutation carriers: an international case-control study. *J Clin Oncol* 2005;23:7491-7496. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16234515>.
296. Xiao YL, Wang K, Liu Q, et al. Risk reduction and survival benefit of risk-reducing salpingo-oophorectomy in hereditary breast cancer: meta-analysis and systematic review. *Clin Breast Cancer* 2019;19:e48-e65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30470623>.
297. Domchek SM, Friebel TM, Neuhausen SL, et al. Mortality after bilateral salpingo-oophorectomy in BRCA1 and BRCA2 mutation carriers: a prospective cohort study. *Lancet Oncol* 2006;7:223-229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16510331>.
298. Domchek SM, Friebel TM, Singer CF, et al. Association of risk-reducing surgery in BRCA1 or BRCA2 mutation carriers with cancer risk and mortality. *JAMA* 2010;304:967-975. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20810374>.
299. Metcalfe K, Lynch HT, Foulkes WD, et al. Effect of oophorectomy on survival after breast cancer in BRCA1 and BRCA2 mutation carriers. *JAMA Oncol* 2015;1:306-313. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26181175>.
300. Heemskerk-Gerritsen BA, Seynaeve C, van Asperen CJ, et al. Breast cancer risk after salpingo-oophorectomy in healthy BRCA1/2 mutation carriers: revisiting the evidence for risk reduction. *J Natl Cancer Inst* 2015;107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25788320>.
301. Chai X, Domchek S, Kauff N, et al. RE: Breast cancer risk after salpingo-oophorectomy in healthy BRCA1/2 mutation carriers: revisiting the evidence for risk reduction. *J Natl Cancer Inst* 2015;107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26264690>.
302. Terry MB, Daly MB, Phillips KA, et al. Risk-reducing oophorectomy and breast cancer risk across the spectrum of familial risk. *J Natl Cancer Inst* 2019;111:331-334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30496449>.
303. Kotsopoulos J, Huzarski T, Gronwald J, et al. Bilateral oophorectomy and breast cancer risk in BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst* 2017;109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27601060>.
304. Stjepanovic N, Villacampa G, Nead KT, et al. Association of premenopausal risk-reducing salpingo-oophorectomy with breast cancer risk in BRCA1/2 mutation carriers: maximising bias-reduction. *Eur J Cancer* 2020;132:53-60. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32325420>.
305. Choi YH, Terry MB, Daly MB, et al. Association of risk-reducing salpingo-oophorectomy with breast cancer risk in women with BRCA1 and BRCA2 pathogenic variants. *JAMA Oncol* 2021;7:585-592. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33630024>.
306. Marchetti C, De Felice F, Boccia S, et al. Hormone replacement therapy after prophylactic risk-reducing salpingo-oophorectomy and breast cancer risk in BRCA1 and BRCA2 mutation carriers: A meta-analysis. *Crit Rev Oncol Hematol* 2018;132:111-115. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30447915>.
307. Gordhandas S, Norquist BM, Pennington KP, et al. Hormone replacement therapy after risk reducing salpingo-oophorectomy in patients



with BRCA1 or BRCA2 mutations; a systematic review of risks and benefits. *Gynecol Oncol* 2019;153:192-200. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30661763>.

308. Chlebowski RT, Prentice RL. Menopausal hormone therapy in BRCA1 mutation carriers: uncertainty and caution. *J Natl Cancer Inst* 2008;100:1341-1343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18812547>.

309. Garber JE, Hartman AR. Prophylactic oophorectomy and hormone replacement therapy: protection at what price? *J Clin Oncol* 2004;22:978-980. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14981100>.

310. McAlpine JN, Hanley GE, Woo MM, et al. Opportunistic salpingectomy: uptake, risks, and complications of a regional initiative for ovarian cancer prevention. *Am J Obstet Gynecol* 2014;210:471.e471-411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24412119>.

311. Findley AD, Siedhoff MT, Hobbs KA, et al. Short-term effects of salpingectomy during laparoscopic hysterectomy on ovarian reserve: a pilot randomized controlled trial. *Fertil Steril* 2013;100:1704-1708. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23993887>.

312. Daly MB, Drescher CW, Yates MS, et al. Salpingectomy as a means to reduce ovarian cancer risk. *Cancer Prev Res (Phila)* 2015;8:342-348. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25586903>.

313. Chlebowski RT, Rohan TE, Manson JE, et al. Breast cancer after use of estrogen plus progestin and estrogen alone: analyses of data from 2 Women's Health Initiative randomized clinical trials. *JAMA Oncol* 2015;1:296-305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26181174>.

314. College of American Pathologists (CAP). Protocol for the Examination of Specimens From Patients With Carcinoma of the Ovary. 2009. Available at: [http://www.cap.org/apps/docs/committees/cancer/cancer\\_protocols/2009/Ovary\\_09protocol.pdf](http://www.cap.org/apps/docs/committees/cancer/cancer_protocols/2009/Ovary_09protocol.pdf). Accessed March 2011.

315. McLaughlin JR, Risch HA, Lubinski J, et al. Reproductive risk factors for ovarian cancer in carriers of BRCA1 or BRCA2 mutations: a case-control study. *Lancet Oncol* 2007;8:26-34. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17196508>.

316. Narod SA, Risch H, Moslehi R, et al. Oral contraceptives and the risk of hereditary ovarian cancer. Hereditary Ovarian Cancer Clinical Study Group. *N Engl J Med* 1998;339:424-428. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9700175>.

317. DeSantis CE, Ma J, Gaudet MM, et al. Breast cancer statistics, 2019. *CA Cancer J Clin* 2019;69:438-451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31577379>.

318. Torre LA, Trabert B, DeSantis CE, et al. Ovarian cancer statistics, 2018. *CA Cancer J Clin* 2018;68:284-296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29809280>.

319. Domchek SM, Robson ME. Update on genetic testing in gynecologic cancer. *J Clin Oncol* 2019;37:2501-2509. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31403865>.

320. Apostolou P, Fostira F. Hereditary breast cancer: the era of new susceptibility genes. *Biomed Res Int* 2013;2013:747318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23586058>.

321. Marabelli M, Cheng SC, Parmigiani G. Penetrance of ATM gene mutations in breast cancer: a meta-analysis of different measures of risk. *Genet Epidemiol* 2016;40:425-431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27112364>.

322. Easton DF, Pharoah PD, Antoniou AC, et al. Gene-panel sequencing and the prediction of breast-cancer risk. *N Engl J Med* 2015;372:2243-2257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26014596>.

323. Couch FJ, Shimelis H, Hu C, et al. Associations between cancer predisposition testing panel genes and breast cancer. *JAMA Oncol* 2017;3:1190-1196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28418444>.





324. Kurian AW, Hughes E, Handorf EA, et al. Breast and ovarian cancer penetrance estimates derived from germline multiple-gene sequencing results in women. *JCO Precision Oncology* 2017;1:1-12. Available at: <http://ascopubs.org/doi/abs/10.1200/PO.16.00066>.

325. Lu HM, Li S, Black MH, et al. Association of breast and ovarian cancers with predisposition genes identified by large-scale sequencing. *JAMA Oncol* 2019;5:51-57. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30128536>.

326. Hauke J, Horvath J, Gross E, et al. Gene panel testing of 5589 BRCA1/2-negative index patients with breast cancer in a routine diagnostic setting: results of the German Consortium for Hereditary Breast and Ovarian Cancer. *Cancer Med* 2018;7:1349-1358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29522266>.

327. Bernstein JL, Haile RW, Stovall M, et al. Radiation exposure, the ATM Gene, and contralateral breast cancer in the women's environmental cancer and radiation epidemiology study. *J Natl Cancer Inst* 2010;102:475-483. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20305132>.

328. Lilyquist J, LaDuca H, Polley E, et al. Frequency of mutations in a large series of clinically ascertained ovarian cancer cases tested on multi-gene panels compared to reference controls. *Gynecol Oncol* 2017;147:375-380. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28888541>.

329. Kurian AW, Ward KC, Howlander N, et al. Genetic testing and results in a population-based cohort of breast cancer patients and ovarian cancer patients. *J Clin Oncol* 2019;37:1305-1315. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30964716>.

330. Grant RC, Selander I, Connor AA, et al. Prevalence of germline mutations in cancer predisposition genes in patients with pancreatic cancer. *Gastroenterology* 2015;148:556-564. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25479140>.

331. Thompson ER, Rowley SM, Li N, et al. Panel testing for familial breast cancer: calibrating the tension between research and clinical care. *J Clin Oncol* 2016;34:1455-1459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26786923>.

332. Weber-Lassalle N, Borde J, Weber-Lassalle K, et al. Germline loss-of-function variants in the BARD1 gene are associated with early-onset familial breast cancer but not ovarian cancer. *Breast Cancer Res* 2019;21:55. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31036035>.

333. Slavin TP, Maxwell KN, Lilyquist J, et al. The contribution of pathogenic variants in breast cancer susceptibility genes to familial breast cancer risk. *NPJ Breast Cancer* 2017;3:22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28649662>.

334. Carter NJ, Marshall ML, Susswein LR, et al. Germline pathogenic variants identified in women with ovarian tumors. *Gynecol Oncol* 2018;151:481-488. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30322717>.

335. Ramus SJ, Song H, Dicks E, et al. Germline mutations in the BRIP1, BARD1, PALB2, and NBN genes in women with ovarian cancer. *J Natl Cancer Inst* 2015;107. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26315354>.

336. Rafnar T, Gudbjartsson DF, Sulem P, et al. Mutations in BRIP1 confer high risk of ovarian cancer. *Nat Genet* 2011;43:1104-1107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21964575>.

337. Kaurah P, MacMillan A, Boyd N, et al. Founder and recurrent CDH1 mutations in families with hereditary diffuse gastric cancer. *JAMA* 2007;297:2360-2372. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17545690>.

338. Pharoah PD, Guilford P, Caldas C. Incidence of gastric cancer and breast cancer in CDH1 (E-cadherin) mutation carriers from hereditary diffuse gastric cancer families. *Gastroenterology* 2001;121:1348-1353. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11729114>.

339. Hansford S, Kaurah P, Li-Chang H, et al. Hereditary diffuse gastric cancer syndrome: CDH1 mutations and beyond. *JAMA Oncol* 2015;1:23-32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26182300>.
340. Xicola RM, Li S, Rodriguez N, et al. Clinical features and cancer risk in families with pathogenic CDH1 variants irrespective of clinical criteria. *J Med Genet* 2019;56:838-843. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31296550>.
341. Jacobs MF, Dust H, Koeppe E, et al. Outcomes of endoscopic surveillance in individuals with genetic predisposition to hereditary diffuse gastric cancer. *Gastroenterology* 2019;157:87-96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30935944>.
342. Frebourg T, Oliveira C, Hochain P, et al. Cleft lip/palate and CDH1/E-cadherin mutations in families with hereditary diffuse gastric cancer. *J Med Genet* 2006;43:138-142. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15831593>.
343. Friedrichsen DM, Malone KE, Doody DR, et al. Frequency of CHEK2 mutations in a population based, case-control study of breast cancer in young women. *Breast Cancer Res* 2004;6:R629-635. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15535844>.
344. Iniesta MD, Gorin MA, Chien LC, et al. Absence of CHEK2\*1100delC mutation in families with hereditary breast cancer in North America. *Cancer Genet Cytogenet* 2010;202:136-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20875877>.
345. Kuusisto KM, Bebel A, Vihinen M, et al. Screening for BRCA1, BRCA2, CHEK2, PALB2, BRIP1, RAD50, and CDH1 mutations in high-risk Finnish BRCA1/2-founder mutation-negative breast and/or ovarian cancer individuals. *Breast Cancer Res* 2011;13:R20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21356067>.
346. Cybulski C, Wokolorczyk D, Jakubowska A, et al. Risk of breast cancer in women with a CHEK2 mutation with and without a family history of breast cancer. *J Clin Oncol* 2011;29:3747-3752. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21876083>.
347. Weischer M, Bojesen SE, Ellervik C, et al. CHEK2\*1100delC genotyping for clinical assessment of breast cancer risk: meta-analyses of 26,000 patient cases and 27,000 controls. *J Clin Oncol* 2008;26:542-548. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18172190>.
348. Naslund-Koch C, Nordestgaard BG, Bojesen SE. Increased risk for other cancers in addition to breast cancer for CHEK2\*1100delC heterozygotes estimated from the Copenhagen General Population Study. *J Clin Oncol* 2016;34:1208-1216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26884562>.
349. CHEK2\*1100delC and susceptibility to breast cancer: a collaborative analysis involving 10,860 breast cancer cases and 9,065 controls from 10 studies. *Am J Hum Genet* 2004;74:1175-1182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15122511>.
350. Schmidt MK, Hogervorst F, van Hien R, et al. Age- and tumor subtype-specific breast cancer risk estimates for CHEK2\*1100delC carriers. *J Clin Oncol* 2016;34:2750-2760. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27269948>.
351. Han FF, Guo CL, Liu LH. The effect of CHEK2 variant I157T on cancer susceptibility: evidence from a meta-analysis. *DNA Cell Biol* 2013;32:329-335. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23713947>.
352. Boland CR, Goel A. Microsatellite instability in colorectal cancer. *Gastroenterology* 2010;138:2073-2087 e2073. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20420947>.
353. Rumilla K, Schowalter KV, Lindor NM, et al. Frequency of deletions of EPCAM (TACSTD1) in MSH2-associated Lynch syndrome cases. *J Mol Diagn* 2011;13:93-99. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21227399>.
354. Kempers MJ, Kuiper RP, Ockeloen CW, et al. Risk of colorectal and endometrial cancers in EPCAM deletion-positive Lynch syndrome: a cohort study. *Lancet Oncol* 2011;12:49-55. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21145788>.



355. Bonadona V, Bonaiti B, Olschwang S, et al. Cancer risks associated with germline mutations in MLH1, MSH2, and MSH6 genes in Lynch syndrome. JAMA 2011;305:2304-2310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21642682>.

356. Kohlmann W, Gruber S. Lynch Syndrome. GeneReviews at GeneTests: Medical Genetics Information Resource 2014. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK1211/>.

357. Lindor NM, Petersen GM, Hadley DW, et al. Recommendations for the care of individuals with an inherited predisposition to Lynch syndrome: a systematic review. JAMA 2006;296:1507-1517. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17003399>.

358. Watson P, Vasen HF, Mecklin JP, et al. The risk of extra-colonic, extra-endometrial cancer in the Lynch syndrome. Int J Cancer 2008;123:444-449. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18398828>.

359. Møller P, Seppälä TT, Bernstein I, et al. Cancer risk and survival in path\_MMR carriers by gene and gender up to 75 years of age: a report from the Prospective Lynch Syndrome Database. Gut 2018;67:1306-1316. Available at: <https://pubmed.ncbi.nlm.nih.gov/28754778/>.

360. Ten Broeke SW, van der Klift HM, Tops CMJ, et al. Cancer risks for PMS2-associated Lynch syndrome. J Clin Oncol 2018;36:2961-2968. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30161022>.

361. Dominguez-Valentin M, Sampson JR, Seppala TT, et al. Cancer risks by gene, age, and gender in 6350 carriers of pathogenic mismatch repair variants: findings from the Prospective Lynch Syndrome Database. Genet Med 2020;22:15-25. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31337882>.

362. Espenschied CR, LaDuca H, Li S, et al. Multigene panel testing provides a new perspective on Lynch syndrome. J Clin Oncol 2017;35:2568-2575. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28514183>.

363. Chen LM, Yang KY, Little SE, et al. Gynecologic cancer prevention in Lynch syndrome/hereditary nonpolyposis colorectal cancer families. Obstet Gynecol 2007;110:18-25. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17601891>.

364. Schmeler KM, Lynch HT, Chen LM, et al. Prophylactic surgery to reduce the risk of gynecologic cancers in the Lynch syndrome. N Engl J Med 2006;354:261-269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16421367>.

365. Stuckless S, Green J, Dawson L, et al. Impact of gynecological screening in Lynch syndrome carriers with an MSH2 mutation. Clin Genet 2013;83:359-364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22775459>.

366. Syngal S, Brand RE, Church JM, et al. ACG clinical guideline: Genetic testing and management of hereditary gastrointestinal cancer syndromes. Am J Gastroenterol 2015;110:223-262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25645574>.

367. Stoffel EM, Mangu PB, Gruber SB, et al. Hereditary colorectal cancer syndromes: American Society of Clinical Oncology Clinical Practice Guideline endorsement of the familial risk-colorectal cancer: European Society for Medical Oncology Clinical Practice Guidelines. J Clin Oncol 2015;33:209-217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25452455>.

368. Auranen A, Joutsiniemi T. A systematic review of gynecological cancer surveillance in women belonging to hereditary nonpolyposis colorectal cancer (Lynch syndrome) families. Acta Obstet Gynecol Scand 2011;90:437-444. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21306348>.

369. Jarvinen HJ, Renkonen-Sinisalo L, Aktan-Collan K, et al. Ten years after mutation testing for Lynch syndrome: cancer incidence and outcome in mutation-positive and mutation-negative family members. J Clin Oncol 2009;27:4793-4797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19720893>.





370. Renkonen-Sinisalo L, Butzow R, Leminen A, et al. Surveillance for endometrial cancer in hereditary nonpolyposis colorectal cancer syndrome. *Int J Cancer* 2007;120:821-824. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17096354>.

371. Rijcken FE, Mourits MJ, Kleibeuker JH, et al. Gynecologic screening in hereditary nonpolyposis colorectal cancer. *Gynecol Oncol* 2003;91:74-80. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14529665>.

372. Dove-Edwin I, Boks D, Goff S, et al. The outcome of endometrial carcinoma surveillance by ultrasound scan in women at risk of hereditary nonpolyposis colorectal carcinoma and familial colorectal carcinoma. *Cancer* 2002;94:1708-1712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11920532>.

373. Harkness EF, Barrow E, Newton K, et al. Lynch syndrome caused by MLH1 mutations is associated with an increased risk of breast cancer: a cohort study. *J Med Genet* 2015;52:553-556. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26101330>.

374. Goldberg M, Bell K, Aronson M, et al. Association between the Lynch syndrome gene MSH2 and breast cancer susceptibility in a Canadian familial cancer registry. *J Med Genet* 2017;54:742-746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28779004>.

375. Wimmer K, Kratz CP, Vasen HF, et al. Diagnostic criteria for constitutional mismatch repair deficiency syndrome: suggestions of the European consortium 'care for CMMRD' (C4CMMRD). *J Med Genet* 2014;51:355-365. Available at: <http://jmg.bmj.com/content/51/6/355.full.pdf>.

376. Zhang B, Beeghly-Fadiel A, Long J, Zheng W. Genetic variants associated with breast-cancer risk: comprehensive research synopsis, meta-analysis, and epidemiological evidence. *Lancet Oncol* 2011;12:477-488. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21514219>.

377. Zhang G, Zeng Y, Liu Z, Wei W. Significant association between Nijmegen breakage syndrome 1 657del5 polymorphism and breast cancer

risk. *Tumour Biol* 2013;34:2753-2757. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23765759>.

378. Suszynska M, Klonowska K, Jasinska AJ, Kozlowski P. Large-scale meta-analysis of mutations identified in panels of breast/ovarian cancer-related genes - providing evidence of cancer predisposition genes. *Gynecol Oncol* 2019;153:452-462. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30733081>.

379. Uusitalo E, Rantanen M, Kallionpaa RA, et al. Distinctive cancer associations in patients with neurofibromatosis type 1. *J Clin Oncol* 2016;34:1978-1986. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26926675>.

380. Rosenfeld A, Listernick R, Charrow J, Goldman S. Neurofibromatosis type 1 and high-grade tumors of the central nervous system. *Childs Nerv Syst* 2010;26:663-667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19937438>.

381. Nishida T, Tsujimoto M, Takahashi T, et al. Gastrointestinal stromal tumors in Japanese patients with neurofibromatosis type I. *J Gastroenterol* 2016;51:571-578. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26511941>.

382. Walker L, Thompson D, Easton D, et al. A prospective study of neurofibromatosis type 1 cancer incidence in the UK. *Br J Cancer* 2006;95:233-238. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16786042>.

383. Stewart DR, Korf BR, Nathanson KL, et al. Care of adults with neurofibromatosis type 1: a clinical practice resource of the American College of Medical Genetics and Genomics (ACMG). *Genet Med* 2018;20:671-682. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30006586>.

384. Sharif S, Moran A, Huson SM, et al. Women with neurofibromatosis 1 are at a moderately increased risk of developing breast cancer and should be considered for early screening. *J Med Genet* 2007;44:481-484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17369502>.





385. Evans DG. Are we ready for targeted early breast cancer detection strategies in women with NF1 aged 30-49 years? *Am J Med Genet A* 2012;158a:3054-3055. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22987630>.

386. Seminog OO, Goldacre MJ. Age-specific risk of breast cancer in women with neurofibromatosis type 1. *Br J Cancer* 2015;112:1546-1548. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25742481>.

387. Ferner RE, Huson SM, Thomas N, et al. Guidelines for the diagnosis and management of individuals with neurofibromatosis 1. *J Med Genet* 2007;44:81-88. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17105749>.

388. Casadei S, Norquist BM, Walsh T, et al. Contribution of inherited mutations in the BRCA2-interacting protein PALB2 to familial breast cancer. *Cancer Res* 2011;71:2222-2229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21285249>.

389. Cybulski C, Kluzniak W, Huzarski T, et al. Clinical outcomes in women with breast cancer and a PALB2 mutation: a prospective cohort analysis. *Lancet Oncol* 2015;16:638-644. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25959805>.

390. Song H, Dicks EM, Tyrer J, et al. Population-based targeted sequencing of 54 candidate genes identifies PALB2 as a susceptibility gene for high-grade serous ovarian cancer. *J Med Genet* 2021;58:305-313. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32546565>.

391. Tischkowitz M, Xia B. PALB2/FANCN: recombining cancer and Fanconi anemia. *Cancer Res* 2010;70:7353-7359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20858716>.

392. Loveday C, Turnbull C, Ruark E, et al. Germline RAD51C mutations confer susceptibility to ovarian cancer. *Nat Genet* 2012;44:475-476; author reply 476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22538716>.

393. Loveday C, Turnbull C, Ramsay E, et al. Germline mutations in RAD51D confer susceptibility to ovarian cancer. *Nat Genet* 2011;43:879-882. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21822267>.

394. Song H, Dicks E, Ramus SJ, et al. Contribution of germline mutations in the RAD51B, RAD51C, and RAD51D genes to ovarian cancer in the population. *J Clin Oncol* 2015;33:2901-2907. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26261251>.

395. Yang X, Song H, Leslie G, et al. Ovarian and breast cancer risks associated with pathogenic variants in RAD51C and RAD51D. *J Natl Cancer Inst* 2020;112:1242-1250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32107557>.

396. Li N, McInerney S, Zethoven M, et al. Combined tumor sequencing and case-control analyses of RAD51C in breast cancer. *J Natl Cancer Inst* 2019;111:1332-1338. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30949688>.

397. Pilie PG, Johnson AM, Hanson KL, et al. Germline genetic variants in men with prostate cancer and one or more additional cancers. *Cancer* 2017;123:3925-3932. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28657667>.

398. Lang SH, Swift SL, White H, et al. A systematic review of the prevalence of DNA damage response gene mutations in prostate cancer. *Int J Oncol* 2019;55:597-616. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31322208>.

399. Ewing CM, Ray AM, Lange EM, et al. Germline mutations in HOXB13 and prostate-cancer risk. *N Engl J Med* 2012;366:141-149. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22236224>.

400. Antonarakis ES, Lu C, Luber B, et al. Germline DNA-repair gene mutations and outcomes in men with metastatic castration-resistant prostate cancer receiving first-line abiraterone and enzalutamide. *Eur Urol* 2018;74:218-225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29439820>.



401. Schweizer MT, Cheng HH, Tretiakova MS, et al. Mismatch repair deficiency may be common in ductal adenocarcinoma of the prostate. *Oncotarget* 2016;7:82504-82510. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27756888>.

402. Risbridger GP, Taylor RA, Clouston D, et al. Patient-derived xenografts reveal that intraductal carcinoma of the prostate is a prominent pathology in BRCA2 mutation carriers with prostate cancer and correlates with poor prognosis. *Eur Urol* 2015;67:496-503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25154392>.

403. Robson M, Im SA, Senkus E, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med* 2017;377:523-533. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28578601>.

404. Tutt ANJ, Garber JE, Kaufman B, et al. Adjuvant olaparib for patients with BRCA1- or BRCA2-mutated breast cancer. *N Engl J Med* 2021;384:2394-2405. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34081848>.

405. Litton JK, Rugo HS, Ettl J, et al. Talazoparib in patients with advanced breast cancer and a germline BRCA mutation. *N Engl J Med* 2018;379:753-763. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30110579>.

406. Moore KN, Secord AA, Geller MA, et al. Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol* 2019;20:636-648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30948273>.

407. Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. *J Clin Oncol* 2015;33:244-250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25366685>.

408. Penson RT, Valencia RV, Cibula D, et al. Olaparib versus nonplatinum chemotherapy in patients with platinum-sensitive relapsed ovarian cancer and a germline BRCA1/2 mutation (SOLO3): a randomized

phase III trial. *J Clin Oncol* 2020;38:1164-1174. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32073956>.

409. Swisher EM, Lin KK, Oza AM, et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. *Lancet Oncol* 2017;18:75-87. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27908594>.

410. de Bono J, Mateo J, Fizazi K, et al. Olaparib for metastatic castration-resistant prostate cancer. *N Engl J Med* 2020;382:2091-2102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32343890>.

411. Abida W, Patnaik A, Campbell D, et al. Rucaparib in men with metastatic castration-resistant prostate cancer harboring a BRCA1 or BRCA2 gene alteration. *J Clin Oncol* 2020;38:3763-3772. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32795228>.

412. Golan T, Hammel P, Reni M, et al. Maintenance olaparib for germline BRCA-mutated metastatic pancreatic cancer. *N Engl J Med* 2019;381:317-327. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31157963>.

413. Gabai-Kapara E, Lahad A, Kaufman B, et al. Population-based screening for breast and ovarian cancer risk due to BRCA1 and BRCA2. *Proc Natl Acad Sci U S A* 2014;111:14205-14210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25192939>.

414. Manchanda R, Loggenberg K, Sanderson S, et al. Population testing for cancer predisposing BRCA1/BRCA2 mutations in the Ashkenazi-Jewish community: a randomized controlled trial. *J Natl Cancer Inst* 2015;107:379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25435541>.

415. Best AF, Tucker MA, Frone MN, et al. A pragmatic testing-eligibility framework for population mutation screening: the example of BRCA1/2. *Cancer Epidemiol Biomarkers Prev* 2019;28:293-302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30692095>.

416. Metcalfe KA, Poll A, Royer R, et al. A comparison of the detection of BRCA mutation carriers through the provision of Jewish population-based



genetic testing compared with clinic-based genetic testing. Br J Cancer 2013;109:777-779. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23778531>.

417. Lieberman S, Tomer A, Ben-Chetrit A, et al. Population screening for BRCA1/BRCA2 founder mutations in Ashkenazi Jews: proactive recruitment compared with self-referral. Genet Med 2017;19:754-762.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27929526>.

418. Manahan ER, Kuerer HM, Sebastian M, et al. Consensus guidelines on genetic testing for hereditary breast cancer from the American Society of Breast Surgeons. Ann Surg Oncol 2019;26:3025-3031. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31342359>.

419. Beitsch PD, Whitworth PW, Hughes K, et al. Underdiagnosis of hereditary breast cancer: are genetic testing guidelines a tool or an obstacle? J Clin Oncol 2019;37:453-460. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30526229>.

420. Yang S, Axilbund JE, O'Leary E, et al. Underdiagnosis of hereditary breast and ovarian cancer in Medicare patients: genetic testing criteria miss the mark. Ann Surg Oncol 2018;25:2925-2931. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29998407>.

421. Sun L, Brentnall A, Patel S, et al. A cost-effectiveness analysis of multigene testing for all patients with breast cancer. JAMA Oncol 2019;5:1718-1730. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31580391>.

422. Kurian AW, Bernhisel R, Larson K, et al. Prevalence of pathogenic variants in cancer susceptibility genes among women with postmenopausal breast cancer. JAMA 2020;323:995-997. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32154851>.

423. Chavarri-Guerra Y, Hendricks CB, Brown S, et al. The burden of breast cancer predisposition variants across the age spectrum among 10 000 patients. J Am Geriatr Soc 2019;67:884-888. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31012959>.

424. Yadav S, Hu C, Hart SN, et al. Evaluation of germline genetic testing criteria in a hospital-based series of women with breast cancer. J Clin Oncol 2020;38:1409-1418. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32125938>.

425. Desai NV, Yadav S, Batalini F, et al. Germline genetic testing in breast cancer: rationale for the testing of all women diagnosed by the age of 60 years and for risk-based testing of those older than 60 years. Cancer 2021;127:828-833. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33146899>.

426. Boddicker NJ, Hu C, Weitzel JN, et al. Risk of late-onset breast cancer in genetically predisposed women. J Clin Oncol 2021;39:3430-3440. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34292776>.

427. Weitzel JN, Kidd J, Bernhisel R, et al. Multigene assessment of genetic risk for women for two or more breast cancers. Breast Cancer Res Treat 2021;188:759-768. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33826040>.

428. Maxwell KN, Wenz BM, Kulkarni A, et al. Mutation rates in cancer susceptibility genes in patients with breast cancer with multiple primary cancers. JCO Precis Oncol 2020;4. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32954205>.

429. Antoniou AC, Hardy R, Walker L, et al. Predicting the likelihood of carrying a BRCA1 or BRCA2 mutation: validation of BOADICEA, BRCAPRO, IBIS, Myriad and the Manchester scoring system using data from UK genetics clinics. J Med Genet 2008;45:425-431. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18413374>.

430. Parmigiani G, Chen S, Iversen ES, Jr., et al. Validity of models for predicting BRCA1 and BRCA2 mutations. Ann Intern Med 2007;147:441-450. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17909205>.

431. Lindor NM, Johnson KJ, Harvey H, et al. Predicting BRCA1 and BRCA2 gene mutation carriers: comparison of PENN II model to previous study. Fam Cancer 2010;9:495-502. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20512419>.





432. Murphy CD, Lee JM, Drohan B, et al. The American Cancer Society guidelines for breast screening with magnetic resonance imaging: an argument for genetic testing. *Cancer* 2008;113:3116-3120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18932252>.
433. Panchal SM, Ennis M, Canon S, Bordeleau LJ. Selecting a BRCA risk assessment model for use in a familial cancer clinic. *BMC Med Genet* 2008;9:116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19102775>.
434. Terry MB, Liao Y, Whittemore AS, et al. 10-year performance of four models of breast cancer risk: a validation study. *Lancet Oncol* 2019;20:504-517. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30799262>.
435. Archer S, Babb de Villiers C, Scheibl F, et al. Evaluating clinician acceptability of the prototype CanRisk tool for predicting risk of breast and ovarian cancer: a multi-methods study. *PLoS One* 2020;15:e0229999. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32142536>.
436. Sidransky D, Tokino T, Helzlsouer K, et al. Inherited p53 gene mutations in breast cancer. *Cancer Res* 1992;52:2984-2986. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1581912>.
437. Gonzalez KD, Noltner KA, Buzin CH, et al. Beyond Li Fraumeni Syndrome: clinical characteristics of families with p53 germline mutations. *J Clin Oncol* 2009;27:1250-1256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19204208>.
438. Lalloo F, Varley J, Ellis D, et al. Prediction of pathogenic mutations in patients with early-onset breast cancer by family history. *Lancet* 2003;361:1101-1102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12672316>.
439. Lane DP. Cancer. p53, guardian of the genome. *Nature* 1992;358:15-16. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1614522>.
440. Levine AJ. p53, the cellular gatekeeper for growth and division. *Cell* 1997;88:323-331. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9039259>.
441. Garber JE, Goldstein AM, Kantor AF, et al. Follow-up study of twenty-four families with Li-Fraumeni syndrome. *Cancer Res* 1991;51:6094-6097. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1933872>.
442. Nichols KE, Malkin D, Garber JE, et al. Germ-line p53 mutations predispose to a wide spectrum of early-onset cancers. *Cancer Epidemiol Biomarkers Prev* 2001;10:83-87. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11219776>.
443. Siddiqui R, Onel K, Facio F, et al. The TP53 mutational spectrum and frequency of CHEK2\*1100delC in Li-Fraumeni-like kindreds. *Fam Cancer* 2005;4:177-181. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15951970>.
444. Mai PL, Best AF, Peters JA, et al. Risks of first and subsequent cancers among TP53 mutation carriers in the National Cancer Institute Li-Fraumeni syndrome cohort. *Cancer* 2016;122:3673-3681. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27496084>.
445. Birch JM, Hartley AL, Tricker KJ, et al. Prevalence and diversity of constitutional mutations in the p53 gene among 21 Li-Fraumeni families. *Cancer Res* 1994;54:1298-1304. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8118819>.
446. Krutilkova V, Trkova M, Fleitz J, et al. Identification of five new families strengthens the link between childhood choroid plexus carcinoma and germline TP53 mutations. *Eur J Cancer* 2005;41:1597-1603. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15925506>.
447. Li FP, Fraumeni JF, Jr. Soft-tissue sarcomas, breast cancer, and other neoplasms. A familial syndrome? *Ann Intern Med* 1969;71:747-752. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/5360287>.
448. Li FP, Fraumeni JF, Jr., Mulvihill JJ, et al. A cancer family syndrome in twenty-four kindreds. *Cancer Res* 1988;48:5358-5362. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3409256>.
449. Malkin D, Li FP, Strong LC, et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms.



Science 1990;250:1233-1238. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/1978757>.

450. Varley JM, Evans DG, Birch JM. Li-Fraumeni syndrome--a molecular and clinical review. Br J Cancer 1997;76:1-14. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/9218725>.

451. Masciari S, Dewanwala A, Stoffel EM, et al. Gastric cancer in individuals with Li-Fraumeni syndrome. Genet Med 2011;13:651-657. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21552135/>.

452. Holmfeldt L, Wei L, Diaz-Flores E, et al. The genomic landscape of hypodiploid acute lymphoblastic leukemia. Nat Genet 2013;45:242-252. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23334668>.

453. Kamihara J, Rana HQ, Garber JE. Germline TP53 mutations and the changing landscape of Li-Fraumeni syndrome. Hum Mutat 2014;35:654-662. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24706533>.

454. Curiel-Lewandrowski C, Speetzen LS, Cranmer L, et al. Multiple primary cutaneous melanomas in Li-Fraumeni syndrome. Arch Dermatol 2011;147:248-250. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/21339461>.

455. Giavedoni P, Ririe M, Carrera C, et al. Familial melanoma associated with Li-Fraumeni Syndrome and Atypical Mole Syndrome: total-body digital photography, dermoscopy and confocal microscopy. Acta Derm Venereol 2017;97:720-723. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/28218344>.

456. Melhem-Bertrandt A, Bojadzieva J, Ready KJ, et al. Early onset HER2-positive breast cancer is associated with germline TP53 mutations. Cancer 2012;118:908-913. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/21761402>.

457. Wilson JR, Bateman AC, Hanson H, et al. A novel HER2-positive breast cancer phenotype arising from germline TP53 mutations. J Med Genet 2010;47:771-774. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20805372>.

458. Packwood K, Martland G, Sommerlad M, et al. Breast cancer in patients with germline TP53 pathogenic variants have typical tumour characteristics: the cohort study of TP53 carrier early onset breast cancer (COPE study). J Pathol Clin Res 2019;5:189-198. Available at:  
<https://www.ncbi.nlm.nih.gov/pubmed/31041842>.

459. Hisada M, Garber JE, Fung CY, et al. Multiple primary cancers in families with Li-Fraumeni syndrome. J Natl Cancer Inst 1998;90:606-611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9554443>.

460. Lustbader ED, Williams WR, Bondy ML, et al. Segregation analysis of cancer in families of childhood soft-tissue-sarcoma patients. Am J Hum Genet 1992;51:344-356. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/1642235>.

461. Birch JM, Blair V, Kelsey AM, et al. Cancer phenotype correlates with constitutional TP53 genotype in families with the Li-Fraumeni syndrome. Oncogene 1998;17:1061-1068. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/9764816>.

462. Chompret A. The Li-Fraumeni syndrome. Biochimie 2002;84:75-82. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11900879>.

463. Chompret A, Abel A, Stoppa-Lyonnet D, et al. Sensitivity and predictive value of criteria for p53 germline mutation screening. J Med Genet 2001;38:43-47. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/11332399>.

464. Eeles RA. Germline mutations in the TP53 gene. Cancer Surv 1995;25:101-124. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/8718514>.

465. Bougeard G, Sesboue R, Baert-Desurmont S, et al. Molecular basis of the Li-Fraumeni syndrome: an update from the French LFS families. J Med Genet 2008;45:535-538. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18511570>.

466. Bougeard G, Renaux-Petel M, Flaman JM, et al. Revisiting Li-Fraumeni syndrome from TP53 mutation carriers. J Clin Oncol



2015;33:2345-2352. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26014290>.

467. Tinat J, Bougeard G, Baert-Desurmont S, et al. 2009 version of the Chompret criteria for Li Fraumeni syndrome. *J Clin Oncol* 2009;27:e108-109; author reply e110. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19652052>.

468. Ginsburg OM, Akbari MR, Aziz Z, et al. The prevalence of germ-line TP53 mutations in women diagnosed with breast cancer before age 30. *Fam Cancer* 2009;8:563-567. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19714488>.

469. Laloo F, Varley J, Moran A, et al. BRCA1, BRCA2 and TP53 mutations in very early-onset breast cancer with associated risks to relatives. *Eur J Cancer* 2006;42:1143-1150. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16644204>.

470. Lee DS, Yoon SY, Looi LM, et al. Comparable frequency of BRCA1, BRCA2 and TP53 germline mutations in a multi-ethnic Asian cohort suggests TP53 screening should be offered together with BRCA1/2 screening to early-onset breast cancer patients. *Breast Cancer Res* 2012;14:R66. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22507745>.

471. Mouchawar J, Korch C, Byers T, et al. Population-based estimate of the contribution of TP53 mutations to subgroups of early-onset breast cancer: Australian Breast Cancer Family Study. *Cancer Res* 2010;70:4795-4800. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20501846>.

472. McCuaig JM, Armel SR, Novokmet A, et al. Routine TP53 testing for breast cancer under age 30: ready for prime time? *Fam Cancer* 2012;11:607-613. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22851211>.

473. Leroy B, Anderson M, Soussi T. TP53 mutations in human cancer: database reassessment and prospects for the next decade. *Hum Mutat*

2014;35:672-688. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24665023>.

474. Kandoth C, McLellan MD, Vandin F, et al. Mutational landscape and significance across 12 major cancer types. *Nature* 2013;502:333-339. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24132290>.

475. Mai PL, Khincha PP, Loud JT, et al. Prevalence of cancer at baseline screening in the National Cancer Institute Li-Fraumeni syndrome cohort. *JAMA Oncol* 2017;3:1640-1645. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28772286>.

476. Kratz CP, Achatz MI, Brugieres L, et al. Cancer screening recommendations for individuals with Li-Fraumeni Syndrome. *Clin Cancer Res* 2017;23:e38-e45. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28572266>.

477. Greer MC, Voss SD, States LJ. Pediatric cancer predisposition imaging: focus on whole-body MRI. *Clin Cancer Res* 2017;23:e6-e13. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28572262>.

478. Ballinger ML, Best A, Mai PL, et al. Baseline surveillance in Li-Fraumeni syndrome using whole-body magnetic resonance imaging: a meta-analysis. *JAMA Oncol* 2017;3:1634-1639. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28772291>.

479. Villani A, Tabori U, Schiffman J, et al. Biochemical and imaging surveillance in germline TP53 mutation carriers with Li-Fraumeni syndrome: a prospective observational study. *Lancet Oncol* 2011;12:559-567. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21601526>.

480. Villani A, Shore A, Wasserman JD, et al. Biochemical and imaging surveillance in germline TP53 mutation carriers with Li-Fraumeni syndrome: 11 year follow-up of a prospective observational study. *Lancet Oncol* 2016;17:1295-1305. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27501770>.

481. Asdahl PH, Ojha RP, Hasle H. Cancer screening in Li-Fraumeni Syndrome. JAMA Oncol 2017;3:1645-1646. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28772307>.
482. Avigad S, Peleg D, Barel D, et al. Prenatal diagnosis in Li-Fraumeni syndrome. J Pediatr Hematol Oncol 2004;26:541-545. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15342977>.
483. Prochazkova K, Foretova L, Sedlacek Z. A rare tumor and an ethical dilemma in a family with a germline TP53 mutation. Cancer Genet Cytogenet 2008;180:65-69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18068537>.
484. Orloff MS, Eng C. Genetic and phenotypic heterogeneity in the PTEN hamartoma tumour syndrome. Oncogene 2008;27:5387-5397. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18794875>.
485. Eng C. PTEN hamartoma tumor syndrome (PTHS). GeneReviews; 2009. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK1488/>.
486. Pilarski R, Stephens JA, Noss R, et al. Predicting PTEN mutations: an evaluation of Cowden syndrome and Bannayan-Riley-Ruvalcaba syndrome clinical features. J Med Genet 2011;48:505-512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21659347>.
487. Varga EA, Pastore M, Prior T, et al. The prevalence of PTEN mutations in a clinical pediatric cohort with autism spectrum disorders, developmental delay, and macrocephaly. Genet Med 2009;11:111-117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19265751>.
488. Nelen MR, Kremer H, Konings IB, et al. Novel PTEN mutations in patients with Cowden disease: absence of clear genotype-phenotype correlations. Eur J Hum Genet 1999;7:267-273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10234502>.
489. Pilarski R, Eng C. Will the real Cowden syndrome please stand up (again)? Expanding mutational and clinical spectra of the PTEN hamartoma tumour syndrome. J Med Genet 2004;41:323-326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15121767>.
490. Bennett KL, Mester J, Eng C. Germline epigenetic regulation of KILLIN in Cowden and Cowden-like syndrome. JAMA 2010;304:2724-2731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21177507>.
491. Hobert JA, Eng C. PTEN hamartoma tumor syndrome: an overview. Genet Med 2009;11:687-694. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19668082>.
492. Starink TM, van der Veen JP, Arwert F, et al. The Cowden syndrome: a clinical and genetic study in 21 patients. Clin Genet 1986;29:222-233. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3698331>.
493. Pilarski R, Burt R, Kohlman W, et al. Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria. J Natl Cancer Inst 2013;105:1607-1616. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24136893>.
494. Bubien V, Bonnet F, Brouste V, et al. High cumulative risks of cancer in patients with PTEN hamartoma tumour syndrome. J Med Genet 2013;50:255-263. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23335809>.
495. Riegert-Johnson DL, Gleeson FC, Roberts M, et al. Cancer and Lhermitte-Duclos disease are common in Cowden syndrome patients. Hered Cancer Clin Pract 2010;8:6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20565722>.
496. Tan MH, Mester JL, Ngeow J, et al. Lifetime cancer risks in individuals with germline PTEN mutations. Clin Cancer Res 2012;18:400-407. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22252256>.
497. Tan MH, Mester J, Peterson C, et al. A clinical scoring system for selection of patients for PTEN mutation testing is proposed on the basis of a prospective study of 3042 probands. Am J Hum Genet 2011;88:42-56. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21194675>.





498. Zbuk KM, Eng C. Hamartomatous polyposis syndromes. *Nat Clin Pract Gastroenterol Hepatol* 2007;4:492-502. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17768394>.

499. Hansen-Kiss E, Beinkampen S, Adler B, et al. A retrospective chart review of the features of PTEN hamartoma tumour syndrome in children. *J Med Genet* 2017;54:471-478. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28526761>.

500. Smith JR, Marqusee E, Webb S, et al. Thyroid nodules and cancer in children with PTEN hamartoma tumor syndrome. *J Clin Endocrinol Metab* 2011;96:34-37. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20962022>.

501. Roche AF, Mukherjee D, Guo SM, Moore WM. Head circumference reference data: birth to 18 years. *Pediatrics* 1987;79:706-712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3575026>.

502. Zhou XP, Waite KA, Pilarski R, et al. Germline PTEN promoter mutations and deletions in Cowden/Bannayan-Riley-Ruvalcaba syndrome result in aberrant PTEN protein and dysregulation of the phosphoinositide-3-kinase/Akt pathway. *Am J Hum Genet* 2003;73:404-411. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12844284>.

503. Zhou XP, Marsh DJ, Morrison CD, et al. Germline inactivation of PTEN and dysregulation of the phosphoinositide-3-kinase/Akt pathway cause human Lhermitte-Duclos disease in adults. *Am J Hum Genet* 2003;73:1191-1198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14566704>.

504. Andres RH, Guzman R, Weis J, et al. Lhermitte-Duclos disease with atypical vascularization--case report and review of the literature. *Clin Neuropathol* 2009;28:83-90. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19353838>.

505. Butler MG, Dasouki MJ, Zhou XP, et al. Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *J Med Genet*

2005;42:318-321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15805158>.

506. Herman GE, Butter E, Enrile B, et al. Increasing knowledge of PTEN germline mutations: Two additional patients with autism and macrocephaly. *Am J Med Genet A* 2007;143:589-593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17286265>.

507. Herman GE, Henninger N, Ratliff-Schaub K, et al. Genetic testing in autism: how much is enough? *Genet Med* 2007;9:268-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17505203>.

508. Orrico A, Galli L, Buoni S, et al. Novel PTEN mutations in neurodevelopmental disorders and macrocephaly. *Clin Genet* 2009;75:195-198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18759867>.

509. Black D, Bogomolny F, Robson ME, et al. Evaluation of germline PTEN mutations in endometrial cancer patients. *Gynecol Oncol* 2005;96:21-24. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15589575>.

510. Nelen MR, Padberg GW, Peeters EA, et al. Localization of the gene for Cowden disease to chromosome 10q22-23. *Nat Genet* 1996;13:114-116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8673088>.

511. Schaffer JV, Kamino H, Witkiewicz A, et al. Mucocutaneous neuromas: an underrecognized manifestation of PTEN hamartoma-tumor syndrome. *Arch Dermatol* 2006;142:625-632. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16702501>.

512. Brownstein MH, Mehregan AH, Bikowski JB, et al. The dermatopathology of Cowden's syndrome. *Br J Dermatol* 1979;100:667-673. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/465314>.

513. Brownstein MH, Mehregan AH, Bilowski JB. Trichilemmomas in Cowden's disease. *JAMA* 1977;238:26. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/577252>.



514. Heald B, Mester J, Rybicki L, et al. Frequent gastrointestinal polyps and colorectal adenocarcinomas in a prospective series of PTEN mutation carriers. *Gastroenterology* 2010;139:1927-1933. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20600018>.
515. Stanich PP, Owens VL, Sweetser S, et al. Colonic polyposis and neoplasia in Cowden syndrome. *Mayo Clin Proc* 2011;86:489-492. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21628613>.
516. Stanich PP, Pilarski R, Rock J, et al. Colonic manifestations of PTEN hamartoma tumor syndrome: case series and systematic review. *World J Gastroenterol* 2014;20:1833-1838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24587660>.
517. Al-Thihli K, Palma L, Marcus V, et al. A case of Cowden's syndrome presenting with gastric carcinomas and gastrointestinal polyposis. *Nat Clin Pract Gastroenterol Hepatol* 2009;6:184-189. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19190598>.
518. Nieuwenhuis MH, Kets CM, Murphy-Ryan M, et al. Cancer risk and genotype-phenotype correlations in PTEN hamartoma tumor syndrome. *Fam Cancer* 2014;13:57-63. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23934601>.
519. Gorlin RJ, Cohen MM, Jr., Condon LM, Burke BA. Bannayan-Riley-Ruvalcaba syndrome. *Am J Med Genet* 1992;44:307-314. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1336932>.
520. Marsh DJ, Coulon V, Lunetta KL, et al. Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation. *Hum Mol Genet* 1998;7:507-515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9467011>.
521. Eng C. Will the real Cowden syndrome please stand up: revised diagnostic criteria. *J Med Genet* 2000;37:828-830. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11073535>.
522. Li S, Shen Y, Wang M, et al. Loss of PTEN expression in breast cancer: association with clinicopathological characteristics and prognosis. *Oncotarget* 2017;8:32043-32054. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28410191>.
523. Feldman R, Gatalica Z, Knezetic J, et al. Molecular profiling of head and neck squamous cell carcinoma. *Head Neck* 2016;38 Suppl 1:E1625-1638. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26614708>.
524. Wise HM, Hermida MA, Leslie NR. Prostate cancer, PI3K, PTEN and prognosis. *Clin Sci (Lond)* 2017;131:197-210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28057891>.
525. SEER Stat Fact Sheets: Thyroid Cancer. 2015. Available at: <http://seer.cancer.gov/statfacts/html/thyro.html>. Accessed May 28, 2015.
526. Schultz KAP, Rednam SP, Kamihara J, et al. PTEN, DICER1, FH, and their associated tumor susceptibility syndromes: clinical features, genetics, and surveillance recommendations in childhood. *Clin Cancer Res* 2017;23:e76-e82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28620008>.
527. Humphris JL, Johns AL, Simpson SH, et al. Clinical and pathologic features of familial pancreatic cancer. *Cancer* 2014;120:3669-3675. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25313458>.
528. Petersen GM. Familial pancreatic cancer. *Semin Oncol* 2016;43:548-553. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27899186>.
529. Yurgelun MB, Chittenden AB, Morales-Oyarvide V, et al. Germline cancer susceptibility gene variants, somatic second hits, and survival outcomes in patients with resected pancreatic cancer. *Genet Med* 2019;21:213-223. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29961768>.
530. Abe T, Blackford AL, Tamura K, et al. Deleterious germline mutations are a risk factor for neoplastic progression among high-risk individuals undergoing pancreatic surveillance. *J Clin Oncol* 2019;37:1070-1080. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30883245>.



531. Roberts NJ, Jiao Y, Yu J, et al. ATM mutations in patients with hereditary pancreatic cancer. *Cancer Discov* 2012;2:41-46. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22585167>.

532. Slater EP, Langer P, Niemczyk E, et al. PALB2 mutations in European familial pancreatic cancer families. *Clin Genet* 2010;78:490-494. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20412113>.

533. Jones S, Hruban RH, Kamiyama M, et al. Exomic sequencing identifies PALB2 as a pancreatic cancer susceptibility gene. *Science* 2009;324:217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19264984>.

534. Hu C, Hart SN, Bamlet WR, et al. Prevalence of pathogenic mutations in cancer predisposition genes among pancreatic cancer patients. *Cancer Epidemiol Biomarkers Prev* 2016;25:207-211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26483394>.

535. Lowery MA, Wong W, Jordan EJ, et al. Prospective evaluation of germline alterations in patients with exocrine pancreatic neoplasms. *J Natl Cancer Inst* 2018;110:1067-1074. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29506128>.

536. Cremin C, Lee MK, Hong Q, et al. Burden of hereditary cancer susceptibility in unselected patients with pancreatic ductal adenocarcinoma referred for germline screening. *Cancer Med* 2020;9:4004-4013. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32255556>.

537. Rainone M, Singh I, Salo-Mullen EE, et al. An emerging paradigm for germline testing in pancreatic ductal adenocarcinoma and immediate implications for clinical practice: a review. *JAMA Oncol* 2020;6:764-771. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32053139>.

538. Klein AP, Brune KA, Petersen GM, et al. Prospective risk of pancreatic cancer in familial pancreatic cancer kindreds. *Cancer Res* 2004;64:2634-2638. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15059921>.

539. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29313949>.

540. Simard EP, Ward EM, Siegel R, Jemal A. Cancers with increasing incidence trends in the United States: 1999 through 2008. *CA Cancer J Clin* 2012;62:118-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22281605>.

541. Vasen H, Ibrahim I, Ponce CG, et al. Benefit of surveillance for pancreatic cancer in high-risk individuals: outcome of long-term prospective follow-up studies from three European expert centers. *J Clin Oncol* 2016;34:2010-2019. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27114589>.

542. Paiella S, Capurso G, Cavestro GM, et al. Results of first-round of surveillance in individuals at high-risk of pancreatic cancer from the AISP (Italian Association for the Study of the Pancreas) registry. *Am J Gastroenterol* 2019;114:665-670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30538291>.

543. Canto MI, Almario JA, Schulick RD, et al. Risk of neoplastic progression in individuals at high risk for pancreatic cancer undergoing long-term surveillance. *Gastroenterology* 2018;155:740-751 e742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29803839>.

544. Goggins M, Overbeek KA, Brand R, et al. Management of patients with increased risk for familial pancreatic cancer: updated recommendations from the International Cancer of the Pancreas Screening (CAPS) Consortium. *Gut* 2020;69:7-17. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31672839>.

545. Weiss FU. Pancreatic cancer risk in hereditary pancreatitis. *Front Physiol* 2014;5:70. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24600409>.

546. Rebours V, Levy P, Ruzsiewicz P. An overview of hereditary pancreatitis. *Dig Liver Dis* 2012;44:8-15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21907651>.



547. Patel MR, Eppolito AL, Willingham FF. Hereditary pancreatitis for the endoscopist. *Therap Adv Gastroenterol* 2013;6:169-179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23503650>.

548. Hasan A, Moscoso DI, Kastrinos F. The role of genetics in pancreatitis. *Gastrointest Endosc Clin N Am* 2018;28:587-603. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30241646>.

549. Canto MI, Hruban RH, Fishman EK, et al. Frequent detection of pancreatic lesions in asymptomatic high-risk individuals. *Gastroenterology* 2012;142:796-804. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22245846>.

Discussion  
update in  
progress