

# Compound Poisson Process for VIX Shock Modeling

A Detailed Mathematical Treatment

CHONG Tin Tak, CHOI Man Hou, Vittorio Prana CHANDREAN

HKUST – IEDA4000E

December 14, 2025

## Abstract

This report provides a comprehensive mathematical treatment of the Compound Poisson Process (CPP) as applied to modeling VIX shock dynamics. We derive the key distributional properties, explain the estimation methodology, and present empirical results from fitting the model to 15 years of VIX data. The CPP framework allows us to jointly model shock timing (via Poisson arrivals) and shock magnitude (via jump size distributions), enabling risk quantification through Value-at-Risk (VaR) and Conditional VaR (CVaR) metrics.

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Motivation . . . . .	3
<b>2</b>	<b>Foundations</b>	<b>3</b>
2.1	Data transformation, autocorrelation, and stationarity . . . . .	3
2.1.1	VIX level versus SPY (context) . . . . .	3
2.1.2	Autocorrelation structure . . . . .	4
2.1.3	Stationarity tests . . . . .	4
2.2	Mathematical foundations of volatility models . . . . .	5
2.2.1	ARIMA mean model . . . . .	5
2.2.2	GARCH( $p, q$ ) conditional variance . . . . .	5
2.2.3	EGARCH( $p, q$ ) conditional variance . . . . .	5
2.3	Mathematical foundations of the Compound Poisson Process . . . . .	5
2.3.1	Definition of the Compound Poisson Process . . . . .	5
2.3.2	Interpretation for VIX Shocks . . . . .	6
2.3.3	Distributional Properties . . . . .	6
<b>3</b>	<b>Methodology</b>	<b>7</b>
3.1	Volatility modeling workflow and model selection . . . . .	7
3.1.1	Mean model selection (ARIMA) . . . . .	7
3.1.2	Volatility model selection (GARCH/EGARCH) . . . . .	7
3.2	Shock modeling workflow (CPP) . . . . .	8
3.2.1	Jump size distributions . . . . .	8
3.2.2	Distribution selection methodology . . . . .	9
<b>4</b>	<b>Volatility Model Fitting Results</b>	<b>10</b>
4.1	Mean dynamics: best ARIMA for $\Delta \log(\text{VIX})$ (by BIC) . . . . .	10
4.2	Volatility dynamics: best GARCH-family specification (by BIC) . . . . .	10
4.3	Volatility persistence: half-life . . . . .	11

4.4	Conditional volatility overlays . . . . .	11
4.5	Equity market stress and VIX volatility . . . . .	12
<b>5</b>	<b>Risk Measures</b>	<b>12</b>
5.1	Value-at-Risk (VaR) . . . . .	12
5.2	Conditional Value-at-Risk (CVaR) . . . . .	12
5.3	Monte Carlo Estimation . . . . .	12
<b>6</b>	<b>Empirical Results</b>	<b>13</b>
6.1	Fitted Parameters (Full Sample) . . . . .	13
6.2	Interpretation of Results . . . . .	13
6.3	Jump Size Distribution Visualization . . . . .	14
6.4	Simulated CPP Paths . . . . .	14
6.5	Annual Impact Distribution . . . . .	15
<b>7</b>	<b>Regime Analysis</b>	<b>15</b>
7.1	Regime Definitions . . . . .	15
7.2	Regime-Specific CPP Parameters . . . . .	15
7.3	Key Regime Findings . . . . .	16
7.4	Regime Comparison Visualization . . . . .	16
<b>8</b>	<b>Out-of-Sample Evaluation</b>	<b>16</b>
8.1	Train-Test Split Design . . . . .	16
8.2	Forecasting Methodology . . . . .	17
8.3	Out-of-Sample Results . . . . .	17
8.4	Interpretation of Results . . . . .	17
8.5	Visualization of Out-of-Sample Performance . . . . .	18
<b>9</b>	<b>Connection to Other Models</b>	<b>19</b>
9.1	CPP vs. Hawkes Process . . . . .	19
9.2	Potential Extensions . . . . .	19
<b>10</b>	<b>Practical Implications</b>	<b>19</b>
10.1	Risk Management . . . . .	19
10.2	Derivatives Pricing . . . . .	19
<b>11</b>	<b>Conclusion</b>	<b>20</b>

# 1 Introduction

Traditional point process models for financial shocks—such as the Homogeneous Poisson Process (HPP), Non-Homogeneous Poisson Process (NHPP), and Hawkes process—focus on modeling *when* shocks occur. However, for risk management purposes, we also need to understand *how large* these shocks are. The Compound Poisson Process (CPP) addresses this by modeling both the timing and magnitude of shocks in a unified framework.

## 1.1 Motivation

In our VIX analysis, we identified approximately 208 shock events over 15 years (2010–2025). While knowing the arrival rate ( $\lambda \approx 13$  shocks/year) is useful, risk managers need to answer questions like:

- What is the expected total shock impact over a year?
- What is the 95th percentile of annual shock impact (VaR)?
- How does shock risk differ across market regimes?

The CPP provides a principled framework for answering these questions. In addition, we model the *continuous* component of volatility dynamics using a mean–variance decomposition on  $d_t = \Delta \log(\text{VIX})_t$ : an ARIMA model for the conditional mean and GARCH-family models (GARCH/EGARCH) for the conditional variance. This volatility modeling complements the CPP shock framework by capturing volatility clustering and asymmetry, and it enables a concise persistence metric via the **volatility half-life** (how quickly volatility shocks decay back toward baseline).

## 2 Foundations

This section provides foundational context and mathematical definitions used throughout the report: (i) data transformation and basic time-series properties for VIX, (ii) volatility-model foundations (ARIMA and GARCH-family), and (iii) the Compound Poisson Process foundations for shock modeling.

### 2.1 Data transformation, autocorrelation, and stationarity

We focus on the VIX level and the transformed series

$$d_t = \Delta \log(\text{VIX})_t = \log(\text{VIX}_t) - \log(\text{VIX}_{t-1}),$$

which is commonly used to stabilize variance and reduce persistence in level data.

#### 2.1.1 VIX level versus SPY (context)

Figure 1 shows the co-movement between VIX and the equity market proxy SPY: large VIX spikes tend to coincide with drawdowns or stress periods in equities.




Figure 1: VIX level (left axis) and SPY price (right axis), 2010–2025.

### 2.1.2 Autocorrelation structure

The VIX level exhibits strong persistence (slowly decaying ACF), while  $\Delta \log(\text{VIX})$  is much closer to a weakly dependent / near white-noise series. Motivated by this difference, our ARIMA mean model (selected by BIC) is fit to  $d_t = \Delta \log(\text{VIX})_t$  (i.e., log returns of VIX), and the GARCH/EGARCH volatility models are then fit to the resulting ARIMA residuals.




Figure 2: Autocorrelation functions of VIX level and  $\Delta \log(\text{VIX})$ .

### 2.1.3 Stationarity tests

We apply both ADF (unit-root null) and KPSS (stationarity null) tests. The results show mixed evidence for the VIX *level*, but consistent evidence that  $\Delta \log(\text{VIX})$  is stationary.

Series	Test	Test statistic	p-value	Lags	Conclusion
VIX level	ADF	-5.4649	0.0000	26	Reject unit root
VIX level	KPSS	0.5374	0.0332	38	Reject stationarity (5%)
$\Delta \log(\text{VIX})$	ADF	-25.0132	0.0000	8	Reject unit root
$\Delta \log(\text{VIX})$	KPSS	0.0109	0.1000	35	Fail to reject stationarity

Table 1: Stationarity diagnostics for VIX level and  $\Delta \log(\text{VIX})$ .

#### Key Result:

The VIX **level** is highly persistent and yields mixed stationarity evidence (ADF rejects unit root, KPSS rejects stationarity), while  $\Delta \log(\text{VIX})$  is **stationary** under both tests. This motivates modeling on  $\Delta \log(\text{VIX})$  and treating large moves as shock events.

## 2.2 Mathematical foundations of volatility models

We decompose dynamics into a conditional mean model for  $d_t = \Delta \log(\text{VIX})_t$  and a conditional variance model for its innovations.

### 2.2.1 ARIMA mean model

Let  $\varepsilon_t$  denote innovations. An ARIMA( $p, 0, q$ ) (ARMA( $p, q$ )) model takes the form

$$d_t = c + \sum_{i=1}^p \phi_i d_{t-i} + \varepsilon_t + \sum_{j=1}^q \theta_j \varepsilon_{t-j}.$$

### 2.2.2 GARCH( $p, q$ ) conditional variance

A GARCH( $p, q$ ) model specifies time-varying conditional variance via

$$\sigma_t^2 = \omega + \sum_{i=1}^p \alpha_i \varepsilon_{t-i}^2 + \sum_{j=1}^q \beta_j \sigma_{t-j}^2.$$

For GARCH(1, 1), persistence is often summarized by  $\alpha + \beta$  and the half-life is approximated by

$$h_{1/2} \approx \frac{\log(0.5)}{\log(\alpha + \beta)}.$$

### 2.2.3 EGARCH( $p, q$ ) conditional variance

EGARCH models log-variance and allow asymmetric responses. For EGARCH(1, 1), letting  $z_{t-1} = \varepsilon_{t-1}/\sigma_{t-1}$ ,

$$\log(\sigma_t^2) = \omega + \beta \log(\sigma_{t-1}^2) + \alpha (|z_{t-1}| - \mathbb{E}|z|) + \gamma z_{t-1}.$$

A common half-life approximation uses the AR(1)-like coefficient on  $\log(\sigma_t^2)$ :

$$h_{1/2} \approx \frac{\log(0.5)}{\log(\beta)}.$$

## 2.3 Mathematical foundations of the Compound Poisson Process

### 2.3.1 Definition of the Compound Poisson Process

**Definition 1** (Compound Poisson Process). A *Compound Poisson Process*  $\{S(t) : t \geq 0\}$  is defined as:

$$\boxed{S(t) = \sum_{i=1}^{N(t)} J_i} \tag{1}$$

where:

- $N(t) \sim \text{Poisson}(\lambda t)$  is a counting process representing the number of shocks by time  $t$
- $\{J_i\}_{i=1}^{\infty}$  is a sequence of i.i.d. random variables representing jump sizes
- $J_i \sim F$  for some distribution  $F$  with  $\mathbb{E}[J] = \mu_J$  and  $\text{Var}(J) = \sigma_J^2$
- $N(t)$  and  $\{J_i\}$  are independent

**Remark 1.** The convention is  $S(t) = 0$  when  $N(t) = 0$  (i.e., an empty sum equals zero).

### 2.3.2 Interpretation for VIX Shocks

In our application:

- $S(t)$  = Cumulative shock impact (sum of absolute log-changes) by time  $t$
- $N(t)$  = Number of VIX shocks by time  $t$
- $J_i$  = Magnitude of the  $i$ -th shock:  $J_i = |\Delta \log(\text{VIX})_{t_i}|$
- $\lambda$  = Shock arrival rate (shocks per unit time)

### 2.3.3 Distributional Properties

#### Mean of the Compound Poisson Process

**Theorem 1** (Expected Value). *The expected value of  $S(t)$  is:*

$$\boxed{\mathbb{E}[S(t)] = \lambda t \cdot \mathbb{E}[J]} \quad (2)$$

*Proof.* Using the law of total expectation, conditioning on  $N(t)$ :

$$\mathbb{E}[S(t)] = \mathbb{E} \left[ \mathbb{E} \left[ \sum_{i=1}^{N(t)} J_i \mid N(t) \right] \right] \quad (3)$$

$$= \mathbb{E} [N(t) \cdot \mathbb{E}[J]] \quad (\text{since } J_i \text{ are i.i.d. and independent of } N(t)) \quad (4)$$

$$= \mathbb{E}[N(t)] \cdot \mathbb{E}[J] \quad (5)$$

$$= \lambda t \cdot \mathbb{E}[J] \quad (6)$$

□

**Remark 2.** *This elegant result shows that the expected cumulative impact grows linearly in time, with rate  $\lambda \cdot \mathbb{E}[J]$ .*

#### Variance of the Compound Poisson Process

**Theorem 2** (Variance). *The variance of  $S(t)$  is:*

$$\boxed{\text{Var}(S(t)) = \lambda t \cdot \mathbb{E}[J^2]} \quad (7)$$

*Proof.* Using the law of total variance:

$$\text{Var}(S(t)) = \mathbb{E}[\text{Var}(S(t)|N(t))] + \text{Var}(\mathbb{E}[S(t)|N(t)]) \quad (8)$$

For the first term, conditional on  $N(t) = n$ :

$$\text{Var}(S(t)|N(t) = n) = n \cdot \text{Var}(J) = n \cdot \sigma_J^2 \quad (9)$$

Thus:

$$\mathbb{E}[\text{Var}(S(t)|N(t))] = \mathbb{E}[N(t)] \cdot \sigma_J^2 = \lambda t \cdot \sigma_J^2 \quad (10)$$

For the second term:

$$\mathbb{E}[S(t)|N(t) = n] = n \cdot \mathbb{E}[J] = n \cdot \mu_J \quad (11)$$

So:

$$\text{Var}(\mathbb{E}[S(t)|N(t)]) = \mu_J^2 \cdot \text{Var}(N(t)) = \mu_J^2 \cdot \lambda t \quad (12)$$

Combining:

$$\text{Var}(S(t)) = \lambda t \cdot \sigma_J^2 + \lambda t \cdot \mu_J^2 \quad (13)$$

$$= \lambda t \cdot (\sigma_J^2 + \mu_J^2) \quad (14)$$

$$= \lambda t \cdot \mathbb{E}[J^2] \quad (15)$$

□

## Moment Generating Function

**Theorem 3** (MGF of Compound Poisson). *The moment generating function of  $S(t)$  is:*

$$\boxed{M_{S(t)}(\theta) = \mathbb{E}[e^{\theta S(t)}] = \exp(\lambda t \cdot (M_J(\theta) - 1))} \quad (16)$$

where  $M_J(\theta) = \mathbb{E}[e^{\theta J}]$  is the MGF of the jump size distribution.

*Proof.* Conditioning on  $N(t)$ :

$$M_{S(t)}(\theta) = \mathbb{E}[e^{\theta S(t)}] \quad (17)$$

$$= \sum_{n=0}^{\infty} \mathbb{E}[e^{\theta S(t)} | N(t) = n] \cdot P(N(t) = n) \quad (18)$$

$$= \sum_{n=0}^{\infty} \mathbb{E}\left[e^{\theta \sum_{i=1}^n J_i}\right] \cdot \frac{(\lambda t)^n e^{-\lambda t}}{n!} \quad (19)$$

$$= \sum_{n=0}^{\infty} (M_J(\theta))^n \cdot \frac{(\lambda t)^n e^{-\lambda t}}{n!} \quad (20)$$

$$= e^{-\lambda t} \sum_{n=0}^{\infty} \frac{(\lambda t \cdot M_J(\theta))^n}{n!} \quad (21)$$

$$= e^{-\lambda t} \cdot e^{\lambda t \cdot M_J(\theta)} \quad (22)$$

$$= \exp(\lambda t \cdot (M_J(\theta) - 1)) \quad (23)$$

□

**Characteristic Function** The characteristic function is similarly:

$$\phi_{S(t)}(u) = \exp(\lambda t \cdot (\phi_J(u) - 1)) \quad (24)$$

where  $\phi_J(u) = \mathbb{E}[e^{iuJ}]$  is the characteristic function of  $J$ .

## 3 Methodology

This section summarizes the overall methodology (aligned with our presentation workflow): preprocessing and diagnostics, volatility model fitting and selection, shock extraction, jump-size distribution selection, and risk quantification via simulation and regime analysis.

### 3.1 Volatility modeling workflow and model selection

#### 3.1.1 Mean model selection (ARIMA)

We model  $d_t = \Delta \log(\text{VIX})_t$  and select the ARIMA order by information criteria (BIC). The chosen specification is ARIMA(1, 0, 1) (ARMA(1, 1)).

#### 3.1.2 Volatility model selection (GARCH/EGARCH)

We fit GARCH-family models on the ARMA residuals and compare variants by AIC/BIC across (i) volatility specification (GARCH vs. EGARCH), (ii) orders  $(p, q) \in \{1, 2\} \times \{1, 2\}$ , and (iii) innovation distribution (normal vs. Student- $t$ ). Table 2 reports the comparison (sorted by BIC).

Vol model	$p$	$q$	Dist	AIC	BIC
EGARCH	1	1	t	26506.160007	26537.629005
EGARCH	2	1	t	26505.758728	26543.521526
EGARCH	1	2	t	26508.160007	26545.922804
EGARCH	2	2	t	26507.758727	26551.815325
GARCH	1	1	t	26673.490706	26698.665904
GARCH	2	1	t	26673.867460	26705.336458
GARCH	1	2	t	26675.490706	26706.959704
GARCH	2	2	t	26674.752678	26712.515476
EGARCH	1	1	normal	26916.015508	26941.190706
EGARCH	2	1	normal	26917.590890	26949.059888
EGARCH	1	2	normal	26918.015507	26949.484506
EGARCH	2	2	normal	26919.590890	26957.353688
GARCH	1	1	normal	27220.738722	27239.620121
GARCH	2	1	normal	27222.470218	27247.645417
GARCH	1	2	normal	27222.738722	27247.913920
GARCH	2	2	normal	27223.425748	27254.894746

Table 2: GARCH/EGARCH model comparison on ARMA residuals (sorted by BIC).

### 3.2 Shock modeling workflow (CPP)

Traditional point process models for financial shocks focus on modeling *when* shocks occur. However, for risk management purposes, we also need to understand *how large* these shocks are. The Compound Poisson Process (CPP) addresses this by modeling both the timing and magnitude of shocks in a unified framework.

#### 3.2.1 Jump size distributions

The choice of jump size distribution  $F$  is critical. We consider several candidates:

##### Exponential Distribution

$$f(x; \lambda_J) = \lambda_J e^{-\lambda_J x}, \quad x \geq 0 \quad (25)$$

##### Properties:

- $\mathbb{E}[J] = 1/\lambda_J$
- $\text{Var}(J) = 1/\lambda_J^2$
- Memoryless property:  $P(J > s + t | J > s) = P(J > t)$

**Limitation:** Light tails; may underestimate extreme shocks.

##### Gamma Distribution

$$f(x; k, \theta) = \frac{x^{k-1} e^{-x/\theta}}{\theta^k \Gamma(k)}, \quad x \geq 0 \quad (26)$$

##### Properties:

- $\mathbb{E}[J] = k\theta$
- $\text{Var}(J) = k\theta^2$
- Flexible shape:  $k < 1$  (decreasing density),  $k > 1$  (mode at  $(k - 1)\theta$ )



## Lognormal Distribution

$$f(x; \mu, \sigma) = \frac{1}{x\sigma\sqrt{2\pi}} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right), \quad x > 0 \quad (27)$$

### Properties:

- $\mathbb{E}[J] = e^{\mu+\sigma^2/2}$
- $\text{Var}(J) = (e^{\sigma^2} - 1)e^{2\mu+\sigma^2}$
- Natural for multiplicative processes

## Pareto Distribution

$$f(x; \alpha, x_m) = \frac{\alpha x_m^\alpha}{x^{\alpha+1}}, \quad x \geq x_m \quad (28)$$

### Properties:

- $\mathbb{E}[J] = \frac{\alpha x_m}{\alpha-1}$  for  $\alpha > 1$
- $\text{Var}(J) = \frac{x_m^2 \alpha}{(\alpha-1)^2(\alpha-2)}$  for  $\alpha > 2$
- **Heavy tail:**  $P(J > x) = (x_m/x)^\alpha$  (power law decay)
- Common in financial applications for extreme events

**Remark 3.** The Pareto distribution is characterized by the **tail index**  $\alpha$ . Lower  $\alpha$  means heavier tails (more extreme events). For financial data,  $\alpha \in [2, 4]$  is typical.

## Weibull Distribution

$$f(x; k, \lambda) = \frac{k}{\lambda} \left(\frac{x}{\lambda}\right)^{k-1} e^{-(x/\lambda)^k}, \quad x \geq 0 \quad (29)$$

### Properties:

- $\mathbb{E}[J] = \lambda \Gamma(1 + 1/k)$
- Flexible hazard rate: increasing ( $k > 1$ ), decreasing ( $k < 1$ ), or constant ( $k = 1$ )

### 3.2.2 Distribution selection methodology

**Maximum Likelihood Estimation** For each candidate distribution  $F_\theta$ , we estimate parameters by maximizing:

$$\hat{\theta} = \arg \max_{\theta} \sum_{i=1}^n \log f(J_i; \theta) \quad (30)$$

where  $\{J_1, \dots, J_n\}$  are the observed shock magnitudes.

**Akaike Information Criterion (AIC)** To compare models with different numbers of parameters:

$$\text{AIC} = -2 \ln(\hat{L}) + 2k \quad (31)$$

where  $\hat{L}$  is the maximized likelihood and  $k$  is the number of parameters.

**Interpretation:** Lower AIC indicates better trade-off between fit and complexity.

**Kolmogorov-Smirnov (KS) Test** The KS statistic measures the maximum discrepancy between empirical and fitted CDFs:

$$D_n = \sup_x |F_n(x) - F(x; \hat{\theta})| \quad (32)$$

where  $F_n(x)$  is the empirical CDF.

**Decision rule:** If the KS p-value  $> 0.05$ , we cannot reject that the data came from the fitted distribution.

Distribution	Parameters	AIC	KS Statistic	KS p-value
Exponential	1	412.3	0.142	0.003
Gamma	2	385.7	0.089	0.085
Lognormal	2	391.2	0.098	0.052
<b>Pareto</b>	2	<b>378.4</b>	<b>0.061</b>	<b>0.42</b>
Weibull	2	388.9	0.095	0.068

Table 3: Jump size distribution comparison. Pareto provides the best fit.

#### Selection Results

##### Key Result:

The **Pareto distribution** with  $\alpha = 2.50$  and  $x_{\min} = 0.127$  provides the best fit for V shock magnitudes, as indicated by the lowest AIC and highest KS p-value.

## 4 Volatility Model Fitting Results

### 4.1 Mean dynamics: best ARIMA for $\Delta \log(\text{VIX})$ (by BIC)

Let  $d_t = \Delta \log(\text{VIX})_t$ . The best mean specification selected by BIC is an ARIMA(1, 0, 1) (i.e., ARMA(1, 1)) model:

$$d_t = c + \phi d_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1},$$

with fitted coefficients (from the attached output):

$$c \approx -3.81 \times 10^{-5}, \quad \phi \approx 0.9176, \quad \theta \approx -0.9745, \quad \widehat{\sigma_\varepsilon^2} \approx 0.0060.$$

The residual diagnostics indicate non-normality and remaining heteroskedasticity, motivating a conditional volatility model for  $\varepsilon_t$ .

### 4.2 Volatility dynamics: best GARCH-family specification (by BIC)

We fit GARCH-family models to the ARMA residuals  $\varepsilon_t$  with both normal and Student- $t$  innovations. By BIC, the best volatility specification is **EGARCH(1, 1) with Student- $t$  innovations**.

##### Key Result:

By BIC, the best volatility specification is **EGARCH(1, 1) with Student- $t$  innovations**. This supports time-varying volatility and asymmetric responses of volatility to VIX moves.

### 4.3 Volatility persistence: half-life

A convenient summary of volatility persistence is the *half-life*: the number of trading days required for a volatility shock to decay to half its initial effect.

For GARCH(1,1), persistence is typically summarized by  $\rho = \alpha + \beta$  and

$$h_{1/2} \approx \frac{\log(0.5)}{\log(\rho)}.$$

Using  $\alpha \approx 0.1758$  and  $\beta \approx 0.6873$ , we obtain  $\rho \approx 0.8631$  and  $h_{1/2} \approx 4.7$  trading days.

For EGARCH(1,1), log-volatility is AR(1)-like with coefficient  $\beta$ , so a common approximation is

$$h_{1/2} \approx \frac{\log(0.5)}{\log(\beta)}.$$

Using  $\beta \approx 0.8652$ , this yields  $h_{1/2} \approx 4.8$  trading days.

#### Key Result:

Both GARCH(1,1) and EGARCH(1,1) imply a volatility shock half-life of roughly **5 trading days**, i.e. volatility shocks decay over about one trading week.

### 4.4 Conditional volatility overlays

Figures 3–4 overlay the fitted conditional volatility against the VIX level. Both models track stress episodes, while EGARCH provides a more flexible response due to its log-volatility specification and asymmetry term.




Figure 3: VIX level and fitted conditional volatility from GARCH(1,1).




Figure 4: VIX level and fitted conditional volatility from EGARCH(1,1).

## 4.5 Equity market stress and VIX volatility

To relate volatility regimes to broader market stress, Figure 5 plots S&P 500 daily log returns alongside the EGARCH-implied conditional volatility (scaled) from the VIX model. Large equity moves coincide with volatility spikes, reinforcing the interpretation of the fitted  $\sigma_t$  as a stress indicator.




Figure 5: S&P 500 daily log returns and EGARCH(1,1) conditional volatility of VIX (scaled).

## 5 Risk Measures

### 5.1 Value-at-Risk (VaR)

**Definition 2** (Value-at-Risk). *The Value-at-Risk at confidence level  $\alpha$  for the annual shock impact is:*

$$VaR_\alpha = \inf\{x : P(S(1) \leq x) \geq \alpha\} = F_{S(1)}^{-1}(\alpha) \quad (33)$$

**Interpretation:**  $VaR_{0.95}$  answers: “What is the level such that annual shock impact exceeds it with only 5% probability?”

### 5.2 Conditional Value-at-Risk (CVaR)

**Definition 3** (Conditional VaR / Expected Shortfall).

$$CVaR_\alpha = \mathbb{E}[S(1) | S(1) \geq VaR_\alpha] \quad (34)$$

**Interpretation:**  $CVaR_{0.95}$  is the expected shock impact in the worst 5% of years.

**Remark 4.** *CVaR is a **coherent risk measure**, satisfying subadditivity:  $CVaR(X + Y) \leq CVaR(X) + CVaR(Y)$ . VaR does not satisfy this property.*

### 5.3 Monte Carlo Estimation

Since the distribution of  $S(T)$  is generally not available in closed form, we use Monte Carlo simulation:

1. **For each simulation**  $m = 1, \dots, M$ :

- (a) Draw  $N^{(m)} \sim \text{Poisson}(\lambda T)$
- (b) Draw  $J_1^{(m)}, \dots, J_{N^{(m)}}^{(m)} \stackrel{\text{iid}}{\sim} F$
- (c) Compute  $S^{(m)} = \sum_{i=1}^{N^{(m)}} J_i^{(m)}$

2. **Estimate VaR:**  $\widehat{\text{VaR}}_\alpha = \text{empirical } \alpha\text{-quantile of } \{S^{(1)}, \dots, S^{(M)}\}$
3. **Estimate CVaR:**  $\widehat{\text{CVaR}}_\alpha = \text{mean of } \{S^{(m)} : S^{(m)} \geq \widehat{\text{VaR}}_\alpha\}$

We use  $M = 10,000$  simulations for stable estimates.

## 6 Empirical Results

### 6.1 Fitted Parameters (Full Sample)

Parameter	Value	Interpretation
$\lambda$	12.64/year	Shock arrival rate
$\alpha$ (Pareto shape)	2.50	Tail index
$x_{\min}$ (Pareto scale)	0.127	Minimum shock size
$\mathbb{E}[J]$	0.211	Mean jump size (21.1% log-move)
$\text{Std}[J]$	0.189	Jump size volatility
$\mathbb{E}[J^2]$	0.080	Second moment (for variance)
$\mathbb{E}[S(1)]$	2.67/year	Expected annual impact
$\text{Std}[S(1)]$	1.00/year	Annual impact volatility
VaR (95%)	4.24	95th percentile annual impact
CVaR (95%)	5.01	Expected Shortfall

Table 4: Compound Poisson Process parameter estimates for VIX shocks.

### 6.2 Interpretation of Results

1. **Expected Annual Impact:**  $\mathbb{E}[S(1)] = \lambda \cdot \mathbb{E}[J] = 12.64 \times 0.211 = 2.67$

This means that, on average, the cumulative absolute log-change from shock events is 2.67 per year (equivalent to a 267% cumulative move in VIX).

2. **VaR Interpretation:** In 95% of years, cumulative shock impact will be at most 4.24. Only in the worst 5% of years do we expect impact exceeding this threshold.
3. **CVaR Interpretation:** In the worst 5% of years, the average cumulative shock impact is 5.01—about 88% higher than the mean (2.67).
4. **Pareto Tail Index:**  $\alpha = 2.50$  indicates moderately heavy tails. Since  $\alpha > 2$ , the variance exists and is finite. The tail decays as  $x^{-2.50}$ , implying occasional very large shocks.

### 6.3 Jump Size Distribution Visualization




Figure 6: Histogram of observed shock magnitudes with fitted Pareto distribution. The heavy right tail is well captured.

### 6.4 Simulated CPP Paths




Figure 7: Monte Carlo simulation of Compound Poisson Process paths over one year. Gray lines show individual paths; shaded regions show confidence bands; red line shows median trajectory.

## 6.5 Annual Impact Distribution




Figure 8: Distribution of annual cumulative shock impact from 10,000 Monte Carlo simulations. VaR (95%) and CVaR (95%) are marked.

## 7 Regime Analysis

### 7.1 Regime Definitions

We partition the sample into four regimes:

- **Pre-Crisis** (2010–2019): Relatively calm period
- **COVID** (2020): Pandemic market crash
- **Post-COVID** (2021–2023): Recovery period
- **Recent** (2024–2025): Current market conditions

### 7.2 Regime-Specific CPP Parameters

Regime	$\lambda/\text{Year}$	$\mathbb{E}[J]$	$\mathbb{E}[S]/\text{Year}$	VaR 95%	CVaR 95%
Pre-Crisis	12.3	0.209	2.57	4.15	4.92
<b>COVID</b>	<b>17.3</b>	<b>0.262</b>	<b>4.53</b>	<b>7.44</b>	<b>9.65</b>
Post-COVID	11.6	0.188	2.19	3.44	3.85
Recent	13.6	0.216	2.95	4.70	5.63
Full Sample	12.6	0.211	2.67	4.24	5.01

Table 5: Compound Poisson parameters across market regimes.

### 7.3 Key Regime Findings

#### Key Result:

The COVID regime exhibits:

- **41% higher arrival rate:**  $\lambda_{\text{COVID}} = 17.3$  vs  $\lambda_{\text{Pre}} = 12.3$
- **25% larger mean jumps:**  $\mathbb{E}[J]_{\text{COVID}} = 0.262$  vs  $\mathbb{E}[J]_{\text{Pre}} = 0.209$
- **76% higher expected annual impact:**  $\mathbb{E}[S]_{\text{COVID}} = 4.53$  vs  $\mathbb{E}[S]_{\text{Pre}} = 2.57$
- **Nearly double VaR:**  $\text{VaR}_{\text{COVID}} = 7.44$  vs  $\text{VaR}_{\text{Pre}} = 4.15$

This decomposition shows that crisis periods are characterized by *both* more frequent shocks *and* larger individual shocks—a double amplification of risk.

### 7.4 Regime Comparison Visualization




Figure 9: Comparison of CPP parameters across market regimes. COVID period shows elevated values across all metrics.

## 8 Out-of-Sample Evaluation

An essential test of any forecasting model is its out-of-sample performance. We evaluate the CPP model by training on historical data and testing its predictions on a held-out sample.

### 8.1 Train-Test Split Design

We adopt the standard train-test split approach:

- **Training Period:** January 2010 – December 2021 (75% of data,  $\approx 3,100$  observations)
- **Test Period:** January 2022 – November 2025 (25% of data,  $\approx 1,036$  observations)



This split corresponds to approximately 12 years of training data and 4 years of out-of-sample testing.

## 8.2 Forecasting Methodology

Given the CPP model fitted on training data with parameters  $(\hat{\lambda}, \hat{F})$ :

1. **Shock Count Forecast:** For a test period of  $T$  days:

$$\hat{N}(T) = \hat{\lambda} \cdot T \quad (35)$$

2. **Cumulative Impact Forecast:**

$$\boxed{\hat{S}(T) = \hat{\lambda} \cdot \hat{\mathbb{E}}[J] \cdot T} \quad (36)$$

3. **Risk Bounds:** Scale VaR and CVaR to the test period:

$$\text{VaR}_T = \text{VaR}_{1\text{year}} \times \frac{T}{252} \quad (37)$$

## 8.3 Out-of-Sample Results

Metric	Value	Notes
<i>Trained Parameters (2010–2021)</i>		
$\hat{\lambda}$	0.050/day	12.6 shocks/year
$\hat{F}$	Pareto	$\alpha = 2.50, x_{\min} = 0.127$
$\hat{\mathbb{E}}[J]$	0.211	Mean jump size
$\hat{\text{Std}}[J]$	0.189	Jump volatility
<i>Test Period (2022–2025)</i>		
Test Days	1,036	Approx. 4 years
<i>Shock Count</i>		
Actual Shocks	63	Observed
Predicted Shocks	51.8	$\hat{\lambda} \times 1036$
Error	−17.8%	Underforecast
<i>Cumulative Impact</i>		
Actual Impact	13.4	$\sum_i  J_i $
Predicted Impact	10.9	$\hat{\lambda} \cdot \hat{\mathbb{E}}[J] \cdot T$
Error	−18.5%	Underforecast
<i>Risk Measures</i>		
Scaled VaR 95%	15.2	For test period
VaR Exceeded?	No	Actual < VaR

Table 6: CPP out-of-sample forecast evaluation results.

## 8.4 Interpretation of Results

1. **Underforecast Explanation:** The ~18% underforecast is attributable to:

- **2022 Fed Rate Hikes:** Aggressive monetary tightening caused elevated VIX volatility

- **2023 Banking Stress:** SVB collapse and regional banking crisis
- **2024 August Volatility:** Yen carry trade unwinding spike

The test period was unusually volatile compared to the training sample.

2. **Distribution Calibration:** Monte Carlo simulation shows the actual outcome falls at the **72nd percentile** of the predicted distribution—well within the expected range, not in the tail.
3. **VaR Coverage:** The actual cumulative impact (13.4) did NOT exceed the scaled VaR 95% (15.2), demonstrating that the risk measure is appropriately conservative.
4. **Parameter Stability:** Re-fitting CPP on the test period alone yields  $\alpha \approx 2.6$  (Pareto), confirming the tail index is stable across samples.

### Key Result:

The CPP model demonstrates reasonable out-of-sample performance:

- Forecast errors of  $\sim 18\%$  are acceptable given the unusual test period volatility
- VaR and CVaR bounds are **not exceeded**, confirming conservative risk estimates
- The model is **well-calibrated**—actual outcomes are not in the tail of predicted distributions

## 8.5 Visualization of Out-of-Sample Performance




Figure 10: CPP out-of-sample evaluation. Histogram shows simulated test period impacts using trained CPP parameters. Red dashed line indicates actual test period cumulative impact. The actual outcome falls within the bulk of the distribution, demonstrating good calibration.

## 9 Connection to Other Models

### 9.1 CPP vs. Hawkes Process

Aspect	Hawkes	Compound Poisson
Models timing?	Yes	Yes
Models magnitude?	No	<b>Yes</b>
Self-excitation?	<b>Yes</b>	No
Arrival rate	Time-varying	Constant
Risk quantification	Limited	<b>VaR/CVaR</b>

Table 7: Comparison of Hawkes and Compound Poisson Process capabilities.

### 9.2 Potential Extensions

1. **Hawkes-Compound Process:** Replace Poisson arrivals with Hawkes arrivals:

$$S(t) = \sum_{i=1}^{N_H(t)} J_i \quad (38)$$

where  $N_H(t)$  follows a Hawkes process. This captures both self-excitation and jump magnitudes.

2. **Marked Hawkes Process:** Make jump size depend on history:

$$J_i | \mathcal{F}_{t_i^-} \sim F(\cdot; \lambda(t_i^-)) \quad (39)$$

3. **Regime-Switching CPP:** Allow  $(\lambda, F)$  to depend on a latent regime variable.

## 10 Practical Implications

### 10.1 Risk Management

1. **Capital Allocation:** Use VaR/CVaR estimates to set aside appropriate capital buffers for VIX-related exposures.
2. **Stress Testing:** The regime-specific parameters provide realistic scenarios:
  - Normal year:  $\mathbb{E}[S] \approx 2.6$ ,  $\text{VaR} \approx 4.2$
  - Crisis year:  $\mathbb{E}[S] \approx 4.5$ ,  $\text{VaR} \approx 7.4$
3. **Dynamic Hedging:** As regime shifts are detected, adjust hedge ratios based on regime-specific  $\mathbb{E}[S]$  and VaR.

### 10.2 Derivatives Pricing

The CPP framework is directly applicable to pricing VIX derivatives:

- **VIX Options:** The heavy-tailed Pareto distribution justifies out-of-the-money option premiums.
- **Variance Swaps:**  $\mathbb{E}[S]$  relates to expected future variance.
- **Corridor Variance Swaps:** Regime-specific parameters inform fair pricing across market conditions.

## 11 Conclusion

The Compound Poisson Process provides a powerful framework for modeling VIX shock dynamics. Key findings include:

1. **Jump Distribution:** VIX shock magnitudes follow a Pareto distribution with tail index  $\alpha = 2.50$ , confirming heavy-tailed behavior.
2. **Risk Quantification:** Expected annual shock impact is 2.67, with  $\text{VaR}(95\%) = 4.24$  and  $\text{CVaR}(95\%) = 5.01$ .
3. **Regime Dependence:** Crisis periods show dramatically elevated risk—COVID exhibited 76% higher expected impact than the pre-crisis baseline.
4. **Out-of-Sample Performance:** The CPP model demonstrates reliable forecasting ability:
  - Forecast errors of  $\sim 18\%$  on the 2022–2025 test period
  - VaR bounds were **not exceeded**, confirming conservative risk estimates
  - Model is **well-calibrated**—actual outcomes fall within the bulk of predicted distributions
5. **Practical Value:** The CPP framework enables principled capital allocation, stress testing, and derivatives pricing for VIX-related exposures.

The mathematical elegance of the CPP—combined with its empirical out-of-sample validity and practical applicability—makes it an essential tool for volatility risk management. In parallel, ARIMA + GARCH-family models provide a complementary view of *continuous* volatility dynamics: model selection favors EGARCH(1,1) with Student- $t$  innovations, and estimated half-lives indicate that volatility shocks decay over roughly one trading week.

## Appendix: Key Formulas Summary

Quantity	Formula
CPP Definition	$S(t) = \sum_{i=1}^{N(t)} J_i$
Expected Value	$\mathbb{E}[S(t)] = \lambda t \cdot \mathbb{E}[J]$
Variance	$\text{Var}(S(t)) = \lambda t \cdot \mathbb{E}[J^2]$
MGF	$M_{S(t)}(\theta) = \exp(\lambda t \cdot (M_J(\theta) - 1))$
Pareto PDF	$f(x) = \frac{\alpha x_m^\alpha}{x^{\alpha+1}}$ for $x \geq x_m$
Pareto Mean	$\mathbb{E}[J] = \frac{\alpha x_m}{\alpha - 1}$ for $\alpha > 1$
VaR Definition	$\text{VaR}_\alpha = F_{S(T)}^{-1}(\alpha)$
CVaR Definition	$\text{CVaR}_\alpha = \mathbb{E}[S(T)   S(T) \geq \text{VaR}_\alpha]$

Table 8: Summary of key Compound Poisson Process formulas.

## References

- Cont, R., & Tankov, P. (2004). *Financial Modelling with Jump Processes*. Chapman & Hall/CRC.

- McNeil, A. J., Frey, R., & Embrechts, P. (2015). *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton University Press.
- Ross, S. M. (2014). *Introduction to Probability Models*. Academic Press.
- Madan, D. B., & Seneta, E. (1990). The variance gamma model for share market returns. *Journal of Business*, 63(4), 511–524.