

**Protocol for estimation of renal threshold for glucose excretion (RTG)
from an oral glucose tolerance test (OGTT)**

1. Discuss the procedure with the subject and obtain informed consent.
2. The subject is instructed to fast for 8 hours before the test.
3. The subject empties their bladder before the test.
4. At the beginning of the test, the subject drinks a glucose solution containing 75g glucose in 5 minutes.
5. Plasma glucose is sampled before (0min) and 30min, 60min, 90min, 120min, 180min, and 240min after glucose ingestion.
6. Each time the subject urinates during the 4-hour test, urine volume is documented and urinary glucose concentration is measured. At the end of the test (t=240min), the subject empties their bladder to collect the last urine sample. Calculate the amount of urinary glucose excretion during the test (UGE_{240min}).
7. Calculate the renal threshold for glucose excretion (RTG) from plasma glucose levels and UGE_{240min} using <https://rtg.renaltubule.com>.

**Data collection sheet for determination of
renal threshold for glucose excretion (RTG)**

Patient Name:

Patient ID:

Age:

Gender:

Height(m):

Weight(kg):

| | | | | | | | |
|---|---|----|-------------------|-------|-----|---------------------------|-----|
| Plasma glucose levels over time | | | | | | | |
| Time(min) | 0 | 30 | 60 | 90 | 120 | 180 | 240 |
| Plasma glucose (mmol/L) | | | | | | | |
| Urinary glucose excretion | | | | | | | |
| Number of urination | 1 | 2 | 3 | 4 | 5 | | |
| Urine volume (ml) | | | | | | | |
| Urinary glucose (mmol/L) | | | | | | | |
| Urinary glucose excretion over the four hours (UGE _{240min})= | | | | | | mmol | |
| Serum creatinine= | | | $\mu\text{mol/L}$ | eGFR= | | ml/min/1.73m ² | |