

Wnt signaling: why is everything so negative?

Jeffrey D Brown* and Randall T Moon†

The Wnt proteins constitute a family of secreted glycoproteins the members of which have essential signaling roles during embryogenesis. The recent identification of several new regulators of this signal transduction pathway have revealed unexpectedly intricate levels of constraint on Wnt-dependent gene activation, and studies in developing embryos and in cell culture systems have allowed a more complete understanding of the functional and biochemical interactions between components of this evolutionarily conserved pathway.

Addresses

University of Washington, Howard Hughes Medical Institute, Department of Pharmacology and Molecular and Cellular Biology Program, Box 357370, Seattle, WA 98195, USA

*e-mail: brownj@u.washington.edu

†e-mail: rtmoon@u.washington.edu

Current Opinion in Cell Biology 1998, 10:182–187

<http://biomednet.com/elecref/0955067401000182>

© Current Biology Ltd ISSN 0955-0674

Abbreviations

APC	adenomatous polyposis coli protein
β-cat	β-catenin
BMP	bone morphogenic protein
CRD	cystein rich domain
CKII	casein kinase II
Dsh	Dishevelled
GSK-3β	glycogen synthase kinase-3β
LEF/TCF	lymphoid enhancer factor/T-cell factor
MBT	mid-blastula transition
PI	phosphatidylinositol
PKC	protein kinase C

Introduction

Signal transduction pathways are elegant systems by which cells and organisms can amplify subtle signals to generate robust responses, but it is essential that signalling pathways are not activated spuriously. To that apparent end the molecule at the heart of the Wnt signaling pathway, β-catenin (or Armadillo, the *Drosophila* homolog), is maintained under constant negative regulation by the constitutively active glycogen synthase kinase-3β (GSK-3β). Only by repressing this perpetual inhibition can a Wnt signal be transduced (reviewed in [1]). In addition, several secreted factors inhibit Wnt signaling [2,3•] (reviewed in [4]), as do other signaling pathways [5,6]. Data accumulated during the past 18 months have filled many gaps in our understanding of signal transduction and have revealed further levels of negative regulation of Wnt signaling. In this review we will concentrate our attention on newly identified components of the pathway as well as on the positive and negative regulation of Wnt signal transduction.

Wnt signaling basics


In the absence of Wnt signaling, GSK-3β causes the free cytosolic pool of β-catenin to be degraded (Figure 1; [7–9], reviewed in [1]). Association of appropriate members of the Wnt (reviewed in [10]) and Frizzled [11] protein families leads to activation of the signaling pathway (Figure 1; [12,13,14•]). The Dishevelled (Dsh) protein, the most proximal intracellular component known in the pathway [15] is somehow activated, phosphorylated in its central domain by casein kinase II (CKII) [16•], and is recruited to cell membranes [13,17]. In spite of changes in localization and phosphorylation state, it is not clear how Dishevelled is activated. Overexpression of either Dishevelled or the *Drosophila* Frizzled-2 protein is sufficient to lead to phosphorylation by CKII, but does not necessarily lead to activation of the protein [16•]. Activation of Dishevelled somehow leads to the inhibition of GSK-3β, possibly involving protein kinase C (PKC) phosphorylation of an amino-terminal serine residue of GSK-3β [18]. As newly translated β-catenin accumulates because of reduced GSK-3β activity, β-catenin can then interact with members of the lymphoid enhancer factor/T-cell factor (LEF/TCF) family of architectural transcription factors [19,20,21••] in the nucleus [22•]. In the nucleus the complex binds consensus LEF/TCF sites in promoters and induces transcription of Wnt-responsive genes (Figure 1; [20,23,24•], reviewed in [25]). Several recent reviews have described the ‘classical’ Wnt-1 signal transduction pathway and the roles of Wnt signaling during development in greater detail [4,10,26,27].

Additional attention has recently been focused upon alternative Wnt and Frizzled signaling pathways. Examples of Wnt/Frizzled signaling pathways that diverge from the model presented in Figure 1 have been observed in *Xenopus* [2] (reviewed in [28]), zebrafish [29•,30••], *Drosophila* [31] (reviewed in [32]), and *Caenorhabditis elegans* [33,34••]. It is not clear whether the divergent pathways in these organisms represent a second cohesive system by which a subset of Wnt signals are transduced, or whether animals with different developmental strategies have tailored components of the well-conserved ‘Wnt-1 pathway’ (Figure 1) to their own needs.

Antagonists abound

Models invoking morphogenic gradients of secreted signaling factors to explain patterning of developing tissues have been a favorite among developmental biologists for a century; however, the recent identification of several secreted antagonists of bone morphogenic protein (BMP), reviewed in [35]) and Wnt (below) signaling add a new dimension to our understanding of these patterning

Figure 1



The 'classical' Wnt-1 signaling pathway. In the absence of Wnt-1 class signals, or if Wnts are bound and inhibited by members of the Frzb family, a complex of GSK-3 β and APC target β -catenin for degradation. Low levels of cytosolic β -cat are insufficient to be translocated to the nucleus and associate with LEF/TCF architectural transcription factors. LEF/TCF are likely bound to promoters in the absence of β -cat and repress basal transcription. Wnt likely binds to the extracellular cysteine-rich domain of Frizzled and leads to the hyperphosphorylation of Dsh by CKII. CKII phosphorylation is not sufficient to activate Dsh; other events are also required. Activated Dsh inhibits GSK-3 β function by a poorly understood mechanism that may involve PKC phosphorylation of GSK-3 β . β -cat levels rise as the result of decreased rates of degradation mediated by GSK-3 β . β -cat appears to be translocated into the nucleus independent of association with LEF/TCF. Inside the nucleus, β -cat may alter the ability of LEF/TCF to bend DNA and may relieve LEF/TCF-dependent transcriptional repression. A carboxy-terminal domain of β -cat then activates transcription.

events. It seems likely that secreted antagonists of other signaling proteins will be revealed in the future.

The Frzb family

The Frzb proteins are expressed in a wide range of tissues in embryos and adults and seem to act by binding to secreted Wnt proteins and preventing productive interactions between Wnt proteins and Frizzled proteins. (reviewed in [4]). The prototypic member of this family, FRZB, was cloned by researchers interested in a chondrogenic activity observed in a purified protein fraction from bovine cartilage [36]. Microsequence analysis of the purified protein followed by degenerate PCR

(polymerase chain reaction) cloning and library screening resulted in the isolation of a full-length cDNA encoding a cysteine-rich domain (CRD) with significant similarity to the extracellular amino-terminal domain of Frizzled receptors [36]—a region now known to be important for binding of Wnt [12,37]. Subsequently, other groups cloned members of the Frzb family by differential screens [38,39,40], isolation of proteins co-purifying with mitogenic fractions from fibroblastic cell lines [41,42], activity screens [43] and by homology to known Frizzled receptors [44]. At least five distinct members of the Frzb family have been cloned to date, with homologs isolated from several vertebrate species.

Members of the Frzb family seem to be surprisingly similar in their specificity for inhibiting Wnt signaling. The three members tested to date, Frzb-1/sFRP-3, sFRP-2, FRP/sFRP-1, have been shown to interact with Wnt-1 [37•,38•,45•], Wnt-8 [45•] or Wingless [44•] by indirect immunofluorescence or coimmunoprecipitation. In various assays for Wnt function in *Xenopus*, Frzb-1/fiz [38•,42,45•] and FRP/sFRP-1 [41,44•] block the activities of Wnt-1, Wnt-8 and Wingless, all members of the 'Wnt-1 functional class' [46,47] (reviewed in [28]). These Frzbs do not efficiently inhibit Wnt-3a activity [41•,48•], in spite of its 'Wnt-1-like' activity in separate assays [46,47]. This result suggests that Frzb proteins discriminate between Wnt proteins that seem functionally similar.

It is interesting to note that although Frzb-1 seems to bind to Wnt-5a it does not seem to interfere with signaling mediated by Wnt-5a [37•,48•]. The observation that association of the Frzb and Wnt proteins does not necessarily inhibit Wnt function was also demonstrated by examination of the activity of various Frzb deletion constructs [37•]. Although mutants of Frzb-1 with small deletions in the CRD bind Wnt-1 in co-immunoprecipitation assays, they do not seem to efficiently interfere with Wnt-1 signaling [37•].

Cerberus antagonism: is it all in your head?

Cerberus is a secreted factor expressed in the *Xenopus* gastrula organizer and is thought to have a role in head induction [49]—a process inhibited by ectopic Wnt-8 signaling in the gastrula dorsal mesoderm [50]. Cerberus has recently been shown to be a potent inhibitor of Wnt signaling in early *Xenopus* embryos, though the mechanism of inhibition is not understood [3•]. As the *Cerberus* expression domain overlaps that of *Frzb-1*, it seems that the gastrula organizer is a source of multiple inhibitors of Wnt signaling, which likely serve to sharpen the boundaries of cell fate (reviewed in [4]; see also Note added in proof).

The enemy within: Xwnt-5a?

The Frzbs proteins may be lurking impostors of Frizzled receptors lying in wait for stray Wnt proteins, but the embryonic responses to some Wnt proteins may also be blocked by members of their own family. It has become clear that the Wnt and Frizzled proteins can transduce at least two distinct types of signal [2,29•,30•,34•,46,47] (reviewed in [28,32]). In various assays, Wnt proteins can be assigned to the 'Wnt-1 class', which transduces signals via the pathway in Figure 1, or to the Wnt-5a class [47], which activates the phosphatidylinositol (PI) cycle via a G-protein coupled pathway [30•]. Members of the Wnt-5a class can inhibit embryonic responses to members of the Wnt-1 class, at least in early *Xenopus* embryos [2]. This inhibition is likely indirect and can be mimicked by stimulating the PI cycle and calcium signaling by a serotonin receptor [29•]. It is worth noting that although Wnt-5a and related Wnt proteins do not

stabilize β -catenin (β -cat) in *Xenopus* embryos before the mid-blastula transition (MBT; the onset of zygotic transcription), Wnt-5a signaling after MBT leads to the stabilization of β -catenin [51•]. This shift in signaling identity is probably the result of the expression of a distinct member of the Frizzled family after MBT, perhaps Frizzled-5 [14•], which can bind Wnt-5a class ligands and transduce a Wnt-1-like signal [51•].

The nuts and bolts of β -catenin regulation

Modest increases in levels of β -catenin are sufficient to activate transcription of Wnt-responsive genes in embryos [24•,51•,52•]. Phosphorylation of β -catenin on conserved amino-terminal serine and threonine residues by GSK-3 β [7] may be all that is required to target the protein for ubiquitination and proteasome degradation [9]; however, others have suggested that β -catenin/Armadillo may be a poor substrate for GSK-3 β *in vitro* [8,53] and suggest an alternative mechanism involving the adenomatous polyposis coli protein (APC). APC has been implicated, at least in tumor cell lines, in the regulation of β -catenin levels [8], however, *Drosophila* embryos lacking zygotic APC regulate Armadillo appropriately [54] (reviewed in [55]). A new player, Axin, may also have a role in this part of the pathway, as described below.

As mutations at the *Fused* locus (as distinct from *Drosophila fused* mutants) were known to cause twinning of the axis in homozygous mouse embryos (reviewed in [56•]), investigators were interested in determining if Axin, the protein encoded at the locus, was involved in regulation of signaling pathways known to cause axis duplications in *Xenopus* embryos [56•]. Overexpression of Axin in cleavage-stage *Xenopus* embryos inhibited the ability of Wnt, Dishevelled, or a dominant-negative mutant of GSK-3 β to promote formation of an ectopic dorsal axis. In contrast, Axin did not block axis induction in response to β -catenin or Siamois, a homeodomain-containing protein induced by Wnt signaling in the early *Xenopus* embryo [13,56•,57,58]. Other factors known to cause axis duplication in *Xenopus* such as Noggin, Activin and a dominant negative form of the BMP receptor were also unaffected by Axin. A mutant of Axin lacking a carboxy-terminal domain with identity to RGS proteins (Axin Δ RGS, regulators of G-protein signaling, reviewed in [59]), behaves like a dominant negative and results in axial duplication when ectopically expressed. Co-expression of cadherin, which may inhibit Wnt signaling by sequestering β -catenin from the free cytoplasmic signaling pool, blocks axis induction by this mutant, suggesting that Axin Δ RGS causes axis duplication by increasing the free signaling pool of β -catenin [56•]. The mechanism of Axin action is not known, and biochemical studies will certainly add to our understanding of its regulation of β -catenin.

LEF/TCF: architects of transcription

Although the nuclear translocation of β -catenin had been thought to correlate with its signaling function in the

Wnt pathway, it has only recently become clear how this activity is transduced. Two-hybrid screens revealed interactions between the LEF/TCF family of architectural transcription factors and β -catenin [19,20]. Complexes of LEF/TCF and β -catenin/Armadillo have been shown to bind to consensus LEF/TCF sites in promoters of several genes [23,24•,60] and in synthetic constructs [19,20]. Interactions are mediated by the conserved amino termini of LEF/TCF family members and arm repeats 3–8 of β -catenin/Armadillo [19,20,22•,23]. In *Drosophila*, *dTCF/pangolin* seems to be required in all *wingless*-dependent signaling events examined [22•,23,61••].

In spite of the sequence similarity that LEF and TCF family members retain in their amino termini and in their highly conserved HMG box DNA-binding domains, these proteins are not functionally equivalent. Overexpression of LEF-1 in *Xenopus* [20] and *Drosophila* [23] results in the activation of Wnt signaling pathways, whereas overexpression of *Xenopus* Tcf-3 [19] or *Drosophila* dTCF/Pangolin [22•] do not.

Association of β -catenin/armadillo with LEF/TCF activates transcription from endogenous and synthetic promoter fragments [19,22•,24•]. This activation could, in principle, be the result of a release from LEF/TCF-mediated transcriptional repression (reviewed in [62]). Indeed, it has been suggested that the LEF/TCF transcription factors are bound to promoters in the absence of β -catenin/Armadillo and repress basal transcription [23,24•]. Also, in an early polarizing event in *C. elegans*, Wnt signaling and LEF/TCF function seem to have strictly antagonistic roles [33,34••] (reviewed in [63]). This model is not sufficient, however, to explain all the details of β -catenin/Armadillo-dependent transcriptional activation in *Drosophila* and vertebrates. Studies in yeast [23] and mammalian cells [22•] have shown that chimeras of a carboxy-terminal region of β -catenin/Armadillo with heterologous DNA-binding domains are sufficient to activate transcription from synthetic minimal promoters. Taken together, these studies suggest that association of β -catenin/Armadillo with LEF/TCF both relieves transcriptional repression by LEF/TCF and activates transcription by the carboxy-terminal β -catenin/Armadillo domain. It is important to note that both promoters studied to date required other factors for maximal expression [23,24•].

Conclusions

Our understanding of the positive and negative regulation of Wnt signaling has broadened tremendously in the past 18 months. Wnt receptors (the Frizzled family), secreted antagonists (Frzb proteins and Cerberus), kinases (CKII, PKC), proteins of less clearly understood function (APC, Axin), and transcription factors (LEF/TCF) have been identified, and the biochemical relationships among these components are being elucidated. There are, no doubt, essential components of these pathways that have yet to

be identified, and the relationship between the 'classical' signal transduction pathway (Figure 1) and divergent Wnt and Frizzled signaling pathways will continue to be clarified. Finally, systematic studies must be undertaken to determine which members of the Wnt and Frizzled families can interact in a productive manner and whether members of the other multi-gene families involved in the transduction of Wnt signals are generally interchangeable.

Note added in proof

An additional secreted antagonist of Wnt signaling has recently been cloned, and found to be expressed in the Spemann Organizer. Its mechanism of action remains unclear [64].

Acknowledgements

We apologize to the many authors whose work was omitted because of space constraints. We would like to thank Minh Sum for assistance in the preparation of Figure 1 and for formatting of this review. We also thank Frank P. Luyten and Luc Leyns for information concerning Frzb genes. JDB is supported by a Public Health Service grant, and RTM is an Investigator of the Howard Hughes Medical Institute.

References and recommended reading

- Papers of particular interest, published within the annual period of review, have been highlighted as:
- of special interest
 - of outstanding interest
1. Miller JR, Moon RT: **Signal transduction through β -catenin and specification of cell fate during embryogenesis.** *Genes Dev* 1996, **10**:2527-2539.
 2. Torres MA, Yang-Snyder JA, Purcell SM, DeMarais AA, McGrew LL, Moon RT: **Activities of the Wnt-1 class of secreted signaling factors are antagonized by the Wnt-5a class and by a dominant negative cadherin in early *Xenopus* development.** *J Cell Biol* 1996, **133**:1123-1137.
 3. Glinka A, Wu W, Onichtchouk D, Blumenstock C, Niehrs C: **Head induction by simultaneous repression of BMP and Wnt signalling in *Xenopus*.** *Nature* 1997, **389**:517-519.
 - Cerberus, a novel secreted protein expressed in the dorsal gastrula (Spemann's) organizer in *Xenopus* blocks Wnt-8 function. This study extends recent findings that Spemann's organizer is the source of multiple secreted antagonists of Wnt and BMP signaling.
 4. Moon RT, Brown JD, Yang-Snyder JA, Miller JR: **Structurally related receptors and antagonists compete for secreted Wnt ligands.** *Cell* 1997, **88**:725-728.
 5. Szuts D, Freeman M, Bienz M: **Antagonism between EGFR and Wingless signaling in the larval cuticle of *Drosophila*.** *Development* 1997, **124**:3209-3219.
 6. Treisman JE, Luk A, Rubin GM, Heberlein U: ***eyelid* antagonizes wingless signaling during *Drosophila* development and has homology to the Bright family of DNA-binding proteins.** *Genes Dev* 1997, **11**:1949-1962.
 7. Yost C, Torres M, Miller JR, Huang E, Kimelman D, Moon RT: **The axis-inducing activity, stability and subcellular distribution of β -catenin is regulated in *Xenopus* embryos by glycogen synthase kinase 3.** *Genes Dev* 1996, **10**:1443-1454.
 8. Rubinfeld B, Souza B, Albert I, Porfiri E, Fiol C, Munemitsu S, Polakis P: **Binding of GSK-3 β to the APC- β -catenin complex and regulation of complex assembly.** *Science* 1996, **272**:1023-1026.
 9. Aberle H, Bauer A, Stappert J, Kispert A, Kemler R: **β -catenin is a target for the ubiquitin-proteasome pathway.** *EMBO J* 1997, **16**:3797-3804.
 10. Parr BA, McMahon AP: **Wnt genes and vertebrate development.** *Curr Opin Genet Dev* 1994, **4**:523-528.
 11. Wang Y, Macke JP, Abella BS, Andreasson K, Worley P, Gilbert DJ, Copeland NG, Jenkins NA, Nathans J: **A large family**

- of putative transmembrane receptors homologous to the product of the *Drosophila* tissue polarity gene *frizzled*. *J Biol Chem* 1996, **271**:4468-4476.
12. Bhanot P, Brink M, Samos CH, Hsieh J-C, Wang Y, Macke JP, Andrew D, Nathans J, Nusse R: **A new member of the frizzled family from *Drosophila* functions as a Wingless receptor.** *Nature* 1996, **382**:225-230.
 13. Yang-Snyder J, Miller JR, Brown JD, Lai CJ, Moon RT: **A frizzled homolog functions in a vertebrate Wnt signaling pathway.** *Curr Biol* 1996, **6**:1302-1306.
 14. He X, Saint-Jeannet J-P, Wang Y, Nathans J, Dawid I, Varmus H: **A member of the frizzled protein family mediating axis induction by Wnt-5a.** *Science* 1997, **275**:1652-1654.
- Frizzled 5 is capable of transducing 'Wnt-1 like' signals in cooperation with Wnt-5a. This suggests that distinct Wnt signaling pathways can be activated by means of the same Wnt interacting with different members of the Frizzled family (see [51]).
15. Klingensmith J, Nusse R, Perrimon N: **The *Drosophila* segment polarity gene *dishevelled* encodes a novel protein required for response to the wingless signal.** *Genes Dev* 1994, **8**:118-130.
 16. Willert K, Brink M, Wodarz A, Varmus H, Nusse R: **Casein kinase 2 associates with and phosphorylates Dishevelled.** *EMBO J* 1997, **16**:3089-3096.
- Overexpression of Dishevelled in an imaginal disc cell line results in hyperphosphorylation of Dishevelled and accumulation of Armadillo. A kinase associated with immunoprecipitated Dishevelled was purified, microsequenced and was determined to be CKII (casein kinase II). Although CKII seems to phosphorylate Dishevelled upon overexpression of Frizzled-2 in the absence of Wingless, this phosphorylation is not sufficient to result in accumulation of Armadillo.
17. Yanagawa S, van Leeuwen F, Wodarz A, Klingensmith J, Nusse R: **The dishevelled protein is modified by wingless signaling in *Drosophila*.** *Genes Dev* 1995, **9**:1087-1097.
 18. Cook D, Fry MJ, Hughes K, Sumathipala R, Woodgett JR, Dale TC: **Wingless inactivates glycogen synthase kinase-3 via an intracellular signalling pathway which involves a protein kinase C.** *EMBO J* 1996, **15**:4526-4536.
 19. Molenaar M, van de Wetering M, Oosterwegel M, Peterson-Maduro J, Godsave S, Korinek V, Roose J, Destree O, Clevers H: **XTcf-3 transcription factor mediates β -catenin-induced axis formation in *Xenopus* embryos.** *Cell* 1996, **86**:391-399.
 20. Behrens J, von Kries JP, Kuhl M, Bruhn L, Wedlich D, Grosschedl R, Birchmeier W: **Functional interaction of β -catenin with the transcription factor LEF-1.** *Nature* 1996, **382**:638-642.
 21. Huber AH, Nelson WJ, Weis WI: **Three-dimensional structure of the armadillo repeat region of β -catenin.** *Cell* 1997, **90**:871-882.
- The structure of the arm repeat region of β -catenin suggests that the protein contains 12, not 13, repeats. The structure also reveals a basic groove in the repeat region that is likely responsible for interaction with acidic regions of LEF/TCF, APC and cadherin.
22. van de Wetering M, Cavallo R, Dooijes D, van Beest M, van Es J, Loureiro J, Ypma A, Hursh D, Jones T, Bejsovec A *et al.*: **Armadillo coactivates transcription driven by the product of the *Drosophila* segment polarity gene *dTCF*.** *Cell* 1997, **88**:789-799.
- A *Drosophila* TCF homolog was cloned and shown to be necessary for transduction of Armadillo signaling. A carboxy-terminal domain of Armadillo and β -catenin is shown to act as a transcriptional activator when bound to a minimal promoter. Interestingly, the authors identify a mutant of Armadillo that cannot associate with dTCF in two-hybrid assays but which still enters the nucleus, suggesting that this translocation occurs independently of association with dTCF.
23. Riese J, Yu X, Munsterlyn A, Eresh S, Hsu S-C, Grosschedl R, Bienz M: **LEF-1, a nuclear factor coordinating signaling inputs from wingless and decapentaplegic.** *Cell* 1997, **88**:777-787.
 24. Brannon M, Gomperts M, Sumoy L, Moon RT, Kimelman D: **A β -catenin-XTcf-3 complex binds to the *siamois* promoter to regulate dorsal axis specification in *Xenopus*.** *Genes Dev* 1997, **11**:2359-2370.
- β -catenin responsiveness of a 0.8 kb fragment of the *siamois* promoter requires consensus LEF/TCF sites. Mutation of those sites results in loss of β -catenin responsiveness and an increase in basal transcription, suggesting that XTcf-3 may be bound to the promoter and repress transcription in the absence of β -catenin.
25. Cavallo R, Rubenstein D, Peifer M: **Armadillo and dTCF: a marriage made in the nucleus.** *Curr Opin Genet Dev* 1997, **7**:459-466.
 26. Nusse R: **A versatile transcriptional effector of wingless signaling.** *Cell* 1997, **89**:321-323.
 27. Orsulic S, Peifer M: **Cell-cell signalling: wingless lands at last.** *Curr Biol* 1996, **6**:1363-1367.
 28. Moon RT, Brown JD, Torres M: **Wnts modulate cell fate and behavior during vertebrate development.** *Trends Genet* 1997, **13**:157-162.
 29. Slusarski DC, Yang-Snyder J, Busa WB, Moon RT: **Modulation of embryonic intracellular Ca^{2+} signaling by Wnt-5a.** *Dev Biol* 1997, **182**:114-120.
- Wnt-5a, but not Wnt-8, can stimulate intracellular Ca^{2+} signaling in zebrafish embryos. This is the first physiological readout for 'Wnt-5a class' responses in vertebrate embryos. Activation of Ca^{2+} signaling by Wnt-5a can be mimicked by stimulation of the PI cycle by activation of a serotonin receptor.
30. Slusarski DC, Corcos VG, Moon RT: **Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling.** *Nature* 1997, **390**:410-413.
- Rat Frizzled-2, but not rat Frizzled-1, can stimulate intracellular Ca^{2+} signaling by G-protein-linked PI signaling and can synergize with Wnt-5a. This study confirms earlier suggestions that some Wnts may activate the PI cycle and places heterotrimeric G-proteins in this pathway.
31. Strutt DJ, Weber U, Mlodzik M: **The role of RhoA in tissue polarity and frizzled signalling.** *Nature* 1997, **387**:292-295.
 32. Adler PA: **The genetic control of tissue polarity in *Drosophila*.** *Bioessays* 1992, **14**:735-741.
 33. Thorpe CJ, Schlesinger A, Carter JC, Bowerman B: **Wnt signaling polarizes an early *C. elegans* blastomere to distinguish endoderm from mesoderm.** *Cell* 1997, **90**:695-705.
 34. Rocheleau CE, Downs WD, Lin R, Wittmann C, Bei Y, Cha Y-H, Ali M, Priess JR, Mello CC: **Wnt signaling and an APC-related gene specify endoderm in early *C. elegans* embryos.** *Cell* 1997, **90**:707-716.
- Several components of the Wnt signal transduction pathway were recovered in screens for excess muscle production in *C. elegans*. Other components were identified by homology with known *Drosophila* and vertebrate genes and were tested by 'RNA-mediated interference', a poorly understood but extremely powerful technique for specifically blocking gene function in *C. elegans*. The elegant combination of genetics, genomics and molecular biology available in this system are demonstrated. In contrast to other systems studied, the endogenous LEF/TCF homolog and Wnt signaling seem to have strictly antagonistic roles.
35. Thomsen GH: **Antagonism within and around the organizer: BMP inhibitors in vertebrate body patterning.** *Trends Genet* 1997, **13**:209-211.
 36. Hoang B, Moos M, Vukicevic S, Luyten FP: **Primary structure and tissue distribution of FRZB, a novel protein related to *Drosophila* frizzled, suggest a role in skeletal morphogenesis.** *J Biol Chem* 1996, **271**:26131-26137.
 37. Lin K, Wang S, Julius MA, Kitajewski J, Moos M, Luyten FP: **The cysteine-rich frizzled domain of frzb-1 is required and sufficient for modulation of Wnt signaling.** *Proc Natl Acad Sci USA* 1997, **94**:11196-11200.
- Deletion mutants of Frzb-1 were used to define the Wnt-binding domain of this antagonist. Although deletion of the domain homologous to Frizzled abrogates interaction with Wnt-1 and the ability of Frzb-1 to block Wnt-1 signaling, smaller deletions in that domain do not disrupt binding to Wnt-1 but do disrupt Wnt-1 antagonism. This study suggests that biochemical interactions of Frzb and Wnt do not necessarily imply inhibition of Wnt function.
38. Leyns L, Bouwmeester T, Kim S-H, Piccolo S, DeRobertis EM: **Frzb-1 is a secreted antagonist of Wnt signaling expressed in the Spemann organizer.** *Cell* 1997, **88**:747-756.
- Frzb-1, a secreted antagonist of Wnt signaling can bind Wnt-1 and block embryonic responses to Wnt-1 and Wnt-8 signaling. A domain of Frzb-1 is closely related to the extracellular ligand-binding domain of Frizzled receptors.
39. Shirozu M, Tada H, Tashiro K, Nakamura T, Lopez ND, Nazarea M, Hamada T, Sato T, Nakano T, Honjo T: **Characterization of novel secreted and membrane proteins isolated by the signal sequence trap method.** *Genomics* 1996, **37**:273-280.
 40. Pfeffer PL, DeRobertis EM, Izpisua-Belmonte JC: **Crescent, a novel chick gene encoding a Frizzled-like cysteine-rich domain, is expressed in anterior regions during early embryogenesis.** *Int J Dev Biol* 1997, **41**:449-458.
 41. Finch PW, He X, Kelley MJ, Uren A, Schaudies RP, Popescu NC, Rudikoff S, Aaronson SA, Varmus HE, Rubin JS: **Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action.** *Proc Natl Acad Sci USA* 1997, **94**:6770-6775.

FRP, a member of the Frzb family, was isolated and shown to inhibit the activity of Wnt-1, Wnt-8 and Wingless, but not Wnt-3a. This result demonstrates that Frzbs can discriminate between Wnts that have similar activities.

42. Mayr T, Deutsch U, Kuhl M, Drexler HCA, Lottspeich F, Deutzmann R, Wedlich D, Risau W: **Fritz: a secreted frizzled-related protein that inhibits Wnt activity.** *Mech Dev* 1997, **63**:109-125.
 43. Salic AN, Kroll KL, Evans LM, Kirschner MW: **Sizzled: a secreted Xwnt8 antagonist expressed in the ventral marginal zone of *Xenopus* embryos.** *Development* 1997, **124**:4739-4748.
 44. Rattner A, Hsieh J-C, Smallwood PM, Gilbert DJ, Copeland NG, Jenkins NA, Nathans J: **A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors.** *Proc Natl Acad Sci USA* 1997, **94**:2859-2863.
- Four *sFRP* genes, members of the Frzb family, are cloned from mouse, revealing that these secreted Wnt antagonists comprise a multi-gene family.
45. Wang S, Krinks M, Lin K, Luyten FP, Moos M: **Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8.** *Cell* 1997, **88**:757-766.
- Xenopus* Frzb-1 was cloned and shown to inhibit Wnt signaling and interact with Wnt proteins by coimmunoprecipitation.
46. Wong GT, Gavin BJ, McMahon AP: **Differential transformation of mammary epithelial cells by Wnt genes.** *Mol Cell Biol* 1994, **14**:6278-6286.
 47. Du SJ, Purcell SM, Christian JL, McGrew LL, Moon RT: **Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos.** *Mol Cell Biol* 1995, **15**:2625-2634.
 48. Wang S, Krinks M, Moos M: **Frzb-1, an antagonist of Wnt-1 and Wnt-8, does not block signaling by Wnts-3a, -5a, or -11.** *Biochem Biophys Res Commun* 1997, **236**:502-504.
- Frzb-1 does not antagonize 'Wnt-5a class' function and discriminates between different Wnts in the 'Wnt-1 class'. Signaling by Wnt-1 and Wnt-8 are blocked, but Wnt-3a activity is not significantly inhibited.
49. Bouwmeester T, Kim S-H, Sasai Y, Lu B, DeRobertis EM: **Cerberus is a head-inducing secreted factor expressed in the anterior endoderm of Spemann's organizer.** *Nature* 1996, **382**:595-601.
 50. Christian J, Moon RT: **Interactions between Xwnt-8 and Spemann organizer signaling pathways generate dorsoventral pattern in the embryonic mesoderm of *Xenopus*.** *Genes Dev* 1993, **7**:13-28.
 51. Larabell CA, Torres M, Rowning BA, Yost C, Miller JR, Wu M, Kimelman D, Moon RT: **Establishment of the dorso-ventral axis in *Xenopus* embryos is presaged by early asymmetries in β -catenin that are modulated by the Wnt signaling pathway.** *J Cell Biol* 1997, **136**:1123-1136.
- Before the onset of zygotic transcription in *Xenopus* embryos, Wnt-5a can block embryonic responses to Wnt-8 (see [2]). After the onset of zygotic transcription, overexpression of Wnt-5a results in the stabilization of β -catenin, suggesting that distinct pathways can be activated by a single Wnt ligand.
52. Miller JR, Moon RT: **Analysis of the signaling activities of localization mutants of β -catenin during axis specification in *Xenopus*.** *J Cell Biol* 1997, **139**:229-243.

Overexpression of a transmembrane-tethered form of β -catenin results in activation of the Wnt pathway in *Xenopus* embryos. This activity is probably the result of overloading the β -catenin degradation pathway which allows the stabilization of the endogenous free β -catenin. This study helps to clarify the mechanism of signaling activity of β -catenin which cannot itself translocate to the nucleus.

53. Pai LM, Orsulic S, Bejsovec A, Peifer M: **Negative regulation of Armadillo, a Wingless effector in *Drosophila*.** *Development* 1997, **124**:2255-2266.
 54. Hayashi S, Rubinfeld B, Souza B, Polakis P, Wieschaus E, Levine AJ: **A *Drosophila* homolog of the tumor suppressor gene *adenomatous polyposis coli* down-regulates β -catenin but its zygotic expression is not essential for the regulation of Armadillo.** *Proc Natl Acad Sci USA* 1997, **94**:242-247.
 55. Moon RT, Miller JR: **The APC tumor suppressor protein in development and cancer.** *Trends Genet* 1997, **13**:256-258.
 56. Zeng L, Fagotto F, Zhang T, Hsu W, Vasicek TJ, Perry WL, Lee JJ, Tilghman SM, Gumbiner BM, Constantini F: **The mouse *Fused* locus encodes axin, an inhibitor of the Wnt signaling pathway that regulates embryonic axis formation.** *Cell* 1997, **90**:181-192.
- Axin seems to be a negative regulator of Wnt signaling, and deletion of an RGS-like domain results in a 'dominant negative' Axin that activates Wnt signaling. Although the mechanism of Axin action is unknown, presence of the RGS domain may suggest the involvement of heterotrimeric G-proteins.
57. Carnac G, Kodjabachian L, Gurdon JB, Lemaire P: **The homeobox gene *siamois* is a target of the Wnt dorsalisation pathway and triggers organiser activity in the absence of mesoderm.** *Development* 1996, **122**:3055-3065.
 58. Brannon M, Kimelman D: **Activation of *siamois* by the Wnt pathway.** *Dev Biol* 1996, **180**:344-347.
 59. Koelle MR: **A new family of G-protein regulators-the RGS proteins.** *Curr Opin Cell Biol* 1997, **9**:143-147.
 60. Huber O, Korn R, McLaughlin J, Ohsugi M, Germann BG, Kemler R: **Nuclear localization of β -catenin by interaction with transcription factor LEF-1.** *Mech Dev* 1996, **59**:3-10.
 61. Brunner E, Peter O, Schweizer L, Basler K: ***pangolin* encodes a LEF-1 homologue that acts downstream of Armadillo to transduce the Wingless signal in *Drosophila*.** *Nature* 1997, **385**:829-833.
- A screen for suppressors of the rough-eye phenotype caused by ectopic Wingless expression resulted in the isolation of mutants in *armadillo* and *pangolin*, a LEF/TCF homologue. Two missense mutations in *pangolin* map to the amino terminus of the protein—the region known to interact with Armadillo. Loss-of-function alleles of *pangolin* result in segment polarity phenotypes which are insensitive to Armadillo overexpression and suggest that Pangolin is absolutely required for Wingless signaling.
62. Klymkowsky MW: **Minireviews, minidogmas and mythinformation.** *BioEssays* 1997, **19**:537-539.
 63. Han M: **Gut reaction to Wnt signaling in worms.** *Cell* 1997, **90**:581-584.
 64. Gilinka A, Wu W, Delius H, Monaghan AP, Blumenstock C, Niehrs C: **Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction.** *Nature* 1998, **391**:357-362.