

Assignment 1 – Discrete Models – Profir Alexandru – ICA 246

– 2

1) Find the solution for the following difference equations:

$$(a) x_{n+1} = \left(\frac{n+1}{n+2} \right)^2 \cdot x_n + \frac{1}{n+2}, x_0 = 1;$$

$$(b) x_{n+3} - 4 \cdot x_{n+2} + x_{n+1} + 6 \cdot x_n = 60 \cdot 4^n, x_0 = 2, x_1 = 12, x_2 = 12;$$

$$(c) x_{n+1} = \frac{2 \cdot x_n}{1 + 4 \cdot x_n}, x_0 = 1 \left(\text{Hint: use substitution } x_n = \frac{1}{y_n} \right);$$

> restart:

```
eq := x(n+1) = ((n+1)/(n+2))^2 * x(n) + 1/(n+2);
ans := factor(simplify(rsolve({eq, x(0)=1}, x(n))));
```

$$eq := x(n+1) = \frac{(n+1)^2 x(n)}{(n+2)^2} + \frac{1}{n+2}$$

$$ans := \frac{n+2}{2(n+1)}$$

(1)

> restart:

```
eq := x(n+3) - 4 * x(n+2) + x(n+1) + 6 * x(n) = 60 * 4^n;
ans := factor(simplify(rsolve({eq, x(0)=2, x(1)=12, x(2)=12}, x(n))));
```

$$eq := x(n+3) - 4x(n+2) + x(n+1) + 6x(n) = 60 \cdot 4^n$$

$$ans := -16 \cdot 3^n + 16 \cdot 2^n - 4 \cdot (-1)^n + 6 \cdot 4^n$$

(2)

> restart:

```
eq := x(n+1) = (2*x(n)) / (1 + 4 * x(n));
```

```
# Subbing x(n) with 1/y(n)
```

```
x0 := 1; y0 := 1/x0;
```

```
eq_y := simplify(subs(x(n)=1/y(n), x(n+1)=1/y(n+1), eq));
```

```
eq_y := y(n+1) = solve(eq_y, y(n+1));
```

```
ans_y := rsolve({eq_y, y(0)=y0}, y(n));
```

```
ans_x := factor(simplify(1/ans_y));
```

$$eq := x(n+1) = \frac{2x(n)}{1+4x(n)}$$

$$x0 := 1$$

$$y0 := 1$$

$$eq_y := \frac{1}{y(n+1)} = \frac{2}{y(n)+4}$$

$$eq_y := y(n+1) = \frac{y(n)}{2} + 2$$

$$ans_y := -3 \left(\frac{1}{2} \right)^n + 4$$

$$ans_x := -\frac{1}{3 \cdot 2^{-n} - 4}$$

(3)

2) Let us consider the difference equation:

$$x_{n+1} = \frac{x_n^2 + 7}{2 \cdot x_n}$$

(a) Find the equilibrium points and study their stability.

(b) Do some numerical simulations.

(a) The equilibrium points can be found using the 'solve' utility of Maple. However, given the non-linearity of the difference equation, the stability of the equilibrium points can be studied using the 'Stability in the first approximation' Theorem, based on 'Taylor Series Expansion'

$$f(x) = f(a^*) + f'(a^*) \cdot (x - a^*) + f''(a^*) \cdot \left(\frac{(x - a^*)^2}{2} \right) + \dots$$

Given the existence of an equilibrium point $a^* = f(a^*)$, and $f'(a^*)$ exists, then:

(i) If $|f'(a^*)| < 1 \Rightarrow a^*$ is locally asymptotically stable

(ii) If $|f'(a^*)| > 1 \Rightarrow a^*$ is unstable

Given the theory support above, and $x_1^* = \sqrt{7}$, $x_2^* = -\sqrt{7}$, and $f(x) = \frac{x^2 - 7}{2x^2}$, both

derivate values in the equilibrium points are < 1 in absolute value, thus both equilibrium points are locally asymptotically stable.

(b) Numerical simulations are done below.

```
> restart;
with(plots):

f := x -> (x^2+7) / (2 * x);

xstar1, xstar2 := solve(x=f(x), x);
evalf(xstar1); evalf(xstar2);

fdx := simplify(D(f)(x));
abs_fdx_xstar1 := abs(subs(x=xstar1, fdx));
abs_fdx_xstar2 := abs(subs(x=xstar2, fdx));
```

```

n := 100:
xxpos[1] := 10:
xxneg[1] := -10:

for i from 1 to n - 1 do
    xxpos[i+1] := evalf(f(xxpos[i]));
    xxneg[i+1] := evalf(f(xxneg[i]));
end do:

ppos := pointplot([seq([k, xxpos[k]], k=1..n)],
    symbol=solidcircle, color=blue):

pneg := pointplot([seq([k, xxneg[k]], k=1..n)],
    symbol=solidcircle, color=red):

display(ppos, pneg, view=[0..n, -10..10]);

```

$$f := x \mapsto \frac{x^2 + 7}{2x}$$

$$xstar1, xstar2 := \sqrt{7}, -\sqrt{7}$$


$$2.645751311$$

$$-2.645751311$$

$$fdx := \frac{x^2 - 7}{2x^2}$$

$$abs_fdx_xstar1 := 0$$

$$abs_fdx_xstar2 := 0$$



3) Let us consider the system of difference equations:

$$\begin{cases} x_{n+1} = x_n - x_n^2 - x_n y_n \\ y_{n+1} = 2y_n - y_n^2 - 3x_n y_n \end{cases}$$

(a) Find the equilibrium points and study their stability.

(b) Do some numerical simulations.

(a) Given the non-linearity of the system, the equilibrium points can be found using the linearized system

resulted from the Jacobian of the system.

$$y_{n+1} = \text{Jacobian}(f)(x^*) y_n$$

Theorem: The following statements hold:

(i) x^* is locally asymptotically stable **if and only if** $|\lambda| < 1$,

for all λ eigenvalues of the Jacobian(f) (x^*);

(ii) x^* is unstable if there exists an eigenvalue λ of the Jacobian(f) (x^*) such that $|\lambda| > 1$.

Equilibrium points

- $x = \frac{1}{2}, y = -\frac{1}{2}; |\lambda_{1,2}| > 1 \Rightarrow \text{unstable};$

- $x = 0, y = 0; |\lambda_2 = 2| > 1 \Rightarrow \text{unstable};$

- $x = 0, y = 1; |\lambda_{1,2}| < 1 \Rightarrow \text{locally asymptotically stable};$

(b) Numerical simulations are done below.

```
> restart:
with(plots): with(linalg):

f1 := (x, y) -> x - x^2 - x*y;
f2 := (x, y) -> 2*y - y^2 - 3*x*y;

ans_sys := solve({f1(x,y)=x, f2(x,y)=y}, {x, y});
J := jacobian([f1(x,y), f2(x,y)], [x,y]);

# 1st solution: {x=1/2, y=-1/2}
A1 := subs(x=1/2, y=-1/2, evalm(J));
lambda1, lambda2 := evalf(eigenvals(A1)):
abs(lambda1); abs(lambda2);

# 2nd solution: {x=0, y=0}
A2 := subs(x=0, y=0, evalm(J));
lambda1, lambda2 := evalf(eigenvals(A2)):
abs(lambda1); abs(lambda2);

# 3rd solution: {x=0, y=1}
A3 := subs(x=0, y=1, evalm(J));
lambda1, lambda2 := evalf(eigenvals(A3)):
abs(lambda1); abs(lambda2);

# simulation
n := 100:
xx[1] := 0.1:
yy[1] := 0.1:

for i from 1 to n-1 do
    xx[i+1] := evalf(f1(xx[i], yy[i]));
    yy[i+1] := evalf(f2(xx[i], yy[i]));
end do:

pts_x := [seq([k, xx[k]], k=1..n)]:
pts_y := [seq([k, yy[k]], k=1..n)]:

p1 := pointplot(pts_x, symbol=circle, color=red):
p2 := pointplot(pts_y, symbol=cross, color=blue):

display(p1, p2, title="x (red), y (blue) vs iteration",
        view=[0..n, -1..2]);
```

$$f1 := (x,y) \mapsto x - x^2 - x\,y$$

$$f2 := (x,y) \mapsto 2\,y - y^2 - 3\,x\,y$$

$$ans_sys := \left\{x = \frac{1}{2}, y = -\frac{1}{2}\right\}, \{x=0, y=0\}, \{x=0, y=1\}$$

$$J := \begin{bmatrix} -2\,x - y + 1 & -x \\ -3\,y & -3\,x - 2\,y + 2 \end{bmatrix}$$

$$A1 := \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{3}{2} & \frac{3}{2} \end{bmatrix}$$


1.224744871
1.224744871

$$A2 := \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

1.
2.

$$A3 := \begin{bmatrix} 0 & 0 \\ -3 & 0 \end{bmatrix}$$

0.
0.



4) Let's consider the simple interest and compound interest models:

$$S_{n+1} = S_n + pS_0$$

$$S_{n+1} = S_n + \frac{p}{r} S_0$$

Company A offers simple interest at an annual rate of 4%. Company B offers compound interest at an annual rate of 3% with a conversion period of one month.

(a) Calculate for the both cases the amount on deposit after 5, 10, 15, and 20 years for principal $S_0 = 1000$.

The results are displayed in the following format: [years, amount_on_deposit, rate_conversion]

- Company A: [5, 1200.00, 1.20], [10, 1400.00, 1.40], [15, 1600.00, 1.60], [20, 1800.00, 1.80];

- Company B: [5, 1161.61, 1.16], [10, 1349.35, 1.34], [15, 1567.43, 1.56], [20, 1820.75, 1.82];

(b) Determine which interest offer maximizes the amount on deposit after 5, 10, 15, and 20 years.

- Company A maximizes the amount on deposit after 5, 10, and 15 years;
- Company B maximizes the amount on deposit after 20 years;

The offer of company A is profitable for acceptable subscription periods, while company B is profitable (and not by a great margin) for longer periods.

The plots provided below confirm the presented conclusion.

```
> restart:
with(plots):

S0 := 1000;
years := [5, 10, 15, 20];

SIM := S(n+1) = S(n) + p * S0;
CIM := S(n+1) = S(n) * (1 + (p/r));

SIM_ans:= unapply(factor(simplify(rsolve({SIM, S(0)=S0}, S(n)))),
p, n);
CIM_ans := unapply(factor(simplify(rsolve({CIM, S(0)=S0}, S(n)))
), p, n, r):
CIM_ans_years := (p, n, r) -> CIM_ans(p, r*n, r);

SIM_A := [seq([y, SIM_ans(0.04, y), SIM_ans(0.04, y) / S0], y in
years)];
CIM_B := [seq([y, CIM_ans_years(0.03, y, 12), CIM_ans_years(0.03,
y, 12) / S0], y in years)];

plot([
[seq([SIM_A[i][1], SIM_A[i][2]], i=1..nops(years))],
[seq([CIM_B[i][1], CIM_B[i][2]], i=1..nops(years))],
],
style = pointline, color = [red, blue, green],
legend = ["Simple 4%", "Compound 3% monthly"],
title = "Comparison of interest growth",
labels = ["Years", "Amount ($)"]);

plot([
[seq([SIM_A[i][1], SIM_A[i][3]], i=1..nops(years))],
[seq([CIM_B[i][1], CIM_B[i][3]], i=1..nops(years))],
],
style = pointline, color = [red, blue],
legend = ["Simple 4%", "Compound 3% monthly"],
title = "Comparison of rate conversion",
labels = ["Years", "Amount (%)"]);
```

$$S_0 := 1000$$

$$\text{years} := [5, 10, 15, 20]$$

$$SIM := S(n+1) = S(n) + 1000 p$$


$$CIM := S(n+1) = S(n) \left(1 + \frac{p}{r} \right)$$

$$SIM_ans := (p, n) \mapsto 1000 n p + 1000$$

$CIM_ans_years := (p, n, r) \mapsto CIM_ans(p, r, n, r)$

$SIM_A := [[5, 1200.00, 1.200000000], [10, 1400.00, 1.400000000], [15, 1600.00, 1.600000000], [20, 1800.00, 1.800000000]]$

$CIM_B := [[5, 1161.616782, 1.161616782], [10, 1349.353547, 1.349353547], [15, 1567.431725, 1.567431725], [20, 1820.754995, 1.820754995]]$



Comparison of rate conversion

