Contents

44	Anin	nal Sensory Systems	3
	44.1	How Do Sensory Organs Convey Information to the Brain?	3
	44.2	Mechanoreception: Sensing Pressure Changes	3
	44.3	Photoreception: Sensing Light	4
	44.4	Chemoreception: Sensing Chemicals	5
	44.5	Other Sensory Systems	6
45	Anin	nal Movement	7
	45.1	How Do Muscles Contract?	-/
	45.2	Muscle Tissues	-/
46	Cher	mical Signals in Animals	10
	46.1	Cell-to-Cell Signaling	10
	46.2	How do Hormones Act on Target Cells?	10
	46.3	What do Hormones Do?	11
	46.4	How is the Production of Hormones Regulated?	11
49	An Ir	ntroduction to Ecology	12
49		ntroduction to Ecology The Levels of ecological study	
49	49.1		12
49	49.1 49.2	The Levels of ecological study	12 12
49	49.1 49.2 49.3	The Levels of ecological study	12 12 12
49	49.1 49.2 49.3 49.4	The Levels of ecological study	12 12 12
49	49.1 49.2 49.3 49.4 49.5	The Levels of ecological study	12 12 13 13
	49.1 49.2 49.3 49.4 49.5 49.6	The Levels of ecological study	12 12 13 13
	49.1 49.2 49.3 49.4 49.5 49.6	The Levels of ecological study	12 12 13 13 14
	49.1 49.2 49.3 49.4 49.5 49.6	The Levels of ecological study	12 12 13 13 13
	49.1 49.2 49.3 49.4 49.5 49.6	The Levels of ecological study	12 12 13 13 14 14
	49.1 49.2 49.3 49.4 49.5 49.6	The Levels of ecological study	12 12 13 13 14 14 14
	49.1 49.2 49.3 49.4 49.5 49.6 Ecos 53.1	The Levels of ecological study What Determines the Distribution and Abundance of Organisms? Abiotic Factors Global Climate Patterns Terrestrial Biomes Aquatic Biomes Systems and Global Ecology How Does Energy Flow through Ecosystems 53.1.1 What Happens to the Biomass of Autorophs? 53.1.2 Energy Transfer between Trophic Levels	12 12 13 13 14 14 14 15
	49.1 49.2 49.3 49.4 49.5 49.6 Ecos 53.1	The Levels of ecological study What Determines the Distribution and Abundance of Organisms? Abiotic Factors Global Climate Patterns Terrestrial Biomes Aquatic Biomes systems and Global Ecology How Does Energy Flow through Ecosystems 53.1.1 What Happens to the Biomass of Autorophs? 53.1.2 Energy Transfer between Trophic Levels 53.1.3 Global Patterns in Productivity	12 12 13 13 14 14 14 14 15 16

54	Biodiversity and Conservation Biology	17
	54.1 What is Biodiversity?	. 17
	54.1.1 Biodiversity Can Be Measured and Analyzed at Sevearl Level	s 17
	54.1.2 Where is Biodiversity Highest?	. 17
	54.2 Threats to Biodiversity	. 18
	54.2.1 Habitat Destruction	. 18
	54.2.2 Estimating the Probability that a Population or Species Will	
	Go Extinct	. 18
	54.3 Benefits of Biodiversity	. 19
50	Behavior Ecology	20
	50.1 Introduction to Behavioral Ecology	. 20
	50.2 What, When, and How to Eat	. 20
	50.4 Choosing a Place to Live	. 20
51	Population Ecology	21
	51.1 Distribution and Abundance	. 21
	51.2 Demography	. 21
	51.3 Population Growth	. 22
52	Community Ecology	23

44 Animal Sensory Systems

44.1 How Do Sensory Organs Convey Information to the Brain?

- ▶ **Transduction**: conversion of an external stimulus to an internal signal in the form of action potentials along sensory neurons.
 - Requires a sensory receptor:
 - 1. *Mechanoreceptor*: respond to changes in pressure.
 - 2. Photoreceptors: respond to particular wavelengths of light.
 - 3. Chemoreceptors: detect specific molecules.
 - 4. Thermoreceptors: respond to changes in temperature.
 - 5. Nociceptors: sense harmful stimuli such as tissue injury.
 - 6. Electroreceptors: detect electric fields.
 - 7. Magnetoreceptors: detect magnetic fields.
 - Frequency of action potential firing rate can indicate the intensity of the stimulus.
- ▶ Transmission: the process of sending the signal to the central nervous system.

44.2 Mechanoreception: Sensing Pressure Changes

- ▶ Statocyst: an fluid filled organ that grabs use to help sense pressure created by gravity.
- ▷ Direct physical pressure causes ion channels to open and close, creating voltage gated action potentials.

> The Mammalian Ear:

 Outer ear: collects incoming pressure waves and funnels them into tube known as the ear canal, which leads to the **tympanic membrane**, or eardrum.

- Middle ear: contains three tiny bones that the eardrum passes
 vibrations to in order to amplify sound. One of the bones, called **stapes**,
 vibrates against a membrane called the **oval window**, which separates
 the middle ear from the inner ear.
- Inner ear: the oval window oscillates in response to vibrations which generates waves in a chamber known as the cochlea. These waves are pressure inputs the hair cells respond to.
- ▶ Hair cells, forming rows that sit in the middle chamber, are embedded in a tissue that sits atop the **basilar membrane**. In addition, the hair cells' stereocilia touch another smaller surface called the **tectorial membrane**.
- ▶ This sandwiching of hair cells produce very specific responses to various frequencies, allowing us to distinguish between them.
- ▶ **Lateral line system**: a mechanoreceptor organ that most fish and larval amphibians use. Consists of embedded gel-like domed structures called cupulae that lay inside canals along the length of the body.

44.3 Photoreception: Sensing Light

▶ The Insect Eye:

- Compound eyes: eyes composed of hundreds of thousands of light-sensing columns called ommatidia.
- Each ommatidium has lens that focuses light into a smaller number of receptor cells—usually four.

> The Vertebrate Eye:

 Simple eye: a structure with a single lens that focuses incoming light onto a layer of many receptor cells.

Structure of the Vertebrate Eye:

- white of the eye: outermost layer of the eye that consists of tough rind of white tissue called the sclera.
- Cornea: a transparent sheet of connective tissue on the front of the

sclera.

- Iris: pigmented, round muscle just inside the cornea that can
 contract or expand to control the amount of light entering the eye.
- Pupil: hole in the center of the iris.
- Lens: works with the cornea to focus incoming light.
- Retina: a layer of photoreceptors and several layers of neurons.
- The photoreceptors are held in place by the pigmented epithelium.
- photoreceptors synapse with an intermediate layer of connecting neurons called **bipolar cells**.
- Bipolar cells synapse with neurons called ganglion cells, which form the innermost layer of the retina.
- The axons of the ganglion cells project to the brian via optic nerve.
- Photoreceptors come in two distinct types:
 - Rods: sensitive to dim light but not to color.
 - Cones: sensitive to different wavelengths of light, but not dim light.
- Opsin: a transparent membrane protein that associates with a molecule of pigment in the retinal.
- **Rhodopsin**: a two molecule complex similar ot opsin in rod cells.

44.4 Chemoreception: Sensing Chemicals

- ▷ Chemoreception occurs when chemicals bind to chemoreceptors, initiating action potentials in sensory neurons.
- ▶ **Gustation**: the sense of taste.
- ▶ Olfaction: sense of smell.
- ▶ Taste buds: clustered structures containing about 100 spindle shaped taste cells that make synapses with sensory neurons.
- ▷ Salty: due to sodium ions dissolved in food.

- ▷ Sourness: due to presence of protons.
- ▶ *Umami*: due to monosodium glutamate.
- Documents: airborne molecules.
- ▶ Olfactory bulb: part of the brain where olfactory signals are processed and interpreted.
- ▶ Pheromone: a chemical that is secreted into the environment that affects the behavior or physiology of animals.

44.5 Other Sensory Systems

- ► Thermoreceptors are located in the central nervous system and also commonly found skin and other outer surfaces of animals.
- ▶ **Ampullae of Lorenzini**: tiny pores scattered across a shark's head contain structures that are responsible for their electroreception. Sharks can detect electrical potentials as small as a nanovolt.
- ▶ Electrogenic fishes have specialized organs near their tails that generate electric fields stronger than those of regular nerves or muscles.
- ▶ Magnetoreception: seen in many organisms, including fungi, invertebrates, and all other vertebrate classes.

45 Animal Movement

45.1 How Do Muscles Contract?

- ▶ Muscle fiber: a muscle fiber is a long, thin muscle cell.
- Within each muscle cell are many threadlike, contractile structures called myofibrils.
- ▶ Myofibrils often look striped or striated due to alternating light and dark units called **sarcomeres**, which repeat along the length of a myofibril.
- ▶ The question of how muscles contract simplifies to how sarcomeres shorten.
- ▶ **Thin filaments** are composed of actin.
- ▶ **Thick filaments** are composed of myosin.
- ▶ **Sliding-filament model**: filaments slide past one another during contraction, with the sarcomere shorting with no change between the lengths of thick and thin filaments themselves.
- ▶ ATP is required for myosin to release from actin once the two molecules have bound to each other.
- ▶ Tropomyosin and Tropnin work together to block the myosin binding sites on actin. When the sites are blocked, the myosin-actin interaction cannot occur, relaxing the muscle.

45.2 Muscle Tissues

> Smooth Muscle:

- Unbranched.
- o Tappered at each end.
- Often organized into thin sheets.
- o Lack sarcomeres, which gave them their smooth appearance.
- o Relatively small and have a single nucleus.

- Essential for the lungs blood vessels, digestive system, urinary bladder, and reproductive system.
- Involuntary

> Cardiac Muscle:

- o Makes up walls of the heart.
- Contain sarcomeres and is satiated.
- Unique branch structure.
- Directly connected end to end via specialized regions called nitercanated discs which are critical to the flow of electrical signals and coordination of the heartbeat.
- Involuntary

> Skeletal Muscle:

- Exceptionally long, unbranched muscle fibers.
- Muitnucleate
- Contains sarcomeres
- Voluntary
- Force depends on:
 - relative proportion of different fiber types.
 - organization of fibers within the muscle.
 - how the muscle is used.

▶ Skeletal Muscle Fiber Types:

- Slow muscle fibers:
 - oxidative.
 - appear red, due to high concentration of myoglobin.
 - myosin hydrolyzes ATP at a slow rate, causing slow contraction.
 - slow to fatigue due to many mitochondria and aerobic respiration.

o Fast muscle fibers:

- glycolytic.
- appear white, due to low myoglobin concentration.
- myosin hydrolyzes ATP at a rapid rate.
- fast to fatigue due to primarily relying on glycolysis.

o Intermediate muscle fibers:

- combination of both fast and slow.

46 Chemical Signals in Animals

46.1 Cell-to-Cell Signaling

- ▶ **Autocrine Signals**: signals that affect the same cell that releases them.
- ▶ Paracrine Signals: diffuse locally and act on target cells near the source cell.
- ▶ **Endocrine Signals**: hormones carried to distinct cells by blood or other body fluids and released by discrete organs called glands.
- ▶ **Neural Signals**, or neurotransmitters, act on other neurons.
- ▶ Neuroincocrine Signals: hormones released by neurons that act on distant cells.
- ▶ Many hormones can act as multiple signal types.
- ▶ All three types of signaling pathway—endocrine, neuron-daocrine, and neuroendocrine-to-endocrine—are regulated by negative feedback, or feedback inhibition.

46.2 How do Hormones Act on Target Cells?

- ▶ There are three chemical classes of hormones:
 - Peptides ane polypeptides.
 - Amino acid derivatives.
 - o Steroids.
- Steroids and thyroid hormones cross plasma membranes much more readily than other types of hormones.
- ▶ Cyclic AMP (cAMP): the key ingredient in the activation of phosphorylase, which catalyzes a reaction that cleaves glucose molecules off glycogen.
- Signal transduction cascade can allow the binding of just a single molecule of epinephrine may trigger the release of millions or even billions of glucose molecules.

▶ Most steroid and thyroid hormones act by inducing a change in gene expression.

46.3 What do Hormones Do?

- ▶ Hormones coordinate the activities of cells in three arenas:
 - o development, growth, and reproduction.
 - o response to environmental challenges.
 - maintenance of homeostasis.
- ▶ A single hormone may affect a wide array of cells and tissues and induce a variety of responses.
- ➤ Xenoestrogens: foreign chemicals that bind to estrogen receptors and induce estrogen like effects.
- ▶ Cortisol triggers the long-term response to stressors by inducing changes that conserve glucose for use by the brain.
- Glucocorticoids make amino acids available for glucose synthesis by promoting the degradation of contractile proteins in muscle. The resulting loss of muscle mass may cause severe weakness.

46.4 How is the Production of Hormones Regulated?

- ▶ In many cases, hormone production is directly or indirectly controlled by the nervous system, where the release of a hormone is regulated by hormones from the anterior pituitary.
- ▶ Hormone-secreting cells in the anterior pituitary are regulated by hormones released by the hypothalamus.
- ▶ Glucocorticoids accomplish feedback inhibition—they suppress their own production.
- ▶ Cushing's disease: the result when feedback inhibition fails.

49 An Introduction to Ecology

49.1 The Levels of ecological study

- ▶ Organisms: study of the morphological, physiological, and behavioral adaptations that allow individuals to live in a particular areal
- ▶ Populations: focus on the number and distribution of individuals in a population and their change over time.
- ▶ Communities: focus on predation, parasitism, and competition, or explore how communities respond to fires, floods, and other disturbances.
- ▷ Ecosystems: study of all the organisms in a particular region along with nonliving components.
- ▶ Biosphere: the sum of all terrestrial and aquatic ecosystems.

49.2 What Determines the Distribution and Abundance of Organisms?

- ▶ For each species, a unique combination of abiotic and biotic factors determines where individuals live and the size of populations
- ▶ Understanding historical events, such as the movement of entire continents, is important to interpreting current patterns of species distributions
- ▷ Abiotic and biotic factors often interact to produce a different effect on species distributions than either type of factor would have on its own.

49.3 Abiotic Factors

- ▶ Air circulation cells such as Hadley cells create bands of wet and dry habitats, and the tilt of Earth's axis causes seasonality in the amount of sunlight that non-equatorial regions receive.
- ➤ The presence of mountains can create local areas of wet or dry habitats, and proximity to an ocean and the direction of ocean currents moderates temperatures in certain terrestrial habitats.

49.4 Global Climate Patterns

▶ The major terrestrial biomes include tropical wet forest, subtropical desert, temperate grassland, temperate forest, boreal forest, and arctic tundra.

49.5 Terrestrial Biomes

49.6 Aquatic Biomes

- ▶ **Intertidal zone**: consists of a rocky shoreline, sandy beach, or mud flat that is exposed to the air at low tide but submerged at high tide.
- ▶ **Neritic zone**: extends from the intertidal zone to depths of about 200 m. Its outermost edge is defined by the end of the continental shelf
- Documental Shelf.
- ▶ **Benthic zone**: bottom of the ocean at all depths.
- ▶ **Photic zone**: intertidal and sunlit regions of the neritic, oceanic, and benthic zones
- > **Aphotic zone**: Areas that do not receive sunlight.

53 Ecosystems and Global Ecology

53.1 How Does Energy Flow through Ecosystems

- ▶ **Primary producer**: a autotroph that allows energy to enter into ecosystems.
- ▶ autotroph: an organism that can synthesize it's own food from inorganic sources.
- ▶ Gross Primary Productivity (GPP): total amount of chemical energy produced in a given area and time period.
- ▶ Net primary productivity (NPP): energy that is invested by primary producers in building new tissue of offspring.
- ▶ NPP = GGP R. R = energy used in cellular respiration or lost.
- ▶ Only 0.8% of solar radiation is captured by plants.
- ▷ It's estimated that 45% of GPP to NPP and the other 55% was used for either cellular respiration or lost to the environment.

53.1.1 What Happens to the Biomass of Autorophs?

- ▶ Primary consumers eat living organisms, secondary eat primary, tertiary eat secondary, and so on.
- ▶ Decomposers: or detritivores, obtain energy from the remains of other organisms or waste products.
- > **Trophic**: the level at which organisms obtain energy from the same source.

53.1.2 Energy Transfer between Trophic Levels

- ▷ All ecosystems share a pattern: the total biomass produced each year declines from lower trophic levels up th the higher levels.
- ightarrow Productivity is measured in g/m²/year
- ▶ Efficiency is a fraction of biomass transferred from one level to the next.
- ▶ Biomass production at each trophic level varies, but generally efficiency is only about 10%.

- ▶ Large mammals are more efficient at producing biomass than small mammals due to less heat loss.
- ▶ Biomass production is more efficient in ectotherms than endotherms, since they primarily rely on heat gained rather than oxidizing sugars.
- ▶ **Biomagnification**: molecules that increase in concentration at higher trophic levels.

53.1.3 Global Patterns in Productivity

- ▶ Productivity is limited by anything that limits the rate of photosynthesis.
- ▶ NPP on land is much higher than it is in oceans.
- ▶ Wet tropics have the highest net productivity.
- Excluding deserts, NPP declines as you move away from the equator due to availability of sunlight and decreasing temperature.
- ▶ Marine productivity is highest along coastlines, due to increased nutrients from rivers and process called upwelling.
- ▶ Tropical wet and dry forests cover less than 5% but account for over 30% of NPP.
- ▶ In aquatic ecosystems, the most productive are algal beds, coral reefs, wetlands, and estuaries.
- ▷ It's estimated that humans are preventing of appropriating 24% of potential NPP.

53.2 How Do Nutrients Cycle through Ecosystems?

53.2.1 Nutrient Cycling within Ecosystems

- ▶ **Biogeochemical cycle**: The path an element takes as it moves from abiotic systems, through biotic, and back.
- ▶ Soil organic matter: the carbon-containing compounds that microscopic decomposers release.
- ▶ **Humus**: result of when detritus becomes completely decayed.

- biogeochemical cycle.
- Decomposition is affected by: abiotic conditions such as oxygen availability, temperature and precipitation; quality of nutrient source fot the fungi, bacteria, and archaea; abundance and diversity of detritivores present.
- ▶ Nutrients can be lost from one ecosystem and exported to another.
- Exported nutrients must be replaced by imports if ecosystems are to function.
- ▶ Mechanisms of nutrient replacement: ions from rocks are released due to weathering; wind or water transport; carbon via photosynthesis; nitrogen via nitrogen-fixing.
- ▶ Vegetation lowers the rate of nutrient export.

53.2.2 Global Biogeochemical Cycles

- ► Human effects on the water cycle: asphalt and concrete limiting water absorption to deep soil; grasslands and forest converted to agricultural fields resulting in lost water in roots, more water off, and less percolates into groundwater; human consumption out pacing recharge rate.
- Consequences of increased nitrogen: algal blooms in aquatic ecosystems, removing oxygen from the area; short term production increase at cost of diversity and long term productivity; acid rain, climate change, depletion of ozone layer.
- ▶ Humans have more than doubled the rate of nitrogen fixation.
- ▶ Humans have quadrupled the amount of phosphorus entering the global biogeochemical system.

54 Biodiversity and Conservation Biology

54.1 What is Biodiversity?

54.1.1 Biodiversity Can Be Measured and Analyzed at Sevearl Levels

- ▶ Genetic diversity: total genetic information contained within all individuals or a population and measured as the number and relative frequencies of all genes in a a group.
- ▶ Species richness: how many species are in an area.
- ▶ Species diversity: both species richness and evenness, of relative abundance of each species.
- ▶ DNA barcoding: use of well characterized gene sequences to distinguish species.
- ► Ecosystem diversity: a measure of horizontal diversity by trophic species richness and vertical diversity by number of trophic levels, plus complex factors with the nonliving environment.
- ▶ **Ecosystem function**: the sum of biological and chemical processes that are characterietic of a given ecosystem.
- ▶ Primary production, nitrogen cycling, decomposition, and carbon storage emerge from the sum of the feeding, growing, moving, respiring, excreting, and decomposing processes.

54.1.2 Where is Biodiversity Highest?

- ▶ Paterns of biodiversity are usually measured by divding geographic area into grids of cells and mapping the biodiversity within each cell.
- ▶ Endemic species: species found in a particular area and no where else.
- ▶ Areas with geographical variation tend to have higher biodiversity.
- ▶ Tropical rainforests are estimated to contain at least 50% of all species, with coral reefs being the most aquatic species rich environment.
- ▶ **Biodiversity hotspot**: areas in urgent need of conservation action.

54.2 Threats to Biodiversity

- ▶ Habitat loss is the most important factor in the decline of species in terrestrial environments.
- ▷ Overexploitation in the dominant problem for marine species, and pollution for freshwater species.
- ▷ Invasive species, native species interactions, and other natural events are additional threats to all environments.

54.2.1 Habitat Destruction

- ▶ human causes: buring, logging, damming, dredging, filling in of estuaries and wetlands, plowing, grazing livestock, mining, and various intfrastructure development.
- ▷ Deforestation of natural areas is the greatest threat.
- ▶ Habitat degradation: a reduction in quality of habitat.
- ▶ **Habitat fragmentation**: a pattern of degradation resulting from the dividing of habitats into smaller isolated fragments.
- ▶ Fragmentation particularly hurts top predators, resulting in trophic cascades due many complex food web interactions and results in often unpredictable consequences.
- Smaller populations have higher chances to be exterminated by catastrophic events.
- ▷ Absence of gene flow results in inbreeding depression.
- Overexploitation: unsustainable removal of organisms for human consumption.

54.2.2 Estimating the Probability that a Population or Species Will Go Extinct

▶ A population is considered viable if the prediction probability of survival for at least 100 years is 95%+.

- > Predictions are made with data from age-specific survivorship and fecundity.
- ▷ geographic structure.
- ▶ rate and severity of habitat disturbance.

54.3 Benefits of Biodiversity

- ▶ Biodiversity increases productivity.
- ▷ Diverse assemblages of plant species make more efficient ues of sunlight, water, and nutrients.
- ▶ Facilitation: some species provide nutrients of other benifits that also make the environment better for other vital species.
- ▶ Higher chance of producing extremely productive species.
- ▶ Resistance: measure of the extent a community remains unchanged by a disturbance.
- ▶ Resilience: measure of how quickly a community recovers following a disturbance.
- ▶ Bioprospecting: exploration of bacteria, archaea, protists, plants, fungi, and animals as novel sources of drugs or ingredients in consumer products.

50 Behavior Ecology

▶ **Behavioral ecology**: a focus a subset of organismal ecology— the behavioral adaptations that have evolved in response to ecological selection pressures.

50.1 Introduction to Behavioral Ecology

- ▶ Proximate causation: explains how actions occur in terms of genetic, neurological, hormonal, and skeletal-muscular mechanisms involved.
- ▶ Ultimate causation: explains why actions occur, based on their evolutionary consequences and history.
- ▶ **Fixed action patterns (FAPs)**: highly inflexible behavior paterns.

50.2 What, When, and How to Eat

 \triangleright **Optimal foraging**: Fitness \propto Feeding efficiency \propto $\frac{\text{benefits}}{\text{costs}}$

50.4 Choosing a Place to Live

- ▶ **Piloting**: the use of familiar landmarks while navigating.
- ▶ **Compass orientation**: movement that is oriented in a specific direction.
- ▶ Map orientation: the ability to locate a specific place on Earth's surface.

51 Population Ecology

- ▶ Population ecology: the study of how and why the number of individuals in a population changes over time and space.
- ▶ **Population thinking**: Understanding variation is key to understanding how populations change through time in their environments.

51.1 Distribution and Abundance

- ▶ **Range**: geographic distribution of different species.
- ▶ Two type of factors determine range:
 - Abiotic: temperature, rainfall, geographical structures, and ongoing historical processes.
 - o Biotic: past and current presence of other species.
- ▶ Ranges are dynamic.
- Dispersion patterns have both proximate and ultimate causes.
- ▶ **Metapopulation**: a population of populations connected by migration.

51.2 Demography

- ▶ Demography: study of factors that determin the size and structure of populations through time.
- ▶ **Age structure**: how many individuals of each age are alive.
- ▶ **Life table**: Summarizes the probability that an individual will survive and reproduce in any given time interval over the course of its lifetime.
- ▶ **Cohort**: a group of the same age that can followed through time.
- ▶ Fecundity: number of female offspring produced by each remale in the population.
- ▶ Net reproductive rate: indicates whether the population is increasing or decreasing.

51.3 Population Growth

- $\triangleright \frac{\Delta N}{\Delta t}$ = Births Deaths + Immigrants Emigrants
- ▶ **per capita rate of increase (r)**: birth rate death rate per individual.
- ▶ **Density independent**: population size does not limit the growth rate.
- ▷ Exponential growth is common colonizing a new habitat or recovery after a disaster.
- ▶ **Carrying capacity, K**: the maximum number of individuals in a population that can be supported in a particular habitat over a sustained period of time.
- ▶ **Logistic population growth**: the decrease in growth rater as a population size approaches the carrying capacity.
- ▷ Density independent factors are usually influenced by abiotic factors.
- ▷ Density dependent facts are usually influenced by biotic factors.

52 Community Ecology