## **Contents**

Animals and Environments	
Introduction	2
Homeostasis	3
Physiology and Time	3
Molecules and Cells in Animal Physiology	
Cell Membrane Review	5
Enzyme Fundamentals	5
Genomics	
Genomics	8
Transcriptomics	
Proteomics	
Metabolomics	
Physiological Development	
Epigenetics	11
Mechanisms of Epigenetic Marking	
Epigenetic Inheritance	
Transport of Solutes and Water	
Passive Transport	12
Concentration gradients	
Electrical gradients	
Biological Aspects of Diffusion	
Active Transport	
Channels and Trapsporters	
Colligative Properties of Aqueous Solutions	
Osmosis	
Osmoregulation	
Nutrition, Feeding, and Digestion	. 14
Hatriton, recallig, and Digestion	1.5

## 1 Animals and Environments

#### Introduction

- ▶ What is physiology?
  - o Form and function of organisms; the study of how organisms work.
- ▶ Central questions of physiology: mechanism and origin.
  - o Mechanism:
    - Refers to the components of living organisms and understanding how components interact to enable the organism to function.
  - Origin:
    - Asks why a mechanism exists, or what is the mechanistic adaptive significance of the mechanism.
  - Mechanism and adaptive significance are distinct concepts; knowing about one doesn't necessarily mean you know anything about the other.

#### Krogh's principle:

"For such a large number of problems there will be some animal of choice or a few such animals on which it can be most conveniently studied."

- This idea is central to disciplines that rely on the comparative method.
- Other key concepts:
  - There is unity in diversity; many organisms are very much alike at the most fundamental levels.
  - The differences are subject to particular niches and often highly specialized that allow for biologist to study more complex systems.
  - The similarities allow us overcome technical limitations via animals that are easier to study.
- ▶ Physiology subdisciplines:
  - Mechanistic: emphasizes the mechanisms by which organisms perform their life functions.
  - Evolutionary: emphasizes evolutionary origins and the adaptive significance of traits.
  - Comparative: emphasizes the way in which diverse phylogenetic groups resemble and differ from each other.

- Environmental: emphasizes the ways in which physiology and ecology interact.
- Integrative: emphasizes the importance of all levels of organization, from genes to proteins and tissues to organs in order to better understand whole physiological systems.

#### **Homeostasis**

- Important ideas to remember:
  - Organisms are structurally dynamic; form stays relatively static while individual cells recycle frequently.
  - Most cells are exposed to the internal environment, not external.
  - o Internal cells may vary or kept constant with the environment.
- > Temperature regulation:
  - Conformity: organism's internal temperature correlates with external temperature in a particular range of temperatures.
  - Regulation: internal environment is held mostly constant using celluar mechanisms.
- ▶ **Homeostasis**: the coordinated physiological processes that maintain a relatively constant state in the organism.
  - Positive feedback: less common in homeostasis due difficulty in regulation; leads to runaway effect easily.
  - Negative feedback: more common in homeostasis due to self correcting nature.
  - **Effector**: executes the change in action that produces an effect, e.g. signals to increase temperature.
  - Sensor: sense changes in environment and sends information to the effector.

## **Physiology and Time**

- ▶ Timeframes of physiological change:
  - Acute: short-term, reversible, and quick to adapt to changes in environment. Usually minutes to hours.
  - **Chronic**: long-term after prolonged exposure to new environments. Changes are usually reversible, but often slower.

- Chronic can be termed acclimation, or phenotypic plasticity/flexibility.
- Repetitive acute responses usually lead to chronic responses.
- **Evolutionary**: changes due to alteration in gene frequencies in populations exposed to new environments.
- ▷ Acclimation is not the same as adaption.
  - Adaption is an evolutionary trait presnet at high frequency in a population due to survival/reproductive advantages.
  - Not all traits are adaptations.
  - The amount of natural variation in a trait must be considered across populations, species etc.

## 2 Molecules and Cells in Animal Physiology

#### **Cell Membrane Review**

- ▶ Major cell memberane structures:
  - **Glycoproteins**: carbohydrate chain attached to a protein.
  - Glycolipids: similar to glycoproteins, but attached to lipid molecues.
  - Glycocalyx: combination of glycoproteins and glycolipids on the surface of cell
  - Integral proteins: embedded in phospholipid bilayer.
  - Peripheral proteins: associated with one side of the bilayer.
- ▶ Unsaturated phospholipid: whey hydrocarbon tails contain double bonds (less hydrogen).
  - Increase membrane fluidity due to extra space created.
- ▶ The fluidity of the cell membrane allows proteins to from complexes and dynamically change shape.

### **Enzyme Fundamentals**

- Enzymes: a protein catalyst that plays two primary roles: accelerating and regulating chemical reactions.
- ▶ Substrates: the initial reactants of the reaction that an enzyme catalyzes.
- Enzyme-substrate-complex (E-S): a combination of enzyme (E) with a molecule of substrate (S) that starts a reaction.
  - Usually stabalized by non-covalent bonds.
  - The substrate is converted to a product by first becomeing an enzyme-product complex (E-P), then dissociates to yield free product and free enzyme.
  - $\circ$   $E + S \rightleftharpoons E-S \rightleftharpoons E-P \rightleftharpoons E+P$
- ▶ Saturation kinetics:
  - $\circ$   $V_{max}$ : the maximum velocity of a reaction and is determined by:
    - The number of active enzyme molecues present relative to substrate.
    - The catalytic effectiveness of each enzyme molecule.
    - These properties usually undergo heavy selection pressure.

- Saturated: all enzymes are occupied by a substrate molecule nearly all the time and now unable to increase reaction velocity.
- Hyperbolic: asymptotically approaches V<sub>max</sub>
  - Tends to happen when enzymes have just one substrate binding site.
  - Or when substrate sites behave independently
- Sigmodal: approaches V<sub>max</sub> with a sigmodal trajectory.
  - When multiple sites influence each other.
- $\circ$  **Turnover number (k<sub>cat</sub>)**: the total effectiveness, expressed as the number of substrate molecules coverted to product per second by each enzyme molecule when saturated.
  - Depends partly on the activation energy of the enzyme-catalyzed reaction.
  - Activation energy: the energy required for the substrate to enter the transition state.
  - Transition state: the intermediate chemical state between substrate and product.
  - Enzymes lower the activation energy required to enter transition state.
- ▶ **Enzyme-substrate affinity**: The proclivity of the enzyme to form a complex with the substrate when they meet.
  - Likely complex formation results in high-affinity.
  - Unlikely complex formation results in low-affinity.
  - Affinity affects the shape of the reaction velocity.
    - Higher affinity produces a steeper velocity, and a lower affinity produces a more linear result.
    - Enzyme concentration is not changed.
  - **Half-saturation constant, K**<sub>m</sub>: the substrate concentration required to attain one-half maximum reaction velocity.
    - $K_m$  and enzyme-substrate affinity are inversely related.
    - i.e. low-affinity enzyme has a greater  $K_m$ .
- Molecular Flexibility:
  - o **Conformation**: the three-dimensional shape of a protein.

- Stabalized mostly by weak, noncovalent bonds—hydrogen, van der Waals, hydrophobic, electrostatic, etc.
- Weak interactions allow for easy yet stable conformational changes.
- Enzyme molecules composed of two, three or our proteins are called dimeric, trimeric, or tetrameric respectively.
- - This is because they accelrate the approach towards equilibrium (principles of mass action).
- ▶ **Ligand**: any molecule that selectively binds by noncovalent bonds to structurally and complementary sites on a specific protein.
- ▶ Cooperativity: the interactions between multiple binding sites that may facilitate or inhibit the binding of other sites.
  - Can either positive or negative; facilitating or inhibiting binding on the same molecule.
  - o Homotropic cooperativity: facilitation or inhibition of the same ligand.
  - Heterotropic cooperativity: influences on the binding of other ligands.
  - Interactions occur at a distance, resulting in delayed, or rippling responses.
  - **Allosteric modulation**: the modulation of the catalytic properties.
    - Allosteric sites: nonsubstrate-binding regulatory sites for nonsubstrate ligands that modulate the catalytic properties.
    - Allosteric modulators: the nonsubstrate ligands.
    - Allosteric activation:increases and inhibition:impairs affinity, thus the catalytic activity.
- ▶ **Isozymes**: enzymes that catalyze the same chemical reaction but differ in amino acid sequence.
- ▶ Interspecific enzyme homologs: different molecular forms of an enzyme coded by homologous gen loci in different species.
  - Isozymes and interspecific enzyme homologs often differ in their catalytic and regulatory properties.
  - Functional differences often prove to be adaptive in different environments.

Week 2 3 Genomics

## 3 Genomics

#### **Genomics**

▶ Genomics: study of the genomes—the full set of genetic material—of organisms.

- Metods of genomics:
  - Computational biology and bioinformatics use various computational methods to process large amount of genomic data.
  - High-throughput: methods of analyzing large data with out much human attention and mostly computation.
  - **Annotation**: laborious direct human interpretation.
- ▶ The overarching goals of genomics is to elucidate the evolution and the current functioning of genes and genomes.
- ▶ Gene families: genes that share distinctive DNA base sequences and tend to code for functionally similar proteins.
- ▶ **Postgenomic era**: the study of species after genome is sequenced.

## **Transcriptomics**

- ▶ Transcriptomics: the study of which genes are transcribed to make mRNA and the rates at which they are transcribed.
  - aka transcription profiling.
  - Implies the study of great numbers of mRNAs.
- ► **Transcriptome**: a species full set set of mRNA molecules. It represents the full complement of genes being transcribed at any given time.
  - Time is emphasized; it's a snapshot transcription activity during the observed period.
  - Very useful in comparative methods.
- ▶ Methods of transcriptomics:
  - DNA midroarrays: aka gene chips; a high throughput method tht allows simultaneous analysis of large number of mRNAs.
  - mRNA sequencing: aka RNA-Seq; similar to microarrays, but can identify both known and novel transcripts.
    - More sensitive than microarrays.

Week 2 3 Genomics

- Readily applicable across wide range of species.
- Gene manipulation: studies that permit the direct assessment of gene function by directly altering its expression.
  - Gene deletion: aka gene knockout; breaking or disurbing function of an animal's gene to interfere with proteins, creating deficient or inferior phenotypic traits.
    - **Forced overexpression**: inverse of gene deletion; experimentally increasing synthesis of the mRNA.
    - **Compensation**: phenotypic alterations of that tend to make up for the manipulation done by forced expression or gene deletion.
  - RNA interference (RNAi): allows specific mRNA targets to be silenced in animals with *normal* genomes.
    - Normal genomes: wild type that is not artificially manipulated.
    - · RNAi is reversible.
  - **CRISPR/Cas**: used to edit nuclear DNA in eukaryotic cells.
    - · Can be used to insert sequences that then can be transcribed and tranlasted, providing insights on protein function.

#### **Proteomics**

- ▶ **Proteomics**: the study of proteins being synthesized by cells and tissues.
  - Implies simultaneous study of large numbers of proteins.
  - Predicting proteins from gene transcription is still very difficult;
    transcription, translation, and post-translational processing are all
    regulated dynamically and independently.
- Two-dimensional gel electrophoresis: the primary proteomics method that separates complex mixtures of samples using two different protein properties.
  - Separated by isoelectric points and then molecular weights.

#### **Metabolomics**

- ▶ **Metabolomics**: study of organic compounds in the cells and tissues other than macromolecules coded by the genome.
  - Metabolites: compounds currently being processed by metabolism and the majority of metabolmics focus of study.

Week 2 3 Genomics

- e.g. sugars, amino acids, and fatty acids.

▶ **Nuclear magnetic resonance (NMR)**: primary method of metabolomics that is capable of detecting and quantifying a large variety of compounds through identification of unique signatures in the NMR spectrum.

## 4 Physiological Development

## **Epigenetics**

- ▶ **Epigenetics**: modifications in gene expression with no change in DNA sequence that are transmitted when genes replicate.
- ▶ Marked: aka tagged; when DNA is modified in way to alter expression.

#### **Mechanisms of Epigenetic Marking**

- DNA methylation: addition of methyl groups to cytosine residues in DNA.
  - Generally represses of silences the gene.
  - DNA methyltransferase 1 (DNMT1): an enzyme acts to perpetuate the pattern of methylation in daughter cells.
  - **Methylome**: the set of all methylated sites.
- **Histone modification**: modified histones that that can make DNA more or less accessbile for transcription.
  - Can be modified by methylation, acetylation, phosphorylation, or other covalent bonding of chemical groups at specific sites.
  - Also has mechanisms for perpetuation, e.g. small RNA molecules play a role.
- ▶ **Epigenome**: the global summary of marks or a set of epigenetic marks in a cell.

### **Epigenetic Inheritance**

- Mitotic inheritance: aka somatic; perpetuation of marks during the process of cell division by mitosis within an individual.
- Meiotic inheritance: aka transgenerational; perpetuation of marks during meiosis that results in passing of marks to offspring.
- ▶ Research is continuing to provide strong evidence that epigenetics can radically alter physiology.
- ▷ Epigenetic marking may also play large roles in lifelong effects due early-life and prenatal environments.

## 5 Transport of Solutes and Water

### **Passive Transport**

- ▶ **Equilibrium**: the state at which a of minimum capacity to do work under locally prevailing conditions.
  - A change toward equilibrium is always in the direction of decreasing work potential.

### **Concentration gradients**

- General definition: the difference in concentration between two solutions or regions.
- Nore accurately:  $\frac{\Delta C}{X}$  where X is the distance separating (boundary layer) the regions of concentraion.
- $\triangleright \text{ Fick diffusion equation: } J = D \frac{\Delta C}{X}$ 
  - J is the net number of solute molecules passing into the low-concentration region from the high-concentraion region.
  - **Diffusion coefficient (D)**: proportionality factor determined by the permeability of the membrane or epithelium as well as the temperature.
- ▶ Each solute diffuses according to its own concentraion gradient.
- ▶ **Simple diffusion**: aka diffusion; moves solute from an area of high solute concentration to an area of low solution concentration.
  - Does not use energy as it can only move material in the direction of the concentration gradient and towards equilibrium.

#### **Electrical gradients**

- ▶ **Electrical gradient**: difference in charge across a membrane.
- ▶ Many solutes bear electrical charge that affects the diffusion of such solutes.
- ▶ Bulk solution: solution not in contact with with a membrane.
  - Has a net charge of zero, this regions do not differe in charge.
  - Lack of net charge does not affect diffusion in the bulk solution, though does affect diffusion across the cell membranes of epithelia.
  - Bulk flow: physical kinetic movement of fluid, typically due to pressure.
- ▶ **Electrochemical gradient**: gradient consisting of the chemical gradient (concentration gradient) and the electrical gradient.

### **Biological Aspects of Diffusion**

- ▶ **Ion channels**: integral membrane protein that permits the passive transport of inorganic ions by diffusion through the membrane.
  - Some can be selective for certain ions, such as Na<sup>+</sup>, Cl<sup>-</sup>, and K<sup>+</sup>
  - Even the least selective discriminate between anions and cations
  - Gated channels: ion channels that can open and close due to the proteins allowing for conformational changes.
    - Voltage-gated: responds to voltage change.
    - Stretch-gated: aka tension gated: responds to physical tensions.
    - Phosphorylation-gated: responds due to changes in protein phosphorylation.
    - Ligan-gated: responds due to extracelluar signaling.
- ▶ **Permeability**: the ease at which the solute can move through the membrane by diffusion.
  - Changed by use and quantity of ion channels
- ▶ **Faciliated diffusion**: moves with or against concentration gradient, though it uses energy due to use of channel or protein transporters.
  - Instead it always occurs in the direction of electrochemical equilibrium.
  - o Solutes are transported much faster than they are in simple diffusion.
  - Solutes must bind reversibly with biding sites on transporter proteins.

## **Active Transport**

D

## **Channels and Trapsporters**

 $\triangleright$ 

## **Colligative Properties of Aqueous Solutions**

 $\triangleright$ 

#### **Osmosis**

 $\triangleright$ 

## Osmoregulation

Excerpt from Chapter 26: Water and Salt in Physiology  $\mapsto$ 

 $\triangleright$ 

# 7 Nutrition, Feeding, and Digestion

 $\triangleright$