

# Contents

<b>1</b>	<b>The Brain and Behavior</b>	<b>6</b>
1.1	The Brain Has Distinct Functional Regions . . . . .	6
<b>2</b>	<b>Nerve Cells, Neural Circuitry, and Behavior</b>	<b>7</b>
2.1	The Nervous System Has Two Classes of Cells . . . . .	7
<b>3</b>	<b>Genes and Behavior</b>	<b>9</b>
<b>4</b>	<b>The Cells of the Nervous System</b>	<b>10</b>
<b>5</b>	<b>Ion Channels</b>	<b>11</b>
5.1	Rapid Signaling in the Nervous System Depends on Ion Channels . .	11
5.2	Ion Channels are Proteins That Span the Cell Membrane . . . . .	11
5.3	Ion Channels in ALL Cells Share Several Characteristics . . . . .	12
<b>6</b>	<b>Electrical Properties of the Neuron</b>	<b>13</b>
<b>7</b>	<b>Propagated Signaling</b>	<b>14</b>
7.1	Personal Notes . . . . .	14
<b>8</b>	<b>Overview of Synaptic Transmission</b>	<b>15</b>
8.1	Synapses Are Either Electrical or Chemical . . . . .	15
8.2	Electrical Synapses Provide Instantaneous Signal Transmission . . .	15
8.3	Chemical Synapses Can Amplify Signals . . . . .	16
<b>9</b>	<b>Directly Gated Transmission</b>	<b>17</b>
<b>10</b>	<b>Synaptic Integration in CNS</b>	<b>18</b>
10.1	Central Neurons Receive Excitatory and Inhibitory Inputs . . . . .	18
10.2	Inhibitory Synaptic Action . . . . .	18
<b>11</b>	<b>Second Messengers</b>	<b>19</b>
<b>12</b>	<b>Transmitter Release</b>	<b>20</b>
<b>13</b>	<b>Neurotransmitters</b>	<b>21</b>
13.1	Four Criteria of a Neurotransmitter . . . . .	21

13.2 Only a Few Small-Molecule Substances Act as Transmitters . . . . .	21
13.3 Small-Molecule Transmitters Are Actively Taken up into Vesicles . . .	24
13.4 Removal of Transmitter from the Synaptic Cleft Terminates Synaptic Transimission . . . . .	24
<b>14 Diseases of the Nerve and Motor Unit</b>	<b>25</b>
<b>15 Organization of the Central Nervous System</b>	<b>26</b>
15.1 The Central Nervous System Consists of the Spinal Cord and the Brain	26
15.2 The Major Functional Systems Are Similarly Organized . . . . .	26
15.3 The Cerebral Cortex is Concerned with Cognition . . . . .	27
15.4 Subcortical Regions of the Brain are Functionally Organized into Nuclei	28
15.5 Modulatory Systems in the Brain Influence Motivation, Emotion, and Memory . . . . .	29
15.6 The Peripheral Nervous System is Anatomically Distinct from the Central Nervous System . . . . .	29
<b>16 Organization of Perception and Movement</b>	<b>30</b>
16.1 Sensory Information Processing is Illustrated in the Somatosensory System . . . . .	30
16.2 The Thalamus is an Essential Link Between Sensory Receptors and the Cerebral Cortex for All Modalities Except Olfaction . . . . .	31
16.3 Sensory Information Processing Culminates in the Cerebral Cortex .	32
16.4 Voluntary Movement is Mediated by Direct Connections Between the Cortex and Spinal Cord . . . . .	33
<b>17 Representation of Space and Action</b>	<b>34</b>
17.1 The Brain has an Orderly Representation of Personal Space . . . . .	34
17.2 The Internal Representation of Personal Space can be Modified by Experience . . . . .	34
17.3 Is Consciousness Accessible to Neurobiological Analysis? . . . . .	34
<b>18 Organization of Cognition</b>	<b>36</b>
18.1 Functionally Related Areas of Cortex Lie Close Together . . . . .	36
18.2 Sensory Information is Processed in the Cortex in Serial Pathways .	36
18.3 Goal-Directed Motor Behavior Is Controlled in the Frontal Lobe . . . .	38

18.4 Limbic Association Cortex is a Gateway to the Hippocampal Memory System . . . . .	38
<b>19 Functions of the Premotor Systems</b>	<b>40</b>
19.1 Direct Connections Between the Cerebral Cortex and Spinal Cord Play a Fundamental Role in the Organization of Voluntary Movements	40
19.2 The Four Premotor Areas of the Primate Brain Also Have Direct Connections in the Spinal Cord . . . . .	41
19.3 Motor Circuits Involved in Voluntary Actions are Organized to Achieve Specific Goals . . . . .	41
19.4 The Hand Has a Critical Role in Primate Behavior . . . . .	41
19.5 The Joint Activity of Neurons in the Parietal and Premotor Cortex Encodes Potential Motor Acts . . . . .	42
<b>20 Functional Imaging of Cognition</b>	<b>43</b>
20.1 Functional Imaging Reflects the Metabolic Demand of Neural Activity	43
<b>21 Sensory Coding</b>	<b>44</b>
21.1 Psychophysics Relates the Physical Properties of Stimuli to Sensations . . . . .	44
21.2 Physical Stimuli are Represented in the Nervous System by Means of the Sensory Code . . . . .	44
21.3 Modality-Specific Pathways Extend to the Central Nervous System .	45
<b>22 Somatosensory System</b>	<b>46</b>
22.1 The Primary Sensory Neuron of the Somatosensory System are Clustered in the Dorsal Root Ganglia . . . . .	46
22.2 Peripheral Somatosensory Nerve Fibers Conduct Action Potentials at Different Rates . . . . .	46
22.3 Many Specialized Receptors Are Employed by the Somatosensory System . . . . .	47
<b>23 Touch</b>	<b>48</b>
23.1 Active and Passive Touch Evoke Similar Responses in Mechanoreceptors . . . . .	48
23.2 The Hand Has Four Major Types of Mechanoreceptors . . . . .	48

23.3 Tactile Information is Processed in the Central Touch System . . . . .	48
<b>24 Pain</b>	<b>50</b>
24.1 Noxious Insults Activate Nociceptors . . . . .	50
<b>25 Visual Processing</b>	<b>51</b>
25.1 Visual Perception is a Constructive Process . . . . .	51
25.2 Form, Color, Motion, and Depth are Processed in Discrete Areas of the Cerebral Cortex . . . . .	51
25.3 The Receptive Fields of Neurons at Successive Relays in an Afferent Pathway Provide Clues to How the Brain Analyzes Visual Form . . . . .	52
25.4 The Visual Cortex is Organized into Columns of Specialized Neurons	52
25.5 Intrinsic Cortical Circuits Transform Neural Information . . . . .	53
<b>26 Low-Level Visual Processing</b>	<b>54</b>
26.1 The Photoreceptor Layer Samples the Visual Image . . . . .	54
26.2 Ganglion Cells Transmit Neural Images to the Brain . . . . .	54
26.3 A Network of Interneurons Shapes the Retinal Output . . . . .	55
<b>27 Intermediate-Level Visual Processing</b>	<b>56</b>
27.1 Internal Models of Object Geometry Help the Brain Analyze Shapes .	56
27.2 Depth Perception Helps Segregate Objects from Background . . . . .	56
27.3 Local Movement Cues Define Object Trajectory and Shape . . . . .	57
27.4 Context Determines the Perception of Visual Stimuli . . . . .	57
27.5 Cortical Connections, Functional Architecture, and Perception are In- timately Related . . . . .	57
<b>28 High-Level Visual Processing</b>	<b>59</b>
28.1 The Inferior Temporal Cortex is the Primary Center for Object Per- ception . . . . .	59
28.2 Object Recognition Relies on Perceptual Constancy . . . . .	59
28.3 Visual Memory is a Component of High-Level Visual Processing . . .	60
28.4 Associative Recall of Visual Memories Depends on Top-Down Acti- vation of the Cortical Neurons that Process Visual Stimuli . . . . .	60
<b>29 Visual Processing and Action</b>	<b>61</b>

29.1 Successive Fixations Focus Our Attention in the Visual Field . . . . . 61

29.2 The Visual Scene Remains Stable Despite Continual Shifts in the  
Retinal Image . . . . . 61

29.3 Vision Lapses During Saccades . . . . . 61

# 1 The Brain and Behavior

## 1.1 The Brain Has Distinct Functional Regions

### The Central Nervous System Has Seven Main Parts

- ▷ **Spinal cord:** most caudal part of the central nervous system. It is subdivided into cervical, thoracic, lumbar, and sacral regions.
- ▷ **Brain stem:** consists of the medulla oblongata, pons, and midbrain. Relays input from the spinal cord and back, and controls input to and from the head.
- ▷ **Medulla oblongata:** rostral to spinal cord and includes several centers responsible for vital autonomic functions.
- ▷ **Pons:** rostral to medulla and conveys information about movement.
- ▷ **Cerebellum:** lies behind pons, modulates force and range of movement, and involved in learning motor skills.
- ▷ **Diencephalon:** lies rostral to midbrain and contains two structures, thalamus (processes information reaching cerebral cortex) and hypothalamus (regulates autonomic, endocrine, and visceral functions).
- ▷ **Cerebrum:** comprises two cerebral hemispheres, each consisting of wrinkled outer layer (the cerebral cortex), and three deep lying structures (basal ganglia, the hippocampus, and the amygdaloid nuclei).
- ▷ **Cerebral cortex:** divided into four distinct lobes— frontal, parietal, occipital, and temporal. The frontal lobe is largely concerned with short-term memory and planning, as well as movement; the parietal lobe with somatic sensation, forming a body image, and relating it to extrapersonal space; the occipital lobe with vision; and the temporal lobe with hearing—combined with deeper structures—with learning, memory, and emotion.

## 2 Nerve Cells, Neural Circuitry, and Behavior

### 2.1 The Nervous System Has Two Classes of Cells

- ▷ There are two main classes of cells in the nervous system: nerve cells, or neurons, and glial cells, or glia.
- ▷ A neuron has four defined regions:
  - **Cell body**: or **soma**, is the metabolic center of the cell, containing normal cell organelles.
  - **Dendrites**: branch out in tree-like fashion and are main apparatus for receiving signals.
  - **Axon**: extends some distance from a cell and carries signals to other neurons.
  - **Presynaptic terminals**: specialized enlarged regions of its axon's branches and is responsible for transfer of signals.
- ▷ **Principle of dynamic polarization**: electrical signals only flow in one direction in neurons.
- ▷ **Connectional specificity**: nerve cells do not connect randomly with one another in these formation of networks.
- ▷ Neurons are classified into three groups:
  - **Unipolar**: simplest due to single primary process, which gives rise to many branches. One branch as axon and others as receiving structures. These cells predominate invertebrates; they occur in the autonomic nervous system in vertebrates.
  - **Bipolar**: oval soma that gives rise to two processes: a dendritic structure that receives signals and an axon that carries information towards the central nervous system. Many sensory cells are bipolar, and pain receptors are pseudo-unipolar.
  - **Multipolar**: predominate nervous system of vertebrates and vary greatly in shape; typically containing a single neuron and many dendritic points emerging from various points around the cell body.

- ▷ Glial cells support nerve cells and greatly outnumber neurons.
- ▷ Glial cells surround the cell bodies, axons, and neurons and can be divided into two major classes:
  - **Microglia**: immune system cells that become phagocytes during injury, infection, or degenerative diseases.
  - There are three main types of **macroglia**: oligodendrocytes, Schwann cells, and astrocytes. About 80% of all brain cells are macrogalia.



### **3 Genes and Behavior**

This chapter has been intentionally left blank, see genetic notes for more information.

## 4 The Cells of the Nervous System

This chapter was intentionally left blank, see cell biology notes for more information.

## 5 Ion Channels

### 5.1 Rapid Signaling in the Nervous System Depends on Ion Channels

- ▷ Up to 100 million ions can pass through a single channel each second, comparable to the turnover rate of the fastest enzymes, catalase and carbonic anhydrase.
- ▷ Each channel allows only one or a few types of ions to pass.
- ▷ Many open and close, however, some remain open resulting in significant contribution to resting potential.
- ▷ Ion pumps maintain gradients and are 100 to 100,000 times slower than channels.
- ▷ Questions for this chapter:
  - Why do nerve cells have channels?
  - How can channels conduct ions at such high rates and still be selective?
  - How are channels gated?
  - How are properties of these channels modified by various intrinsic and extrinsic conditions?

### 5.2 Ion Channels are Proteins That Span the Cell Membrane

- ▷ Cells have channels in order to transport ions across lipid bilayer easily and eliminate the need to be stripped of waters of hydration.
- ▷ The smaller the ion, the greater attraction to water, and the lower its mobility. This partially explains selection, but how does the inverse selection, that selecting of lower mobility, occur?
- ▷ Some ions bind to proteins that can transport them, but this is far too slow for some cases.

- ▷ An extension of pore theory says that channels have narrow regions that act as molecular sieves, where the ion sheds most of its water and only is let through by a binding to a specifically charged selectivity filter.

### 5.3 Ion Channels in ALL Cells Share Several Characteristics

- ▷ The opening and closing of a channel involve conformational changes.
- ▷ **Gating**: the transition of a channel between these stable functional states.
- ▷ Three major gating mechanisms:
  - Ligand: binding of chemical ligands known as agonists at either cellular site; transmitters on the extracellular; others that activate signaling cascades; and more.
  - Voltage-gated: changes in electrochemical changes as often as temperature sensors.
  - Mechanical stretch or physical changes in the membrane.

## 6 Electrical Properties of the Neuron

This chapter was intentionally left blank. No alternative notes, but may need to review chemistry and physics if this chapter is needed.

## 7 Propagated Signaling

### 7.1 Personal Notes

- ▷ The variety of voltage-sensitive ion channels and the influence of cytoplasmic factors may be analogous to bias/weights or other hyperparameters in neural networks.
- ▷ Genetic changes thus change these networks and may be how transfer learning takes place instead of starting from complete scratch for every organism.
- ▷ Innate abilities could represent earlier layers in the network and heavily genetically programmed, while later layers are given more time to develop based on environment.
- ▷ Can epigenetics have a relatively fast acting change on inherited intelligence?

The rest of this chapter has been left blank, as it is more of an extension of molecular genetics, which I may need to return to later to answer questions like those listed above.

## 8 Overview of Synaptic Transmission

### 8.1 Synapses Are Either Electrical or Chemical

- ▷ Average neuron forms and receive several thousand synaptic connections each, with the Purkinje cell of the cerebellum receiving up to 100,000.
- ▷ Both forms of transmission can be enhanced or diminished by cellular activity.
- ▷ Electrial synapses are used to send rapid stereotyped depolarizing signals.
- ▷ Chemical synapses are capable of more complex behaviors due to vairable signaling.
- ▷ Most synapses are chemical.

### 8.2 Electrical Synapses Provide Instantaneous Signal Transmission

- ▷ Presynaptic terminals must be big enough for its membrane to contain many ion channels to trigger initial depolarization.
- ▷ Postsynaptic terminals must be relatively to small in due to Ohm's law.
- ▷ Even weak subthreshold depolarizing currents can be carried to the postsynaptic neuron and depolarize it.
- ▷ Electrical synapses have a specialized region of contact called the gap junction, with seperation of only 4 nm, bridged by gap-junction channels specialized to conduct ionic current.
- ▷ Electrical transmission can be used to orchestrate actions or large groups of neurons.
- ▷ Groups of electrically coupled cells allows for explosive reactions.
- ▷ Gap junctions are formed between glial cells as well as neurons.

### 8.3 Chemical Synapses Can Amplify Signals

- ▷ Chemical synapses are used to amplify or inhibit signals.
- ▷ The synaptic cleft is 20-40 nm wide and depend on the diffusion of neurotransmitters to carry out signaling.
- ▷ Neurotransmitters are clustered at speclized regions called **active zones**, which allow for selective activate of nearby postsynaptic receptors, which lead to the opening or closing of ion channels.
- ▷ Chemical synapses can be as short as 0.3 ms but often last several ms.
- ▷ Weak activations of chemical synapses can activate larger electrial synapses.
- ▷ The action of a transmitter depends on the properties of the postsynaptic receptor, not the chemical properties of itself.
- ▷ Neurotransmitters control the opening of ion channels in the postsynaptic cell either directly or indirectly.
- ▷ Indirect effects tend to last seconds to minutes and often modulate behavior due to alterations in the excitability of neurons and their synaptic connections.



## 9 Directly Gated Transmission

This chapter has been intentionally left blank. Unlikely to return to this chatper.

## 10 Synaptic Integration in CNS

### 10.1 Central Neurons Receive Excitatory and Inhibitory Inputs

- ▷ Generation of an action potential often requires the near-synchronous firing of a number of sensor neurons.
- ▷ Small inhibitory postsynaptic potential (IPSP), if strong enough, can counteract the sum of the excitatory actions and prevent membrane potential from reaching threshold potential.
- ▷ **Sculpting** function of synaptic inhibition that exerts control over action potentials in neurons that are spontaneously active due to intrinsic pacemaker channels, often completely shaping the firing patterns of cells.
- ▷ Most transmitters usually are inhibitory or excitatory despite being able to be either type.

### 10.2 Inhibitory Synaptic Action

- ▷ Inhibitory synapses play an essential role in the nervous system both by preventing too much excitation and by helping coordinate activity among networks of neurons.
- ▷ Inhibitory inputs that hyperpolarize the cell perform subtraction on the excitatory inputs, where those that **shunt** perform division.
- ▷ Adding or removing nonshunting inhibitory inputs results in summation, while the combination or excitatory with the removal of inhibitory shunt produces a multiplication.

## 11 Second Messengers

This chapter has been intentionally left blank. Molecular focused chapters may be reviewed later when a more narrow question needs to be answered.

## 12 Transmitter Release

This chapter has been intentionally left blank. More on the chemistry and molecular function of transmission. Such information May answer questions in the future, but it's not the focus of inquiry at the moment.

## 13 Neurotransmitters

### 13.1 Four Criteria of a Neurotransmitter

- ▷ Four steps of synaptic transmission:
  1. Synthesis and storage of a transmitter.
  2. Release of the transmitter.
  3. Interaction of the transmitter with receptors and postsynaptic membrane.
  4. removal of the transmitter from the synaptic cleft.
- ▷ First approximation of a neurotransmitter can be defined as a substance released by a neuron that affects a specific target in a specific manner.
- ▷ Neurotransmitters typically act on targets other than the releasing neuron itself, unlike autacoids.
- ▷ Neurotransmitter interaction with receptors is typically transient, lasting from milliseconds to minutes.
- ▷ General criteria for neurotransmitters:
  - It is synthesized in the presynaptic neuron.
  - It is present in the presynaptic terminal and is released in amounts sufficient to exert a defined action on the postsynaptic neuron or effector organ.
  - When administered exogenously in a reasonable concentration it mimics the action of the endogenous transmitter.
  - A specific mechanism usually exists for removing the substance from the synaptic cleft.

### 13.2 Only a Few Small-Molecule Substances Act as Transmitters

- ▷ **Acetylcholine (ACh):**

- Only low weight amine transmitter substance that is not an amino acid or derived directly from one.
- Nervous tissue cannot synthesize choline, which limits ACh biosynthesis due to choline acetyltransferase being the only enzymatic reaction.
- ACh is released by spinal motor neurons.
- In the autacoids nervous system it is the transmitter for all preganglionic neurons and for parasympathetic postganglionic neurons as well.
- ACh is the principle neurotransmitter of the reticular activating system, which modulates arousal, sleep, wakefulness, and other critical aspects of human consciousness.

▷ **Biogenic Amines:**

- **Catecholamine Transmitters:**

- **Dopamine** — Tyrosine
- **Norepinephrine** — Tyrosine
- **Epinephrine** — Tyrosine
- Tyrosine hydroxylase is the rate-limiting for synthesis of both dopamine and norepinephrine.
- $\beta$ -hydroxylase converts dopamine to norepinephrine and is membrane-associated.
- Norepinephrine is the only transmitter synthesized within vesicles.
- In order for epinephrine to be formed, then its immediate precursor, norepinephrine, must exit from vesicles into the cytoplasm.
- In order to be released, epinephrine must be taken up into vesicles.
- Three of four dopaminergic nerve tracts arise in the midrain, with the last arising in the arcuate nucleus of the hypothalamus.

- Synthesis of biogenic amines is highly regulated and can be rapidly increased.

- **Serotonin** — Tryptophan

- **Melatonin** — Serotonin

- Tryptophan hydroxylase is the limiting reaction and the first enzyme in the pathway.
- Cell bodies with serotonergic neurons are found around the midline raphe nuclei of the brain stem and are involved in regulating attention.
- Productions of serotonergic cells are widely distributed throughout the brain and spinal cord.
- Antidepressant medication inhibit the uptake of serotonin, norepinephrine, and dopamine.

- **Histamine** — Histidine

- Long been recognized as an autacoid, active when released from mast cells in the inflammatory reaction.
- Concentrated in the hypothalamus.

▷ **Amino Acid Transmitters:**

- **Aspartate** — Oxaloacetate

- **$\gamma$ -Aminobutyric acid (GABA)** — Glutamine:

- Present at high concentrations throughout the central nervous system and detectable in other tissues.

- **Glutamate** — Glutamine:

- Most frequently used at excitatory synapses throughout the central nervous system.

- **Glycine** — Serine:

- Major transmitter used by inhibitory interneurons of the spinal cord.

▷ **ATP and Adenosine**

- Can act as transmitters at some synapses.
- Adenosine has an inhibitory effect in the central nervous system.
- Caffeine's stimulatory effect depends on inhibition of adenosine binding to its receptors.
- ATP released by tissue damage acts to transmit pain sensation in some cases.

### 13.3 Small-Molecule Transmitters Are Actively Taken up into Vesicles

- ▷ **Tranmitter** glutamate must be kept separate from **metabolic** glutamate; this is done through compartmentalization in synaptic vesicles.
- ▷ Drugs that are sufficiently similar to the normal transmitter substance can act as false transmitters, tho they often bind weakly decreasing efficacy of of transmission.

### 13.4 Removal of Transmitter from the Synaptic Cleft Terminates Synaptic Transimission

- ▷ If transmitter molecules released in one synaptic action were allowed to remain in the cleft after release, then they would prevent ner signlas from getting through, the synapse would become refractory due to desensitization.
- ▷ Transmitters are removed by three mechanisms: diffusion, enzymatic degradation, and reuptake.
- ▷ Degradation is only used by cholinergic synapses.
- ▷ Degradation allows for single use signaling and for the lingering choline to be reused.



## 14 Diseases of the Nerve and Motor Unit

This chapter has been intentionally left blank. Might revisit, reading comprehension was low.

## 15 Organization of the Central Nervous System

### 15.1 The Central Nervous System Consists of the Spinal Cord and the Brain

- ▷ The spinal cord is divided into a core of central gray matter and surrounding white matter.
- ▷ The gray matter is divided into **dorsal** and **ventral** horns.
- ▷ **Dorsal horn**: contains orderly sensory relay neurons that receive input from periphery.
- ▷ **Ventral horn**: contains group of motor neurons and interneurons that regulate motor neural firing patterns.
- ▷ The brain stem(**Medulla, pons, and midbrain**) has five distinct functions:
  1. Spinal cord mediate sensation and motor control of trunk and limbs, but the brain stem control the head, neck and face.
  2. Site of entry for information from several specialized sites such as hearing, balance, and taste.
  3. Mediation of parasympathetic reflexes, such out cardiac output, pupil constriction, and more.
  4. Contains ascending and descending pathways that carry sensory and motor information to other parts of the CNS.
  5. Contains the **reticular formation**, which receives a summary of incoming sensory information and regulates alertness and arousal.

### 15.2 The Major Functional Systems Are Similarly Organized

- ▷ The central nervous system consists of several functional systems that are relatively autonomous and much work together using numerous interconnected anatomical sites throughout the brain.
- ▷ Information is transformed at each synaptic relay, with the output rarely being the same as the input.

- ▷ Neurons at each synaptic relay are organized into a neural map of the body.
- ▷ Most sensory systems inputs are arranged topographically through out successive stages of processing.
- ▷ Each functional system is hierarchically organized.
- ▷ **Decussations:** crossing of second order fiber from the brain stem and the spinal cord.

### 15.3 The Cerebral Cortex is Concerned with Cognition

- ▷ Increasing the surface area due to sulci and gyri allow for greater number of cortical neurons which provide a greater capacity for information processing.
- ▷ Neurons in the cerebral cortex are organized in layers and columns which helps computational efficiency.
- ▷ The neocortex receives inputs from the thalamus, other cortical regions on both sides of the brain, and other structures then output to other various regions.
- ▷ The input-output relation is organized into orderly layering of cortical neurons, with most containing six layers.
  - Layer I: the **molecular layer**, is occupied by dendrites of cells located in deeper layers and axons that make connections to other areas of the cortex.
  - Layer II: the **external granule cell layer**, one of two layers that contain small spherical neurons.
  - Layer III: the **external pyramidal cell layer**, second layer of small spherical neurons, typically larger than layer II.
  - Layer IV: the **internal granule cell layer**, contains much larger number of spherical neurons and is main recipient of sensory input from the thalamus.
  - Layer V: the **internal pyramidal cell layer**, contains pyramidal neurons

that are also larger than its external layer. These neurons give rise to major output pathways of the cortex.

- Layer VI: the **multiform layer**, a blend of neurons into white matter that forms the deep limit of the cortex and carries axons to and from areas of the cortex.
- ▷ Thickness of the layers varies throughout the cortex.
- ▷ The cerebral cortex has a large variety of neurons, more than 40 different types based only on the distribution of their dendrites and axons.
- ▷ Most neurons are either principal (projection) neurons or local interneurons.

#### 15.4 Subcortical Regions of the Brain are Functionally Organized into Nuclei

- ▷ Three major structures lie deep within the cerebral hemisphere: the basal ganglia, the hippocampal formation, the amygdala, and the basal ganglia. These subcortical structures act to regulate the cortical activity.
- ▷ Basal ganglia regulates movement and certain cognitive functions such as learning of motor skills.
- ▷ The basal ganglia has five major functional subcomponents: the caudate nucleus, putamen, globus pallidus, subthalamic nucleus, and substantia nigra.
- ▷ The hippocampal formation includes the hippocampus, dentate gyrus, and subiculum. Together these structures are responsible for the formation of long-term memories episodic memories, but not responsible for storage.
- ▷ The amygdala is involved in analyzing the emotional significance of sensory stimuli.

### 15.5 Modulatory Systems in the Brain Influence Motivation, Emotion, and Memory

- ▷ Some brain areas are neither sensory nor motor, but instead modify specific functions.
- ▷ Distinct modulatory systems within the brain stem modulate attention and arousal.

### 15.6 The Peripheral Nervous System is Anatomically Distinct from the Central Nervous System

- ▷ The peripheral nervous system supplies the central nervous system with a continuous stream of information about both external and internal environments. It is split into two divisions.
- ▷ The **somatic division** includes the sensory neurons that receive information from skin, muscles, and joints and provide information about position and pressure.
- ▷ The **autonomic division** mediates visceral sensation as well as motor control of the viscera, vascular, and exocrine glands. It consists of sympathetic (response to stress), parasympathetic (restores homeostasis), and enteric (controls smooth muscle of the gut) systems.

## 16 Organization of Perception and Movement

### 16.1 Sensory Information Processing is Illustrated in the Somatosensory System

- ▷ Complex behaviors require the integrated action of several nuclei and cortical regions, processed in a hierarchical fashion, and becomes increasingly complex.
- ▷ Complex processing results in a light touch or painful prick in the skin being mediated by often very different pathways.
- ▷ Somatosensory information from the trunk and limbs is conveyed to the spinal cord.
- ▷ The spinal cord is divided into four major regions: cervical, thoracic, lumbar, and sacral.
- ▷ Spinal nerves at the cervical level are involved with sensory perceptions and motor function of the back of the head, neck, and arms.
- ▷ Thoracic nerves innervate the upper trunk.
- ▷ Lumbar and sacral nerves innervate the lower trunk, back, and legs.
- ▷ Each of the four regions of the spinal cord contains several segments, despite the lack of appearance of segmentation of mature spinal cords.
- ▷ The spinal cord varies in size and shape due to two organizational features.
- ▷ First, the relatively few sensory axons enter the cord at the sacral level, with number of entering axons increasing progressively at higher levels.
- ▷ Most descending axons from the brain terminate at cervical levels.
- ▷ Second, the variation in the size of the ventral and dorsal horns.
- ▷ The number of ventral motor neurons dedicated to the body region roughly parallels the dexterity of movements of that region.
- ▷ **lumbosacral** and **cervical enlargements**: regions of the spinal cord where fibers enter the cord due demands of sensory neurons for finer tactile discrimination in limbs.

- ▷ The primary sensory neurons of the trunk and limbs are clustered in the dorsal root ganglia. These neurons are pseudo-unipolar in shape and have bifurcated axon with central and peripheral branches.
- ▷ Local branches activate local reflex circuits while ascending branches carrying information to the brain that give rise to the perception.
- ▷ The central axons of dorsal root ganglion neurons are arranged to produce a map of the body surface.
- ▷ Each somatic submodality is processed in a distinct subsystem from the periphery to the brain.

## 16.2 The Thalamus is an Essential Link Between Sensory Receptors and the Cerebral Cortex for All Modalities Except Olfaction

- ▷ The thalamus conveys sensory input to the primary sensory areas of the cerebral cortex and additionally acts as a gatekeeper depending on the behavioral state of the animal.
- ▷ The thalamus is a good example of a brain region made up of several well-defined nuclei.
- ▷ Some nuclei receive information specific to a sensory modality and project to a specific area of the neocortex.
- ▷ The nuclei of the thalamus are most commonly classified into four groups:
  - **anterior group:** receives most input from the mammillary nuclei of the hypothalamus and presubiculum of the hippocampal formation. The role of this region is uncertain, but thought to be related to memory and emotion.
  - **Medial group:** consists mostly of the mediodorsal nucleus. It receives input from the basal ganglia, amygdala, and midbrain and is been implicated in memory.
  - **ventral group:** important for motor control and carry information from basal ganglia and cerebellum to the motor cortex.

- **posterior group:** includes the medial and lateral geniculate nucleus (component of auditory system), lateral posterior nucleus (component of the retina and visual cortex in the occipital lobe), and the pulvinar (involved in the parietal-occipital-temporal cortex).
- ▷ **Reticular nucleus:** a unique sheet-like structure covering the thalamus.
- ▷ Neurons of the reticular nucleus are not interconnected with the neocortex, instead the axons terminate on the other nuclei of the thalamus.
- ▷ Thus, the reticular nucleus modulates activity in other thalamic nuclei based on its monitoring of the entirety of the thalamocortical stream.
- ▷ The thalamus not only relays information but is a crucial step and adds substantial degree of information processing.

### 16.3 Sensory Information Processing Culminates in the Cerebral Cortex

- ▷ Parts of the body are represented in the cortex somatotopically, but the area of the cortex is not proportional to its mass. Instead, it is proportional to the density of innervation.
- ▷ The cortical areas involved in the early stages of sensory processing are concerned primarily with a single modality.
- ▷ Unimodal association areas converge on multimodal association areas of the cortex concerned with combining sensory modalities.
- ▷ Multimodal associational areas are heavily interconnected with the hippocampus and appear to be important for unified percept and representation of the percept in memory.
- ▷ There is a close linkage between the somatosensory and motor functions of the cortex.



## **16.4 Voluntary Movement is Mediated by Direct Connections Between the Cortex and Spinal Cord**

- ▷ The human corticospinal tract consists of approximately one million axons, with 40% originating from the motor cortex.
- ▷ Most of the corticospinal fibers cross the midline in the medulla at a location known as the pyramidal decussation.
- ▷ 10% of those fibers do not cross until they reach the local where they terminate.
- ▷ The motor information carried in the corticospinal tract is significantly modulated by the sensory information and information from other motor regions.
- ▷ The cerebellum is thought to be part of an error-correcting mechanism for movements because it can compare movement commands from the cortex with somatic sensory information about what actually happened.

## 17 Representation of Space and Action

### 17.1 The Brain has an Orderly Representation of Personal Space

- ▷ Internal representation can be thought of as a certain pattern of neural activity that has at least two aspects:
  - The pattern of activation within a particular population of neurons.
  - the pattern of firing in individual cells.
- ▷ The cortex has a map of the sensory receptive surface for each sensory modality.
- ▷ Cortical maps of the body are the basis of accurate clinical neurological examinations.
- ▷ There is a direct relationship between the anatomical organization of the functional pathways in the brain and specific perceptual and motor behaviors.

### 17.2 The Internal Representation of Personal Space can be Modified by Experience

- ▷ Details of sensory maps vary considerably from one individual to another.
- ▷ Lost connections can be taken over by existing nearby connections.

### 17.3 Is Consciousness Accessible to Neurobiological Analysis?

- ▷ John Searle and Thomas Nagel have defined three essential features of self-awareness:
  - Subjectivity, or the awareness of a self that is the center of experience.
  - Unity, or the fact that our experience of the world at any given moment is felt as a single unified experience.

- Intentionality, or the the experience that connets successive moments and the sense that the successive moments are directed to some goal.
- ▷ Crick and Koch argue that our efforts should be focused on visual perceptionand in particular on two phenomena: binocular rivalry and selective attention.
- ▷ Sensory input alone does not give rise to consciousness; higher-level interpretation of that input is needed.

## 18 Organization of Cognition

### 18.1 Functionally Related Areas of Cortex Lie Close Together

- ▷ The cortex of each cerebral hemisphere is a continuous sheet of gray matter.
- ▷ At the coarsest level, it consists of five lobes, with each lobe further subdivided.
- ▷ Functional areas are distinguished by cellular structure, connectivity, and the physiological response properties of neurons.
- ▷ Precepts that govern the organization of functional areas in the macaque (old world monkey) cerebral cortex:
  1. All areas fall into a few major functional groups.
  2. Areas in a given category occupy a discrete, continuous portion of the cortical sheet.
  3. Functionally related areas occupy neighboring sites.

### 18.2 Sensory Information is Processed in the Cortex in Serial Pathways

- ▷ Cortical areas communicate with each other through bundles of axons traveling together in identifiable tracts.
- ▷ **Primary sensory areas** possess four properties characteristic of their role in the early stages of information processing:
  - Inputs from thalamic sensory relay nuclei.
  - Neurons in a primary sensory area have small receptive fields and are arranged to form a precise somatotopic map of the sensory receptor surface.
  - Injury to a part of the map causes a simple sensory loss confined to the corresponding part of the contralateral sensory receptor surface.

- Connections to other cortical areas are limited, mostly to nearby areas that process information in the same modality.
- ▷ Higher-order sensory areas have a different set of properties important to their role in the later stages of information processing:
  - Inputs arise from other thalamic nuclei and lower-order areas of sensory cortex instead of sensory relay nuclei.
  - Large receptive fields and imprecise maps of the array of receptors in the periphery.
  - Injury results in abnormalities of perception, but does not impair ability to detect sensory stimuli.
  - Connected to distant areas in the frontal and limbic nodes as well as nearby unimodal areas.
- ▷ Sensory information is processed serially, but not exclusively; higher-order areas project back to lower-order areas which can modulate the activity of neurons in lower-order areas.
- ▷ **Association cortex:** regions of the cortex where injury causes cognitive deficits that cannot be explained by impairment of sensory or motor function alone.
- ▷ Large regions of association cortex are contained within each of the four lobes:
  - **parietal:** critical for sensory guidance of motor behavior and spatial awareness.
  - **temporal:** recognition of sensory stimuli and for storage of semantic (factual) knowledge.
  - **frontal:** key role in organizing behavior in working memory.
  - **limbic:** complex functions related to emotion and episodic (autobiographical) memory.
- ▷ Association areas have much more extensive input and output connections than do lower-order sensory and motor areas.

- ▷ All association lobes are densely interconnected network of pathways.

### **18.3 Goal-Directed Motor Behavior Is Controlled in the Frontal Lobe**

- ▷ All areas of the frontal lobe participate in the control of motor behavior and are connected in a series of functional hierarchy.
- ▷ Neuronal activity in the premotor cortex, adjacent to the primary motor cortex, reflects global aspects of motor behavior.
- ▷ Dorsolateral prefrontal cortex contributes to cognitive control of behavior.
- ▷ The orbital-ventromedial prefrontal cortex, connected to the dorsolateral prefrontal cortex (then premotor), is involved with emotional processes associated with executive control of behavior.
- ▷ Information flows from higher-order areas in the frontal lobe to primary order cortex, contrasting sensory's periphery first flow.
- ▷ Prefrontal cortex is important for the executive control of behavior.
- ▷ The orbital-ventromedial prefrontal cortex is linked strongly to the hypothalamus and amygdala, receives input from every sensory system, and projects to the dorsolateral prefrontal cortex.
- ▷ Thus, the above pathway allows for response to emotional and sensory inputs and allows the trigger of appropriate behavior.

### **18.4 Limbic Association Cortex is a Gateway to the Hippocampal Memory System**

- ▷ The limbic (limbus—edge) association cortex forms a ring that is visible in the medial view of the hemisphere.
- ▷ Previously it was thought to make up an entire system in combinations with other areas, but some divisions of the limbic lobe have other functions, with some not yet well understood.



## 19 Functions of the Premotor Systems

### 19.1 Direct Connections Between the Cerebral Cortex and Spinal Cord Play a Fundamental Role in the Organization of Voluntary Movements

- ▷ Individual muscles and joints are represented in the cortex multiple times in a complex mosaic.
- ▷ Each muscle joint is represented by a column of neurons whose axons branch and terminate in several functionally related spinal motor nuclei.
- ▷ Movement can also be elicited by stimulation of premotor areas.
- ▷ Neurons in the primary motor cortex fire in connection with a variety of goal-directed movements.
- ▷ There are three pathways from the premotor and motor areas to the motor neurons in the spinal cord: a direct corticospinal and two indirect, the medial and lateral brain stem systems.
- ▷ The pathways together make up the **corticospinal system**.
- ▷ **Medial brain stem system**: receives information from the cortex and other motor centers for the control of posture and locomotion.
- ▷ **Lateral brain stem system**: similar to medial but is involved in control of arm and hand movements.
- ▷ Reflex circuits can generate stereotyped movements without descending commands; new patterns can be generated through direct action on motor neurons.
- ▷ The cortical motor areas receive feedback from the cerebellum and basal ganglia in order for smooth execution of skilled movements in motor learning.



## 19.2 The Four Premotor Areas of the Primate Brain Also Have Direct Connections in the Spinal Cord

- ▷ **Lateral ventral premotor area:** concerned the "what" in visual preception and controls mostly mouth and hand movements.
- ▷ **Lateral dorsal premotor area:** concerned with the "where" in visual preception and controls direction or movements.
- ▷ **Supplementary motor area:**
- ▷ **Cingulate motor areas:** a group of areas in the cingulate sulcus.
- ▷ The four areas are connected to the primary motor cortex.

## 19.3 Motor Circuits Involved in Voluntary Actions are Organized to Achieve Specific Goals

- ▷ The parietal lobe contains more than one representation of space and each one is dependent on motor activity.
- ▷ Neurons that respond to objects in peripersonal space are located mostly in the inferior parietal lobe where hand and mouth movements are represented, whereas neurons that respond to futher away objects are found where eye movements are represented.

## 19.4 The Hand Has a Critical Role in Primate Behavior

- ▷ Investigation of the **anterior intraparietal area (AIP)** show that neurons fall into three main classes: motor-dominant, visual-dominant, and visual-motor combination.
- ▷ Futher suggestion shows that these neurons are involved in transferring sensory representations of objects into motor representations.
- ▷ **Canonical neurons:** neurons that fire in response to visual observation (and actual grasping) of graspable objects of certain size, shape, and orientation.
- ▷ Canonical neurons are thought to translate objects physical properties into *potential motor acts*.

## 19.5 The Join Activity of Neurons in the Parietal and Premotor Cortex Encodes Potential Motor Acts

- ▷ Studies of the parietal and premotor canonical neurons show that some neurons encode the possibilities for interaction with an object.
- ▷ **Mirror neurons:** discharge during specific motor acts, but also fire when the individual observation action being done by another.
- ▷ Mirror neurons may help us understand the intention of others.
- ▷ Potential motor acts are suppressed or released by motor planning centers.
- ▷ Neurons in the supplementary motor area are involved in the planning, generation, and control of sequential motor actions.
- ▷ After long periods of practice, when the behavior becomes automatic, activity in the presupplementary motor area ceases.

## 20 Functional Imaging of Cognition

### 20.1 Functional Imaging Reflects the Metabolic Demand of Neural Activity

- ▷ A large amount of neuron's total energy metabolism, about one-half, is devoted to maintaining resting potential.
- ▷ The other half is for other biochemical processes, including all molecular reactions for normal function.
- ▷ fMRI responses have been highly correlated with neural spiking.
- ▷ Though, some fMRI measurements fail to capture subthreshold modulatory inputs.

Interesting chapter, but each section had one-two random notes. Omitted due to lack of conclusiveness.

## 21 Sensory Coding

### 21.1 Psychophysics Relates the Physical Properties of Stimuli to Sensations

- ▷ **Psychophysics:** relationship between the physical characteristics of a stimulus and attributes of sensory experience.
- ▷ **Sensory physiology:** examination of neural consequences of a stimulus.
- ▷ **Sensory threshold:** the lowest stimulus strength a subject can detect.
- ▷ Sensory thresholds can be altered by emotional or psychological factors.
- ▷ Sensations are quantified using probabilistic statistics.
- ▷ Reaction times are correlated with cognitive processes.

### 21.2 Physical Stimuli are Represented in the Nervous System by Means of the Sensory Code

- ▷ Neural coding of sensory information is better understood at the early stages than later.
- ▷ Sensory receptors are responsive to a single type of stimulus energy.
- ▷ **Stimulus transduction:** the time it takes to convert a stimulus response into an electrical signal.
- ▷ Multiple subclasses of sensory receptors are found in each sense organ.
- ▷ There are rapid and slowly adapting sensors that illustrate a major principal of decoding: *contrast*.
- ▷ The timing of action potentials between neurons has a profound effect on long-term potentiation and depression at synapses.
- ▷ The receptive field of a sensory neuron conveys spatial information.
- ▷ Fragmentation of a stimulus into component parts, each encoded by an individual neuron, is the initial step in sensory processing.

### 21.3 Modality-Specific Pathways Extend to the Central Nervous System

- ▷ Activity of sensory neurons are more variable than that of neurons in the periphery.
- ▷ Central sensory neurons fire irregularly before and after stimulation, even during times of no stimulation.
- ▷ The variability is a result of: alertness, attention, previous experience, and recent activation by similar stimuli.
- ▷ The receptor surface is represented topographically in central nuclei.
- ▷ Feedback regulates sensory coding and top-down learning mechanisms influences sensory processing.

## 22 Somatosensory System

- ▷ The somatosensory system serves three major functions:
  - **Proprioception:** the sense of oneself. Skeletal muscle, joint capsules, and the skin allow for awareness of our own body.
  - **Exteroception:** the sense of direct interaction with the external world. Touch, contact, pressure, stroking, temperature, pain, motion, vibration are used to identify objects.
  - **Interoception:** sense of major organ systems of the body and its internal state. Most information does not appear conscious, but plays a major role. Primarily consist of chemoreceptors.
- ▷ All somatic senses are mediated by the dorsal root ganglion neurons.

### 22.1 The Primary Sensory Neuron of the Somatosensory System are Clustered in the Dorsal Root Ganglia

- ▷ Dorsal root ganglion neurons are pseudo-unipolar cells.
- ▷ The central branches terminate in the spinal cord or brain stem, forming the first synapses in somatosensory pathways.
- ▷ **Primary afferent fiber:** the axon of each dorsal root ganglion cell serves as a single transmission line from receptor to central nervous system.
- ▷ **Peripheral nerves:** individual primary afferent fibers group that are grouped together, and also include motor axons innervating nearby muscles, blood vessels, glands, or viscera.

### 22.2 Peripheral Somatosensory Nerve Fibers Conduct Action Potentials at Different Rates

- ▷ Difference in peripheral nerve's diameter and conduction velocity mediate somatic sensation.
- ▷ Larger diameter tends to relay faster, not accounting for degree myelinated fibers.

- ▷ Electrical stimulation of whole nerves is also used to classify peripheral nerve fibers.
- ▷ **Compound action potential:** summed action potential of all nerve fibers excited by a stimulus pulse and is roughly proportional to the total number of active nerve fibers.
- ▷ The conduction velocity throughout the nervous system is correlated with the need to maintain synchrony.

### 22.3 Many Specialized Receptors Are Employed by the Somatosensory System

- ▷ The receptor class expressed in the nerve terminal of a sensory neuron determines the type of stimulus detected.
- ▷ Mechanoreceptors mediate touch and proprioception.
- ▷ The skin has eight types of mechanoreceptors that are responsible for touch.
- ▷ Proprioceptors measure muscle activity and joint positions.
- ▷ Nociceptors mediate pain.
- ▷ Thermal receptors detect changes in skin temperature.
- ▷ Itch is a distinctive cutaneous sensation.
- ▷ Visceral sensations represent the status of various internal organs.

## 23 Touch

### 23.1 Active and Passive Touch Evoke Similar Responses in Mechanoreceptors

- ▷ Active and passive modes of tactile stimulation excite the same population of receptors in the skin and evoke similar responses in afferent fibers.
- ▷ Passive touch is used for naming objects or describing sensations.
- ▷ Active touch is used when the hand manipulates objects.

### 23.2 The Hand Has Four Major Types of Mechanoreceptors

- ▷ **Merkel cell**: tips of epidermal sweat ridges; detects edges, points; slow adaption to sustained indentation.
- ▷ **Meissner corpuscle**: close to skin surface; detects lateral motion; no adaption to sustained indentation.
- ▷ **Ruffini ending**: located in dermis; senses skin stretching; slow adaption to sustained indentation.
- ▷ **Pacinian corpuscle**: located deep in dermis; senses vibration; no adaption to sustained indentation.
- ▷ Receptive fields define the zone of tactile sensitivity.
- ▷ There are two types of receptive fields: one with highly specialized fields; the other with broader fields with a central hotspot.
- ▷ Slowly adapting fibers detect object pressure and from.
- ▷ Rapidly adapting fibers detect motion and vibration.
- ▷ Combination of slow and rapidly adapting fibers contribute to grip control.

### 23.3 Tactile Information is Processed in the Central Touch System

- ▷ Cortical receptive fields integrate information from neighboring receptors.



- ▷ Neurons in the somatosensory cortex are organized into functionally specialized columns.
- ▷ All neurons within a column receive inputs from the same local area of the receptor sheet and respond to the same class(es) of receptors.
- ▷ Columns share a common center that is clearly evident in layer IV.
- ▷ Horizontal connections within layers II and III link neurons in neighboring columns, sharing information when activated by the same stimulus.
- ▷ Cortical columns are organized somatotopically.
- ▷ **Cortical magnification:** the amount of cortical area devoted to a unit of area of skin. This varies by more than a hundredfold across different body surfaces.

## 24 Pain

### 24.1 Noxious Insults Activate Nociceptors

- ▷ Most nociceptors are simply the free nerve endings of primary sensory neurons.
- ▷ There are three main classes of nociceptors:
  - Thermal
  - Mechanical
  - Polymodal— high-intensity mechanical, chemical, or thermal.
- ▷ There is a less understood fourth class: silent nociceptors.
- ▷ The three main classes are widely distributed in the skin and deep tissues and are often coactivated.
- ▷ Silent nociceptors are found in viscera; activated by inflammation and various chemical agents.
- ▷ **Allodynia**: pain in response to stimuli that are normally innocuous.
- ▷ **Hyperalgesia**: an exaggerated response to noxious stimuli, typically persistent even in absence of sensory stimulation.
- ▷ **Nociceptive pain**: activation of nociceptors and normally from accompanying inflammation.
- ▷ **Neuropathic pain**: direct injury to nerves in peripheral or central nervous system and is accompanied by burning or electric sensation.

## 25 Visual Processing

### 25.1 Visual Perception is a Constructive Process

- ▷ The brain guesses at scene presented to the eyes based on past experience.
- ▷ The modern view of perception is based on the gestalt psychology—the perceptual interpretation we make of any visual object depends not just on the properties of the stimulus, but also the context.
- ▷ An important step in object recognition separating figures from the background.
- ▷ The brain analyzes a scene at three levels:
  - Low: local contrast, orientation, color, and movement.
  - Intermediate: analysis of the layout, surfaces, parsing global contours, and depth.
  - High: object recognition.
- ▷ Motion, depth, form, and color occur in a unified percept due to interacting neural pathways.

### 25.2 Form, Color, Motion, and Depth are Processed in Discrete Areas of the Cerebral Cortex

- ▷ Visual areas of the cortex can be differentiated either by a visuotopic map, or by functional properties of the neurons.
- ▷ Visual areas are organized into two hierarchical pathways: ventral, involved in object recognition; and dorsal, dedicated to the use of visual information guiding movements.
- ▷ Pathways are interconnected so that information is shared and each connection is reciprocal— each area sends information back to areas from which it receives input.
- ▷ The shared connections provide information about cognitive functions, spatial attention, stimulus expectation, and emotional content, to earlier

levels of visual processing.

### 25.3 The Receptive Fields of Neurons at Successive Relays in an Afferent Pathway Provide Clues to How the Brain Analyzes Visual Form

- ▷ **On-center:** cells that fire when a spot of light is turned on within a circular central region.
- ▷ **Off-center:** cells that fire inversely to on-center.
- ▷ If both cells are stimulated with diffuse light, then there is little to no response. This allows them to distinguish borders and contours very well and leads to the encoding of contrast.
- ▷ **Eccentricity:** size of the retina's receptive field, which varies in relative to the fovea and the position of neurons along the visual pathway.

### 25.4 The Visual Cortex is Organized into Columns of Specialized Neurons

- ▷ The dominant feature of the functional organization of the primary visual cortex is the visuotopic organization of the of its cells: the visual field is systematically represented across the surface of the cortex.
- ▷ Columns reflect the functional role of that area in vision.
- ▷ Orientation and ocular dominance columns have embedded clusters of neurons that have strong color preferences.
- ▷ These clusters specialize to provide information about surfaces rather than edges.
- ▷ **Serial processing:** processing in successive connections between cortical areas that run from the back of the brain forward.
- ▷ **Parallel processing:** occurs simultaneously in subsets of fibers that process different submodalities such as from, color, movement.

## 25.5 Intrinsic Cortical Circuits Transform Neural Information

- ▷ Each area of the visual cortex transforms information gathered by the eyes.
- ▷ Principal input to the primary visual cortex comes from two parallel pathways that originate in the parvocellular and magnocellular layers of the lateral geniculate nucleus.
- ▷ Neurons in different layers have distinctive receptive-field properties, with superficial layers have smaller fields while deeper layers tend to have larger ones.
- ▷ Feedback projections are thought to provide a means where higher centers in a pathway can influence lower ones.
- ▷ Feedback projection is still largely unknown.

## 26 Low-Level Visual Processing

### 26.1 The Photoreceptor Layer Samples the Visual Image

- ▷ Ocular optics limit the quality of the retinal image.
- ▷ The density of photoreceptors, bipolar cells, and ganglion cells is highest at the fovea(center of eye).
- ▷ There are two types of photoreceptors: rods and cones.
- ▷ Rods: very sensitive—low light, no color.
- ▷ Cones: less sensitive—for daylight, multiple types, faster response time.
- ▷ Central fovea has an absence of rods.

### 26.2 Ganglion Cells Transmit Neural Images to the Brain

- ▷ Optic nerve has only 1% as many axons as there are receptor cells, so the retinal circuit must edit information before it is conveyed to the brain.
- ▷ The two major ganglion cells are binary, ON or OFF cells.
- ▷ Many ganglion cells fire regardless of current lighting condition, but ON cells fire more rapidly with increasing light, while OFF slows or stops. The inverse is also true when going from light to dark.
- ▷ Many ganglion cells respond strongly to edges in the image.
- ▷ Output produced by ganglion cells enhance regions of contrast, while reducing homogenous illumination.
- ▷ Ganglion output also emphasizes temporal changes in stimuli through transient (burst response) and sustained (steady) neurons.
- ▷ Retinal output emphasizes moving objects.
- ▷ Several ganglion cell types project to the brain through parallel pathways.
- ▷ About 20 ganglion cells have been described, which allow the optic nerve to convey about 20(?) different representations of the world based on polarity

(on/off, fine/coarse, sustained/transient, motion, spectral filtering... and more?)

### 26.3 A Network of Interneurons Shapes the Retinal Output

- ▷ Parallel pathways originate in bipolar cells.
- ▷ Most retinal processing is accomplished with graded membrane potentials via the *ribbon synapse*.
- ▷ Action potentials occur only in certain amacrine and ganglion cells.
- ▷ Stimulus representation in ganglion cell population originates in dedicated bipolar cell pathways that are differentiated by their selective connections to photoreceptors and postsynaptic targets.
- ▷ Spatial filtering is accomplished by lateral inhibition.
- ▷ Amacrine cells are axonless neurons with dendrites that ramify in the inner plexiform layer, generally producing an inhibitory network.
- ▷ Temporal filtering occurs in synapses and feedback circuits.
- ▷ Color vision begins in cone-selective circuits.
- ▷ Rod and cone circuits merge in the inner retina.

## 27 Intermediate-Level Visual Processing

### 27.1 Internal Models of Object Geometry Help the Brain Analyze Shapes

- ▷ **Visual Primitives:** local features in a visual scene: contrast, line orientation, brightness, color, movement, and depth.
- ▷ In the visual cortex neurons respond selectively to lines of particular orientations, which reflects the arrangement of inputs from cells in the lateral geniculate.
- ▷ There are two types of orientation-selective neurons:
  - **Simple cells:** receptive fields are divided into binary subregions, firing when an light enters the ON region.
  - Simple cells are cells highly selective for the position of a line or edge in space.
  - **Complex cells:** lack discrete binary subregions. Instead they fire continuously as line or edge stimulus traverses.
- ▷ Visual cortex neurons do not respond to an image that is stabilized on the retina.
- ▷ **Contextual modulation:** visual cortex neuron that is modulated by stimuli outside the receptive field's core, allowing for complex selectivity.

### 27.2 Depth Perception Helps Segregate Objects from Background

- ▷ **Plane of fixation:** the point where corresponding positions are displayed on both retinas.
- ▷ Visual cortex neurons can be selective for objects lying on, in front, or behind the plane of fixation.
- ▷ Depth is also determined through monocular cues such as size, perspective, occlusion, brightness, and movement.



### **27.3 Local Movement Cues Define Object Trajectory and Shape**

- ▷ The primary visual cortex determines the direction of movement of objects.
- ▷ Determining the direction of motion requires resolving multiple cues.
- ▷ An important determinant of perceived direction is scene segmentation.

### **27.4 Context Determines the Perception of Visual Stimuli**

- ▷ The visual system attempts to measure the surface characteristics of objects by comparing the light arriving from different parts of the visual field.
- ▷ Transient and variable stimulation of the retina construct representations of a stable, three-dimensional world.
- ▷ Color and brightness depend heavily on nearby contextual cues.
- ▷ Receptive-field properties also depend on context.
- ▷ Contextual influences are pervasive in contour integration, scene segmentation, and determination of object shape and surface properties.

### **27.5 Cortical Connections, Functional Architecture, and Perception are Intimately Related**

- ▷ Intermediate-level visual processing requires sharing information from throughout the visual field.
- ▷ Cortical circuits make horizontal connections that link distant locations in the visual field suggest connections between orientation columns of similar specificity have a role in contour integration over a large area of the visual cortex.
- ▷ Perceptual learning requires plasticity in cortical connections.
- ▷ Perceptual learning involves repeating a discrimination task many times, does not require error feedback, and improvement manifests itself as decrease in threshold in discrimination of smaller differences.

- ▷ Visual search relies on the cortical representation of visual attributes and shapes.
- ▷ Certain objects pop out from others in a complex image due to visual system processes in parallel pathways—features of target and the features of surrounding distractors.
- ▷ Cognitive processes influence visual perception and segmentation, such as top-down spatial attention, visual expectation, or the particular perceptual task at hand.

## 28 High-Level Visual Processing

### 28.1 The Inferior Temporal Cortex is the Primary Center for Object Perception

- ▷ Object perception is the nexus between vision and cognition, and is what high-level visual processing is concerned with.
- ▷ The hierarchy of synaptic relays in the cortical system extend from the primary visual cortex to the temporal lobe.
- ▷ Temporal lobe is a site of convergence of many types of visual information.
- ▷ Clinical evidence identifies the inferior temporal cortex as essential for object recognition.
- ▷ Neurons in the inferior temporal cortex encode complex visual stimuli, such as specific patterns or features.
- ▷ Neurons in the inferior temporal cortex are functionally organized in columns.
- ▷ Face-selective cells constitute a highly specialized class of neurons.
- ▷ Object recognition is intertwined with visual categorization, visual memory, and emotion; all of which the inferior temporal cortex contribute to.

### 28.2 Object Recognition Relies on Perceptual Constancy

- ▷ For object recognition to take place, various invariant attributes must be represented independently of other image properties. Doing this correctly is termed **Perceptual constancy**
- ▷ Perceptual constancy has many forms, from size, position, rotation, viewing angle, likeness, and more.
- ▷ A more general type of constancy is the perception of objects as belonging to the same semantic category.

### **28.3 Visual Memory is a Component fo High-Level Visual Processing**

- ▷ Visual memory can influence the processing of new incoming visual information.
- ▷ Implicit visual learning leads to changes in the selectivity of neuronal responses.
- ▷ Explicit visual learning depends on linkage of the visual system and declarative memory formation.
- ▷ The hippocampus and neocortical areas of the medial temporal lobe are essential both for the acquisition of visual associative memories and for the functional plasticity of the inferior temporal cortex.

### **28.4 Associative Recall or Visual Memories Depends on Top-Down Activation of the Cortical Neurons that Process Visual Stimuli**

- ▷ Sensory experience of an image and recall of the same image are subjectively similar.
- ▷ Sensory experience depends on bottom-up flow of visual information, while the latter is top-down flow.
- ▷ The distinction is important, but under normal conditions both signal pathways collaborate to yield visual experience.

## 29 Visual Processing and Action

### 29.1 Successive Fixations Focus Our Attention in the Visual Field

- ▷ A saccade usually lasts less than 40 ms, occurs several times per second, and redirects center of sight in the visual field.
- ▷ Attention selects objects for further visual examination.
- ▷ Voluntary attention is closely linked to saccades.
- ▷ Activity in the parietal lobe correlates with attention paid to objects.
- ▷ The locus of attention can be ascertained only by examining the entire salience map and choosing its peak; it cannot be identified by monitoring activity at one point alone.

### 29.2 The Visual Scene Remains Stable Despite Continual Shifts in the Retinal Image

- ▷ Neurons shift their receptive fields from one part of the visual field to another before the saccade occurs.
- ▷ The parietal neurons must have advanced information about the saccade; either from feedback from the peripheral proprioceptors in the eye muscles directly (unlikely) or from the motor system that controls the movement of the eyes.

### 29.3 Vision Lapses During Saccades

- ▷ An object can be seen during a saccade if it is moving as fast and in the same direction of the eye.
- ▷ Visual scene is thought to be blurred, but not unconscious during a saccade.
- ▷ Visual masking may hide the lower contrast, blurred image if two images are close enough, which would happen often during a saccade.

- ▷ Extravisual input, such as corollary discharge, must also be present. This would reduce sensitivity during a saccade if the input is not there (*Saccadic suppression*).
- ▷ Saccadic suppression due to corollary discharge is most pronounced for stimuli with low spatial frequency and high contrast, which are most effective for activating neurons in the magnocellular pathway of the visual system.
- ▷ There is evidence that visual masking and corollary discharge must act together to reduce disruption of vision during saccades.