

Lab Practical One

Week 1: Methods and Organelles

Background	3
Microscopic Techniques	4
Organelles and Cytoplasmic Inclusions	7
Labeled Diagram of a Eukaryotic Cell	9
Mitotic Phases	10
Apoptosis Events	12
Ross and Paulina Atlas Plates	13

Week 2: Epithelial Connective Bone Cartilage

Epithelial Tissue	16
Examples	16
Review	17
Connective Tissue Proper	19
Examples	19
Review	20
Cartilage and Bone	22
Examples	22
Review	22
Bone Examples	23

Week 3: Adipose Blood Muscle Nerve

Adipose	26
White Adipose	26
Brown Adipose	27
White Adipose Examples	28
Brown Adipose Examples	29
Blood	30
Muscle	32
Skeletal Muscle	32
Cardiac Muscle	34
Colon	35
Uterus	37
Central Nervous System	38
Spinal Cord	38
Cerebellum	40
Sympathetic Nerve	41
Enteric Nervous System	42

Week 4: Cardiovascular

The Heart	43
------------------	-----------

Blood Vessels	45
Arteries	45
Veins	47
Arterioles and Venules	48
Capillaries	49
Continuous Capillaries	49
Fenestrated Capillaries	50
Sinusoidal (Discontinuous) Capillaries	51

Week 1: Methods and Organelles

Background

- **Describe the steps in tissue preparation for microscopy:**

- **Fixation:** typically the first step in the preparation of histological sections in where tissues samples are treated with fixatives as a means to preserve and protect from biological decay.
 - **Fixatives:** solutions, compounds, or others means meant to either disable degradative enzymes, induce cross-linking (stabilizing proteins), or protect from extrinsic damage.
- **Embedding:** the process of placing tissues in a harder medium (e.g., paraffin and plastic resins) as a means to allow for thin slicing of tissue.
 - Embedding occurs later in the process of preparation; once a tissue is fixed it must first undergo a series of steps:
 - **Dehydration:** the removal of water using ethanol.
 - **Clearing:** replacement of an organic solvent miscible with both alcohol and the embedding medium, giving a translucent appearance.
 - **Infiltration:** evaporation of the clearing solvent via exposure to heat (50–60 °C) promoting the final embedding of tissue into the medium.
- **Staining:** used as a means to increase contrast in tissue or specific features of tissue that are of interest as most biology tissue has very little inherent contrast.
 - **Basophilic:** dyes that have an affinity for **anionic** (net negative charge) cells parts.
 - E.g., hematoxylin, toluidine blue, alcian blue, and methylene blue.
 - **Acidophilic:** dyes that have an affinity for **cationic** (net positive charge) cell parts.
 - E.g., eosin, orange G, and acid fuchsin

- **What does H & E Stain?**

- **Hematoxylin (H)** and **eosin (E)** stains are among the most commonly used stains.
 - As mentioned above, hematoxylin acts as a **basophilic dye**, turning negatively charged organelles like the cell nucleus, RNA-rich regions of cytoplasm, cartilage, anywhere from **blue → purple**.
 - Eosin acts as an **Acidophilic dye**, typically turning **cationic structures pink**; sometimes it is considered to be a **counterstain**, i.e., typically a secondary dye that is meant to distinguish features.

- **What does PAS Stain?**

- **Periodic acid-Schiff (PAS)** utilizes hexose rings of polysaccharides and other carbohydrate rich structures to stain macromolecules **purple → magenta**.

- **Describe Enzyme Histochemistry.**

- Enzyme histochemistry is a method for localizing cellular structures using specific enzymatic activity in such structures.
- Preservation of enzymes often requires non-fixed or mildly fixed tissue and generally adhere to the following steps:
 1. Tissues sections are immersed in solution containing the substrate of the enzyme to be localized.
 2. The enzyme is exposed to and allowed to act on the substrate.
 3. A marker compound is introduced and reacted with the product from step 2.
 4. Location is determined via precipitation of the insoluble product, which must be visible a light or electron microscopy, over the site of the enzyme.
- Phosphatase, dehydrogenase, and peroxidase are common examples of enzymes detected with histochemistry.

- **How does Immunohistochemistry work?**

- Immunohistochemistry (IHC): the use of labeled antibodies and antigens to identify and localize many proteins and macromolecules that lack specific enzymatic activity.
- Visualization of such interactions are commonly accomplished with either:
 - Chromogenic immunohistochemistry (CIH): use of antibodies conjugated to an enzyme that catalyzes a color-producing reaction.
 - Immunofluorescence: tagging of a fluorophore (fluorescein, rhodamine) to an antibody.
- Common used in diagnosis of abnormal cells such as those in cancerous tumors.

Microscopic Techniques


- **Bright field:** a very common method that uses ordinary light and stained structures to discern differences and cell structures.




- **Phase contrast:** uses differences in refractive index of natural cell and tissue components; allows for observation of living cells.



- **Confocal:** use of scans at successive focal planes with a more focused light beam, usually from a laser, to construct a 3D image.



- **Fluorescent:** use of longer wavelengths emitted via specific perturbations of cellular substances using a specific wavelength (usually UV). Useful for identification of cells and components that have affinity for specific fluorescent compounds.



- **Transmission Electron Microscopy (TEM)**: a high resolution (3 nm) that allows for particles to be magnified many times via transmission of electrons through a specimen; very thin tissue sections are used and a flat image is made from the intersection of the electron beam and the sample.




- **Scanning Electron Microscopy (SEM)**: used to provide a high resolution view of the surface of cells, tissues, and organs. Unlike TEM, SEM does not intersect with the specimen, instead the electrons are reflected by a coating and then collected, analyzed, and used to generate a 3D view.




Organelles and Cytoplasmic Inclusions


- **Nucleus:** a large membrane bound organelle that contains chromatin, the nucleolus, and nucleoplasm.
Size: **5–20 μm** , the largest organelle.




- **Nucleolus:** large, dense structure within the nucleus that functions in the synthesis of ribosomes.
Size: **0.5–5 μm** , smaller than nucleus.




- **Plasma Membrane:** phospholipid bilayer containing various elements, acts as a physical barrier to regulate internal environment; maintains charge and functions in cell communication.
Size: **5–10 nm**, very thin—3 orders of magnitude less than width of nucleus.




- **Rough ER:** interconnect membrane that modifies, transports, and stores proteins produced by attached ribosomes.
Size: **5–8 nm thick, 20–30 nm wide**, membrane of rER thin, while the lumen is wide and surrounds nucleus (often).




- **Smooth ER:** Like rER, but lacking ribosomes; synthesizes, transports, and stores lipids; metabolizes carbohydrates, toxins (of various types); and forms vesicles and peroxisomes.
Size: lumen is **30–60 nm** thick, larger than rER; overall fraction of cell varies.




- **Golgi Apparatus:** modifies, packages, and sorts materials that arrive from the ER; forms secretory vesicles/lysosomes.
Size: **2–5 mm**, stacks of flat membranes, can occupy large portion of cell.




- **Vesicles:** transports cellular material
Size: 30–100 nm, large range varies among different types of vesicles, still relatively small.




- **Mitochondria:** *the powerhouse of the cell.*
Size: 0.5–1 μm, decently sized, but still smaller than other organelles in the cell.




- **Endosomes:** collection of intracellular sorting organelles, originating from trans Golgi network that move molecules and ligands to lysosomes (L), or recycled back to cell membrane.
Size: 1–50 μm, depends on stage; (E)rly forms dynamic tubular network, can be long. (M)vbs (late) lack tubes.
TEM, not light microscope below*




- **Lysosomes:** spherical-shaped from Golgi, contains digestive enzymes to break down microbes and materials.
Size: 0.5–1 μm, rather small, TEM used below instead of light microscope.



- **Peroxisomes:** formed via ER or fission; contains enzymes to break down specific harmful substances, also used for beta oxidation of fatty acids.
Size: 0.1–1 μm, typically smaller than lysosomes, can be similar sized.




- **Cytoskeleton:** organized network of {actin, micro-, intermediate} filaments, and microtubules and other proteins.
Size: 7 nm, varies, depending on structure and filaments used.




- **Ribosomes:** composed of protein and rRNA and engage in protein synthesis; can be on rER, in plasma membrane, in lysosomes, and free in cell.

Size: 20–30 nm, varies, organized into both large and small subunits.




- **Glycogen granules:** aggregates of carbohydrate polymer in which glucose is stored, notably in liver cells

Size: 20–30 μm, many can cluster.




- **Lipid droplets:** accumulations of lipid-filling adipocytes.



Size: 20–40 nm to 100 μm, can be removed, can take up majority of cell.




Labeled Diagram of a Eukaryotic Cell



Mitotic Phases







- **Interphase:** The long period between mitosis (the G_1 , S , and G_2 phases) that takes place before division is able to occur again.
- The **mitotic phase** consists of:
 - **Prophase** ± 1 hr: the first stage where the nucleolus disappears, replicated chromatin condenses and each sister chromatid joins with the centromere.
 - **Prometaphase:** not always presented as a distinct section from prophase, instead it meant to describe processes typically found in later parts of prophase—i.e., where nuclear lamina and pore complexes disperse in cytoplasmic membrane vesicles—and early parts of metaphase—i.e., the attachment to kinetochores, which help with alignment, and other processes.
 - **Metaphase** < 1 hr: further condensation of chromosomes, the cell is more spherical, and microtubules move the chromosomes towards the equatorial plate.
 - **Anaphase** $\leq \frac{1}{2}$ hr: the chromosomes separate towards opposite spindle poles and reach maximum condensation.
 - **Telophase** (minutes): the final stage, where effects of prophase and prometaphase are reversed, and a contractile ring of actin filament forms to allow for constriction and thus cytokinesis to occur.

Apoptosis Events


- **DNA fragmentation:** endonucleases are activated, cleaving DNA into fragments.




- **Decrease of cell volume:** destruction of the cytoskeleton and chromatin causing dense (dark purple stained) pyknotic nuclei.



- **Membrane blebbing:** shrinking of the cell causes the plasma membrane to undergo dramatic shape changes, producing “blebbing” structures as proteins are degraded and lipid mobility increases.




- **Formation of apoptotic bodies (and phagocytic removal):** membrane-bound remnants of cytoplasm and nucleus into even smaller fragments, forming apoptotic bodies. Phospholipids on these bodies induce the phagocytosis by neighboring cells (or white blood cells).




Ross and Paulina Atlas Plates


- **Stratified squamous epithelium (plate 2)**: deeper cells, especially those in the basal layer, are small with little cytoplasm making the nuclei appear closely packed.
 - Larger cells flatten out and form disc-like squames (squamous; square-flat).
 - The deeper cells may be columnar or cuboidal.
 - No intracellular space and flexibility around size allows for constant abrasion; useful for mouth, esophagus, and vagina.




- **Simple cuboidal epithelium (plate 2)**: a single layer of flat of cuboidal (cube-like) cells, with large spherical central nuclei.
 - Being on the surface layer allows cells perform secretion and absorption.
 - Found on surface of ovaries, lining of nephrons, walls of renal tubules, and parts of the eye, thyroid, and salivary glands.




- **Skeletal muscle (plate 22)**: one of the three major muscle types that forms striated muscle tissue and generally under control of the somatic nervous system.
 - Below is a cross-section (left) of a muscle fascicle and a longitudinal-section (right) of a gluteraldehyde.
 - Muscle fibers (MF) exhibit polygonal shape and vary only slightly in width.
 - Muscle fiber nuclei (MFN) are embedded within the extreme periphery of the fiber.
 - Fibroblast nuclei (FN) lie outside the muscle fiber; they both smaller and more dense than the MFN.
 - Capillaries (C) are present between the muscle fibers.



- **Cardiac muscle (plate 24)**: another major muscle type that constitutes the main tissues of the walls of the heart; it is involuntary, striated muscle.
 - Below are a longitudinal-section (1) and cross-sections (2), each with an increased resolution respectively.
 - Intercalated discs (ID) very pronounced cross-bands, if they are stained for; they oppose cell-to-cell contacts which promotes end-to-end alignment of cells.
 - Connective tissue cells (CT): surrounds bundles of fibers, and more delicate CT surrounds individual MF.



- **Smooth muscle (plate 26):** the last of the major muscle types; it is non-striated, divided into two subgroups (unitary and multi-unitary).
 - Found in hollow organs such as the stomach, intestines, urinary bladder, and uterus. Also found in passageways such as in arteries and veins, as well as tracts in the respiratory, urinary and reproductive system.
 - In the eye its found as ciliary muscle, a specific type of smooth muscle that dilates and contracts the iris.
 - Smooth muscle (SM) in the skin also helps hair stain erect.
 - Below are cross-sectional bundles (CS) as well as longitudinally sections bundles (LS).
 - Dense connective tissue (DCT) separates the smooth muscle bundles, while dense irregular connective tissue (DICT) help bundle smooth muscle into circular profiles.
 - High resolution shows smooth muscle cells (SMC), which wavy forms here indicate partial contraction.




Week 2: Epithelial Connective Bone Cartilage


Epithelial Tissue

Examples


- Simple squamous epithelium ↑




- Cell nuclei
- Basement membrane (BM)




- Stratified squamous epithelium






- Cell nuclei
- BM



- Ciliated pseudostratified columnar epithelium (PsEp)



- Goblet c.
- Cilia
- BM



- Simple cuboidal epithelium




- BM (maybe?)




- **Simple columnar epithelium**



- Microvillous border



- Goblet cells



Review


- **Simple squamous:** wider than their height, hence flat and scale like (squamous).
 - Locations: in mouth, esophagus, blood vessels (endothelium), alveoli of lungs, lymphatic vessels, lining of cavities (mesothelium).
 - Function: facilitates movements of viscera via diffusion and filtration, active transport by pinocytosis, secretion of molecules and lubricating substances.
- **Simple cuboidal:** cells with height and width that are approximately the same, hence cuboidal.
 - Location: covering the ovary and thyroid, in kidney tubules, and other secretory portions of small glands.
 - Function: covering, secretion, absorption.
- **Simple columnar:** cells that are taller than they are wide, i.e., column shaped, hence columnar.
 - Location: ciliated tissues are in bronchi, uterine tubes, and uterus; smooth line intestine and gallbladder.
 - Function: protection, lubrication, absorption, and secretion.
- **Pseudostratified columnar:** cells with nuclei that appear at different heights, leading the pseudo impression of stratified columnar cells when viewed in cross-section.
 - Location: ciliated tissues lines the trachea and much of the upper respiratory tract, including the nasal cavity.
 - Function: protection, secretion (mostly mucus), cilia-mediated transport of particles trapped in mucus.

- **Stratified squamous:** like simple squamous, but multilayered. Cells generally become more squamous as they become more apical.
 - Location: lines the esophagus, mouth, vagina, anal canal, larynx.
 - Function: multi layers protects against abrasion, prevents water loss.
- **Stratified cuboidal:** again, same as simple cuboidal, but multi layered.
 - Location: sweat, salivary, and mammary glands, also in developing ovarian follicles.
 - Function: Protection, secretion.
- **Stratified columnar:** multilayered columnar...
 - Location: the male urethra and the ducts of some glands, and the conjunctiva (mucus membrane in front eye and inside eyelid).
 - Function: protection, secretion of mucus.
- **Transitional:** cells that can change from squamous to cuboidal, depending on amount of tension in the epithelium.
 - Location: bladder, ureters, urethra, and renal calyces (chambers in kidney through which urine passes).
 - Function: protection, distensibility (ability to swell from inside), stretch.


Connective Tissue Proper

Examples


- Dense regular connective tissue:




- Collagenous tissue




- Loose connective tissue




- Nuclei




- Matrix




- Elastin fibers




- Collagen fibers




- Reticular tissue




- Reticular fibers




- Collagen (type I)



- Lymph node reticular fibers




- Muscle tendon junctions




- Collagen I



- Fibroblasts



- Thin skin



- Thick collagenous fibers type I



Review


- **Mesenchyme:** connective tissue found mostly during embryonic development and is generally undifferentiated, sparse, and uniformly distributed in the matrix with sparse collagen fibers.
 - Location: mesodermal layer of early embryo, develops into tissues of lymphatic and circulatory system; can migrate easily. Organized into adherent sheets.
 - Function: contains stem/progenitor cells for all adult connective tissue cells.
- **Areolar:** or loose connective tissue, is the most widely distributed connective tissue type, has much ground substance (gel like), many cells, little collagen, and randomly distributed.
 - Location: can be found in the skin and in places that connect the epithelium to other tissues; found beneath the dermis layer and under epithelial tissue of body systems that have external openings. Also surrounds blood vessels and nerves and a component of mucus membranes in the digestive, respiratory, reproductive and urinary systems.
 - Function: supports microvasculature, nerves, immune defense cells, holds organs in place, and serves as reservoir of water and salts.

- **Loose CT reticular:** a network of reticular fibers (synthesized by fibroblasts called reticular cells) made up of type III collagen.
 - Location: bone marrow, spleen, liver, kidney, lymph nodes (not thymus), adrenal glands.
 - Function: supports blood-forming cells, many secretory cells, and lymphocytes in most lymphoid organs.
- **Dense regular:** filled with mostly parallel bundles of collagen, few fibroblasts, is aligned with collagen, and very strong. Slow to heal due to poor blood supply.
 - Location: tendons, ligaments, aponeuroses (layers of flat broad tendons), and corneal stroma.
 - Function: provide strong connections within musculoskeletal system and strong resistance to force, especially in one direction.
- **Dense elastic fibers:** an essential component of the extracellular matrix composed of bundles of protein (elastin) produced by fibroblasts, endothelial, smooth muscle, and airway epithelial cells. Can stretch many times their length and return to back to original length.
 - Location: found in the skin, lungs, arteries, veins, connective tissue proper, elastic cartilage, periodontal ligaments, fetal tissues and more; basically anything that must undergo stretching.
 - Function: provides the elastic properties to connective tissues.
- **Dense irregular:** fibers not arranged in parallel bundles as in dense regular connective tissue. Still consists of mostly collagen fibers, but has less ground substance than loose connective tissue. Fibroblasts are scattered sparsely across the tissue.
 - Location: found mostly in the reticular layer of the dermis, but also in the sclera in the deeper skin layers. Found in the submucosal layer of the digestive tract.
 - Function: mainly provides strength, similar to regular tissue. Protects organs and helps resist tearing.


Cartilage and Bone

Examples

- **Hyaline cartilage**




- Incomplete rings???
- Outer Perichondrium




- Inner Perichondrium
- Chondrocytes



- **Elastic cartilage**



- **Fibrocartilage**




Review

- **Hyaline Cartilage:** homogeneous with type II collagen, most common of the cartilage types. Covered externally by fibrous membrane known as perichondrium.
 - Location: components of the upper respiratory tract; in the larynx, trachea, bronchi, sternal ends of the ribs, and articular ends and epiphyseal plates of long bones.
 - Function: gives structures (particularly respiratory) a definite but pliable form, low friction surface to joints.


- **Elastic Cartilage:** similar to hyaline cartilage, but contains many yellow elastic fibers in a solid matrix.
 - Location: external ear, external acoustic meatus (ear canal), auditory tube; epiglottis (leaf shape flap in throat), and more.
 - Function: provides flexible shape and support of soft tissues.
- **Fibrocartilage:** a mixture of white fibrous tissue and cartilaginous tissue in various proportions; contains type I collagen in addition to type II.
 - Location: many joints (shoulder, hip, knee), as well secondary joints (pubic symphysis, manubriosternal joint), intervertebral discs, and insertions of tendons.
 - Function: provides cushioning, tensile strength, and resistance to tearing and compression.
- **Bone:** calcified connective tissue also consisting of cells, fibers, and ground substances. Structurally rigid using calcium phosphate salts within its matrix. There are multiple types of bone, i.e., cortical (hard, outer) and cancellous (trabecular, spongy, internal).
 - Location: like, everywhere. Spooky Scary Skeletons.
 - Function: provides solid support for the body, protects organs, encloses internal cavities, serve as reservoir of calcium, phosphate, and other ions, form systems of levers that multiply forces generated during skeletal muscle contraction, and transforms energy into bodily movement.

Bone Examples


- **Ground bone cross-section and longitudinal-section**



- Osteons canal (HC)




- Lacunae (L)




- Axial canals




- Canaliculi ↑



- Decalcified bone




- Compact bone




- Spongy bone




- Osteoblasts




- Osteoclasts (Ocl)




- Growth at the epiphyseal plate




- Resting zone




- Zone of proliferation





- Zone of hypertrophy



- Zone of calcification




- Zone of ossification



Week 3: Adipose Blood Muscle Nerve


Adipose

White Adipose



- Location: subcutaneous layer, mammary gland, greater omentum, mesenteries, retroperitoneal space, visceral pericardium, orbits, bone marrow cavity.
- Function: energy storage, insulation, cushioning, hormone production, source of metabolic water.
- Adipocyte Morphology: unilocular, spherical, flatten nuclear, rim of cytoplasm, large diameter (15–150 µm)
- Factors inducing differentiation: PPAR- γ /RXR
- UCP-1 gene expression: None
- Mitochondria: Few, poorly developed
- Response to environmental stress (cold): Decreased lipogenesis, increased lipoprotein lipase activity.
- Growth and differentiation: entire life, starting from stromal-vascular cells


Brown Adipose




- Location: large amounts in newborn; in adults: retroperitoneal space, deep cervical and supraclavicular regions of neck, interscapular, paravertebral regions of the back, mediastinum.
- Function: thermogenesis.
- Adipocyte Morphology: multiocular, spherical, round eccentric nucleus, smaller diameter (10–25 µm).
- Factors inducing differentiation: PRDM16/PGC-1
- UCP-1 gene expression: Yes (unique to brown fat)
- Mitochondria: many, well-developed.
- Response to environmental stress (cold): increased lipogenesis, decreased lipoprotein lipase activity.
- Growth and differentiation: only during fetal period, decreases in adult life, except those with pheochromocytoma and hibernoma.

White Adipose Examples


- **Adipocyte:** the largest component of loose connective tissue.




- **EM of adipose (blue)**




- **Reticular Fibers (purple)**




- **Small blood vessel (red)**



- **Nerve Fiber (cyan)**




- **Nuclei**




Brown Adipose Examples


- Brown adipose tissue



- Brown adipocyte



- Contrast between white unilocular and brown multilocular adipocyte




Blood


- **RBC (erythrocytes):** light red, biconcave discs without nuclei that carry oxygen and return CO₂. Size: $\approx 7.5 \mu\text{m}$



- **Platelet:** small, basophilic discs that help prevent bleeding via blood clotting. Size: 2–4 μm



- **Eosinophil:** contain distinctive large, eosinophilic granules and usually nuclei with two lobes. Aids in anti-parasitic and bacterial activity, allergic reactions, and modulating inflammation.




- **Neutrophil:** have distinctive nuclei with 2 to 5 lobes and azurophilic granules that form essential part of innate immune system with various functions.




- **Basophil:** rare cells with distinctive large, basophilic granules. Aids immune responses by releasing cytokines, leukotrienes, and histamine.



- **Monocyte:** large cells with "kidney-shaped" or notched nuclei. Precursors of macrophages and other cells of the mononuclear phagocyte system.




- **Lymphocyte:** occur in a range of sizes (small:left, large:right). Have many functions relating to immune system, as well as subtypes (T and B), helps produce antibodies.




Muscle

Skeletal Muscle


- Tendon



- Skeletal Muscle:




- Skeletal Muscle Cross-Section




- Fibroblasts Nuclei and Sacromere




- Sarcomere (Em)



- Fascicles (F)




- **Perimysium:** thick layer of connective tissue that surrounds a group of muscle cells to form fascicles. Blood and nerves are found in this layer.




- **Endomysium:** thin layer of connective tissue that surrounds each muscle cell.




- **Epimysium:** dense connective tissue that surrounds the entire muscle and is usually continuous with a tendon.



- **Satellite Cells:** occur on the surface of muscle cells






- **Capillaries:** a collapsed capillary containing flattened red blood cells is present on the left edge of the center muscle cell.





Cardiac Muscle

- **Cross-striations:** cardiac muscles cells have rounded cross-sections ($< 25 \mu\text{m}$) with a centrally located nucleus.
- **Branched cells:** cells are joined end-to-end and are often branched





- **Longitudinal section:** cardiac muscle cells are smaller than skeletal muscle.

- **Straight intercalated disks:** specialized junctions that connect the individual cells




- **Stepped intercalated disks**




Colon


- Smooth muscle in small intestine:



- Inner layer: individual cells vary in diameter depending on their location within the cell. Cross-sections through the middle of cells have centrally located nuclei, usually surrounded by an unstained region.




- Outer layer: in relaxed smooth muscle, the nuclei are elongated with rounded ends. When contracted, the nuclei spiral, kink, or twist. The cytoplasm is pink, unstriated and with little detail.





Uterus

- **Myometrium (left) Fascicles of smooth muscle (right)**




- **Blood vessels within myometrium**




Central Nervous System

Spinal Cord


- Gray matter




- Ventral structure




- White matter




- Motor neurons in anterior horn




- Dorsal structure




- Perikaryon, Nucleus of ventral horn cell, and neuropil



- Surrounding meninges




- Central canal of spinal cord




Cerebellum


- Cerebellum




- Molecular (Mol) and granular (Gr) layer




- Granular nuclei count



- Molecular nuclei count



- Purkinje cells: large neurons found at the interface between the two layers of the cerebellum.




- Extension of Purkinje axons:




Sympathetic Nerve


- **Satellite cells:** glial cells with small nuclei at the periphery of nerve cell bodies.




- **Ganglion Cells:** large, nerve cell bodies with prominent nuclei and nucleoli.



- **Nerves**




- **Lipofuscin:** yellow-brown pigment located in the cytoplasm. It is end-stage lysosomes that accumulates with age.




Enteric Nervous System


- **Mucosa**




- **Epithelium**




- **Submucosa:** dense irregular connective tissue.




- **Meissner's Plexus:** provides secretory innervation of goblet cells and motor innervation of the muscularis mucosae.



- **Muscularis Externa:** two orthogonal layers of smooth muscle.




- **Auerbach's Plexus:** provides motor innervation of the muscularis externa.





Week 4: Cardiovascular

The Heart


- **Epicardium:** outer surface of mesothelial cells supported by dense irregular connective tissue.
- **Right Atrium:** receives venous blood from the systemic circulation.



- **Myocardium:** thickest, middle layer of cardiac muscle responsible for the pumping action of the heart; contains blood vessels, nerves and adipose cells.



- **Endocardium:** inner surface of a simple squamous epithelium supported by a thin layer of dense irregular connective tissue.




- **Collagen fibers (Fibrous rings):** support heart valves and contain bands of collagen and elastic fibers that appear dark red.


- **Intercalated discs:** thin, dark stained linear structures dividing adjacent cells that are perpendicular to the direction of the muscle fiber




- **Pectinate muscles:** bundles of muscle that protrude from the surface that are common in the right atrium.



- **Ventricle:** receives blood from the right atrium and pumps it into the lungs via the pulmonary arteries.



- **Ventricle myocardium:** receives blood from the right atrium and pumps it into the lungs via the pulmonary arteries.




- **Is the myocardium thicker in the atrium or ventricle? Why?**

- The **ventricles'** myocardium are thicker, since it does more work **pumping** blood, as the atrium mainly **receives** blood and only pumps it to the ventricles. Since pumping increases size of muscles, then the myocardium housing said muscles, will be larger.


Blood Vessels

Arteries


- **Elastic Artery (Carotid Artery)**: conveys blood from the heart to systemic and pulmonary circulations.




Tunica intima





Tunica media




Tunica adventitia



- **Vasa Vasorum:** blood vessels that supply the tunica adventitia and tunica media.
- **Nerves in adventitia:** responsible for regulation of the contraction of the smooth muscle.



- **Comparison of Muscular Artery (Top) and Medium Vein (Bottom):**



Veins


- **Large Vein (Brachiocephalic Vein)**: has thinner walls than arteries with less distinct layers, as well as the adventitia instead of media being the thickest portion.




Tunica intima




Tunica media




Tunica adventitia




Vasa Vasorum



Nerves




White Adipose




Arterioles and Venules


- **Arteriole:** small diameter blood vessels that allow blood to flow into and out of capillary beds, respectively.
- **Venules:** small diameter blood vessel that allows blood to return from capillary beds to veins.



- **Two layers of smooth muscle:**



- **Arterioles vs. Venules**



Capillaries


Continuous Capillaries

- Continuous capillaries have many, tight well-developed occluding junctions between slightly overlapping endothelia cells. They are the most common capillary that provide well-regulated metabolic exchange across the cells.

Longitudinal




Cross-section




Fenestrated Capillaries

- Similar to continuous, but with endothelia cells that are penetrated by numerous small circular openings. Found in organs with rapid interchange of substances between tissue and the blood. The pore can only be seen with electron microscopy.

SEM




TEM



Sinusoidal (Discontinuous) Capillaries

- These capillaries allow for maximal exchange of macromolecules as well as easier movement of cells between tissues and blood.
- The endothelium has large perforations without diaphragms and irregular intercellular clefts.
- Unlike other capillaries, sinusoids have highly discontinuous basement membranes and larger diameters, slowing blood flow.

Network of sinusoids



Sinusoid surrounded by endocrine cells

