Calculus

Limits and Continuity

Limits

Limit | Thomas' Calculus (2.2–2.4)

- **Limit** $\lim\limits_{x\to c}$: the value of a function (or sequence) approaches as the input (or index) approaches some value (informal definition)
  - Limits are used to define continuity, derivatives, and integrals

Limits of a Functions and Sequences

Limit of a function | Limit of a sequence

- **Limit of a function**: a fundament concept in calculus and analysis concerning the behavior L of a function near a particular input c, i.e.,

$$\lim_{x\to c} f(x) = L$$

  - Reads as "f of x tends to L as x tends to c"
- ε, δ **Limit of function**: a formalized definition, wherein $f(x)$ is defined on an open interval, except possibly at c itself, leading to above definition, if and only if:
  - For every real measure of closeness $\varepsilon > 0$, there exists a real corresponding $\delta > 0$, such that for all existing further approaches there exist a smaller ε, i.e.,

$$f : \mathbb{R} \to \mathbb{R},\ c, L \in \mathbb{R} \Rightarrow \lim_{x\to c} f(x) = L$$
$$\Updownarrow$$
$$\forall \varepsilon > 0\, (\exists \delta > 0 : \forall x, 0 < |x - c| < \delta \Rightarrow |f(x) - L| < \varepsilon)$$

- Functions do not have a limit when the function:
  - has a unit step, i.e., it "jumps" at a point;
  - is not bounded, i.e., it tends towards infinity;
  - or does not stay close to any single number, i.e., it oscillates too much.
- **Limit of a sequence**: the value that the terms of a sequence (x_n) "tends to" (and not to any other) as n approaches infinity (or some point), i.e.,

$$\lim_{n\to\infty} x_n = x$$

- ε **Limit of sequence**: for every measure of closeness ε, the sequence's term eventually converge to the limit, i.e.,

$$\forall \varepsilon > 0\, (\exists N \in \mathbb{N}\, (\forall n \in \mathbb{N}\, (n \geq N \Rightarrow |x_n - x| < \varepsilon)))$$

  - **Convergent**: when a limit of a sequence exists.
  - **Divergent**: a sequence that does not converge.

Properties of Limits

List of limits | Squeeze theorem

- **Operations on a single known limit**: if $\lim_{x\to c} f(x) = L$ then:
  - $\lim_{x\to c}[f(x) \pm \alpha] = L \pm \alpha$
  - $\lim_{x\to c} \alpha f(x) = \alpha L$
  - $\lim_{x\to c} f(x)^{-1} = L^{-1}, L \neq 0$
  - $\lim_{x\to c} f(x)^{n} = L^{n}, n \in \mathbb{N}$
  - $\lim_{x\to c} f(x)^{n^{-1}} = L^{n^{-1}}, n \in \mathbb{N}, \text{ if } n \in \mathbb{N}_e \Rightarrow L > 0$
- **Operations on two known limits**: if $\lim_{x\to c} f(x) = L_1$ and $\lim_{x\to c} g(x) = L_2$
  - $\lim_{x\to c}[f(x) \pm g(x)] = L_1 \pm L_2$
  - $\lim_{x\to c}[f(x)g(x)] = L_1 L_2$
  - $\lim_{x\to c} f(x)g(x)^{-1} = L_1 {L_2}^{-1}$
- **Squeeze theorem**: used to confirm the limit of a function via comparison with two other functions whose limits are easily known or computed.
  - Let I be an interval having the point a as a limit point.
  - Let g, f, and h, be functions defined on I, except possibly at a itself.
  - Suppose that $\forall x \in I \wedge \neq a \Rightarrow g(x) \leq f(x) \leq h(x)$
  - And suppose that $\lim_{x\to a} g(x) = \lim_{x\to a} h(x) = L$
  - Then, $\lim_{x\to a} f(x) = L$
  - Essentially, the hard to compute limit of the "middle function" is found by finding two other easy functions that "squeeze" the middle function at that point.

One-Sided Limit

One-Sided Limit

-

Continuity

- Sources:

  ### Continuity at a Point

  -

  ### Continuous Functions

  -

  ### Intermediate Value Theorem

  -

Limits Involving Infinity

- Sources:

  ## Limits at Infinity

  -

  ## Infinite Limits

  -

Derivatives

Integrals