

Calculus



Limits and Continuity

Limits	2
Limits of a Functions and Sequences.....	2
Properties of Limits.....	3
One-Sided Limit.....	3
Continuity	4
Continuous Functions	4
Intermediate Value Theorem.....	4
Extreme Value Theorem	4
Limits Involving Infinity	5
Limits at Infinity.....	5
Infinite Limits	5

Derivatives

Applications of Derivatives

Integrals

Applications of Integrals

Transcendental Functions

Techniques of Integration

Infinite Sequences and Series

First-Order Differential Equations

Parametric Equations and Polar Coordinates

Vectors and Vector-Valued Functions

Partial Derivatives

Multiple Integrals

Vector Calculus

Second-Order Differential Equations

Limits and Continuity



Limits

🌐 Limit 🌐 | Thomas' Calculus (2.2–2.4) 🌐

- **Limit** $\lim_{x \rightarrow c}$: the value of a function (or sequence) approaches as the input (or index) approaches some value (informal definition)
 - Limits are used to define **continuity** ↓, **derivatives** ↓, and **integrals** ↓

Limits of a Functions and Sequences

🌐 Limit of a function 🌐 | Limit of a sequence 🌐

- **Limit of a function**: a fundamental concept in calculus and analysis concerning the behavior L of a function near a particular input c , i.e.,

$$\lim_{x \rightarrow c} f(x) = L$$

- Reads as “ f of x tends to L as x tends to c ”
- ϵ, δ **Limit of function**: a formalized definition, wherein $f(x)$ is defined on an open interval I , except possibly at c itself, leading to above definition, if and only if:
 - For every real measure of **closeness** $\epsilon > 0$, there exists a real **corresponding** $\delta > 0$, such that for all existing further approaches there exist a smaller ϵ , i.e.,

$$f : \mathbb{R} \rightarrow \mathbb{R}, c, L \in \mathbb{R} \Rightarrow \lim_{x \rightarrow c} f(x) = L$$



$$\forall \epsilon > 0 (\exists \delta > 0 : \forall x \in I (0 < |x - c| < \delta \Rightarrow |f(x) - L| < \epsilon))$$

- Functions do not have a limit when the function:
 - has a unit step, i.e., it “jumps” at a point;
 - is not bounded, i.e., it tends towards infinity;
 - or does not stay close to any single number, i.e., it oscillates too much.
- **Limit of a sequence**: the value that the terms of a sequence (x_n) “tends to” (and not to any other) as n approaches infinity (or some point), i.e.,

$$\lim_{n \rightarrow \infty} x_n = x$$

- ϵ **Limit of sequence**: for every measure of closeness ϵ , the sequence’s term eventually converge to the limit, i.e.,

$$\forall \epsilon > 0 (\exists N \in \mathbb{N} (\forall n \in \mathbb{N} (n \geq N \Rightarrow |x_n - x| < \epsilon)))$$

- **Convergent**: when a limit of a sequence **exists**.
- **Divergent**: a sequence that **does not** converge.

Properties of Limits

📌 List of limits 📌 | Squeeze theorem 📌

- **Operations on a single known limit:** if $\lim_{x \rightarrow c} f(x) = L$ then:
 - $\lim_{x \rightarrow c} [f(x) \pm \alpha] = L \pm \alpha$
 - $\lim_{x \rightarrow c} \alpha f(x) = \alpha L$
 - $\lim_{x \rightarrow c} f(x)^{-1} = L^{-1}, L \neq 0$
 - $\lim_{x \rightarrow c} f(x)^n = L^n, n \in \mathbb{N}$
 - $\lim_{x \rightarrow c} f(x)^{n^{-1}} = L^{n^{-1}}, n \in \mathbb{N}, \text{ if } n \in \mathbb{N}_e \Rightarrow L > 0$
- **Operations on two known limits:** if $\lim_{x \rightarrow c} f(x) = L_1$ and $\lim_{x \rightarrow c} g(x) = L_2$
 - $\lim_{x \rightarrow c} [f(x) \pm g(x)] = L_1 \pm L_2$
 - $\lim_{x \rightarrow c} [f(x)g(x)] = L_1 L_2$
 - $\lim_{x \rightarrow c} f(x)g(x)^{-1} = L_1 L_2^{-1}$
- **Squeeze theorem:** used to confirm the limit of a function via comparison with two other functions whose limits are easily known or computed.
 - Let I be an interval having the point a as a limit point.
 - Let g, f , and h , be functions defined on I , except possibly at a itself.
 - Suppose that $\forall x \in I \wedge x \neq a \Rightarrow g(x) \leq f(x) \leq h(x)$
 - And suppose that $\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L$
 - Then, $\lim_{x \rightarrow a} f(x) = L$
 - Essentially, the hard to compute limit of the “middle function” is found by finding two other easy functions that “squeeze” the middle function at that point.

One-Sided Limit

📌 One-Sided Limit 📌

- **One-sided limit:** one of two limits of $f(x)$ as x approaches a specified point from either the **left** or from the **right**.
 - From the **left**: $\lim_{x \rightarrow c^-} = L$
 - From the **right**: $\lim_{x \rightarrow c^+} = L$
- If the left and right limits exist and are equal, then

$$\lim_{x \rightarrow c} f(x) = L \Leftrightarrow \lim_{x \rightarrow c^-} f(x) = L \wedge \lim_{x \rightarrow c^+} f(x) = L$$

- Limits can still exist, even if the function is defined at a different point, as long as both one-sided limits approach the same value near the given input.

Continuity

🌐 Continuous function 🌐 | Classification of discontinuities 🌐 | Thomas' Calculus (2.5) 🌐

- Continuity of functions is one of the core concepts of topology, however, there are definitions in terms of limits that prove useful; the following is only a primer.

Continuous Functions

- **Continuous function:** a function that does not have any abrupt changes in value.
 - I.e., a function is continuous if and only if arbitrarily small changes in its output can be assured by restricting to sufficiently small changes in its input.
- **Discontinuous:** when a function is not continuous at a point in its domain, leading to a discontinuity; there are three classifications:
 - **Removable:** when both **one-sided limits** \uparrow exist, are finite, and are equal, but the actual value of $f(x)$ is not equal to the limit and equal to some other value.
 - The discontinuity can be removed to regain continuity.
 - Sometimes the term *removable discontinuity* is mistaken for *removable singularity*, or a "hole" in the function (the point is not defined elsewhere).
 - **Jump:** when a single limit does not exist because the one-sided limits exist and are finite, but not equal; points can be defined at the discontinuity, but the function can not be made continuous.
 - **Essential:** when at least one of two one-sided limits doesn't exist; can be the result of oscillating or unbounded functions.

Intermediate Value Theorem

🌐 Intermediate Value Theorem 🌐

◦

Extreme Value Theorem

🌐 Intermediate Value Theorem 🌐

◦

Limits Involving Infinity

- Sources:

Limits at Infinity

-

Infinite Limits

-

Derivatives



Applications of Derivatives



Integrals



Applications of Integrals



Transcendental Functions



Techniques of Integration



Infinite Sequences and Series



First-Order Differential Equations



Parametric Equations and Polar Coordinates



Vectors and Vector-Valued Functions



Partial Derivatives



Multiple Integrals



Vector Calculus



Second-Order Differential Equations

