Covasim | Candidate hypotheses

67 nodes / 454 edges

Upward: solid | Downward: dashed | Same scale: dotted

Meso
N28 Exposures generated per silent infectious person-step by group
N47 Layer-level degree suppression ratio under quarantine
N12 Within-layer degree dispersion
N37 Per-index contact load within a layer
N19 Household dyad attributed secondary exposure count
N13 Layer consecutive active-step run length
N62 Layer attributed-exposure rate ratio after versus at the trough
N54 Layer share of attributed new exposure transitions
N38 Cross-layer dispersion of contact load
N63 Age-band exposure rate ratio after versus at the trough
N55 Age-band exposure share drift from first active step
N3 Age-band exposure hazard per susceptible person-step
N21 Fraction of household dyads with any attributed exposure
N5 Cross-band dispersion of exposure hazard
N11 Layer exposure incidence per contact edge
N53 Age-band share of new exposure transitions
N36 Layer known-contact incidence per contact edge
N27 Silent share of infectious person-steps by quarantine-status group
N4 Age-band cumulative attack ratio
N20 Dispersion of attributed exposures across household dyads
N46 Dispersion of contact degree among quarantined agents
Micro
N45 Agent full-contact-suppression indicator under quarantine
N60 Agent exposure-step offset from the city infectious trough
N33 Agent known-contact state indicator
N52 Agent exposure-to-diagnosis interval
N25 Agent cumulative silent-infectious duration
N35 Agent quarantine state indicator
N43 Agent contact degree ratio in and out of quarantine
N1 Agent new-exposure transition indicator
N17 Agent source-attributed exposure indicator
N26 Agent symptomatic-but-undiagnosed indicator
N61 Agent exposure transition after the infectious trough
N10 Agent per-layer contact weight load
N16 Agent degree in the persistent household layer
N34 Agent known-contact to quarantine latency
N8 Agent contact degree within a contact layer
N18 Agent household-attributed exposure indicator
N2 Agent cumulative ever-exposed indicator
N9 Agent exposure transition attributed to a layer
N44 Agent absolute contact-degree change under quarantine
N24 Agent silent-infectious indicator
N51 Agent first observed exposure step index
Macro
N30 Silent share of current infectious agents
N7 Exposure-share versus susceptible-person-time mismatch
N6 Age-band attack ratio range
N42 Known-contact prevalence share
N15 Normalised entropy of layer exposure shares
N67 Share of layers with post-trough incidence increase
N64 Mean post-trough log growth of infectious agents
N65 Post-trough rebound amplitude of infectious count
N50 Population-level quarantine contact suppression ratio
N49 Share of quarantined agents at or below the non-quarantined median degree
N66 Susceptible share at the infectious trough
N29 New diagnoses per infectious person-step
N41 Quarantine coverage of observed agent-steps
N40 Mean per-agent tracing latency at quarantine start
N39 Known-contact transitions per new diagnosis
N32 Mean recorded symptom-to-diagnosis interval
N56 Total-variation drift of age composition of exposures
N57 Total-variation drift of layer composition of attributed exposures
N59 Median age of agents with new exposure transitions
N23 Gini concentration of household dyad attributed exposures
N22 Household share of all observed exposure transitions
N31 Cumulative never-diagnosed to ever-diagnosed infection ratio
N14 Layer edge-set Jaccard turnover
N48 Relative step-to-step change in total contact volume
N58 Contrast of exposure growth before and after first quarantine onset