

3.2 Exploratory analysis of a theoretical model

Sean Bittner and John Cunningham

September 1, 2019

Dynamical models with two populations (excitatory (E) and inhibitory (I) neurons) of visual processing have been used to reproduce a host of experimentally documented phenomena in V1. When an inhibition stabilized network (ISN, the I population stabilizes an otherwise unstable E population), these models exhibit the paradoxical effect [1], selective amplification [2], surround suppression [3], and sensory integrative properties [4]. Since almost all I neurons fall into one of three classes (parvalbumin (P)-, somatostatin (S)-, and vasointestinal peptide (V)-expressing neurons) [5, 6], theoretical neuroscientists look to extend these dynamical models to four populations [7]. A current challenge in theoretical neuroscience is understanding the distributed role of inhibition stabilization across these inhibitory subtypes.

These four populations exhibit neuron-type specific connectivity (Fig. 1A) [8], in which some populations do not project to others. Since S and V are the only populations that mutually inhibit each other, a popular conceptualization is that S and V have winner-take-all dynamics. In fact, evidence in mice suggests that V silences S when presented with large stimuli, and S silences V for small stimuli [9]. Here, we use DSNs to understand the possible sources of inhibition stabilization in this V1 model, when either S or V is inactive, selecting the weight matrix parameters as the free parameters of the DSN. The behavior of the DSN sampled models is constrained to produce two things: 1.) a mean-zero distribution of ISN coefficients $\gamma(W) = 1 - f'(f^{-1}(r_E(W)))W_{EE}$ with some variance, and 2.) α -population silencing $r_\alpha(W) = 0$, for $\alpha \in \{S, V\}$. When $\gamma < 0$ the network is ISN, and not ISN otherwise. Constraining the DSN behavior to a zero-mean distribution of ISN coefficients gives us samples of both ISN and non-ISN networks, optimized to have greatest variety of stabilization motifs.

The DSN posteriors are blah blah blah




Figure 1: A.) Model of primary visual cortex (V1) Neurons: E - excitatory (black), P - parvalbumin- (blue), S - somatostatin- (red), and V - VIP-expressing (green). Parameters: weights of the dynamics matrix W . B.) The DSNs are conditioned on population-silenced ISN criticality. C.) DSN distribution of the parameters of the V1 model conditioned on population-silenced ISN criticality. D.) Eigenmodes of the hessian of each DSN ordered by eigenvalue. E.) Behavioral sensitivity of the model along each mode of the hessian.

References

- [1] Misha V Tsodyks, William E Skaggs, Terrence J Sejnowski, and Bruce L McNaughton. Paradoxical effects of external modulation of inhibitory interneurons. *Journal of neuroscience*, 17(11):4382–4388, 1997.
- [2] Brendan K Murphy and Kenneth D Miller. Balanced amplification: a new mechanism of selective amplification of neural activity patterns. *Neuron*, 61(4):635–648, 2009.
- [3] Hirofumi Ozeki, Ian M Finn, Evan S Schaffer, Kenneth D Miller, and David Ferster. Inhibitory stabilization of the cortical network underlies visual surround suppression. *Neuron*, 62(4):578–592, 2009.
- [4] Daniel B Rubin, Stephen D Van Hooser, and Kenneth D Miller. The stabilized supralinear network: a unifying circuit motif underlying multi-input integration in sensory cortex. *Neuron*, 85(2):402–417, 2015.
- [5] Henry Markram, Maria Toledo-Rodriguez, Yun Wang, Anirudh Gupta, Gilad Silberberg, and Caizhi Wu. Interneurons of the neocortical inhibitory system. *Nature reviews neuroscience*, 5(10):793, 2004.
- [6] Bernardo Rudy, Gordon Fishell, SooHyun Lee, and Jens Hjerling-Leffler. Three groups of interneurons account for nearly 100% of neocortical gabaergic neurons. *Developmental neurobiology*, 71(1):45–61, 2011.
- [7] Ashok Litwin-Kumar, Robert Rosenbaum, and Brent Doiron. Inhibitory stabilization and visual coding in cortical circuits with multiple interneuron subtypes. *Journal of neurophysiology*, 115(3):1399–1409, 2016.
- [8] Carsten K Pfeffer, Mingshan Xue, Miao He, Z Josh Huang, and Massimo Scanziani. Inhibition of inhibition in visual cortex: the logic of connections between molecularly distinct interneurons. *Nature neuroscience*, 16(8):1068, 2013.
- [9] Mario Dipoppa, Adam Ranson, Michael Krumin, Marius Pachitariu, Matteo Carandini, and Kenneth D Harris. Vision and locomotion shape the interactions between neuron types in mouse visual cortex. *Neuron*, 98(3):602–615, 2018.