



题目




O=C(C=C1OC)C(C)=CC1=O>C=C/C=C/C.CCl[Sn](Cl)(Cl)Cl>[A], 反应于 -20°C 进行



O=C(C=C1OC)C(C)=CC1=O>C=C/C=C/C.FB(F)(O(CC)CC)F>[B], 反应于 -20°C 进行


请分别给出以上两个反应最有可能得到的产物A和B（不考虑立体化学）

A.



O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O,产物A


产物A



O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O,产物B


产物B

B.



O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O,产物A


产物A



O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O,产物B


产物B

C.



O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O,产物A


产物A



O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O,产物B


产物B

D.



O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O,产物A

产物A



O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O,产物B

产物B

答案

正确答案: **D**

详细解析

SnCl_4 可以同时与羰基和甲氧基络合, 从而活化下面的羰基

CHECKPOINT

1 PTS

SnCl_4 可以同时与羰基和甲氧基络合, 从而活化下面的羰基

而 BF_3 只能四配位, 优先与上方富电子羰基络合, 活化上方的羰基


CHECKPOINT

1 PTS

而 BF_3 只能四配位, 优先与上方富电子羰基络合, 活化上方的羰基


对双烯体而言, 烷基给电子, 因此与烷基连接一侧为正电性, 另一端为负电性, 与亲双烯体的相应位点反应可以得到产物A和B:

A.



O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O,产物A

B.



O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O,产物B

因此选项D正确。

CHECKPOINT

1 PTS

产物A为O=C(C=C1OC)C(CC=CC2C)(C)C2C1=O

CHECKPOINT

1 PTS

产物B为O=C(C=C1OC)C(C(C)C=CC2)(C)C2C1=O