

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/333951232>

Quick calculation of TWO-WAY and ONE-WAY ANOVA (Experiment finding paper)

Experiment Findings · June 2019

DOI: 10.13140/RG.2.2.35083.75041

CITATIONS

0

READS

4,069

1 author:



[Aung Pyae](#)

Asia Pacific University of Technology and Innovation

9 PUBLICATIONS 1 CITATION

SEE PROFILE

Quick calculation for TWO-WAY and ONE-WAY ANOVA

(Experiment finding paper)

Experiment finding paper about calculation of one-way and two-way ANOVA in fast and effective way using a scientific calculator. The paper starts explaining from two-way ANOVA because it may difficult or complex to study one-way ANOVA without knowing two-way first. But if the researcher knows clearly about two-way, one-way is also clearly known and no need to add any new knowledge to study one-way.

The two-way ANOVA compares the mean differences between groups that have been split on two independent variables (called factors). The primary purpose of a two-way ANOVA is to understand if there is an interaction between the two independent variables on the dependent variable. It is very easy to calculate ANOVA using computer, but it is incredibly complex with number by hand. If some numerical calculation is wrong in complex steps like finding variance, the whole calculation answer will go wrong. However, ANOVA calculation needs to use scientific calculator and this method is an effective and good also for time consuming.

One-way ANOVA is to determine whether there are any statistically significant differences between the means of two or more independent groups. This method is used when there is a minimum of three or rather than two groups.

Key ideas

ANOVA is a statistical procedure for analyzing the results of an experiment with multiple groups. It is the extension of similar procedures for A/B test, used to access whether the overall variation among groups is within the range of chance variation. A useful outcome of an ANOVA is the identification of variance components associated with group treatments, interaction effects and errors.

All calculator instructions are made using **Canon F-789SGA** scientific calculator.



Canon F-789SGA scientific calculator

Shop here: <http://tiny.cc/348h8y>

Please note that all calculator calculation instructions made in this paper are done with above illustrated calculator brand. Button and instructions may vary with other brands but generally all scientific calculator has similar instructions.

Contents

TWO-WAY ANOVA	3
ONE-WAY ANOVA	11
Two-way ANOVA in SPSS	18
One-way ANOVA in SPSS	22
Two-way ANOVA in SAS	26
One-way ANOVA in SAS	29
Datasets	32

TWO-WAY ANOVA

Assumptions

1. The populations from which the samples were obtained must be normally or approximately normally distributed.
2. The samples must be independent.
3. The variances of the populations must be equal.
4. The groups must have the same sample size.

Two-way ANOVA calculation steps

- Step 1: Define hypothesis
- Step 2: Find the means for Row and Column
- Step 3: Frame the ANOVA summary table
- Step 4: Calculate DF (Degree of freedom)
- Step 5: Calculate SS (Sum of squares)
- Step 6: Calculate MS (Mean squares)
- Step 7: Calculate F (F value)
- Step 8: Calculate F-critical values
- Step 10: Compare F and F-critical for each effect and conclusion

Hypothesis

- There are three sets of hypotheses with the two-way ANOVA.
- The null hypotheses for each of the sets are given below.
- The population means of the first factor are equal.
- The population means of the second factor are equal.
- There is no interaction between the two factors.

$$F > F\text{-critical} = H1$$

$$F < F\text{-critical} = H0$$

Formula

$$SS = A \text{ Variance}^2 \times N$$

$$SS \text{ total} = \text{Standard deviation}^2 \times (N - 1)$$

$$MS = SS \div DF$$

$$F = MS \div MS \text{ residual}$$

Two-way ANOVA summary table

	DF	SS	MS	F	F-critical
Main effect A	A – 1	A Variance ^{2 x N}	SS _A ÷ DF _A	MS _A ÷ MS _r	DF _A , DF _r
Main effect B	B – 1	B Variance ^{2 x N}	SS _B ÷ DF _B	MS _B ÷ MS _r	DF _B , DF _r
Residual (Error)	Total – (A + B)	SS _{total} – (SS _A + SS _B)	SS _r ÷ DF _r	N/A	N/A
Total	N – 1	Standard deviation ^{2 x (N – 1)}	N/A	N/A	N/A

Where:

- DF = Degree of freedom
- SS = Sum of squares
- MS = Mean square
- F = F values
- F-critical = Critical statistics F

Example Question

Three variety of potatoes are being compared for yield. The experiment was carried out by assigning each variety at random to four different plots. One variety of potato in each of the 4 locations. Potato type (3 levels) and location (4 levels). The plot is as follows.

		Potato type		
		P-1	P-2	P-3
Yield	Y-1	18	13	12
	Y-2	20	23	21
	Y-3	14	12	9
	Y-4	11	17	10

Answer**Step 1: Define hypothesis**

Set 1

$H_0: \mu_1 = \mu_2 = \mu_3$ (The population means of the yields are equal.)

$H_1: \mu_1 \neq \mu_2 \neq \mu_3$ (At least one mean is different.)

Set 2

$H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$ (The population means of the potato types are equal.)

$H_1: \mu_1 \neq \mu_2 \neq \mu_3 \neq \mu_4$ (At least one mean is different.)

Step 2: Find the means for Row and Column

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put Row data (for each yield type). Click AC to go back main screen and data will be automatically saved inside.
3. Click APPS to enter statistics mode and select S-VAR.
4. Select \bar{X} and press = to show mean result.
5. Repeat this process to find mean for each row and column of dataset. The answer will obtain as follows. G mean (grand mean) is the mean of overall dataset.

		Potato type (B)			
		P-1	P-2	P-3	Mean (\bar{X})
Yield (A)	Y-1	18	13	12	14.33
	Y-2	20	23	21	21.33
	Y-3	14	12	9	11.66
	Y-4	11	17	10	12.66
Mean (\bar{X})		15.75	16.25	13	15 (G mean)

Step 3: Frame the ANOVA summary table

	DF	SS	MS	F	F-critical
Yield (A)					
Potato (B)					
Residual					
Total					

Step 4: Calculate DF (Degree of freedom)

DF for yield	Number of rows – 1 = 4 – 1 = 3
DF for potato	Number of columns – 1 = 3 – 1 = 2
DF total	Total number of observations in dataset – 1 = 12 – 1 = 11
DF residual	DF total – (DF yield + DF potato) = 11 – 5 = 6

Step 5: Calculate SS (Sum of squares)

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put row **mean** data (**calculated in step 2**). Click AC to go back main screen and data will be automatically saved inside.
3. Click APPS to enter statistics mode and select S-VAR.
4. Select **xσn** for variance.
5. Put square and multiply with total number of observations in dataset. This will produce SS for Yield. Final formula on calculator screen will like this.

$$\mathbf{x\sigma n^{2 \times 12}}$$

Therefore, SS for yield is like

$$\mathbf{SS \text{ yield} = \text{Variance} (14.33, 21.33, 11.66, 12.66)^{2 \times 12} = 171.44}$$

Repeat same process of Potato and result is like

$$\mathbf{SS \text{ potato} = \text{Variance} (15.75, 16.25, 13)^{2 \times 12} = 24.5}$$

After these two SSs for main effects are complete, the next process is to calculate SS total.

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put **ALL** observation values in dataset. Click AC to go back main screen and data will be automatically saved inside.

3. Click APPS to enter statistics mode and select S-VAR.
4. Select **xσn-1** for standard deviation of dataset.
5. Put square and multiply with DF total which is already calculated in **Step 4**. This will produce SS total. Final formula on calculator screen will like this.

$\mathbf{x\sigma n-1^2 \times (12-1)}$
--

Therefore, SS total result is like

$$\mathbf{SS\ total = Standard\ deviation^2 \times 11 = 238}$$

After 3 SS values are complete, calculation is simple to find SS residual values by

$$\mathbf{SS\ residual = SS\ total - (SS\ yield + SS\ potato) = 238 - (171.44 + 24.5) = 42.39}$$

After these DF and SS processes are complete, put the values in ANOVA frame.

	DF	SS	MS	F	F-critical
Yield (A)	3	171.44			
Potato (B)	2	24.5			
Residual	6	42.17			
Total	11	238			

Step 6: Calculate MS (Mean squares)

The formula of MS is as follows

$$\mathbf{MS = SS \div DF}$$

Therefore, MS for each row is calculated as follows.

$$\mathbf{MS\ yield = SS\ yield \div DF\ yield = 171.44 \div 3 = 57.14}$$

$$\mathbf{MS\ potato = SS\ potato \div DF\ potato = 24.5 \div 2 = 12.25}$$

$$\mathbf{MS\ residual = SS\ residual \div DF\ residual = 42.17 \div 6 = 7.03}$$

Note: There is no need to find MS total.

Place the MS values in frame.

	DF	SS	MS	F	F-critical
Yield (A)	3	171.44	57.14		
Potato (B)	2	24.5	12.25		
Residual	6	42.17	7.03		
Total	11	238	-		

Step 7: Calculate F (F value)

F values for yield and potato are calculated as follows.

$$\mathbf{F \text{ yield} = MS \text{ yield} \div MS \text{ residual} = 57.14 \div 7.03 = 8.12}$$

$$\mathbf{F \text{ potato} = MS \text{ potato} \div MS \text{ residual} = 12.55 \div 7.03 = 1.74}$$

Update the table with F values

	DF	SS	MS	F	F-critical
Yield (A)	3	171.44	57.14	8.12	
Potato (B)	2	24.5	12.25	1.74	
Residual	6	42.17	7.03	-	
Total	11	238	-	-	

Step 8: Calculate F-critical values

Check the critical F values in F table ($p=0.05$) with calculated **DFs**.

$$\mathbf{F\text{-critical} = F \text{ 0.05, DF effect (F table column), DF residual (F table row)}}$$

Therefore,

$$\mathbf{F\text{-critical} \text{ yield} = F_{0.05, 3, 6} = 4.76}$$

$$\mathbf{F\text{-critical} \text{ potato} = F_{0.05, 2, 6} = 5.14}$$

Update the table (Final table)

	DF	SS	MS	F	F-critical
Yield (A)	3	171.44	57.14	8.12	4.76
Potato (B)	2	24.5	12.25	1.74	5.14
Residual	6	42.17	7.03	-	-
Total	11	238	-	-	-

Step 10: Compare F and F-critical for each effect and conclusion

F yield > F-critical yield (H1)

Reject null hypothesis and accept alternative hypothesis. Yield has significant effect on the dependent variable. The result is significant at $p < .05$

F potato < F-critical potato (H0)

Accept null hypothesis and reject alternative hypothesis. Potato has no significant effect on the dependent variable. The result is not significant at $p < .05$

More investigation

We can convert F-critical to P values from following link.

<https://www.socscistatistics.com/pvalues/fdistribution.aspx>

The result is as follows.

	DF	SS	MS	F	F-critical	P value
Yield	3	171.44	57.14	8.12	4.76 (3,6)	.049938
Potato	2	24.5	12.25	1.74	5.14 (2,6)	.05006
Residual	6	42.17	7.03	7.03	-	-
Total	11	238	-	-	-	-

ONE-WAY ANOVA

Assumptions

1. The populations from which the samples were obtained must be normally or approximately normally distributed.
2. The samples must be independent.
3. The variances of the populations must be equal.

One-way ANOVA calculation steps

- Step 1: Define hypothesis
- Step 2: Find the means of each group
- Step 3: Frame the ANOVA summary table
- Step 4: Calculate DF (Degree of freedom)
- Step 5: Calculate SS (Sum of squares)
- Step 6: Calculate MS (Mean squares)
- Step 7: Calculate F (F value)
- Step 8: Calculate F-critical values
- Step 10: Compare F and F-critical then conclusion

Hypothesis

The null hypothesis will be that all population means are equal, the alternative hypothesis is that at least one mean is different.

$$F > F\text{-critical} = H1$$

$$F < F\text{-critical} = H0$$

Formula

$$SS = \text{Variance}^2 \times N$$

$$SS_{\text{total}} = \text{Standard deviation}^2 \times (N - 1)$$

$$MS = SS \div DF$$

$$F = MS \div MS_{\text{residual}}$$

Two-way ANOVA summary table

	DF	SS	MS	F	F-critical
Main effect A	A – 1	A Variance ^{2 x N}	SS _A ÷ DF _A	MS _A ÷ MS _r	DF _A , DF _r
Residual (Error)	Total – A	SS _{total} – SS _A	SS _r ÷ DF _r	N/A	N/A
Total	N – 1	Standard deviation ^{2 x (N – 1)}	N/A	N/A	N/A

Where:

- DF = Degree of freedom
- SS = Sum of squares
- MS = Mean square
- F = F values
- F-critical = Critical statistics F

Example Question

Students of different countries were randomly selected and their height (in meter) were recorder as below:

African 1.7, 1.5, 1.6, 1.6, 1.8

Asian 1.5, 1.5, 1.7, 1.7, 1.6

European 1.6, 1.8, 1.6, 1.7, 1.8

Use one-way ANOVA to test at 0.05 level of significant whether there is difference among the mean of heights for the three nationalities. The heights for all nationalities follow normal distribution.

Answer

Dataset in clear format

African	1.7	1.5	1.6	1.6	1.8
Asian	1.5	1.5	1.7	1.7	1.6
European	1.6	1.8	1.6	1.7	1.8

Step 1: Define hypothesis

H0: All mean heights are equal

H1: Not all mean heights are equal

Step 2: Find the means of each group

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put data (for each ethnicity type). Click AC to go back main screen and data will be automatically saved inside.

3. Click APPS to enter statistics mode and select S-VAR.
4. Select \bar{X} and press = to show mean result.
5. You need to repeat this process **3 times** for each ethnicity.

The result will be obtained as follows;

						Mean (\bar{X})
African	1.7	1.5	1.6	1.6	1.8	1.64
Asian	1.5	1.5	1.7	1.7	1.6	1.6
European	1.6	1.8	1.6	1.7	1.8	1.7
					Grand mean = 1.64	

Grand mean is the mean of all observations in dataset.

Step 3: Frame the ANOVA summary table

	DF	SS	MS	F	F-critical
Main effect					
Residual					
Total					

Step 4: Calculate DF (Degree of freedom)

DF for main effect **Number of columns – 1 = 3 – 1 = 2**

DF total **Total number of observations in dataset – 1 = 15 – 1 = 14**

DF residual **DF total – DF main effect = 14 – 2 = 12**

Arrange the values in ANOVA frame.

	DF	SS	MS	F	F-critical
Main effect	2				
Residual	12				
Total	14				

Step 5: Calculate SS (Sum of squares)

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put row **mean** data (**calculated in step 2**). Click AC to go back main screen and data will be automatically saved inside.
3. Click APPS to enter statistics mode and select S-VAR.
4. Select **xσn** for variance.
5. Put square and multiply with total number of observations in dataset. This will produce SS for main effect (Ethnicity). Final formula on calculator screen will like this.

$$x\sigma n^{2 \times 15}$$

Therefore, SS for main effect (ethnicity) is like

$$SS \text{ main} = \text{Variance } (16.4, 1.6, 1.7)^{2 \times 15} = 0.0253$$

After SS for main effects is complete, the next process is to calculate SS total.

Calculator setup

1. Enter Mode and select STAT to enter statistical calculation mode.
2. Choose SD and put **ALL** observation values in dataset. Click AC to go back main screen and data will be automatically saved inside.
3. Click APPS to enter statistics mode and select S-VAR.
4. Select **xσn-1** for standard deviation of dataset.
5. Put square and multiply with DF total which is already calculated in **Step 4**. This will produce SS total. Final formula on calculator screen will like this.

$$x\sigma n-1^{2 \times (15-1)}$$

Therefore, SS total result is like

$$\text{SS total} = \text{Standard deviation}^2 \times 14 = 0.1573$$

After SS main and total are calculated, SS residual is easy to find by

$$\text{SS residual} = \text{SS total} - \text{SS residual} = 0.1573 - 0.0253 = 0.132$$

After SS calculations are complete, update the ANOVA frame.

	DF	SS	MS	F	F-critical
Main effect	2	0.0253			
Residual	12	0.132			
Total	14	0.1573			

Step 6: Calculate MS (Mean squares)

The formula of MS is as follows

$$\text{MS} = \text{SS} \div \text{DF}$$

Therefore, MS for each row is calculated as follows.

$$\text{MS main} = \text{SS main} \div \text{DF main} = 0.0253 \div 2 = 0.0126$$

$$\text{MS residual} = \text{SS residual} \div \text{DF residual} = 0.132 \div 12 = 0.011$$

Note: There is no need to find MS total.

Place the MS values in frame.

	DF	SS	MS	F	F-critical
Main effect	2	0.0253	0.0126		
Residual	12	0.132	0.011		
Total	14	0.1573			

Step 7: Calculate F (F value)

Using the formula, F values for main effect is calculated as follows.

$$\mathbf{F\ main = MS\ main \div MS\ main = 0.0126 \div 0.011 = 1.145}$$

Update the table with F values

	DF	SS	MS	F	F-critical
Main effect	2	0.0253	0.0126	1.145	
Residual	12	0.132	0.011		
Total	14	0.1573			

Step 8: Calculate F-critical values

Check the critical F values in F table ($p=0.05$) with calculated **DF**.

$$\mathbf{F\text{-}critical = F\ 0.05,\ DF\ main\ (F\ table\ column),\ DF\ residual\ (F\ table\ row)}$$

Therefore,

$$\mathbf{F\text{-}critical\ main = F_{0.05,\ 2,\ 12} = 3.89}$$

Update the table (Final table)

	DF	SS	MS	F	F-critical
Main effect	2	0.0253	0.0126	1.145	3.89
Residual	12	0.132	0.011		
Total	14	0.1573			

Step 10: Compare F and F-critical then conclusion

$$\mathbf{F\ main < F\text{-}critical\ (H_0)}$$


Null hypothesis cannot be rejected. The difference among the mean heights are not significant.

The result is not significant at $p < .05$


Two-way ANOVA in SPSS

	Yield	Potato	Result
1	Y1	P1	18.00
2	Y1	P2	13.00
3	Y1	P3	12.00
4	Y2	P1	20.00
5	Y2	P2	23.00
6	Y2	P3	21.00
7	Y3	P1	14.00
8	Y3	P2	12.00
9	Y3	P3	9.00
10	Y4	P1	11.00
11	Y4	P2	17.00
12	Y4	P3	10.00


Step 1: Insert data into SPSS data frame



Step 2: Select Analyze → General Linear Model → Univariate



Step 3: Arrange variables into their correct order. Next 3 steps to configure are “Plots”, “Post Hoc”, and “Options” buttons.



Univariate: Profile Plots

Factors:
Yield
Potato

Horizontal Axis:
Yield

Separate Lines:
Potato

Separate Plots:

Plots:
Add Change Remove

Yield*Potato


Chart Type:
☒ Line Chart
☐ Bar Chart

Error Bars
☐ Include Error bars
☒ Confidence Interval (95.0%)
☐ Standard Error Multiplier: 2

☐ Include reference line for grand mean
☐ Y axis starts at 0

Continue Cancel Help

Step 4: Click “Plots” button and configure above settings. Click “Add” button and select chart type. Then click “Continue”. This will lead to previous screen.



Univariate: Post Hoc Multiple Comparisons for Observed Means

Factor(s):
Yield
Potato


Post Hoc Tests for:
Yield
Potato

Equal Variances Assumed
☐ LSD ☐ S-N-K ☐ Waller-Duncan
☐ Bonferroni ☒ Tukey Type I/Type II Error Ratio: 100
☐ Sidak ☐ Tukey's-b ☐ Dunnett
☐ Scheffe ☐ Duncan Control Category: Last
☐ R-E-G-W-F ☐ Hochberg's GT2 Test
☐ R-E-G-W-Q ☐ Gabriel ☒ 2-sided ☐ < Control ☐ > Control

Equal Variances Not Assumed
☐ Tamhane's T2 ☐ Dunnett's T3 ☐ Games-Howell ☐ Dunnett's C

Continue Cancel Help

Step 5: Click “Post Hoc” button and configure above settings. Click “Continue” for next step.




The image shows the 'Univariate: Options' dialog box in SPSS. The 'Display' section has the following checked options: Descriptive statistics, Estimates of effect size, Homogeneity tests, and Parameter estimates. Under 'Heteroskedasticity Tests', the 'Modified Breusch-Pagan test' is selected. The 'Significance level' is set to .05, and 'Confidence intervals are 95.0 %'. The 'Continue' button is highlighted.

Step 6: Click “Options” and check for above values. Click “Continue” and this will close that form. ANOVA setup is completed and click “OK” to start analysis.


Result

Tests of Between-Subjects Effects						
Dependent Variable:						
Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared
Corrected Model	238.000 ^a	11	21.636			1.000
Intercept	2700.000	1	2700.000			1.000
Yield	171.333	3	57.111			1.000
Potato	24.500	2	12.250			1.000
Yield * Potato	42.167	6	7.028			1.000
Error	0.000	0				
Total	2938.000	12				
Corrected Total	238.000	11				

a. R Squared = 1.000 (Adjusted R Squared = .)



If you select “Bar Chart” in step 4, the plot will produce like this.




One-way ANOVA in SPSS


On-way process is same with two-way.

	Ethnicity	Result
1	African	1.7
2	African	1.5
3	African	1.6
4	African	1.6
5	African	1.8
6	Asian	1.5
7	Asian	1.5
8	Asian	1.7
9	Asian	1.7
10	Asian	1.6
11	European	1.6
12	European	1.8
13	European	1.6
14	European	1.7
15	European	1.8


Step 1: Insert data into SPSS data frame



Step 2: Select Analyze → General Linear Model → Univariate



Step 3: Arrange variables into their correct order. Next 3 steps to configure are “Plots”, “Post Hoc”, and “Options” buttons.



Univariate: Profile Plots

Factors: Ethnicity

Horizontal Axis: []

Separate Lines: []

Separate Plots: []

Plots: Add Change Remove

Plots: Ethnicity

Chart Type: ☒ Line Chart ☐ Bar Chart

Error Bars ☐ Include Error bars


☒ Confidence Interval (95.0%) ☐ Standard Error Multiplier: 2

☐ Include reference line for grand mean

☐ Y axis starts at 0

Continue Cancel Help

Step 4: Click “Plots” button. Drag Ethnicity to Horizontal Axis and click Add. Select chart type then click “Continue”. This will lead to previous screen.



Univariate: Post Hoc Multiple Comparisons for Observed Means

Factor(s): Ethnicity

Post Hoc Tests for: Ethnicity

Equal Variances Assumed

☐ LSD ☐ S-N-K ☐ Waller-Duncan

☐ Bonferroni ☒ Tukey Type I/Type II Error Ratio: 100

☐ Sidak ☐ Tukey's-b ☐ Dunnett

☐ Scheffe ☐ Duncan Control Category: Last

☐ R-E-G-W-F ☐ Hochberg's GT2 Test


☐ R-E-G-W-Q ☐ Gabriel ☒ 2-sided ☐ < Control ☐ > Control

Equal Variances Not Assumed

☐ Tamhane's T2 ☐ Dunnett's T3 ☐ Games-Howell ☐ Dunnett's C

Continue Cancel Help

Step 5: Click “Post Hoc” button and configure above settings. Click “Continue” for next step.




The image shows the 'Univariate: Options' dialog box in SPSS. The 'Display' section has the following checked options: 'Descriptive statistics', 'Estimates of effect size', 'Homogeneity tests', and 'Spread vs. level plot'. Other options like 'Observed power', 'Parameter estimates', 'Contrast coefficient matrix', 'Residual plot', 'Lack of fit', and 'General estimable function' are unchecked. The 'Heteroskedasticity Tests' section has 'Modified Breusch-Pagan test' and 'Breusch-Pagan test' checked, with 'F test' and 'White's test' unchecked. There are 'Model...' buttons next to the checked tests. The 'Parameter estimates with robust standard errors' section has radio buttons for HC0, HC1, HC2, HC3, and HC4, all of which are currently unselected. At the bottom, the 'Significance level' is set to .05 and 'Confidence intervals are 95.0 %'. There are 'Continue', 'Cancel', and 'Help' buttons at the bottom.

Step 6: Click “Options” and check for above values. Click “Continue” and this will close that form. ANOVA setup is completed and click “OK” to start analysis.


Result

Tests of Between-Subjects Effects						
Dependent Variable:						
Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared
Corrected Model	.025 ^a	2	0.013	1.152	0.349	0.161
Intercept	40.673	1	40.673	3697.515	0.000	0.997
Ethnicity	0.025	2	0.013	1.152	0.349	0.161
Error	0.132	12	0.011			
Total	40.830	15				
Corrected Total	0.157	14				

a. R Squared = .161 (Adjusted R Squared = .021)



Bar chart



Two-way ANOVA in SAS


SAS supports GUI setup for two-way ANOVA. Example codes are shown below.

```
DATA Potato;
INPUT Yield$ Potato$ Result;
DATALINES;
Y1 P1 18
Y1 P2 13
Y1 P3 12
Y2 P1 20
Y2 P2 23
Y2 P3 21
Y3 P1 14
Y3 P2 12
Y3 P3 9
Y4 P1 11
Y4 P2 17
Y4 P3 10
;
RUN;
PROC PRINT DATA=Potato;
RUN;
```


Obs	Yield	Potato	Result
1	Y1	P1	18
2	Y1	P2	13
3	Y1	P3	12
4	Y2	P1	20
5	Y2	P2	23
6	Y2	P3	21
7	Y3	P1	14
8	Y3	P2	12
9	Y3	P3	9
10	Y4	P1	11
11	Y4	P2	17
12	Y4	P3	10

```
proc glm data=WORK.POTATO plot(only)=(intplot);
class Yield Potato;
model Result=Yield Potato Yield*Potato / ss3;
lsmeans Yield Potato / adjust=tukey pdiff=all alpha=0.05 cl plots=();
quit;
```

Source	DF	Type III SS	Mean Square	F Value	Pr > F
Yield	3	171.3333333	57.1111111	-	-
Potato	2	24.5000000	12.2500000	-	-
Yield*Potato	6	42.1666667	7.0277778	-	-



SAS setup



Settings | Code/Results | Split | [Icons]

DATA MODEL OPTIONS

DATA

WORK.POTATO

Filter: (none)

ROLES


*Dependent variable: (1 item)

Result

*Factors:

Yield

Potato



DATA MODEL OPTIONS

MODEL EFFECTS

Model

Model Effects


Intercept

Yield

Potato

Yield*Potato

Edit



Settings | Code/Results | Split | [Icons]

DATA MODEL OPTIONS

STATISTICS

Select statistics to display:

Default statistics

PLOTS

Select plots to display:

Selected plots

☒ Interaction plot

☐ LSMeans plot

☐ Mean difference plot

☐ Diagnostic plots

Maximum number of plot points:

Default(5,000)

One-way ANOVA in SAS

SAS supports GUI setup for one-way ANOVA. Example codes are shown below.

```
data Ethnicity;
input Nationality$ Height;
datalines;
African 1.7
African 1.5
African 1.6
African 1.6
African 1.8
Asian 1.5
Asian 1.5
Asian 1.7
Asian 1.7
Asian 1.6
European 1.6
European 1.8
European 1.6
European 1.7
European 1.8
;
run;
proc print data=Ethnicity;
run;
```

Obs	Nationality	Height
1	African	1.7
2	African	1.5
3	African	1.6
4	African	1.6
5	African	1.8
6	Asian	1.5
7	Asian	1.5
8	Asian	1.7
9	Asian	1.7
10	Asian	1.6
11	European	1.6
12	European	1.8
13	European	1.6
14	European	1.7
15	European	1.8

```
Title 'ONE-WAY ANOVA';
ods noproctitle;
ods graphics / imagemap=on;


proc glm data=WORK.ETHNICITY plots(only);
  class Nationality;
  model Height=Nationality;
  means Nationality / welch plots=none;
  lsmeans Nationality / adjust=tukey pdiff alpha=.05 plots=(diffplot);
run;
quit;
```

Dependent Variable: Height					
Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	0.02533333	0.01266667	1.15	0.3488
Error	12	0.13200000	0.01100000		
Corrected Total	14	0.15733333			


R-Square	Coeff Var	Root MSE	Height Mean
0.161017	6.369285	0.104881	1.646667

Source	DF	Type I SS	Mean Square	F Value	Pr > F
Nationality	2	0.02533333	0.01266667	1.15	0.3488

Source	DF	Type III SS	Mean Square	F Value	Pr > F
Nationality	2	0.02533333	0.01266667	1.15	0.3488



SAS setup



Settings Code/Results Split

DATA OPTIONS OUTPUT INFORMATION

DATA

WORK.ETHNICITY

Filter: (none)


ROLES

*Dependent variable: (1 item)

Result

*Categorical variable: (1 item)

Nationality



Settings Code/Results Split

DATA OPTIONS OUTPUT INFORMATION

HOMOGENEITY OF VARIANCE

Test:

None

☒ Welch's variance-weighted ANOVA

COMPARISONS

Comparisons method:

Tukey

Significance level: 0.05

PLOTS

Display plots:

Selected plots

☐ Box plot

☐ Means plot

☒ LS-mean difference plot

☐ Diagnostics plot

Maximum number of plot points:

Default(5,000)

Datasets

Data table for two-way ANOVA

Yield	Potato	Result
Y1	P1	18
Y1	P2	13
Y1	P3	12
Y2	P1	20
Y2	P2	23
Y2	P3	21
Y3	P1	14
Y3	P2	12
Y3	P3	9
Y4	P1	11
Y4	P2	17
Y4	P3	10

Data table for one-way ANOVA

Ethnicity	Height
African	1.7
African	1.5
African	1.6
African	1.6
African	1.8
Asian	1.5
Asian	1.5
Asian	1.7
Asian	1.7
Asian	1.6
European	1.6
European	1.8
European	1.6
European	1.7
European	1.8

Reference websites

[1]: <https://statistics.laerd.com/spss-tutorials/two-way-anova-using-spss-statistics.php>

[2]: <https://people.richland.edu/james/lecture/m170/ch13-2wy.html>

[3]: <https://www.socscistatistics.com/pvalues/fdistribution.aspx>

[4]: <https://www.spss-tutorials.com/spss-two-way-anova-basics-tutorial/>

[5]: <http://vassarstats.net/anova1u.html>

[6]: <http://vassarstats.net/anova2u.html>

[7]: <https://people.richland.edu/james/lecture/m170/ch13-1wy.html>

PROFILE**AUNG PYAE**

Studies M.SC in Data Science and Business Analytics at Asia Pacific University of Technology and Innovation (Kuala Lumpur, Malaysia). Does researches in Data science, Statistics, Machine learning and AI, Operation research, Business intelligence.

Website: <http://aungpyae.me>