

CYCLE LOG

CYCLE NO. _____ START DATE _____ TARGET _____ H ONE SHEET PER CYCLE

DAY	WEIGHT	WATER L	SALT TSP	K ⁺	MG	B1	WALK	ENERGY, SYMPTOMS, NOTES
P-3				☐	☐	☐	☐	
P-2				☐	☐	☐	☐	
P-1				☐	☐	☐	☐	
F1				☐	☐	☐	☐	
F2				☐	☐	☐	☐	
F3				☐	☐	☐	☐	
F4				☐	☐	☐	☐	
R1				☐	☐	☐	☐	First meal:
R2				☐	☐	☐	☐	

KEY P = prime · F = fast · R = refeed · shaded rows are the fast itself · tick each box when done

DAILY TARGETS DURING THE FAST

3,000–5,000 mg Sodium — 1¼-2 tsp fine salt, divided, in water
1,000–3,000 mg Potassium — small doses only, never one large dose
300–500 mg Magnesium — glycinate or malate, at night
50–100 mg Thiamine (B1) — refeeding insurance, daily
2–3 L Water — salted; plain volume alone risks hyponatremia
30–60 min Easy walking, days 1–2, tapering after

BREAK THE FAST IMMEDIATELY IF

- Heart palpitations, irregular heartbeat, or chest pain
 - Fainting, or near-fainting that doesn’t resolve on sitting
 - Confusion, slurred speech, or disorientation
 - Persistent vomiting, or you can’t keep water down
 - Visual disturbance, or severe weakness (can’t climb stairs)
 - Numbness or tingling that spreads
- None of these are “push through” symptoms. Eat something, salt it, reassess. A cycle abandoned early costs almost nothing.