

# AMATYC J-4

Curtis Bradley

July 24, 2024

## 1 Problem

Start with a regular hexagon of side 1 as level 1. At each level  $n - 1$ , form the regular hexagon of level  $n$  by joining consecutive midpoints of the one for level  $n - 1$ . Find the lowest level at which the associated regular hexagon has a perimeter under 1.

## 2 Solution

Construct a regular hexagon with side length  $S_k$ , such that the perimeter  $P_k = 6S_k$ . Construct a second hexagon inscribed in the first with vertices intersecting the first's midpoints. Observe that six identical triangles are made between both hexagons with side lengths of the triangle of  $\frac{S_k}{2}$  separated by an angle of  $120^\circ$  (the interior angle of the hexagon). Using law of cosines, the side length of the inscribed hexagon can be found to be  $S_{k+1} = \frac{\sqrt{3}}{2} S_k$ . We can construct an equation for  $P_k$  as such:

$$P_k = 6S_1 \cdot \left( \frac{\sqrt{3}}{2} \right)^{k-1}$$

Setting  $P_k$  to 1 and  $S_1$  to 1, it can be found  $k = 1 + \frac{\ln 6}{\ln \frac{2\sqrt{3}}{3}} \approx 13.45$ . Therefore the smallest integer level  $n$  where  $P_n \leq 1$  is  $n = \lceil k \rceil = 14$  where  $P_{14} \approx 0.92$ .  $\square$




Diagram of hexagon with side lengths  $S_k$  with inscribed hexagon of side lengths  $S_{k+1}$