

# Computational topology: Lecture 4

Alberto Paoluzzi

March 14, 2019

1 Simplicial Complexes

2 Cellular Complexes

3 Chain Complexes

# Simplicial Complexes

# Join operation

The **join** of two sets  $P, Q \subset \mathbb{E}^n$  is the set of **convex combination** of their points:

$$PQ = \{\alpha \mathbf{x} + \beta \mathbf{y} \mid \mathbf{x} \in P, \mathbf{y} \in Q\},$$

where  $\alpha, \beta \in \mathbb{R}$ ,  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

# Join operation

The **join** of two sets  $P, Q \subset \mathbb{E}^n$  is the set of **convex combination** of their points:

$$PQ = \{\alpha \mathbf{x} + \beta \mathbf{y} \mid \mathbf{x} \in P, \mathbf{y} \in Q\},$$

where  $\alpha, \beta \in \mathbb{R}$ ,  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

The join operation is associative and commutative.

# Join operation

The **join** of two sets  $P, Q \subset \mathbb{E}^n$  is the set of **convex combination** of their points:

$$PQ = \{\alpha \mathbf{x} + \beta \mathbf{y} \mid \mathbf{x} \in P, \mathbf{y} \in Q\},$$

where  $\alpha, \beta \in \mathbb{R}$ ,  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

The join operation is associative and commutative.

## Implementation

How ?

# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

A  $d$ -simplex can be seen as a  $d$ -dimensional triangle: a 0-simplex is a **point**, a 1-simplex is a **segment**, a 2-simplex is a **triangle**, a 3-simplex is a **tetrahedron**, and so on.



# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

A  $d$ -simplex can be seen as a  $d$ -dimensional triangle: a 0-simplex is a **point**, a 1-simplex is a **segment**, a 2-simplex is a **triangle**, a 3-simplex is a **tetrahedron**, and so on.

The set  $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d\}$  of vertices of  $\sigma_d$  is called the **0-skeleton** of  $\sigma_d$ .

# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

A  $d$ -simplex can be seen as a  $d$ -dimensional triangle: a 0-simplex is a **point**, a 1-simplex is a **segment**, a 2-simplex is a **triangle**, a 3-simplex is a **tetrahedron**, and so on.

The set  $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d\}$  of vertices of  $\sigma_d$  is called the **0-skeleton** of  $\sigma_d$ .

The  $s$ -simplex generated from **any** subset of  $s + 1$  vertices ( $0 \leq s \leq n$ ) of  $\sigma_d$  is called an  **$s$ -face** of  $\sigma_d$ .

# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

A  $d$ -simplex can be seen as a  $d$ -dimensional triangle: a 0-simplex is a **point**, a 1-simplex is a **segment**, a 2-simplex is a **triangle**, a 3-simplex is a **tetrahedron**, and so on.

The set  $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d\}$  of vertices of  $\sigma_d$  is called the **0-skeleton** of  $\sigma_d$ .

The  $s$ -simplex generated from **any** subset of  $s + 1$  vertices ( $0 \leq s \leq n$ ) of  $\sigma_d$  is called an  **$s$ -face** of  $\sigma_d$ .

Let us notice, from the definition, that a simplex may be considered both as a purely **combinatorial object** and as a **geometric object**, i.e. as the compact point set defined by the convex hull of a discrete set of points.

# Simplex

A  $d$ -simplex  $\sigma_d \subset \mathbb{E}^n$  ( $0 \leq d \leq n$ ) may be defined as the repeated join of  $d + 1$  affinely independent points, called **vertices**.

A  $d$ -simplex can be seen as a  $d$ -dimensional triangle: a 0-simplex is a **point**, a 1-simplex is a **segment**, a 2-simplex is a **triangle**, a 3-simplex is a **tetrahedron**, and so on.

The set  $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d\}$  of vertices of  $\sigma_d$  is called the **0-skeleton** of  $\sigma_d$ .

The  $s$ -simplex generated from **any** subset of  $s + 1$  vertices ( $0 \leq s \leq n$ ) of  $\sigma_d$  is called an  **$s$ -face** of  $\sigma_d$ .

Let us notice, from the definition, that a simplex may be considered both as a purely **combinatorial object** and as a **geometric object**, i.e. as the compact point set defined by the convex hull of a discrete set of points.

## Implementation

How ?

# Simplicial complex

## Definition

A set  $\Sigma$  of simplices is called a **triangulation**.

# Simplicial complex

## Definition

A set  $\Sigma$  of simplices is called a **triangulation**.

A **simplicial complex**, often simply denoted as **complex**, is a triangulation  $\Sigma$  that verifies the following conditions:

- 1 if  $\sigma \in \Sigma$ , then any face of  $\sigma$  belongs to  $\Sigma$ ;
- 2 if  $\sigma, \tau \in \Sigma$ , then either  $\sigma \cap \tau = \emptyset$ , or  $\sigma \cap \tau$  is a face of both  $\sigma$  and  $\tau$ .

# Simplicial complex

## Definitions

A simplicial complex can be considered a “well-formed” triangulation.

# Simplicial complex

## Definitions

A simplicial complex can be considered a “well-formed” triangulation.

Such kind of triangulations are widely used in engineering analysis, e.g., in topography or in finite element methods.



# Simplicial complex

## Definitions

A simplicial complex can be considered a “well-formed” triangulation.

Such kind of triangulations are widely used in engineering analysis, e.g., in topography or in finite element methods.

The **order** of a complex is the maximum order of its simplices. A complex  $\Sigma_d$  of order  $d$  is also called a  **$d$ -complex**.

# Simplicial complex

## Definitions

A simplicial complex can be considered a “well-formed” triangulation.

Such kind of triangulations are widely used in engineering analysis, e.g., in topography or in finite element methods.

The **order** of a complex is the maximum order of its simplices. A complex  $\Sigma_d$  of order  $d$  is also called a  **$d$ -complex**.

A  $d$ -complex is said to be **regular** or **pure** if **each simplex is a face of a  $d$ -simplex**.

A **regular  $d$ -complex** is homogeneously  $d$ -dimensional.

# Simplicial complex

## Definitions

A simplicial complex can be considered a “well-formed” triangulation.

Such kind of triangulations are widely used in engineering analysis, e.g., in topography or in finite element methods.

The **order** of a complex is the maximum order of its simplices. A complex  $\Sigma_d$  of order  $d$  is also called a  **$d$ -complex**.

A  $d$ -complex is said to be **regular** or **pure** if **each simplex is a face of a  $d$ -simplex**.

A **regular  $d$ -complex** is homogeneously  $d$ -dimensional.

## Implementation

How ?

# Combinatorial boundary

The **combinatorial boundary**  $\Sigma_{d-1} = \partial\sigma_d$  of a simplex  $\sigma_d$  is a simplicial complex consisting of all proper  $s$ -faces ( $s < d$ ) of  $\sigma_d$ .

# Combinatorial boundary

The **combinatorial boundary**  $\Sigma_{d-1} = \partial\sigma_d$  of a simplex  $\sigma_d$  is a simplicial complex consisting of all proper  $s$ -faces ( $s < d$ ) of  $\sigma_d$ .

Two simplices  $\sigma$  and  $\tau$  in a complex  $\Sigma$  are called  **$s$ -adjacent** if they have a common  $s$ -face.

# Combinatorial boundary

The **combinatorial boundary**  $\Sigma_{d-1} = \partial\sigma_d$  of a simplex  $\sigma_d$  is a simplicial complex consisting of all proper  $s$ -faces ( $s < d$ ) of  $\sigma_d$ .

Two simplices  $\sigma$  and  $\tau$  in a complex  $\Sigma$  are called  **$s$ -adjacent** if they have a common  $s$ -face.

With some abuse of language, we call (combinatorial)  **$s$ -skeletons** the sets  $\mathcal{K}_s$  ( $s \leq d$ ). **Geometric carrier**  $|\Sigma|$ , also called the **support space**, is the point-set union of simplices in  $\Sigma$ .

## Implementation

How ?

```
help?> Lar.simplexFacets
```

# Orientation

The **ordering** of the 0-skeleton of a simplex implies an **orientation** of it. The simplex can be oriented according to the **even or odd permutation class** of its 0-skeleton.

The two **opposite orientations** of a simplex will be denoted as  $+\sigma$  and  $-\sigma$ .

# Orientation

The **ordering** of the 0-skeleton of a simplex implies an **orientation** of it. The simplex can be oriented according to the **even or odd permutation class** of its 0-skeleton.

The two **opposite orientations** of a simplex will be denoted as  $+\sigma$  and  $-\sigma$ .

Two simplices are **coherently oriented** when their common faces have opposite orientation. A complex is **orientable** when all its simplices can be coherently oriented.



# Orientation

The **ordering** of the 0-skeleton of a simplex implies an **orientation** of it. The simplex can be oriented according to the **even or odd permutation class** of its 0-skeleton.

The two **opposite orientations** of a simplex will be denoted as  $+\sigma$  and  $-\sigma$ .

Two simplices are **coherently oriented** when their common faces have opposite orientation. A complex is **orientable** when all its simplices can be coherently oriented.

It is assumed that:

- 1 The two orientations of a simplex may represent its relative interior and exterior;
- 2 the two orientations of an orientable simplicial complex analogously represent the relative interior and exterior of the complex, respectively;
- 3 the boundary of a complex maintains the same orientation of the complex.

# Orientation and volume

The **volume** associated with an orientation of a simplex (or complex) is positive, while the one associated with the opposite orientation has the same absolute value and opposite sign.

## Orientation and volume

The **volume** associated with an orientation of a simplex (or complex) is positive, while the one associated with the opposite orientation has the same absolute value and opposite sign.

It is assumed that the bounded object has positive volume.

It is also assumed that either a minus sign or a multiplying factor  $-1$  denote a complementation, i.e. an opposite orientation of the simplex, which can be explicitly obtained by swapping two vertices in its ordered 0-skeleton:

# Orientation and volume

The **volume** associated with an orientation of a simplex (or complex) is positive, while the one associated with the opposite orientation has the same absolute value and opposite sign.

It is assumed that the bounded object has positive volume.

It is also assumed that either a minus sign or a multiplying factor  $-1$  denote a complementation, i.e. an opposite orientation of the simplex, which can be explicitly obtained by swapping two vertices in its ordered 0-skeleton:

$$+\sigma_3 = \langle \mathbf{v}_0, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \rangle$$

$$-\sigma_3 = \langle \mathbf{v}_1, \mathbf{v}_0, \mathbf{v}_2, \mathbf{v}_3 \rangle$$

## Implementation

How ?

```
help?> Lar.surface
```

## Face extraction

The oriented facets  $\sigma_{d-1,(i)}$  ( $0 \leq i \leq d$ ) of the oriented  $d$ -simplex  $\sigma_d = +\langle \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d \rangle$  are obtained by removing the  $i$ -th vertex  $\mathbf{v}_i$  from the 0-skeleton of  $\sigma_d$ :

## Face extraction

The oriented facets  $\sigma_{d-1,(i)}$  ( $0 \leq i \leq d$ ) of the oriented  $d$ -simplex  $\sigma_d = +\langle \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d \rangle$  are obtained by removing the  $i$ -th vertex  $\mathbf{v}_i$  from the 0-skeleton of  $\sigma_d$ :

$$\sigma_{d-1,(i)} = (-1)^i(\sigma_d - \langle \mathbf{v}_i \rangle), \quad 0 \leq i \leq d. \quad (1)$$

where the minus sign denotes a subtraction between 0-skeletons.

## Face extraction

The oriented facets  $\sigma_{d-1,(i)}$  ( $0 \leq i \leq d$ ) of the oriented  $d$ -simplex  $\sigma_d = +\langle \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d \rangle$  are obtained by removing the  $i$ -th vertex  $\mathbf{v}_i$  from the 0-skeleton of  $\sigma_d$ :

$$\sigma_{d-1,(i)} = (-1)^i (\sigma_d - \langle \mathbf{v}_i \rangle), \quad 0 \leq i \leq d. \quad (1)$$

where the minus sign denotes a subtraction between 0-skeletons.

The 0-skeleton of  $\sigma_{d-1,(i)}$  is therefore obtained by removing the  $i$ -th vertex from the 0-skeleton of  $\sigma_d$  and either by swapping a pair of vertices or, better, by inverting the simplex sign, when  $i$  is odd.

# Face extraction

The oriented facets  $\sigma_{d-1,(i)}$  ( $0 \leq i \leq d$ ) of the oriented  $d$ -simplex  $\sigma_d = +\langle \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d \rangle$  are obtained by removing the  $i$ -th vertex  $\mathbf{v}_i$  from the 0-skeleton of  $\sigma_d$ :

$$\sigma_{d-1,(i)} = (-1)^i(\sigma_d - \langle \mathbf{v}_i \rangle), \quad 0 \leq i \leq d. \quad (1)$$

where the minus sign denotes a subtraction between 0-skeletons.

The 0-skeleton of  $\sigma_{d-1,(i)}$  is therefore obtained by removing the  $i$ -th vertex from the 0-skeleton of  $\sigma_d$  and either by swapping a pair of vertices or, better, by inverting the simplex sign, when  $i$  is odd.

## Implementation

How ?

```
help?> Lar.simplexFacets
```



# Facet extraction




Figure 1: Coherent orientation of the facets of a 3-simplex

# Facet extraction




Figure 1: Coherent orientation of the facets of a 3-simplex

Implementation? See:

```
function Lar.simplexFacets(simplices::Cells)::Cells
```

# Simplicial prism

The prism over a simplex  $\sigma_d = \langle \mathbf{v}_0, \dots, \mathbf{v}_d \rangle$ , defined as the set  $P_{d+1} := \sigma_d \times [a, b]$ , with  $[a, b] \subset \mathbb{E}$ , will be called **simplicial  $(d + 1)$ -prism**.

# Simplicial prism

The prism over a simplex  $\sigma_d = \langle \mathbf{v}_0, \dots, \mathbf{v}_d \rangle$ , defined as the set  $P_{d+1} := \sigma_d \times [a, b]$ , with  $[a, b] \subset \mathbb{E}$ , will be called **simplicial  $(d+1)$ -prism**.

An oriented complex which triangulates  $P_{d+1}$  can be defined combinatorially, by using a closed form formula for its  $\mathcal{K}_{d+1}$  skeleton:

# Simplicial prism

The prism over a simplex  $\sigma_d = \langle \mathbf{v}_0, \dots, \mathbf{v}_d \rangle$ , defined as the set  $P_{d+1} := \sigma_d \times [a, b]$ , with  $[a, b] \subset \mathbb{E}$ , will be called **simplicial  $(d+1)$ -prism**.

An oriented complex which triangulates  $P_{d+1}$  can be defined combinatorially, by using a closed form formula for its  $\mathcal{K}_{d+1}$  skeleton:

$$\mathcal{K}_{d+1} = \{ \sigma_{d+1, (i)} = (-1)^{id} \langle \mathbf{v}_i^a, \mathbf{v}_{i+1}^a, \dots, \mathbf{v}_d^a, \mathbf{v}_0^b, \mathbf{v}_1^b, \dots, \mathbf{v}_i^b \rangle \mid 0 \leq i \leq d \}$$

where  $\mathbf{v}_i^a = (\mathbf{v}_i, a)$  and  $\mathbf{v}_i^b = (\mathbf{v}_i, b)$ .

## Exercise: extract the boundary of implicit prism

extract the boundary of the simplicial cube

- write dimension-independent code

### Hint

See and modify the code of `Lar.simplexGrid`

## Exercise: extract the boundary of implicit prism

extract the boundary of the simplicial cube

- write dimension-independent code
- apply to a grid of such decomposed cubes

Hint

See and modify the code of `Lar.simplexGrid`

# Cellular Complexes



# Preliminary definitions

Let  $X$  be a topological space, and  $\Lambda(X) = \bigcup \Lambda_p$  ( $p \in \{0, 1, \dots, d\}$ ) a partition of  $X$ , with  $\Lambda_p$  a set of  $\{(\text{relatively}) \text{ open}\}$ , connected, and manifold  $p$ -cells.

# Preliminary definitions

Let  $X$  be a topological space, and  $\Lambda(X) = \bigcup \Lambda_p$  ( $p \in \{0, 1, \dots, d\}$ ) a partition of  $X$ , with  $\Lambda_p$  a set of  $\{(relatively) \text{ open}\}$ , connected, and manifold  $p$ -cells.

A **CW-structure** on the space  $X$  is a filtration  $\emptyset = X_{-1} \subset X_0 \subset X_1 \subset \dots \subset X_{d-1} \subset X = \bigcup_p X_p$ , such that, for each  $p$ , the **skeleton**  $X_p$  is homeomorphic to a space obtained from  $X_{p-1}$  by attachment of  $p$ -cells in  $\Lambda_p = \Lambda_p(X) \sim [?]$ .

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

A cellular complex is **finite** when it contains a finite number of cells.

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

A cellular complex is **finite** when it contains a finite number of cells.

A **{regular}**  $d$ -complex is a complex where each closed  $d$ -cell equates the closure of its interior, and where  $p$ -cells ( $p < d$ ) are contained in the boundary of a  $d$ -cell.

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

A cellular complex is **finite** when it contains a finite number of cells.

A **{regular}**  $d$ -complex is a complex where each closed  $d$ -cell equates the closure of its interior, and where  $p$ -cells ( $p < d$ ) are contained in the boundary of a  $d$ -cell.

- Two  $d$ -cells are **coherently oriented** when their common  $(d - 1)$ -cells have opposite orientations.

## Implementation

How ?

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

A cellular complex is **finite** when it contains a finite number of cells.

A **{regular}**  $d$ -complex is a complex where each closed  $d$ -cell equates the closure of its interior, and where  $p$ -cells ( $p < d$ ) are contained in the boundary of a  $d$ -cell.

- Two  $d$ -cells are **coherently oriented** when their common  $(d - 1)$ -cells have opposite orientations.
- A cellular  $d$ -complex  $X$  is **orientable** when its  $d$ -cells can be coherently oriented.

## Implementation

How ?

# Cellular complex

A **CW-complex** is a space  $X$  endowed with a CW-structure, and is also called a **cellular complex**.

A cellular complex is **finite** when it contains a finite number of cells.

A **{regular}**  $d$ -complex is a complex where each closed  $d$ -cell equates the closure of its interior, and where  $p$ -cells ( $p < d$ ) are contained in the boundary of a  $d$ -cell.

- Two  $d$ -cells are **coherently oriented** when their common  $(d - 1)$ -cells have opposite orientations.
- A cellular  $d$ -complex  $X$  is **orientable** when its  $d$ -cells can be coherently oriented.
- The **support space**  $|\sigma|$  of a cell  $\sigma$  is its compact point-set.

## Implementation

How ?



# Chain Complexes

# Chain groups

aaaaa

Chains are defined by attaching coefficients to cells.

# Chain groups

aaaaa

Chains are defined by attaching coefficients to cells.

Since one wishes to add chains, one has to pick coefficients from a set endowed with the structure of a commutative group, or stronger.

Let  $(G, +, 0)$  be a nontrivial commutative group, whose identity element is denoted 0.

# Chain groups

aaaaa

Chains are defined by attaching coefficients to cells.

Since one wishes to add chains, one has to pick coefficients from a set endowed with the structure of a commutative group, or stronger.

Let  $(G, +, 0)$  be a nontrivial commutative group, whose identity element is denoted 0.

A  $p$ -chain of  $X$  with coefficients in  $G$  is a mapping  $c_p : X \rightarrow G$  such that, for each  $\sigma \in X_p$ , reversing a cell orientation changes the sign of the chain value:

$$c_p(-\sigma) = -c_p(\sigma).$$

# Chain groups

aaaaa

Chain addition is defined by addition of chain values: if  $c_p^1, c_p^2$  are  $p$ -chains, then  $(c_p^1 + c_p^2)(\sigma) = c_p^1(\sigma) + c_p^2(\sigma)$ , for each  $\sigma \in X_p$ .

The resulting group is denoted  $C_p(X; G)$ . When clear from the context, the group  $G$  is often left implied, writing  $C_p(X)$ .

# Chain groups

aaaaa

Chain addition is defined by addition of chain values: if  $c_p^1, c_p^2$  are  $p$ -chains, then  $(c_p^1 + c_p^2)(\sigma) = c_p^1(\sigma) + c_p^2(\sigma)$ , for each  $\sigma \in X_p$ .

The resulting group is denoted  $C_p(X; G)$ . When clear from the context, the group  $G$  is often left implied, writing  $C_p(X)$ .

Let  $\sigma$  be an oriented cell in  $X$  and  $g \in G$ . The **elementary chain** whose value is  $g$  on  $\sigma$ ,  $-g$  on  $-\sigma$  and 0 on any other cell in  $X$  is denoted  $g\sigma$ .

# Chain groups

aaaaa

Chain addition is defined by addition of chain values: if  $c_p^1, c_p^2$  are  $p$ -chains, then  $(c_p^1 + c_p^2)(\sigma) = c_p^1(\sigma) + c_p^2(\sigma)$ , for each  $\sigma \in X_p$ .

The resulting group is denoted  $C_p(X; G)$ . When clear from the context, the group  $G$  is often left implied, writing  $C_p(X)$ .

Let  $\sigma$  be an oriented cell in  $X$  and  $g \in G$ . The **elementary chain** whose value is  $g$  on  $\sigma$ ,  $-g$  on  $-\sigma$  and 0 on any other cell in  $X$  is denoted  $g\sigma$ .

Each chain can be written in a unique way as a sum of elementary chains.

# Chain groups

aaaaa

Chains are often thought of as attaching orientation and/or multiplicity to cells



# Chain groups

aaaaa

Chains are often thought of as attaching orientation and/or multiplicity to cells

If coefficients are taken from the group  $G = (\{-1, 0, 1\}, +, 0)$ , then cells can only be discarded or selected, possibly inverting their orientation

# Chain groups

aaaaa

Chains are often thought of as attaching orientation and/or multiplicity to cells

If coefficients are taken from the group  $G = (\{-1, 0, 1\}, +, 0)$ , then cells can only be discarded or selected, possibly inverting their orientation

A  $p$ -cycle is a closed  $p$ -chain, i.e., a  $p$ -chain without boundary.

# Chain orientation

It is useful to select a **conventional orientation** to orient cells **automatically**.

# Chain orientation

It is useful to select a **conventional orientation** to orient cells **automatically**.

- **0-cells** are considered all **positive**.

# Chain orientation

It is useful to select a **conventional orientation** to orient cells **automatically**.

- **0-cells** are considered all **positive**.
- Closed  $p$ -cells can be given a **coherent (internal) orientation** according with the orientation of the first  $(p - 1)$ -cell in their **canonical representation** (sorted on indices) of their  $(p - 1)$ -cycles.

# Chain orientation

It is useful to select a **conventional orientation** to orient cells **automatically**.

- **0-cells** are considered all **positive**.
- Closed  $p$ -cells can be given a **coherent (internal) orientation** according with the orientation of the first  $(p - 1)$ -cell in their **canonical representation** (sorted on indices) of their  $(p - 1)$ -cycles.
- Finally, a  **$d$ -cell** may be oriented as the **sign of its oriented volume**.

# Chain spaces

aaaaa

To allow not only for chain addition, but also for linear combination of chains, coefficients should be taken from a set endowed with the structure of a field, such as  $(\mathbb{F}, +, \times, 0, 1)$ ,

where 0 and  $1 \neq 0$  denote, respectively, the additive and multiplicative identities.

# Chain spaces

aaaaa

To allow not only for chain addition, but also for linear combination of chains, coefficients should be taken from a set endowed with the structure of a field, such as  $(\mathbb{F}, +, \times, 0, 1)$ ,

where 0 and  $1 \neq 0$  denote, respectively, the additive and multiplicative identities.

**Unit chains** are elementary chains whose value is  $u = 1\sigma$  for some cell  $\sigma$ .



# Chain spaces

aaaaa

To allow not only for chain addition, but also for linear combination of chains, coefficients should be taken from a set endowed with the structure of a field, such as  $(\mathbb{F}, +, \times, 0, 1)$ ,

where 0 and  $1 \neq 0$  denote, respectively, the additive and multiplicative identities.

**Unit chains** are elementary chains whose value is  $u = 1\sigma$  for some cell  $\sigma$ .

Each chain can be written in a **unique way** as a **linear combination** of unit chains  $u \in U$ , if the outer cell is not taken into account.

# Chain spaces

## Canonical representation and standard basis

Hence, the space of  $p$ -chains  $C_p$  is endowed with a **standard (or natural) basis**, comprised of **all the independent unit  $p$ -chains**.

# Chain spaces

## Canonical representation and standard basis

Hence, the space of  $p$ -chains  $C_p$  is endowed with a **standard (or natural) basis**, comprised of **all the independent unit  $p$ -chains**.

The **standard basis** may be the one associated with the **canonical representation** . . .

# Chain spaces

## Canonical representation and standard basis

Hence, the space of  $p$ -chains  $C_p$  is endowed with a **standard (or natural) basis**, comprised of **all the independent unit  $p$ -chains**.

The **standard basis** may be the one associated with the **canonical representation** . . .

In particular,  $\#U_d = \#\Lambda_d - 1$ .

# Chain spaces

## Canonical representation and standard basis

Hence, the space of  $p$ -chains  $C_p$  is endowed with a **standard (or natural) basis**, comprised of **all the independent unit  $p$ -chains**.

The **standard basis** may be the one associated with the **canonical representation** ...

In particular,  $\#U_d = \#\Lambda_d - 1$ .

Often, with some **abuse of notation**, one does not distinguish between a  $p$ -cell and the corresponding unit  $p$ -chain.

# Characteristic matrices

Given a set  $S = \{s_j\}$ , the **characteristic function**  $\chi_A : S \rightarrow \{0, 1\}$  takes value 1 for all elements of  $A \subseteq S$  and 0 at all elements of  $S$  not in  $A$ .

# Characteristic matrices

Given a set  $S = \{s_j\}$ , the **characteristic function**  $\chi_A : S \rightarrow \{0, 1\}$  takes value 1 for all elements of  $A \subseteq S$  and 0 at all elements of  $S$  not in  $A$ .

We call **characteristic matrix**  $M$  of a collection of subsets  $A_i \subseteq S$  ( $i = 1, \dots, n$ ) the binary matrix  $M = (m_{ij})$ , with  $m_{ij} = \chi_{A_i}(s_j)$ .

A matrix  $M_p$ , whose rows are indexed by unit  $p$ -chains and columns are indexed by unit 0-chains, provides a useful representation of a basis for the linear space  $C_p$ .

# Characteristic matrices

Given a set  $S = \{s_j\}$ , the **characteristic function**  $\chi_A : S \rightarrow \{0, 1\}$  takes value 1 for all elements of  $A \subseteq S$  and 0 at all elements of  $S$  not in  $A$ .

We call **characteristic matrix**  $M$  of a collection of subsets  $A_i \subseteq S$  ( $i = 1, \dots, n$ ) the binary matrix  $M = (m_{ij})$ , with  $m_{ij} = \chi_{A_i}(s_j)$ .

A matrix  $M_p$ , whose rows are indexed by unit  $p$ -chains and columns are indexed by unit 0-chains, provides a useful representation of a basis for the linear space  $C_p$ .

Permuting (**reindexing**) either rows or columns provides a **different basis**.



# Characteristic matrices

Given a set  $S = \{s_j\}$ , the **characteristic function**  $\chi_A : S \rightarrow \{0, 1\}$  takes value 1 for all elements of  $A \subseteq S$  and 0 at all elements of  $S$  not in  $A$ .

We call **characteristic matrix**  $M$  of a collection of subsets  $A_i \subseteq S$  ( $i = 1, \dots, n$ ) the binary matrix  $M = (m_{ij})$ , with  $m_{ij} = \chi_{A_i}(s_j)$ .

A matrix  $M_p$ , whose rows are indexed by unit  $p$ -chains and columns are indexed by unit 0-chains, provides a useful representation of a basis for the linear space  $C_p$ .

Permuting (**reindexing**) either rows or columns provides a **different basis**.

While chains are mostly presented as formal sums of cells, in the actual implementation their **signed coordinate vectors** are used as **sparse arrays**, and in particular as CSC (Compressed Sparse Column) maps :

$\mathbb{N} \rightarrow \{-1, 0, 1\}$ .

# Characteristic matrices

## Implementation

# Characteristic matrices

## Implementation

### Implementation

How ?

```
help?> Lar.characteristicMatrix
```

## WHY Characteristic matrices are useful

Consider the **incidence queries** that arise in a **cellular decomposition**  $\Lambda(X)$  of a 2D space, such as **2D triangulation** or the **boundary** of a **manifold 3-solid**.

## WHY Characteristic matrices are useful

Consider the **incidence queries** that arise in a **cellular decomposition**  $\Lambda(X)$  of a 2D space, such as **2D triangulation** or the **boundary** of a **manifold 3-solid**.

There are **9 incidence relations** between pairs of cells in such a  $V, E, F$  decomposition, and traditional **graph-based representations** are chosen by optimizing the trade-off between **space requirements** and **efficiency** of queries.

With LAR, all such **queries are sparse matrix–vector multiplications** without any additional **space requirements**.

## WHY Characteristic matrices are useful

Consider the **incidence queries** that arise in a **cellular decomposition**  $\Lambda(X)$  of a 2D space, such as **2D triangulation** or the **boundary** of a **manifold 3-solid**.

There are **9 incidence relations** between pairs of cells in such a  $V, E, F$  decomposition, and traditional **graph-based representations** are chosen by optimizing the trade-off between **space requirements** and **efficiency** of queries.

With LAR, all such **queries are sparse matrix–vector multiplications** without any additional **space requirements**.

The chain operators corresponding to the incidence relations are

$$\begin{array}{lll}
 \mathcal{V}\mathcal{V} : C_0 \rightarrow C_0, & \mathcal{E}\mathcal{V} : C_0 \rightarrow C_1, & \mathcal{F}\mathcal{V} : C_0 \rightarrow C_2; \\
 \mathcal{V}\mathcal{E} : C_1 \rightarrow C_0, & \mathcal{E}\mathcal{E} : C_1 \rightarrow C_1, & \mathcal{F}\mathcal{E} : C_1 \rightarrow C_2; \\
 \mathcal{V}\mathcal{F} : C_2 \rightarrow C_0, & \mathcal{E}\mathcal{F} : C_2 \rightarrow C_1, & \mathcal{F}\mathcal{F} : C_2 \rightarrow C_2.
 \end{array}$$

## WHY Characteristic matrices are useful

Consider the **incidence queries** that arise in a **cellular decomposition**  $\Lambda(X)$  of a 2D space, such as **2D triangulation** or the **boundary** of a **manifold 3-solid**.

There are **9 incidence relations** between pairs of cells in such a  $V, E, F$  decomposition, and traditional **graph-based representations** are chosen by optimizing the trade-off between **space requirements** and **efficiency** of queries.

With LAR, all such **queries are sparse matrix–vector multiplications** without any additional **space requirements**.

The chain operators corresponding to the incidence relations are

$$\begin{array}{lll}
 \mathcal{V}\mathcal{V} : C_0 \rightarrow C_0, & \mathcal{E}\mathcal{V} : C_0 \rightarrow C_1, & \mathcal{F}\mathcal{V} : C_0 \rightarrow C_2; \\
 \mathcal{V}\mathcal{E} : C_1 \rightarrow C_0, & \mathcal{E}\mathcal{E} : C_1 \rightarrow C_1, & \mathcal{F}\mathcal{E} : C_1 \rightarrow C_2; \\
 \mathcal{V}\mathcal{F} : C_2 \rightarrow C_0, & \mathcal{E}\mathcal{F} : C_2 \rightarrow C_1, & \mathcal{F}\mathcal{F} : C_2 \rightarrow C_2.
 \end{array}$$

# Characteristic matrices: Examples

aaaaa

The corresponding **CSR matrices** are readily computed:

$$\mathcal{V}\mathcal{V} = \mathcal{V}\mathcal{E} \circ \mathcal{E}\mathcal{V} = \mathcal{E}\mathcal{V}^\top \circ \mathcal{E}\mathcal{V} \Rightarrow [\mathcal{V}\mathcal{V}] = M_1^t M_1$$

$$\mathcal{V}\mathcal{E} = \mathcal{E}\mathcal{V}^\top \Rightarrow [\mathcal{V}\mathcal{E}] = M_1^t$$

$$\mathcal{V}\mathcal{F} = \mathcal{F}\mathcal{V}^\top \Rightarrow [\mathcal{V}\mathcal{F}] = M_2^t$$

$$\mathcal{E}\mathcal{V} \quad [\mathcal{E}\mathcal{V}] = M_1$$

$$\mathcal{E}\mathcal{E} = \mathcal{E}\mathcal{V} \circ \mathcal{V}\mathcal{E} = \mathcal{E}\mathcal{V} \circ \mathcal{E}\mathcal{V}^\top \Rightarrow [\mathcal{E}\mathcal{E}] = M_1 M_1^t$$

$$\mathcal{E}\mathcal{F} = \mathcal{E}\mathcal{V} \circ \mathcal{V}\mathcal{F} = \mathcal{E}\mathcal{V} \circ \mathcal{F}\mathcal{V}^\top \Rightarrow [\mathcal{E}\mathcal{F}] = M_1 M_2^t$$

$$\mathcal{F}\mathcal{V} \quad [\mathcal{F}\mathcal{V}] = M_2$$

$$\mathcal{F}\mathcal{E} = \mathcal{F}\mathcal{V} \circ \mathcal{V}\mathcal{E} = \mathcal{F}\mathcal{V} \circ \mathcal{E}\mathcal{V}^\top \Rightarrow [\mathcal{F}\mathcal{E}] = M_2 M_1^t$$

$$\mathcal{F}\mathcal{F} = \mathcal{F}\mathcal{V} \circ \mathcal{V}\mathcal{F} = \mathcal{F}\mathcal{V} \circ \mathcal{F}\mathcal{V}^\top \Rightarrow [\mathcal{F}\mathcal{F}] = M_2 M_2^t.$$