$\begin{array}{c} \text{group theory} \\ \text{birdtracks, Lie's, and exceptional groups} \\ a \ blog \end{array}$

Predrag Cvitanović

January 5, 2023

1.7 Quantum marginal problem

The marginal distribution of a subset of a collection of random variables is the probability distribution of the variables contained in the subset. It gives the probabilities of various values of the variables in the subset without reference to the values of the other variables.

Marginal variables are those variables in the subset of variables being retained. These concepts are "marginal" because they can be found by summing values in a table along rows or columns, and writing the sum in the margins of the table.

Quantum marginal problem (QMP) : Find a pure quantum state $|\psi\rangle$ given some marginals ρ_{ℓ} .

2023-01-02 Erik Aurell The quantum marginal problem: is there a globally pure state which has a set of reduced density matrices (in Fraser's notation): Let there be

- (a) a joint Hilbert space $J = X_1 \otimes X_2 \otimes X_k$,
- (b) a family of index sets $S_1,S_2,\cdots S_m$ with cardinalities $k_1,k_2,....k_m$ labelling Hilbert spaces $X_{S_i}=X_{i,j_1}\otimes\cdots\otimes X_{i,j_{k_1}}$, and
- (c) for each Hilbert space X_{S_i} a density matrix ρ_i .

Question: does there exist a pure state $|\psi\rangle$ on J such that the partial trace of $|\psi\rangle\langle\psi|$ over the Hilbert space complementary to X_{S_i} equals ρ_i ? That is, in standard notation, is there a $|\psi\rangle$ such that ${\rm Tr}_{J\backslash S_i}[|\psi\rangle\langle\psi|] = \rho_i$ for each i?

I have a paper with Pawel Horodecki [11] where we use some results in this field.

This problem was solved about 10 years ago in the case when the marginals are disjoint (those are the results we used with Horodecki) by Alexander Klyachko [72, 73] Quantum marginal problem and representations of the symmetric group, arXiv:quant-ph/0409113; Quantum marginal problem and N-representability, arXiv:quant-ph/0511102. In this case only the eigenvalues of marginals ρ_ℓ matter. In principle. Relies on algebraic geometry, not straightforward to apply. One defines sets of marginals of compatible states, then checks whether they contain any pure states. Testing for purity is a nonlinear problem. Trick: take a convex hull of copies of compatible states. Pure states have traces equal to 1; this reduces the problem to study of symmetric products of the copies / separable states.

Aside: Klyachko was born in 1946, in Vladivostok.

Disjoint marginals mean questions like if there exists a pure state over N particles with given 1-particle reduced density matrices. There is no general solution for overlapping marginals. Overlapping marginals means questions like if there exists a pure state over N particles with a set of given 2-particle reduced density matrices, where the connectivity graph of particles linked by these given 2-particle reduced density matrices is large.

Thomas C. Fraser [49] A sufficient family of necessary inequalities for the compatibility of quantum marginals, arXiv:2211.00685, gives a general (but ab-

stract) solution to QMP. Here is Thomas' (a graduate student) short video on this work:

Thomas Fraser: Simultaneous Spectral Estimation and the Quantum Marginal Problem (2021)

I have yet to decide if the solution in this paper is practical in any way. It is however clear that it uses properties of symmetric tensors a lot. There is an appendix explaining Predrag's birdtrack notation, with calculations (which I did not try to do yet). Predrag's book is cited.

Maybe something new that we could think of together?

2023-01-02 Predrag Fraser [49] *Appendix E: Diagrammatics* is taken directly from my *Group Theory* [37]. So is most of 'tensor networks' literature, which never cites me but cites Penrose instead, so Fraser is being very kind. The new and key operation, not in my book, is eq. (E15) for operations on 'multi-partite Hilbert spaces', i.e., multiple copies of a tensor.

Appendix F: The n=1 Case I can follow step-by-step, but do not understand the significance of the results.

2023-01-03 Erik Aurell .

Otfried Gühne: The Quantum Marginal Problem (2022)

is up to 32 min at least partly about other (but related) problems. I have read

Huber, Gühne and Siewert [59] Absolutely maximally entangled states of seven qubits do not exist (2017)

Huber, Eltschka, Siewert and Gühne [58] Bounds on absolutely maximally entangled states from shadow inequalities, and the quantum MacWilliams identity, (2018)

Yu, Simnacher, Wyderka, Nguyen and Gühne [127] A complete hierarchy for the pure state marginal problem in quantum mechanics (2021)

that he is referring to (though not in detail). I know other work by his postdoc Nguyen, who coauthored a very good classical stat mech review;

Nguyen, Zecchina and Berg [89] *Inverse statistical problems: From the inverse Ising problem to data science* (2017), arXiv:1702.01522.

At 32 min Gühne starts on the problem as I see it. The (unpublished) Klyachko [73] paper he refers to is a classic in the field, though almost incomprehensible (at least to me). It concerns the problem where the marginals are non-overlapping. In that setting there is a much easier later paper for Gaussian states:

Eisert, Tyc, Rudolph and Sanders [43] *Gaussian quantum marginal problem* (2008), arXiv:quant-ph/0703225.

At around 43minutes or so comes the construction of Gühne which is in the same ballpark as Fraser's. It considers more and more copies of the system, and then conditions on those N copies of the system such that the quantum marginal

problem is solvable by a technique called semi-definite programming, standard in the field. I never bothered to learn exactly what it is because (1) there are already many others who know it very well and use it and (2) I am not convinced it is in the end so practical to consider these problems of N copies of the original problem, where N eventually tends to infinity. I may of course be wrong on this point (2). After 44 minutes I think Gühne again looks at a special case, maybe not generally useful.

The last part of the talk is I think a bit inconclusive (or not quite up-to-date). The case "AME(4,6)" was solved in Suhail Ahmad Rather, Adam Burchardt, Wojciech Bruzda, Grzegorz Rajchel-Mieldzioć, Arul Lakshminarayan, Karol Życzkowski [100] *Thirty-six entangled officers of Euler: Quantum solution to a classically impossible problem* (2022).

I am not sure if there is not some almost-cheating behind this. Życzkowski was one of the co-authors of the list of open problems Gühne refers to. Maybe Życzkowski knew that he could solve that problem when he put together the list of open problems together with Pawel Horodecki. I do not know.

References

- [1] G. 't Hooft, "On the phase transition towards permanent quark confinement", Nucl. Phys. B **138**, 1–25 (1978).
- [2] G. 't Hooft, "On the convergence of planar diagram expansions", Comm. Math. Phys. **86**, 449–464 (1982).
- [3] G. 't Hooft, "Rigorous construction of planar diagram field theories in four dimensional Euclidean space", Comm. Math. Phys. **88**, 1–25 (1983).
- [4] G. 't Hooft, "Planar diagram field theories", in *Progress in Gauge Field Theory*, edited by G. 't Hooft, A. Jaffe, H. Lehmann, P. K. MitterI, M. Singer, and R. Stora (Springer, 1984), pp. 271–335.
- [5] G. 't Hooft, "Counting planar diagrams with various restrictions", Nucl. Phys. B **538**, 389–410 (1999).
- [6] G. 't Hooft, "A planar diagram theory for strong interactions", Nucl. Phys. B 72, 461–473 (1974).
- [7] J. Alcock-Zeilinger and H. Weigert, "Compact Hermitian Young projection operators", J. Math. Phys. **58**, 051702 (2017).
- [8] J. Alcock-Zeilinger and H. Weigert, "Simplification rules for birdtrack operators", J. Math. Phys. **58**, 051701 (2017).
- [9] J. Alcock-Zeilinger and H. Weigert, "Transition operators", J. Math. Phys. **58**, 051703 (2017).
- [10] J. Alcock-Zeilinger and H. Weigert, "A simple counting argument of the irreducible representations of SU(N) on mixed product spaces", J. Algebr. Comb., 10.1007/s10801-018-0853-z (2018).
- [11] E. Aurell, M. Eckstein, and P. Horodecki, "Hawking radiation and the quantum marginal problem", J. Cosmol. Astropart. Phys. **2022**, 014 (2022).
- [12] J. C. Baez and J. Dolan, "Categorification", in *Higher Category Theory*, edited by E. Getzler and M. Kapranov (Amer. Math. Soc., Providence R.I., 1998), pp. 1–36.

[13] D. Binosi and L. Theußl, "JaxoDraw: a graphical user interface for drawing Feynman diagrams", Comput. Phys. Commun. **161**, 76–86 (2004).

- [14] J. F. Blinn, "Quartic discriminants and tensor invariants", IEEE Computer Graphics and Appl. 22, 86–91 (2002).
- [15] C. J. Bomhof, P. J. Mulders, and F. Pijlman, "The construction of gauge-links in arbitrary hard processes", Euro. Phys. J. C 47, 147–162 (2006).
- [16] R. Brauer, "On algebras which are connected with the semisimple continuous groups", Ann. Math. **38**, 857 (1937).
- [17] E. Brézin, C. Itzykson, G. Parisi, and J. B. Zuber, "Planar diagrams", Comm. Math. Phys. **59**, 35–51 (1978).
- [18] J. Brundan, "On the definition of Heisenberg category", Algebraic Combinatorics 1, 523–544 (2018).
- [19] P. Butera, G. M. Cicuta, and M. Enriotti, "Group weight and vanishing graphs", Phys. Rev. D **21**, 972–978 (1980).
- [20] L.-Q. Cai, B.-S. Lin, and K. Wu, "A diagrammatic categorification of q-boson and q-fermion algebras", Chin. Phys. B **21**, 020201 (2012).
- [21] G. P. Canning, "Diagrammatic group theory in quark models", Phys. Rev. D 18, 395–410 (1978).
- [22] A. Cayley, "XXVIII. On the theory of the analytical forms called trees", The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science **13**, 172–176 (1857).
- [23] S. Chmutov, S. Duzhin, and J. Mostovoy, *Introduction to Vassiliev Knot Invariants* (Cambridge Univ. Press, Cambridge UK, 2012).
- [24] G. M. Cicuta, "Vanishing graphs, planarity, and Reggeization", Phys. Rev. Lett. 43, 826–829 (1979).
- [25] W. K. Clifford, "Extract of a letter to Mr. Sylvester from Prof. Clifford of University College, London", Amer. J. Math. 1, 126–128 (1878).
- [26] B. Coecke and A. Kissinger, *Picturing Quantum Processes: A First Course in Quantum Theory and Diagrammatic Reasoning* (Cambridge Univ. Press, Cambridge UK, 2017).
- [27] B. Coecke and A. Kissinger, *Categorical Quantum Mechanics I: Causal Quantum Processes* (Oxford Univ. Press, 2018).
- [28] B. Cooper and M. Hogancamp, "An exceptional collection for Khovanov homology", Algebraic & Geometric Topology 15, 2659–2707 (2015).
- [29] M. S. Costa and T. Hansen, "Conformal correlators of mixed-symmetry tensors", J. High Energy Phys. 2015, 151 (2015).
- [30] M. S. Costa, T. Hansen, J. Penedones, and E. Trevisani, "Projectors and seed conformal blocks for traceless mixed-symmetry tensors", J. High Energy Phys. **2016**, 018 (2016).

- [31] F. Cougoulic, Nuclear effects in high-energy proton-nucleus collisions: transverse momentum broadening of energetic parton systems and soft anomalous dimension matrices, PhD thesis (Ecole Nationale Supérieure Mines-Télécom Atlantique, 2018).
- [32] F. Cougoulic and S. Peigné, "Nuclear p_⊥-broadening of an energetic parton pair", J. High Energy Phys. **2018**, 203 (2018).
- [33] C. W. Curtis, "The four and eight square problem and division algebras", in *Studies in Modern Algebra*, edited by A. A. Albert (Math. Assn. of America, New York, 1963), pp. 100–125.
- [34] P. Cvitanović, "Group theory for Feynman diagrams in non-Abelian gauge theories", Phys. Rev. D 14, 1536–1553 (1976).
- [35] P. Cvitanović, Classical and exceptional Lie algebras as invariance algebras, Oxford Univ. preprint 40/77, unpublished, 1977.
- [36] P. Cvitanović, *Field Theory*, Notes prepared by E. Gyldenkerne (Nordita, Copenhagen, 1983).
- [37] P. Cvitanović, *Group Theory: Birdtracks, Lie's and Exceptional Groups* (Princeton Univ. Press, Princeton NJ, 2008).
- [38] P. Cvitanović, P. G. Lauwers, and P. N. Scharbach, "Gauge invariance structure of Quantum Chromodynamics", Nucl. Phys. B **186**, 165–186 (1981).
- [39] P. Cvitanović, P. G. Lauwers, and P. N. Scharbach, "The planar sector of field theories", Nucl. Phys. B **203**, 385–412 (1982).
- [40] A. K. Cyrol, M. Mitter, and N. Strodthoff, "FormTracer. A mathematica tracing package using FORM", Comput. Phys. Commun. **219**, 346–352 (2016).
- [41] Y. L. Dokshitzer, R. Scheibl, and C. Slotta, Perturbative QCD (and beyond), in *Lectures on QCD: Applications*, edited by F. Lenz, H. Grießhammer, and D. Stoll (Springer, Berlin, 1997), pp. 87–135.
- [42] Y.-J. Du, M. Sjödahl, and J. Thorén, "Recursion in multiplet bases for tree-level MHV gluon amplitudes", J. High Energy Phys. **2015**, 119 (2015).
- [43] J. Eisert, T. Tyc, T. Rudolph, and B. C. Sanders, "Gaussian quantum marginal problem", Comm. Math. Phys. **280**, 263–280 (2008).
- [44] H. Elvang, P. Cvitanović, and A. Kennedy, "Diagrammatic Young projection operators for u(n)", J. Math. Phys. **46**, 043501 (2005).
- [45] G. Falcioni, E. Gardi, M. Harley, M. Magnea, and C. D. White, "Multiple gluon exchange webs", J. High Energy Phys. **2014**, 010 (2014).
- [46] F. Feng, Y.-F. Xie, Q.-C. Zhou, and S.-R. Tang, HepLib: A C++ library for high energy physics, 2021.
- [47] B. Fiol, J. Martínez-Montoya, and A. Rios Fukelman, "Wilson loops in terms of color invariants", J. High Energy Phys. **2019**, 10.1007/jhep05 (2019) 202 (2018).
- [48] S. Fomin and P. Pylyavskyy, "Tensor diagrams and cluster algebras", Adv. Math. 300, 717–787 (2016).

[49] T. C. Fraser, A sufficient family of necessary inequalities for the compatibility of quantum marginals, 2022.

- [50] W. Fulton, *Young Tableaux*, with Applications to Representation Theory and Geometry (Cambridge Univ. Press, Cambridge, 1997).
- [51] S. Garoufalidis and I. Popescu, "Analyticity of the planar limit of a matrix model", Ann. Henri Poincaré 14, 499–565 (2012).
- [52] B. Geyer and M. Lazar, "Twist decomposition of nonlocal light-cone operators II: general tensors of 2nd rank", Nucl. Phys. B **581**, 341–390 (2000).
- [53] N. S. González, "Categorical Bernstein operators and the Boson-Fermion correspondence", Selecta Math. **26**, 1–64 (2020).
- [54] J. A. Gracey, " F_4 symmetric ϕ^3 theory at four loops", Phys. Rev. D **95**, 065030 (2017).
- [55] J. Gu and H. Jockers, "A note on colored HOMFLY polynomials for hyperbolic knots from WZW models", Comm. Math. Phys. **338**, 393–456 (2015).
- [56] M. Hamermesh, *Group Theory and its Applications to Physical Problems* (Dover, New York, 1962).
- [57] D. Hill and J. Sussan, "A categorification of twisted Heisenberg algebras", Adv. Math. **295**, 368–420 (2016).
- [58] F. Huber, C. Eltschka, J. Siewert, and O. Gühne, "Bounds on absolutely maximally entangled states from shadow inequalities, and the quantum MacWilliams identity", J. Phys. A **51**, 175301 (2018).
- [59] F. Huber, O. Gühne, and J. Siewert, "Absolutely maximally entangled states of seven qubits do not exist", Phys. Rev. Lett. **118**, 200502 (2017).
- [60] I. P. Ivanov, C. C. Nishi, J. P. Silva, and A. Trautner, "Basis-invariant conditions for CP symmetry of order four", Phys. Rev. D **99**, 015039 (2019).
- [61] B. R. Judd, *Operator Techniquesin Atomic Spectroscopy* (Princeton Univ. Press, 1963).
- [62] A. B. Kempe, "On the application of Clifford's graphs to ordinary binary quantics", Proceedings of the London Mathematical Society **s1-17**, 107–123 (1885).
- [63] A. D. Kennedy, Group algebras, Lie algebras, and Clifford algebras, Colloquium (Moscow State University), 1997.
- [64] S. Keppeler, "Birdtracks for SU(n)", SciPost Phys. Lect. Notes, 3 (2018).
- [65] S. Keppeler and M. Sjödahl, "Orthogonal multiplet bases in $SU(N_c)$ color space", J. High Energy Phys. **2012**, 124 (2012).
- [66] S. Keppeler and M. Sjödahl, "Hermitian Young operators", J. Math. Phys. 55, 021702 (2014).
- [67] M. Khovanov, "Heisenberg algebra and a graphical calculus", Fundam. Math. **225**, 169–210 (2014).
- [68] J.-H. Kim, M. S. H. Oh, and K.-Y. Kim, "Boosting vector calculus with the graphical notation", Am. J. Phys. **89**, 200–209 (2019).

- [69] A. A. Kirillov, "Family algebras", Electron. Res. Announc. Amer. Math. Soc. 6, 7–20 (2000).
- [70] A. A. Kirillov, "Introduction to family algebras", Moscow Math. J. 1, 49–63 (2001).
- [71] A. Kissinger, Abstract tensor systems as monoidal categories, in *Categories and Types in Logic, Language, and Physics: Essays Dedicated to Jim Lambek on the Occasion of His 90th Birthday*, edited by C. Casadio, B. Coecke, M. Moortgat, and P. Scott (Springer, Berlin, 2014), pp. 235–252.
- [72] A. Klyachko, Quantum marginal problem and representations of the symmetric group, 2004.
- [73] A. A. Klyachko, "Quantum marginal problem and N-representability", J. Phys.: Conf. Ser. **36**, 72–86 (2006).
- [74] B. Kol and R. Shir, "Color structures and permutations", J. High Energy Phys. **2014**, 020 (2014).
- [75] J. Koplik, A. Neveu, and S. Nussinov, "Some aspects of the planar perturbation series", Nucl. Phys. B **123**, 109–131 (1977).
- [76] J. Kuipers, T. Ueda, and J. A. M. Vermaseren, "Code optimization in FORM", Comput. Phys. Commun. **189**, 1–19 (2015).
- [77] T. Lappi, A. Ramnath, K. Rummukainen, and H. Weigert, "JIMWLK evolution of the odderon", Phys. Rev. D **94**, 054014 (2016).
- [78] A. Lascoux, "Young's representations of the symmetric group", in *Symmetry and Structural Properties of Condensed Matter*, edited by T. Lulek, B. Lulek, and A. Wal (World Scientific, Singapore, 2001).
- [79] A. Licata, D. Rosso, and A. Savage, "A graphical calculus for the Jack inner product on symmetric functions", J. Combin. Theory A **155**, 503–543 (2018).
- [80] A. Licata and A. Savage, A survey of Heisenberg categorification via graphical calculus, 2011.
- [81] A. Licata and A. Savage, "Hecke algebras, finite general linear groups, and Heisenberg categorification", Quantum Topol. 4, 125–185 (2013).
- [82] A. Lifson, C. Reuschle, and M. Sjödahl, "Introducing the chirality-flow for-malism", Acta Phys. Pol. B 51, 1547 (2020).
- [83] I. Lindgren and J. Morrison, *Atomic Many-Body Theory* (Springer, Berlin, 1982).
- [84] R. Lipshitz, P. Ozsváth, and D. Thurston, Bordered Heegaard Floer homology: invariance and pairing, 2008.
- [85] T. Liu and N. Zerf, Irreducible tensor basis and general Fierz relations for Bhabha scattering like amplitudes, 2016.
- [86] C. Marquet and H. Weigert, "New observables to test the Color Glass Condensate beyond the large- N_c limit", Nucl. Phys. A **843**, 68–97 (2010).

[87] A. Y. Morozov, A. A. Morozov, and A. V. Popolitov, "Matrix model and dimensions at hypercube vertices", Theor. Math. Phys. **192**, 1039–1079 (2017).

- [88] S. Morse and E. Peterson, "Trace diagrams, signed graph colorings, and matrix minors", Involve 3, 33–66 (2010).
- [89] H. C. Nguyen, R. Zecchina, and J. Berg, "Inverse statistical problems: From the inverse Ising problem to data science", Adv. Phys. **66**, 197–261 (2017).
- [90] S. Okubo and J. Patera, "General indices of simple Lie algebras and symmetrized product representations", J. Math. Phys. **24**, 2722–2733 (1983).
- [91] H. Osborn and A. Stergiou, "Seeking fixed points in multiple coupling scalar theories in the ϵ expansion", J. High Energy Phys. **2018**, 10.1007/jhep05 (2018) 051 (2016).
- [92] Y. Pang, J. Rong, and N. Su, " ϕ^3 theory with F_4 flavor symmetry in $6-2\epsilon$ dimensions: 3-loop renormalization and conformal bootstrap", J. High Energy Phys. **2016**, 057 (2016).
- [93] G. Parisi, "A simple expression for planar field theories", Phys. Lett. B 112, 463–464 (1982).
- [94] R. Penrose, "Applications of negative dimensional tensors", in *Combinatorial mathematics and its applications*, edited by D. J. A. Welsh (Academic, New York, 1971), pp. 221–244.
- [95] R. Penrose, *The Road to Reality: A Complete Guide to the Laws of the Universe* (A. A. Knopf, New York, 2005).
- [96] E. Peterson, Trace diagrams, representations, and low-dimensional topology, PhD thesis (Univ. of Maryland, 2006).
- [97] E. Peterson, A not-so-characteristic equation: the art of linear algebra, 2007.
- [98] E. Peterson, On a diagrammatic proof of the Cayley-Hamilton theorem, 2009.
- [99] E. Peterson, Unshackling linear algebra from linear notation, 2009.
- [100] S. A. Rather, A. Burchardt, W. Bruzda, G. Rajchel-Mieldzioć, A. Lakshminarayan, and K. Życzkowski, "Thirty-six entangled officers of Euler: Quantum solution to a classically impossible problem", Phys. Rev. Lett. 128, 080507 (2022).
- [101] F. Rejon-Barrera and D. Robbins, "Scalar-vector bootstrap", J. High Energy Phys. 2016, 139 (2016).
- [102] J. Richter-Gebert and P. Lebmeir, "Diagrams, tensors and geometric reasoning", Discr. Comp. Geometry **42**, 305–334 (2009).
- [103] T. van Ritbergen, A. N. Schellekens, and J. A. Vermaseren, "Group theory factors for Feynman diagrams", Int. J. Mod. Phys. A **14**, 41–96 (1999).
- [104] A. Roy and T. Quella, "Chiral Haldane phases of SU(N) quantum spin chains", Phys. Rev. B **97**, 155148 (2018).
- [105] D. Rutherford, Substitutional Analysis (Dover, New York, 1948).
- [106] A. Savage, String diagrams and categorification, 2018.

- [107] D. S. Silver, "The new language of mathematics", Amer. Sci. 105, 364 (2017).
- [108] B. Simon, *Representations of Finite and Compact Groups* (Amer. Math. Soc., Providence, RI, 1996).
- [109] M. Sjödahl, "ColorMath a package for color summed calculations in SU(N)", Euro. Phys. J. C **73**, 2310 (2013).
- [110] M. Sjödahl, "ColorFull: a C++ library for calculations in $SU(N_c)$ ", Euro. Phys. J. C **75**, 236 (2015).
- [111] M. Sjödahl and S. Keppeler, "Tools for calculations in color space", in *Proceedings*, 21st International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2013) (PoS, 2013), p. 0166.
- [112] M. Sjödahl and J. Thorén, "Decomposing color structure into multiplet bases", J. High Energy Phys. **2015**, 055 (2015).
- [113] M. Sjödahl and J. Thorén, "QCD multiplet bases with arbitrary parton ordering", J. High Energy Phys. **2018**, 198 (2018).
- [114] G. E. Stedman, *Diagram Techniques in Group Theory* (Cambridge Univ. Press, Cambridge, 1990).
- [115] J. J. Sylvester, "On an application of the new atomic theory to the graphical representation of the invariants and covariants of binary quantics, with three appendices", Amer. J. Math. 1, 64–125 (1878).
- [116] L. Tagliacozzo, A. Celi, and M. Lewenstein, "Tensor networks for lattice gauge theories with continuous groups", Phys. Rev. X 4, 041024 (2014).
- [117] M. Tai, The classical family algebra of the adjoint representation of sl(n), 2013.
- [118] M. Tai, Family algebras and the isotypic components of $g \otimes g$, PhD thesis (Mathematics, Univ. of Pennsylvania, Philladelphia, 2014).
- [119] J. Thorén, Decomposing colour structures into multiplet bases, MA thesis (Lund Univ., 2013).
- [120] J. Thorén, Multiplet Bases, Recursion Relations and Full Color Parton Showers, PhD thesis (Theoretical Physics Dept., Lund, Sweden, 2018).
- [121] A. Trautner, "Systematic construction of basis invariants in the 2HDM", J. High Energy Phys. **2019**, 208 (2019).
- [122] W.-K. Tung, *Group Theory in Physics* (World Scientific, 1985).
- [123] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).
- [124] H. Weigert, "Non-global jet evolution at finite N_c ", Nucl. Phys. B **685**, 321–350 (2004).
- [125] A. Young, "On quantitative substitutional analysis III", Proc. London Math. Soc. **28**, 255–292 (1928).
- [126] A. Young, "The application of substitutional analysis to invariants III", Phil. Trans. Roy. Soc. Lond. A **234**, 79–114 (1935).

[127] X.-D. Yu, T. Simnacher, N. Wyderka, H. C. Nguyen, and O. Gühne, "A complete hierarchy for the pure state marginal problem in quantum mechanics", Nat. Commun. 12, 1012 (2021).

- [128] A. P. Yutsis, I. B. Levinson, and V. V. Vanagas, *The Theory of Angular Momentum* (Gordon and Breach, New York, 1964).
- [129] A. Zvonkin, "Matrix integrals and map enumeration: An accessible introduction", Math. Comput. Modell. **26**, 281–304 (1997).