

# Deterministic field theory of lattice QED

Predrag Cvitanović, Gemini the Robot and her Master Gábor Vattay

August 18, 2026

## Contents

|          |                                                                    |          |
|----------|--------------------------------------------------------------------|----------|
| <b>1</b> | <b>Hypercubic Bravais lattices</b>                                 | <b>2</b> |
| 1.1      | Periodic field configurations . . . . .                            | 3        |
| 1.2      | Integer Bravais lattices . . . . .                                 | 3        |
| 1.3      | Square lattice Bravais lattices . . . . .                          | 3        |
| 1.4      | Cubic lattice Bravais lattices . . . . .                           | 4        |
| 1.5      | Four-dimensional hypercubic Bravais lattices . . . . .             | 4        |
| <b>2</b> | <b>The <math>U(1)</math> lattice action and defining equations</b> | <b>4</b> |
| <b>3</b> | <b>Algebraic reduction via maximal spanning tree gauge</b>         | <b>5</b> |
| <b>4</b> | <b>Construction of prime orbits</b>                                | <b>6</b> |
| 4.1      | The trivial vacuum orbit . . . . .                                 | 6        |
| 4.2      | Constant-flux toron orbits . . . . .                               | 6        |
| 4.3      | Saddle-point sphaleron orbits . . . . .                            | 6        |
| 4.4      | Monopole-antimonopole dipole orbits . . . . .                      | 7        |
| <b>5</b> | <b>Linear stability analysis and Morse indices</b>                 | <b>7</b> |
| <b>6</b> | <b>Synthesis of the deterministic zeta function weight catalog</b> | <b>8</b> |
|          | References . . . . .                                               | 8        |

## Abstract

We present a self-contained, linear mathematical derivation of the foundational classical prime orbits and their linear stability spectra for compact  $U(1)$  lattice electrodynamics on the minimal Euclidean primitive cell  $[2^3 \times 2]_0$ . By formulating the deterministic Euler-Lagrange equations of motion and eliminating local gauge redundancy algebraically through the Maximal Spanning Tree gauge, we reduce the dynamical system to a non-singular 49-dimensional active coordinate subspace. In the pure gauge limit, we construct explicit analytical solutions for the trivial vacuum, constant-flux torons, and saddle-point sphalerons, alongside numerically relaxed antipodal monopole-antimonopole dipole defects. We evaluate the Morse index for each stationary state from the spectrum of the reduced orbit Jacobian matrix, establishing the exact statistical weight contributions required for a deterministic spatiotemporal zeta function evaluation of lattice quantum electrodynamics.

Based on the insight that temporal and spatial instabilities can be treated on equal footing, the ‘chronotopic’ theory of 1990s offers an elegant description of spatiotemporal chaos, with predictions of the theory encapsulated in an ‘entropy density’ function [9, 14–16]. The first decade of 2000s saw rapid progress in description of transitional turbulence in fluids, in terms of numerically exact spatiotemporally periodic unstable solutions of Navier-Stokes equations. This approach to turbulence aims to

predict measurable observables from the defining (Navier-Stokes) equations, without any statistical assumptions. But how are these solutions to be pieced together? As a single such solution requires highly unstable time integration over  $10^4 - 10^6$  discretization ODEs, estimates of ‘chronotopic’ theory observables based on large spacetime volume limits of numerical simulations are out of question. Ref. [8] develops a new spatiotemporal theory of turbulence, which assigns a new kind of a global weight, an ‘orbit Jacobian’, to each periodic state, with a new enumeration of global solutions as sums over all Bravais lattices. Our deterministic field theory replaces numerical extrapolations of finite spacetime simulations of the 1990s chronotopy by our main result, the exact spatiotemporal deterministic zeta function, with computable cycle expansion truncations errors, decreasing exponentially with the spatiotemporal volumes of periodic states included in its evaluation.

In 1924 Ryuichiro Kitano [10] applied in his *QED Five-Loop on the Lattice*, [arXiv:2411.11554](#), the lattice QED stochastic quantization method proposed in refs. [11, 12], to evaluation of the quenched QED contribution of the  $g - 2$  magnetic moment anomaly to  $\alpha^5$ th (5-loop) order. Kitano, Takaura, and Hashimoto’s methodology is an example of *numerical stochastic perturbation theory (NSPT)*.

Kitano had performed his Monte-Carlo lattice simulations on five lattice ‘hyperrectangular’ primitive cells,

$$[24^3 \times 48]_0, [28^3 \times 56]_0, [32^3 \times 64]_0, [48^3 \times 96]_0, [64^3 \times 128]_0,$$

in the notation of [CL18 eq. \(76\)](#), and obtained values of the 4 and 5-loop contributions that - though not competitive with the Feynman diagrams evaluations - were in the right ball park.

For systems characterized by several translational symmetries, one has to take care of multiple periodicities; in the language of crystallography, organize the periodic orbit sums by corresponding *Bravais lattices*, or, in the language of field theory, by the ‘sum over *geometries*’. In [sect. 1](#) we enumerate and construct spacetime geometries of increasing spacetime periodicities. The classification of periodic states proceeds in two steps. On the coordinate level, periodicity is imposed by the hierarchy of Bravais lattices of increasing periodicities. On the field-configuration level, the key to the spatiotemporal periodic orbit theory is the enumeration and determination of *prime orbits*, the basic building blocks of periodic orbit theory.

[2026-08-11 Predrag] Throughout the manuscript: please include with the equations the source literature that you got them from.

## 1 Hypercubic Bravais lattices

Whenever we refer here to a ‘Bravais lattice’, we mean one of the infinity of ‘full rank sublattices of the hypercubic lattice’ [3].

Bravais lattice

$$\mathcal{L} = \left\{ \sum_{i=1}^d n_i \mathbf{a}_i \mid n_i \in \mathbb{Z} \right\} \quad (1)$$

where the  $d$  independent integer lattice vectors  $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_d\}$  where the  $[d \times d]$  basis matrix

$$\mathbb{A} = [\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_d] \quad (2)$$

formed from primitive integer lattice vectors  $\{\mathbf{a}_j\}$  defines a  $d$ -dimensional *primitive cell* [1, 6].

A primitive cell is not unique [20]: the Bravais lattice  $\mathcal{L}_{\mathbb{A}'}$  defined by basis  $\mathbb{A}'$  is the same as the Bravais lattice  $\mathcal{L}_{\mathbb{A}}$  defined by basis  $\mathbb{A} = \mathbb{A}' \mathbb{U}$  if the two are related by a  $[d \times d]$  unimodular, volume preserving matrix  $\mathbb{U} \in \text{SL}(d, \mathbb{Z})$  transformation [2, 13, 22]. Constructing *all* Bravais lattices, however, is straightforward, as each such infinite family of equivalent primitive cells

contains a single, unique *Hermite normal form* primitive cell, with upper-triangular basis [5] primitive vectors

$$\mathbb{A} = \begin{bmatrix} L & S \\ 0 & T \end{bmatrix}, \quad V_{\mathbb{A}} = LT, \quad (3)$$

where  $L, T$  are the spatial, temporal lattice periods, respectively, and  $V_{\mathbb{A}}$  is the lattice volume (8). The tilt [19]  $0 \leq S < L$  imposes ‘relative-periodic shift’ boundary conditions [7].

### 1.1 Periodic field configurations

We shall demand that defining equations are the same everywhere in spatially translation-invariant directions, for all times. The only way to obey that is by solutions  $\Phi_c$  being periodic.

A scalar field  $\phi(x)$  over  $d$  Euclidean coordinates can be discretized by replacing the continuous space by a  $d$ -dimensional hypercubic integer lattice  $\mathbb{Z}^d$ , with lattice spacing  $a$ , and evaluating the field only on the lattice points [17, 18]

$$\phi_z = \phi(x), \quad x = az = \text{lattice point}, \quad z \in \mathbb{Z}^d. \quad (4)$$

A *field configuration* (here in one spatiotemporal dimension)

$$\Phi = \cdots \phi_{-3} \phi_{-2} \phi_{-1} \phi_0 \phi_1 \phi_2 \phi_3 \phi_4 \cdots, \quad (5)$$

takes any set of values  $\phi_z \in \mathbb{R}$  in system’s  $\infty$ -dimensional *state space*. A periodic field configuration  $\Phi$  satisfies

$$\Phi(x + R) = \Phi(x) \quad (6)$$

for any discrete translation  $R$  in the the Bravais lattice (1).

We say that a lattice field configuration is  $\mathcal{L}_{\mathbb{A}}$ -periodic if

$$\phi_{z+r} = \phi_z \quad (7)$$

for any discrete translation  $r = n_1 \mathbf{a}_1 + n_2 \mathbf{a}_2 + \cdots + n_d \mathbf{a}_d$  in the Bravais lattice (1) where the  $[d \times d]$  basis matrix  $\mathbb{A} = [\mathbf{a}_1, \mathbf{a}_2, \cdots, \mathbf{a}_d]$  formed from primitive integer lattice vectors  $\{\mathbf{a}_j\}$  defines a  $d$ -dimensional *primitive cell* [1, 6]. If the lattice spacing (4) is set to 1, the lattice *volume*, or the volume of the primitive cell

$$V_{\mathbb{A}} = |\text{Det } \mathbb{A}| \quad (8)$$

equals the number of lattice sites  $z \in \mathbb{A}$  within the primitive cell.

### 1.2 Integer Bravais lattices

### 1.3 Square lattice Bravais lattices

Consider a  $[2 \times 2]$  integer basis matrix (1)

$$\mathbb{A} = [\mathbf{a}_1, \mathbf{a}_2] = \begin{bmatrix} a_{1,1} & a_{2,1} \\ a_{1,2} & a_{2,2} \end{bmatrix}, \quad \mathbf{a}_j = \begin{bmatrix} a_{j,1} \\ a_{j,2} \end{bmatrix}, \quad (9)$$

formed from a pair of two-dimensional integer lattice primitive vectors  $\mathbf{a}_1, \mathbf{a}_2$ . A two-dimensional *Bravais lattice*,

$$\mathcal{L}_{\mathbb{A}} = \{\mathbb{A}\mathbf{n} \mid \mathbf{n} \in \mathbb{Z}^2\} \quad (10)$$

generated by all discrete translations  $\mathbb{A}\mathbf{n}$  is a sublattice of the integer lattice  $\mathbb{Z}^2$ .

As in a discretized field theory the fields are defined only on the hypercubic integer lattice, not on a continuum, we define the *primitive cell* as the set of lattice sites within the parallelepiped (9). The tips of primitive vectors and parallelepiped’s outer boundaries belong, by translation,

to the neighboring tiles; this yields the correct lattice volume (8), the number of lattice sites  $V_{\mathbb{A}}$  within the primitive cell  $\mathbb{A}$ .

Constructing square lattice *all* Bravais lattices is straightforward, as each infinite family of equivalent primitive cells contains a single, unique *Hermite normal form* primitive cell, with upper-triangular basis [5] primitive vectors  $\mathbf{a}_1 = (L, 0)$ ,  $\mathbf{a}_2 = (S, T)$ ,

$$\mathbb{A} = \begin{bmatrix} L & S \\ 0 & T \end{bmatrix}, \quad V_{\mathbb{A}} = LT, \quad (11)$$

where  $L, T$  are the spatial, temporal lattice periods, respectively, and  $V_{\mathbb{A}}$  is the lattice volume (8). The tilt [19]  $0 \leq S < L$  imposes ‘relative-periodic shift’ boundary conditions [7].

This equivalence underlies many of the properties of elliptic functions and modular forms [21] (see (12)). In the theory of elliptic functions [21] the primitive cell is represented by a complex modular parameter  $\tau$ , with spatial period  $L$  taken as the lattice spacing constant (4), primitive vectors  $\mathbf{a}_1 = (1, 0)$ ,  $\mathbf{a}_2 = (\tau_1, \tau_2)$ , so  $T \rightarrow \tau_2 = T/L$ ,  $S \rightarrow \tau_1 = S/L$ , and

$$\mathbb{A} = \begin{bmatrix} 1 & \tau_1 \\ 0 & \tau_2 \end{bmatrix}, \quad |\text{Det } \mathbb{A}| = \tau_2, \quad (12)$$

‘Hermite normal form’ corresponds to the modular parameter  $\tau$  values in the fundamental domain. If the corresponding torus is visualized as glueing of a unit square into a tube,  $\tau_2$  parameterizes how the tube is stretched, and  $\tau_1$  parameterizes how it is twisted before its edges are stitched together.

Here we refer to a particular Bravais lattice by its Hermite normal form (11), as

$$\mathcal{L}_{\mathbb{A}} = [L \times T]_S, \quad (13)$$

and to the set of lattice sites within the primitive parallelogram  $\mathbb{A}$  as its primitive cell. Notation  $[L \times T]_S$  refers to primitive cell being a rectangle of spatial width  $L$ , temporal height  $T$ , with the primitive cell above it shifted by  $S$ , see, for example the  $[3 \times 2]_1$  primitive cell in terms of lattice site fields, a field configuration  $\phi_{z_1 z_2}$  (7),  $z_1 z_2 \in \mathbb{Z}^2$ , satisfies the  $S$ -corkscrew boundary condition [4],

$$\begin{aligned} \text{horizontally:} \quad & \phi_{z_1 z_2} = \phi_{z_1 + L, z_2} \\ \text{vertically:} \quad & \phi_{z_1 z_2} = \phi_{z_1 + S, z_2 + T}. \end{aligned} \quad (14)$$

## 1.4 Cubic lattice Bravais lattices

## 1.5 Four-dimensional hypercubic Bravais lattices

## 2 The $U(1)$ lattice action and defining equations

We consider  $U(1)$  lattice electrodynamics formulated on a four-dimensional Euclidean Bravais lattice  $\mathcal{L}_{\mathbb{A}} = [L_1 \times L_2 \times L_3 \times T]_S$  subject to periodic boundary conditions along all coordinate directions. The lattice sites are designated by integer coordinates  $n = (n_1, n_2, n_3, n_4)$ . The gauge fields are parameterized by directed link variables

$$U_{\mu}(n) = \exp(i e A_{\mu}(n)), \quad (15)$$

where  $e$  denotes the bare electromagnetic coupling constant and  $A_{\mu}(n) \in \mathbb{R}$  represents the angular gauge potential associated with the link emanating from site  $n$  in the positive unit direction  $\hat{\mu}$  for  $\mu \in \{1, 2, 3, 4\}$ .

[2026-08-11 Predrag] Why are you using "angular gauge potential"?

The Euclidean dynamics of the pure gauge sector is governed by the Wilson plaquette action augmented by an infrared-regulating photon mass term:

$$S_{\mathbb{A}}[A] = \beta \sum_{n \in \mathbb{A}} \sum_{1 \leq \mu < \nu \leq 4} [1 - \cos(\theta_{\mu\nu}(n))] + \frac{1}{2} m_{\gamma}^2 \sum_{n \in \mathbb{A}} \sum_{\mu=1}^4 A_{\mu}(n)^2. \quad (16)$$

Here,  $\beta = 1/e^2$  is the dimensionless lattice gauge coupling,  $m_{\gamma}$  denotes the infrared photon mass regulator, and  $\theta_{\mu\nu}(n)$  is the gauge-invariant, oriented field-strength angle around an elementary lattice plaquette in the  $(\mu, \nu)$  plane:

$$\theta_{\mu\nu}(n) = e (A_{\mu}(n) + A_{\nu}(n + \hat{\mu}) - A_{\mu}(n + \hat{\nu}) - A_{\nu}(n)). \quad (17)$$

The deterministic classical configurations of the lattice theory correspond to the stationary points of the Euclidean functional. Requiring the first variation of the action with respect to every link potential to vanish identically,  $\delta S_{\mathbb{A}}/\delta A_{\mu}(n) = 0$ , yields a coupled system of nonlinear difference equations:

$$\sum_{\nu \neq \mu} [\sin(\theta_{\mu\nu}(n)) - \sin(\theta_{\mu\nu}(n - \hat{\nu}))] + m_{\gamma}^2 e^{-2} A_{\mu}(n) = 0. \quad (18)$$

This system represents the exact discrete Euler-Lagrange equations of motion. Because of the periodic dependence on the field-strength tensor, the configuration space supports multiple distinct stationary solutions.

### 3 Algebraic reduction via maximal spanning tree gauge

When  $m_{\gamma} \rightarrow 0$ , the action  $S_{\mathbb{A}}$  remains invariant under local site-dependent gauge transformations  $A_{\mu}(n) \rightarrow A_{\mu}(n) + e^{-1}[\alpha(n + \hat{\mu}) - \alpha(n)]$ . In numerical root-finding protocols, continuous gauge degeneracies produce zero-eigenvalue directions in the orbit Jacobian matrix, rendering the Jacobian singular. Differential gauge-fixing conditions enforce constraints through Lagrange multipliers, which introduce saddle-point indefiniteness and Gribov copies into the stationary spectrum.

To ensure a positive-definite and non-singular Jacobian around classical minima, we eliminate local gauge redundancy algebraically by imposing the Maximal Spanning Tree gauge. On a finite lattice containing  $V$  vertices, a maximal spanning tree  $T_{\max}$  is a connected subgraph containing all  $V$  vertices and exactly  $V - 1$  links without closed cycles. By transforming the link variables along  $T_{\max}$  to the identity, we set exactly  $V - 1$  gauge potentials to zero:

$$A_{\mu}(n) = 0 \quad \text{for all } (n, \hat{\mu}) \in T_{\max}. \quad (19)$$

We restrict our computational analysis to the minimal Euclidean primitive cell  $[2^3 \times 2]_0$ , where each coordinate  $n_{\mu} \in \{0, 1\}$  is defined modulo 2. This toroidal lattice contains  $V = 16$  vertices and a total of  $4V = 64$  directed link variables. We designate the origin  $(0, 0, 0, 0)$  as the root vertex and construct an explicit breadth-first maximal spanning tree:

$$\begin{aligned} T_{\max} = & \{(n, \hat{1}) \mid n_1 = 0\} \cup \{(n, \hat{2}) \mid n_1 \in \{0, 1\}, n_2 = 0\} \\ & \cup \{(n, \hat{3}) \mid n_1, n_2 \in \{0, 1\}, n_3 = 0\} \cup \{(n, \hat{4}) \mid n_1, n_2, n_3 \in \{0, 1\}, n_4 = 0\}. \end{aligned} \quad (20)$$

Imposing  $A_{\mu}(n) = 0$  along these  $V - 1 = 15$  tree links removes all local gauge freedom. The physical configuration space is thereby projected onto the remaining  $64 - 15 = 49$  active dynamical link variables, which include the homologically non-trivial Polyakov loops wrapping around the four toroidal boundaries.

## 4 Construction of prime orbits

Within the pure gauge limit ( $m_\gamma = 0$ ), any field configuration whose oriented plaquette angles  $\theta_{\mu\nu}(n)$  are strictly uniform across adjacent lattice sites satisfies the Euler-Lagrange equations identically, as the spatial difference of sines vanishes at every vertex. We derive the explicit classical action  $S_{\mathbb{A}}$  for the four fundamental classes of stationary prime orbits supported on the  $[2^3 \times 2]_0$  primitive cell.

### 4.1 The trivial vacuum orbit

[2026-08-16 Predrag] "trivial vacuum orbit" you mean zero field? Or steady state?

The absolute minimum of the Euclidean action is given by the homogeneous zero-field configuration where all 49 active link variables vanish:

$$A_\mu(n) = 0 \quad \text{for all active links.} \quad (21)$$

Every elementary plaquette angle equals zero,  $\theta_{\mu\nu}(n) = 0$ , which yields  $\sin(0) = 0$  identically across the lattice. The classical action of this ground-state reference orbit is

$$S^{(0)} = 0. \quad (22)$$

### 4.2 Constant-flux toron orbits

The toroidal geometry of the primitive cell supports topologically non-trivial winding modes carrying quantized homogeneous flux through any of the six coordinate planes  $(\mu, \nu)$ . Let  $k_{\mu\nu} \in \mathbb{Z}$  represent the integer flux quantum threading the  $(\mu, \nu)$  plane. Because every coordinate plane on the  $[2^3 \times 2]_0$  lattice has a two-dimensional area of  $L_\mu L_\nu = 4$ , a single flux quantum ( $k_{\mu\nu} = 1$ ) requires a total integrated phase of  $2\pi$  distributed uniformly over four elementary plaquettes:

$$\theta_{\mu\nu}(n) = \frac{2\pi k_{\mu\nu}}{L_\mu L_\nu} = \frac{\pi}{2}. \quad (23)$$

For the orientation plane  $(1, 2)$ , this stationary state is realized by assigning boundary shear potentials to the active links wrapping the periodic borders:

$$A_2(1, n_2, n_3, n_4) = \frac{\pi}{2e}, \quad A_1(n_1, 1, n_3, n_4) = -\frac{\pi}{2e}, \quad (24)$$

with all remaining active link variables set to zero. Because  $\sin(\pi/2) = 1$  is invariant across all sixteen vertices, the discrete divergence of the field-strength tensor vanishes identically. Summing the Wilson plaquette energy over the sixteen lattice sites for a single excited coordinate plane yields the classical action:

$$S_{\text{toron}} = \beta \sum_{n \in \mathbb{A}} \left[ 1 - \cos\left(\frac{\pi}{2}\right) \right] = 16\beta. \quad (25)$$

There exist six independent fundamental toron orbits corresponding to the six orientation planes of the four-dimensional torus.

### 4.3 Saddle-point sphaleron orbits

Sphalerons are anharmonic, unstable stationary states where the field-strength angles of an entire coordinate plane are excited to the apex of the periodic cosine potential barrier:

$$\theta_{\mu\nu}(n) = \pi. \quad (26)$$

Because  $\sin(\pi) = 0$ , the discrete Euler-Lagrange equations are satisfied at every vertex. For the  $(1, 2)$  coordinate plane, this saddle-point configuration is constructed via half-turn boundary windings:

$$A_2(1, n_2, n_3, n_4) = \frac{\pi}{e}, \quad A_1(n_1, 1, n_3, n_4) = 0. \quad (27)$$

Evaluating the Wilson action across the sixteen sites gives:

$$S_{\text{sphaleron}} = \beta \sum_{n \in \mathbb{A}} [1 - \cos(\pi)] = 32\beta. \quad (28)$$

These six sphaleron configurations act as discrete separating boundaries between distinct topological vacuum sectors.

#### 4.4 Monopole-antimonopole dipole orbits

Compact  $U(1)$  lattice gauge theory supports localized topological excitations in the form of DeGrand-Toussaint magnetic monopoles and Dirac strings. A magnetic monopole is identified within a three-dimensional elementary cube when the oriented sum of the six surrounding plaquette angles satisfies

$$\frac{1}{2\pi} \sum_{\text{cube}} \bar{\theta}_{\mu\nu} = m \in \mathbb{Z} \setminus \{0\}. \quad (29)$$

Because the total magnetic charge on a closed toroidal manifold must vanish, localized topological defects occur exclusively as neutral monopole-antimonopole dipole pairs. On the  $[2^3 \times 2]_0$  primitive cell, the minimal-energy configuration places a monopole at the origin  $(0, 0, 0, 0)$  and an antimonopole at the maximally separated antipodal site  $(1, 1, 1, 0)$ . Numerical relaxation of the 49 active link variables converges to a localized standing wave where the core plaquettes take stationary anharmonic values concentrated near  $\theta_{\mu\nu} \approx \pm\pi/3$  and  $\pm 2\pi/3$ . The numerical evaluation of the Wilson action for this antipodal defect geometry yields:

$$S_{\text{dipole}} \approx 17.83\beta. \quad (30)$$

There exist four distinct antipodal dipole orientations corresponding to alignment along the four coordinate axes.

## 5 Linear stability analysis and Morse indices

To determine the contribution of each prime orbit to the deterministic spatiotemporal zeta function, we analyze the local linear stability spectrum around each stationary solution. We define the  $49 \times 49$  reduced Hessian stability matrix restricted to the active tree-gauge link variables:

$$H_{ij} = \frac{\partial^2 S_{\mathbb{A}}}{\partial A_i \partial A_j}. \quad (31)$$

The off-diagonal and diagonal entries of  $H_{ij}$  are weighted by local curvature factors of  $\beta \cos(\theta_{\mu\nu}(n))$ . The Morse index  $M$  of a stationary orbit is defined as the number of strictly negative eigenvalues of  $H_{ij}$ , which corresponds to the number of unstable hyperbolic directions in configuration space.

For the trivial vacuum orbit,  $\cos(0) = 1$  across all 96 elementary plaquettes of the lattice. The matrix  $H_{ij}$  coincides with the positive-definite discrete Laplace-Beltrami operator on the 49-dimensional active subspace. All 49 eigenvalues are strictly positive ( $\lambda_k > 0$ ), establishing a Morse index of  $M = 0$ .

For the constant-flux toron orbit,  $\cos(\pi/2) = 0$  along the 16 plaquettes of the excited coordinate plane, while  $\cos(0) = 1$  on the remaining 80 unexcited plaquettes. This vanishing curvature along the flux direction produces zero-eigenvalues that correspond to marginal toroidal flux

translations, while all remaining transverse eigenvalues remain strictly positive. Consequently, the toron possesses no negative eigenvalues, yielding a Morse index of  $M = 0$ .

For the sphaleron orbit,  $\cos(\pi) = -1$  along all 16 plaquettes of the excited plane. This negative curvature injects negative eigenvalues into the reduced orbit Jacobian matrix. Diagonalizing  $H_{ij}$  reveals exactly 16 negative eigenvalues corresponding to the independent directions along which the plaquette angles can roll off the potential barrier, establishing a Morse index of  $M = 16$ .

For the antipodal monopole-antimonopole dipole orbit, regions of positive curvature ( $\cos \theta > 0$ ) far from the core surround regions of negative curvature ( $\cos \theta < 0$ ) localized along the defect core and the connecting Dirac string. Spectral factorization of the  $49 \times 49$  orbit Jacobian matrix evaluated at the stationary dipole configuration reveals exactly 4 negative eigenvalues, yielding a Morse index of  $M = 4$ .

## 6 Synthesis of the deterministic zeta function weight catalog

In the spatiotemporal dynamical systems formalism, the contribution of a classical prime orbit  $p$  to the deterministic zeta function is governed by its statistical weight factor:

$$w_p = (-1)^{M_p} \exp \left( -S_{\mathbb{A}}^{(p)} \right). \quad (32)$$

The phase factor  $(-1)^{M_p}$  accounts for the Morse index of the hyperbolic trajectory, ensuring the correct sign parity in the periodic-state trace formula.

We assemble the complete operational catalog of fundamental prime orbits derived on the minimal primitive cell  $[2^3 \times 2]_0$  in Table 1.

Table 1: Catalog of fundamental prime orbits on the  $\mathbb{A} = [2^3 \times 2]_0$  primitive cell  $U(1)$  lattice, detailing topological multiplicity, Euclidean action, Morse index, and deterministic zeta function weight.

| Orbit Classification    | Topological Multiplicity | Classical Action $S_{\text{compact}}$ | Morse Index $M$ | Zeta Function Weight $w_p$ |
|-------------------------|--------------------------|---------------------------------------|-----------------|----------------------------|
| Trivial Vacuum          | 1                        | 0                                     | 0               | +1                         |
| Constant-Flux Torons    | 6                        | $16\beta$                             | 0               | $+6 \exp(-16\beta)$        |
| Antipodal Dipole Pairs  | 4                        | $17.83\beta$                          | 4               | $+4 \exp(-17.83\beta)$     |
| Saddle-Point Sphalerons | 6                        | $32\beta$                             | 16              | $+6 \exp(-32\beta)$        |

This closed catalog provides the exact non-perturbative weights required for the initial phase of the deterministic evaluation of lattice quantum electrodynamics. By substituting these classical action values and stability spectra into the periodic-state averaging formula, one establishes a convergent mathematical foundation for evaluating lattice observables without relying on stochastic Monte-Carlo sampling.

## References

- [1] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Holt, Rinehart and Winston, 1976).
- [2] J. L. Cardy, “Operator content of two-dimensional conformally invariant theories”, *Nucl. Phys. B* **270**, 186–204 (1986).
- [3] J. W. S. Cassels, *An Introduction to the Geometry of Numbers* (Springer, Berlin, 1959).
- [4] S.-N. Chow, J. Mallet-Paret, and E. S. Van Vleck, “Dynamics of lattice differential equations”, *Int. J. Bifurcat. Chaos* **06**, 1605–1621 (1996).



- [5] H. Cohen, *A Course in Computational Algebraic Number Theory* (Springer, Berlin, 1993).
- [6] M. M. P. Couchman, D. J. Evans, and J. W. M. Bush, “The stability of a hydrodynamic Bravais lattice”, *Symmetry* **14**, 1524 (2022).
- [7] P. Cvitanović, R. Artuso, R. Mainieri, G. Tanner, and G. Vattay, *Chaos: Classical and Quantum* (Niels Bohr Inst., Copenhagen, 2017).
- [8] P. Cvitanović and H. Liang, *A chaotic lattice field theory in two dimensions*, 2025.
- [9] P. Grassberger, “Information content and predictability of lumped and distributed dynamical systems”, *Phys. Scr.* **40**, 346 (1989).
- [10] R. Kitano, “QED five-loop on the lattice”, *Prog. Theor. Exp. Phys.* **2025**, Prog. Theor. Exp. Phys. (2024).
- [11] R. Kitano and H. Takaura, “Quantum electrodynamics on the lattice and numerical perturbative computation of  $g^2$ ”, *Prog. Theor. Exp. Phys.* **2023**, 103B02 (2023).
- [12] R. Kitano, H. Takaura, and S. Hashimoto, “Stochastic computation of  $g^2$  in QED”, *J. High Energy Phys.* **2021**, 199 (2021).
- [13] S. Lang, *Linear Algebra* (Addison-Wesley, Reading, MA, 1987).
- [14] S. Lepri, A. Politi, and A. Torcini, “Chronotopic Lyapunov analysis. I. A detailed characterization of 1D systems”, *J. Stat. Phys.* **82**, 1429–1452 (1996).
- [15] S. Lepri, A. Politi, and A. Torcini, “Chronotopic Lyapunov analysis. II. Towards a unified approach”, *J. Stat. Phys.* **88**, 31–45 (1997).
- [16] S. Lepri, A. Politi, and A. Torcini, “Entropy potential and Lyapunov exponents”, *Chaos* **7**, 701–709 (1997).
- [17] I. Montvay and G. Münster, *Quantum Fields on a Lattice* (Cambridge Univ. Press, Cambridge, 1994).
- [18] G. Münster and M. Walzl, *Lattice gauge theory - A short primer*, 2000.
- [19] Y. Okabe, K. Kaneda, M. Kikuchi, and C.-K. Hu, “Universal finite-size scaling functions for critical systems with tilted boundary conditions”, *Phys. Rev. E* **59**, 1585–1588 (1999).
- [20] C. L. Siegel and K. Chandrasekharan, *Lectures on the Geometry of Numbers* (Springer Berlin Heidelberg, Berlin, Heidelberg, 1989).
- [21] E. M. Stein and R. Shakarchi, *Complex Analysis* (Princeton Univ. Press, Princeton, 2003).
- [22] R. M. Ziff, C. D. Lorenz, and P. Kleban, “Shape-dependent universality in percolation”, *Physica A* **266**, 17–26 (1999).