

## Important Notice to Authors

Attached is a PDF proof of your forthcoming article in *Physical Review Letters*. The article accession code is LR15587. Your paper will be in the following section of the journal: LETTERS — Nonlinear Dynamics, Fluid Dynamics, Classical Optics, etc.

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, CrossRef, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate PostScript files containing color appear in color in the online journal. However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<http://journals.aps.org/authors/color-figures-print>).

***No further publication processing will occur until we receive your response to this proof.***

## Specific Questions and Comments to Address for This Paper

The numbered items below correspond to numbers in the margin of the proof pages pinpointing the source of the question and/or comment. The numbers will be removed from the margins prior to publication.

- 1 Second proof: Please carefully confirm that all first-proof corrections were addressed and that changes were made accurately.
- 2 Second proof: Please confirm the paper is now ready to be published in its current form.
- 3 Second proof: Please check the updated email address for corresponding author.
- 4 Second proof: Please check the updated acknowledgement section.
- 5 Second proof: Please note that the Ref. [13] was retained as it is. Please check.

## Titles in References

The editors now encourage insertion of article titles in references to journal articles and e-prints. This format is optional, but if chosen, authors should provide titles for *all* eligible references. If article titles remain missing from eligible references, the production team will remove the existing titles at final proof stage.

## Funding Information

Information about an article's funding sources is now submitted to CrossRef to help you comply with current or future funding agency mandates. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. CrossRef's FundRef registry (<http://www.crossref.org/fundref/>) is the definitive registry of funding agencies. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

- National Science Foundation, FundRef ID <http://dx.doi.org/10.13039/1000000001> (United States/Virginia)
- Japan Society for the Promotion of Science, FundRef ID <http://dx.doi.org/10.13039/501100001691> (Japan)
- Ministry of Education, Culture, Sports, Science, and Technology, FundRef ID <http://dx.doi.org/10.13039/501100001700> (Japan)

## Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.
- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- References: Please check to ensure that titles are given as appropriate.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in “x-ray diffraction,” “4-mm-long gas cell,” and “*R*-matrix theory.” However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in “emission by x rays,” “was 4 mm in length,” and “the *R* matrix is tested.”  
Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.
- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

## Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to [aps-robot@luminad.com](mailto:aps-robot@luminad.com). Include the accession code LR15587 in the subject line.
- **Fax:** Return this proof with corrections to +1.855.808.3897.

## If You Need to Call Us

You may leave a voicemail message at +1.855.808.3897. Please reference the accession code and the first author of your article in your voicemail message. We will respond to you via email.

# Estimating the Dimension of an Inertial Manifold from Unstable Periodic Orbits

X. Ding,<sup>1,\*</sup> H. Chaté,<sup>2,3</sup> P. Cvitanović,<sup>1</sup> E. Siminos,<sup>4</sup> and K. A. Takeuchi<sup>5</sup>

<sup>1</sup>*Center for Nonlinear Science, School of Physics, Georgia Institute of Technology, Atlanta, Georgia 30332-0430, USA*

<sup>2</sup>*Service de Physique de l'Etat Condensé, CEA, CNRS, Université Paris-Saclay, 91191 Gif-sur-Yvette, France*

<sup>3</sup>*Beijing Computational Science Research Center, Beijing 100094, China*

<sup>4</sup>*Department of Physics, Chalmers University of Technology, Gothenburg SE-412 96, Sweden*

<sup>5</sup>*Department of Physics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan*

(Received 28 April 2016)

We provide numerical evidence that a finite-dimensional inertial manifold on which the dynamics of a chaotic dissipative dynamical system lives can be constructed solely from the knowledge of a set of unstable periodic orbits. In particular, we determine the dimension of the inertial manifold for the Kuramoto-Sivashinsky system and find it to be equal to the “physical dimension” computed previously via the hyperbolicity properties of covariant Lyapunov vectors.

DOI:

Dynamics in chaotic dissipative systems is expected to land, after a transient period of evolution, on a finite-dimensional object in state space called the inertial manifold [1–5]. This is true even for infinite-dimensional systems described by partial differential equations and offers hope that their asymptotic dynamics may be represented by a finite set of ordinary differential equations, a drastically simplified description. The existence of a finite-dimensional inertial manifold has been established for systems such as the Kuramoto-Sivashinsky, the complex Ginzburg-Landau, and some reaction-diffusion systems [2]. For the Navier-Stokes flows, its existence remains an open problem [3], but dynamical studies, such as the determination of sets of periodic orbits embedded in a turbulent flow [6,7], strengthen the case for a geometrical description of turbulence. However, while mathematical approaches may provide rigorous bounds on the dimensions of inertial manifolds, their constructive description remains a challenge.

Recent progress towards this aim came from numerical investigations of the covariant Lyapunov vectors of spatiotemporally chaotic flows [8,9], made possible by the algorithms developed in Refs. [10–12]. These works have revealed that the tangent space of a generic spatially extended dissipative system is split into two hyperbolically decoupled subspaces: a finite-dimensional subspace of “entangled” or “physical” Lyapunov modes (referred to in what follows as the “physical manifold”), which is presumed to capture all long-time dynamics, and the remaining infinity of transient (“isolated,” “spurious”) Lyapunov modes. Covariant Lyapunov vectors span the Oseledec subspaces [13,14] and thus indicate the intrinsic directions of growth or contraction at every point on the physical manifold. The dynamics of the vectors that span the physical manifold is entangled, with frequent tangencies between them. The transient modes, on the other hand, are damped so strongly that they are isolated—at no time

do they couple by tangencies to the entangled modes. It was conjectured in Refs. [8,9] that the physical manifold provides a local linear approximation to the inertial manifold at any point on the attractor and that the dimension of the inertial manifold is given by the number of entangled Lyapunov modes. Further support for this conjecture was provided by Ref. [15], which verified that the vectors connecting pairs of recurrent points—points on the chaotic trajectory far apart in time but nearby each other in state space—are confined within the local tangent space of the physical manifold.

While these works showed that the physical manifold captures the finite dimensionality of the inertial manifold, they do not tell us much about how this inertial manifold is actually laid out in state space.

In this Letter, we go one important step further and show that the finite-dimensional physical manifold can be precisely embedded in its infinite-dimensional state space, thus opening a path towards its explicit construction. The key idea [16] is to populate the inertial manifold by an infinite hierarchy of unstable time-invariant solutions, such as periodic orbits, an invariant skeleton which, together with the local “tiles” obtained by linearization of the dynamics, fleshes out the physical manifold. Chaos can then be viewed as a walk on the inertial manifold, chaperoned by the nearby unstable solutions embedded in the physical manifold. Unstable periodic orbits have already been used to compute global averages of spatiotemporally chaotic flows [6,17–19].

There are infinitely many unstable orbits, and each of them possesses infinitely many Floquet modes. While in the example that we study here we do not have a detailed understanding of the organization of periodic orbits (their symbolic dynamics), we show that one needs to consider only a finite number of them to tile the physical manifold to a reasonable accuracy. We also show, for the first time, that each local tangent tile spanned by the Floquet vectors of an

unstable periodic orbit splits into a set of entangled Floquet modes and the remaining set of transient modes. Furthermore, we verify numerically that the entangled Floquet manifold coincides locally with the physical manifold determined by the covariant Lyapunov vector approach.

Throughout this Letter, we focus on the one-dimensional Kuramoto-Sivashinsky equation [20,21], chosen here as a prototypical dissipative partial differential equation that exhibits spatiotemporal chaos [22,23]:

$$u_t + u_x u + u_{xx} + u_{xxx} = 0, \quad x \in [0, L], \quad (1)$$

with a real-valued “velocity” field  $u(x, t)$  and the periodic boundary condition  $u(x, t) = u(x + L, t)$ . Following Ref. [19], we fix the size at  $L = 22$ , which is small enough so that unstable orbits are still relatively easy to determine numerically and large enough for the Kuramoto-Sivashinsky equation to exhibit essential features of spatiotemporal chaos [24]. Dynamical evolution traces out a trajectory in the  $\infty$ -dimensional state space,  $\mathbf{x}(t) = f^t(\mathbf{x}(0))$ , with  $\mathbf{x}(t) \equiv u(x, t)$ , where the time-forward map  $f^t$  is obtained by integrating  $\dot{\mathbf{x}} = \mathbf{v}(\mathbf{x})$  up to time  $t$ . The linear stability of the trajectory is described by the Jacobian matrix  $J^t(\mathbf{x}(0)) = \partial \mathbf{x}(t) / \partial \mathbf{x}(0)$ , obtained by integrating  $\dot{J} = AJ$ , where  $A$  is the stability matrix  $A(\mathbf{x}) = \partial \mathbf{v}(\mathbf{x}) / \partial \mathbf{x}$ . We integrate the system (1) numerically, by a pseudospectral truncation [25,26] of  $u(x, t) = \sum_{k=-\infty}^{+\infty} a_k(t) e^{iq_k x}$ ,  $q_k = 2\pi k/L$  to a finite number of Fourier modes. For the numerical accuracy required here, we found 31 Fourier modes (62-dimensional state space) sufficient. The system is invariant under the Galilean transformations  $u(x, t) \rightarrow u(x - ct, t) + c$ , reflection  $u(x, t) \rightarrow -u(-x, t) \equiv \sigma u(x, t)$ , and spatial translations  $u(x, t) \rightarrow u(x + \ell, t) \equiv g(\theta)u(x, t)$ , where  $\theta = 2\pi\ell/L$ . The Galilean symmetry is reduced by setting the mean velocity  $\int dx u(x, t)$ , a conserved quantity, to zero. Because of the  $O(2)$  equivariance of Eq. (1), this system can have two types of relative recurrent orbits (referred to collectively as “orbits” in what follows): preperiodic orbits  $u(x, 0) = \sigma u(x, T_p)$  and relative periodic orbits  $u(x, 0) = g(\theta_p)u(x, T_p)$ , where  $g(\theta_p)$  is the spatial translation by distance  $\ell_p = L\theta_p/2\pi$ . All orbits used here are found by a multiple shooting method together with the Levenberg-Marquardt algorithm (see Ref. [19] for details). In our analysis, we use 200 preperiodic orbits  $\overline{ppo}_{T_p}$  and 200 relative periodic orbits  $\overline{rpo}_{T_p}$ , labeled by their periods  $T_p$ . These are the shortest period orbits taken from the set of over 60 000 determined in Ref. [19] by near-recurrence searches. The method preferentially finds orbits embedded in the long-time attracting set but offers no guarantee that all orbits up to a given period have been found. Floquet multipliers  $\Lambda_j$  and vectors  $\mathbf{e}_j(\mathbf{x})$  are the eigenvalues and eigenvectors of Jacobian matrix  $J_p = \sigma J^{T_p}$  or  $J_p = g(\theta_p)J^{T_p}$  for preperiodic or relative periodic orbits,




FIG. 1. (a) Floquet exponents for  $\overline{ppo}_{10,25}$  (circles),  $\overline{rpo}_{16,31}$  (squares), and Lyapunov exponents of a chaotic trajectory (crosses). The inset shows a close-up of the eight leading exponents.  $\overline{ppo}_{10,25}$  and  $\overline{rpo}_{16,31}$  have, respectively, two and one positive Floquet exponents:  $\lambda_{1,2} = 0.033$  and  $\lambda_1 = 0.328$ . Only one Lyapunov exponent is positive:  $\lambda_1 = 0.048$ . The number of vanishing exponents is always two. The fourth Lyapunov exponent is small but strictly negative:  $\lambda_4 = -0.003$ . (b) Time series of local Floquet exponents  $\lambda_j(\mathbf{x}(t))$  for  $\overline{ppo}_{10,25}$ . (c) Close-up of (b) showing the eight leading exponents. Dashed lines indicate degenerate exponent pairs corresponding to complex Floquet multipliers.

respectively. The Floquet exponents  $\lambda_j$  (if complex, we shall consider only their real parts, with multiplicity 2) are related to multipliers by  $\lambda_j = \ln |\Lambda_j| / T_p$ . A high-accuracy computation of all Floquet exponents and vectors for this finite-dimensional state space (the key to all numerics presented here) has been made possible by the algorithm recently developed in Ref. [27]. For an orbit,  $(\lambda_j, \mathbf{e}_j)$  denotes the  $j$ th Floquet (exponent or vector); for a chaotic trajectory, it denotes the  $j$ th Lyapunov (exponent or vector).

Figure 1(a) shows the Floquet exponent spectra for the two shortest orbits,  $\overline{ppo}_{10,25}$  and  $\overline{rpo}_{16,31}$ , overlaid on the Lyapunov exponents computed from a chaotic trajectory. The basic structure of this spectrum is shared by all 400 orbits used in our study [28]. For chaotic trajectories, hyperbolicity between an arbitrary pair of Lyapunov modes can be characterized by a property called the domination of Oseledec splitting (DOS) [29,30]. Consider a set of finite-time Lyapunov exponents

$$\lambda_j^\tau(\mathbf{x}) \equiv \frac{1}{\tau} \ln \|J^\tau(\mathbf{x}) \mathbf{e}_j(\mathbf{x})\|, \quad (2)$$

with  $L^2$  normalization  $\|\mathbf{e}_j(\mathbf{x})\| = 1$ . A pair of modes  $j < \ell$  is said to fulfill “DOS strict ordering” if  $\lambda_j^\tau(\mathbf{x}) > \lambda_\ell^\tau(\mathbf{x})$  along the entire chaotic trajectory, for  $\tau$  larger than some lower bound  $\tau_0$ . Then this pair is guaranteed not to have tangencies [29,30]. For chaotic trajectories, DOS turned out



to be a useful tool to distinguish entangled modes from hyperbolically decoupled transient modes [8,9]. Periodic orbits are by definition the infinite-time orbits ( $\tau$  can be any repeat of  $T_p$ ), so generically all nondegenerate pairs of modes fulfill DOS. Instead, we find it useful to define, by analogy to the “local Lyapunov exponent” [31], the “local Floquet exponent” as the action of the strain rate tensor [32]  $2D(\mathbf{x}) = A(\mathbf{x})^\top + A(\mathbf{x})$  (where  $A$  is the stability matrix) on the normalized  $j$ th Floquet eigenvector:

$$\lambda_j(\mathbf{x}) = \mathbf{e}_j(\mathbf{x})^\top D(\mathbf{x}) \mathbf{e}_j(\mathbf{x}) = \lim_{\tau \rightarrow 0} \lambda_j^\tau(\mathbf{x}). \quad (3)$$

We find that time series of local Floquet exponents  $\lambda_j(\mathbf{x}(t))$  indicate a decoupling of the leading “entangled” modes from the rest of the strictly ordered, strongly negative exponents [Figs. 1(b) and 1(c)]. Perhaps surprisingly, for every one of the 400 orbits we analyzed, the number of entangled Floquet modes was *always* eight, equal to the previously reported number of the entangled Lyapunov modes for this system [15,28]. This leads to our first surmise: (i) Each individual orbit embedded in the attracting set carries enough information to determine the dimension of the physical manifold.

For an infinite-time chaotic trajectory, hyperbolicity can be assessed by measuring the distribution of minimal principal angles [33,34] between any pair of subspaces spanned by Lyapunov vectors [8–10]. Numerical work indicates that, as the entangled and transient modes are hyperbolically decoupled, the distribution of the angles between these subspaces is bounded away from zero and that observation yields a sharp entangled-transient threshold. This strategy cannot be used for individual orbits, as each one is of a finite period, and the minimal principal angle reached by a pair of Floquet subspaces remains strictly positive. Instead, we measure the angle distribution for a *collection* of orbits and find that the entangled-transient threshold is as sharp as for a long chaotic trajectory: Fig. 2 shows the principal angle distribution between two subspaces  $S_n$  and  $\bar{S}_n$ , with  $S_n$  spanned by the leading  $n$  Floquet vectors and  $\bar{S}_n$  by the rest. As in the Lyapunov analysis of long chaotic trajectories [8], the distributions for small  $n$  indicate strictly positive density as  $\phi \rightarrow 0$ . In contrast, the distribution is strictly bounded away from zero angles for  $n \geq 8$ , the number determined above by the local Floquet exponent analysis. This leads to our second surmise: (ii) The distribution of principal angles for collections of periodic orbits enables us to identify a finite set of *entangled* Floquet modes, the analog of the chaotic trajectories’ entangled covariant Lyapunov vector modes.

It is known, at least for low-dimensional chaotic systems, that a dense set of periodic orbits constitutes the skeleton of a strange attractor [16]. Chaotic trajectories meander around these orbits, approaching them along their stable manifolds and leaving them along their unstable manifolds. If trajectories are indeed confined to a finite-dimensional




FIG. 2. A histogram of the principal angles  $\phi$  between  $S_n$  (the subspace spanned by the  $n$  leading Floquet vectors) and  $\bar{S}_n$  (the subspace spanned by the remaining  $d - n$  Floquet vectors), accumulated over the 400 orbits used in our analysis. (Top panel) For  $n = 1, 2, \dots, 7$  ( $S_n$  within the entangled manifold), the angles can be arbitrarily small. (Bottom panel) For  $n = 8, 10, 12, \dots, 28$  (in the order of the arrow), for which all entangled modes are contained in  $S_n$ , the angles are bounded away from unity.

physical manifold, such shadowing events should take place within the subspace of entangled Floquet modes of the shadowed orbit. To analyze such shadowing, we need to measure the distances between the chaotic trajectories and the invariant orbits. We use *symmetry reduction*, i.e., replacement of a group orbit of states identical up to a symmetry transformation by a single state. Since the translation  $u(x, t) \rightarrow u(x + \ell, t)$  on a periodic domain implies a rotation  $a_k(t) \rightarrow e^{iq_k \ell} a_k(t)$  in Fourier space, we choose to send both trajectories and orbits to the hyperplane  $\text{Im}(a_1) = 0, \text{Re}(a_1) > 0$ , called the first Fourier-mode slice [35], and measure distances therein. This transformation reads

$$\hat{u}(x, t) = g(-\theta(t))u(x, t) \quad (4)$$

with  $\theta(t) = \arg a_1(t)$ . In the slice, both relative periodic orbits and preperiodic orbits are reduced to periodic orbits. From Eq. (4), one easily finds how infinitesimal perturbations  $\delta u(x, t)$  are transformed [36]. This allows us to define the symmetry-reduced tangent space, with the in-slice perturbations  $\delta \hat{u}(x, t)$ , Jacobian matrix  $\hat{J}'(\hat{a})$ , Floquet matrix  $\hat{J}_p(\hat{a})$ , and Floquet vectors  $\hat{e}_j(\hat{a})$ . The dimension of the slice subspace is one less than the full state space: The slice eliminates the marginal translational direction, while the remaining Floquet multipliers  $\Lambda_j$  are unchanged. Therefore, for the system studied here, there are only seven entangled modes, with one marginal mode (time invariance) in the in-slice description, instead of eight and two, respectively, in the full state space description. A shadowing of an orbit  $u_p(x, t')$  by a nearby chaotic

trajectory  $u(x, t)$  is then characterized by the in-slice separation vector

$$\Delta\hat{u}(x, t) \equiv \hat{u}(x, t) - \hat{u}_p(x, t_p), \quad (5)$$

where  $t_p$  is chosen to minimize the in-slice distance  $\|\Delta\hat{u}\|$ .

Now we test whether the  $\Delta\hat{u}(x, t)$  is confined to the tangent space spanned by the entangled in-slice Floquet vectors. To evaluate this confinement, one needs to take into account the nonlinearity of the stable and unstable manifolds for a finite amplitude of  $\Delta\hat{u}(x, t)$ . We decompose the separation vector as

$$\Delta\hat{u}(x, t) = \hat{v}_n(x, t) + \hat{w}_n(x, t), \quad (6)$$

where  $\hat{v}_n(x, t)$  is a vector in the subspace  $\hat{S}_n$  spanned by the leading  $n$  in-slice Floquet vectors and  $\hat{w}_n(x, t)$  is in the orthogonal complement of  $\hat{S}_n$ . If  $n$  is large enough so that  $\hat{S}_n$  contains the local approximation of the inertial manifold, we expect  $\|\hat{w}_n\| \sim \|\hat{v}_n\|^2 \sim \|\Delta\hat{u}\|^2$  because of the smoothness of the inertial manifold; otherwise,  $\|\hat{w}_n\|$  does not vanish as  $\|\Delta\hat{u}\| \rightarrow 0$ . In terms of the angle  $\varphi_n$  between  $\hat{S}_n$  and  $\Delta\hat{u}$ ,  $\sin \varphi_n \sim \|\hat{w}_n\|/\|\hat{v}_n\| \sim \|\Delta\hat{u}\|$  for  $n$  above the threshold, while  $\sin \varphi_n$  remains nonvanishing otherwise.

Following this strategy, we collected segments of a long chaotic trajectory during which it stayed sufficiently close to a specific orbit for at least one period of the orbit. Figure 3(a) illustrates such a shadowing event for  $\overline{ppo}_{33,39}$ . A parametric plot of  $\sin \varphi_n(t)$  vs  $\|\Delta\hat{u}(x, t)\|$  during this event is shown in Fig. 3(b) for  $n = 6, 7, 8$  (blue circles, red squares, and orange triangles, respectively). We can already speculate from such a single shadowing event that  $\sin \varphi_n$  does not necessarily decrease with  $\|\Delta\hat{u}\|$  for  $n < 7$ , while it decreases linearly with  $\|\Delta\hat{u}\|$  for  $n \geq 7$ . This threshold is clearly identified by accumulating data for all the recorded shadowing events with  $\overline{ppo}_{33,39}$  [Fig. 3(c)]:  $\sin \varphi_n$  is confined below a line that depends linearly on  $\|\Delta\hat{u}\|$  if and only if  $n \geq 7$ . Similarly, there is a clear separation in the average of  $\sin \varphi_n$  taken within each bin of the abscissa [Fig. 3(d)]. This indicates that for  $n < 7$  (empty symbols) typical shadowing events manifest a significant deviation of  $\Delta\hat{u}$  from the subspace  $\hat{S}_n$ , whereas for  $n \geq 7$  (solid symbols)  $\Delta\hat{u}$  is always confined to  $\hat{S}_n$ . We therefore conclude that shadowing events are confined to the subspace spanned by the leading seven in-slice Floquet vectors, or equivalently, by all eight entangled Floquet vectors in the full state space. The same conclusion was drawn for  $\overline{ipo}_{34,64}$  [Figs. 3(e) and 3(f)] and five other orbits (not shown). We also verified that, when a chaotic trajectory approaches an orbit, the subspace spanned by all entangled Floquet modes of the orbit coincides with that spanned by all entangled Lyapunov modes of the chaotic trajectory. This implies our third surmise: (iii) The entangled Floquet manifold coincides locally with the entangled Lyapunov manifold, with either capturing the local structure of the inertial manifold.




FIG. 3. (a) Shadowing event between a chaotic trajectory and  $\overline{ppo}_{33,39}$ , drawn over  $2T_p$ . (b) Parametric plot of  $\sin \varphi_n(t)$  vs  $\|\Delta\hat{u}(x, t)\|$  during the single shadowing event shown in (a), for  $n = 6, 7, 8$ . (c) The same as (b), but a total of 230 shadowing events of  $\overline{ppo}_{33,39}$  are used. (d) Average of  $\sin \varphi_n$  in (c), taken within each bin of the abscissa, for  $n = 4, 5, 6, 7, 9, 11, 17, 21, 25$  from top to bottom. (e), (f) The same as (c), (d), respectively, but for 217 shadowing events with  $\overline{ipo}_{34,64}$ . The dashed lines show  $\sin \varphi_n \propto \|\Delta\hat{u}\|$  in all panels.

In summary, we used the Kuramoto-Sivashinsky system to demonstrate by six independent calculations that the tangent space of a dissipative flow splits into entangled vs transient subspaces and to determine the dimension of its inertial manifold. The Lyapunov mode approach of Refs. [8–10,15,28] identifies (i) the entangled Lyapunov exponents, by the dynamics of finite-time Lyapunov exponents [Eq. (2)], and (ii) the entangled tangent manifold, or physical manifold, by measuring the distributions of angles between covariant Lyapunov vectors. The Floquet mode approach [27] developed here shows that (iii) Floquet exponents of each *individual* orbit separate into entangled vs transient (Fig. 1), (iv) for ensembles of orbits, the principal angles between hyperplanes spanned by Floquet vectors separate the tangent space into entangled vs transient (Fig. 2), (v) for a chaotic trajectory shadowing a given orbit, the separation vector lies within the orbit's Floquet entangled manifold (Fig. 3), and (vi) for a chaotic trajectory shadowing a given orbit, the separation vector lies within the covariant Lyapunov vectors' entangled manifold.

All six approaches yield the same inertial manifold dimension, reported in earlier work [15,28]. The approach

with Floquet modes and unstable periodic orbits is constructive, in the sense that periodic points should enable us, in principle (but not attempted in this Letter), to tile the global inertial manifold by local tangent spaces of an ensemble of such points. Moreover, and somewhat surprisingly, our results on individual orbits' Floquet exponents [Figs. 1(b) and 1(c)] and on shadowing of chaotic trajectories (Fig. 3) suggest that *each individual orbit* embedded in the attracting set contains sufficient information to determine the entangled-transient threshold. However, the computation and organization of unstable periodic orbits is still a major undertaking and can currently be carried out only for rather small computational domains [7,19]. The good news is that the approach with entangled Lyapunov modes [8] suffices to determine the inertial manifold dimension, as Lyapunov mode calculations require only averaging over long chaotic trajectories, are much easier to implement, and can be scaled up to much larger domain sizes than  $L = 22$  considered here.

We hope the computational tools introduced in this Letter will eventually contribute to solving outstanding issues of the dynamical system theory, such as the existence of an inertial manifold in the transitional turbulence regime of the Navier-Stokes equations.

**4** We are indebted to R. L. Davidchack for valuable input throughout this project, in particular, for the 60 000 relative periodic orbits that started it, and to M. M. Farazmand for a critical reading of the manuscript. X. D. and P. C. were supported by a grant from G. Robinson, Jr., and by the National Science Foundation under Grants Nos. DMS-1211827 and CMMI-1028133. K. A. T. acknowledges support by JSPS KAKENHI Grant Nos. JP25707033 and JP25103004, as well as the JSPS Core-to-Core Program "Non-equilibrium dynamics of soft matter and information".

\*Corresponding author.

xding@gatech.edu

- [1] P. Constantin, C. Foias, B. Nicolaenko, and R. Temam, *Integral Manifolds and Inertial Manifolds for Dissipative Partial Differential Equations* (Springer, New York, 1989).
- [2] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, 2nd ed. (Springer, New York, 2013).
- [3] R. Temam, *Math. Intell.* **12**, 68 (1990).
- [4] C. Foias, G. R. Sell, and R. Temam, *J. Diff. Equ.* **73**, 309 (1988).
- [5] J. C. Robinson, *Chaos* **5**, 330 (1995).
- [6] J. F. Gibson, J. Halcrow, and P. Cvitanović, *J. Fluid Mech.* **611**, 107 (2008).
- [7] A. P. Willis, K. Y. Short, and P. Cvitanović, *Phys. Rev. E* **93**, 022204 (2016).
- [8] H. L. Yang, K. A. Takeuchi, F. Ginelli, H. Chaté, and G. Radons, *Phys. Rev. Lett.* **102**, 074102 (2009).

- [9] K. A. Takeuchi, H. L. Yang, F. Ginelli, G. Radons, and H. Chaté, *Phys. Rev. E* **84**, 046214 (2011).
- [10] F. Ginelli, P. Poggi, A. Turchi, H. Chaté, R. Livi, and A. Politi, *Phys. Rev. Lett.* **99**, 130601 (2007).
- [11] C. L. Wolfe and R. M. Samelson, *Tellus, Ser. A* **59**, 355 (2007).
- [12] F. Ginelli, H. Chaté, R. Livi, and A. Politi, *J. Phys. A* **46**, 254005 (2013).
- [13] V. I. Oseledec, *Trans. Moscow Math. Soc.* **19**, 197 (1968).
- [14] J.-P. Eckmann and D. Ruelle, *Rev. Mod. Phys.* **57**, 617 (1985).
- [15] H. L. Yang and G. Radons, *Phys. Rev. Lett.* **108**, 154101 (2012).
- [16] P. Cvitanović, R. Artuso, R. Mainieri, G. Tanner, and G. Vattay, *Chaos: Classical and Quantum* (Niels Bohr Institute, Copenhagen, 2016).
- [17] F. Christiansen, P. Cvitanović, and V. Putkaradze, *Non-linearity* **10**, 55 (1997).
- [18] Y. Lan and P. Cvitanović, *Phys. Rev. E* **78**, 026208 (2008).
- [19] P. Cvitanović, R. L. Davidchack, and E. Siminos, *SIAM J. Appl. Dyn. Syst.* **9**, 1 (2010).
- [20] Y. Kuramoto and T. Tsuzuki, *Prog. Theor. Phys.* **55**, 356 (1976).
- [21] G. I. Sivashinsky, *Acta Astronaut.* **4**, 1177 (1977).
- [22] M. C. Cross and P. C. Hohenberg, *Rev. Mod. Phys.* **65**, 851 (1993).
- [23] P. Holmes, J. L. Lumley, and G. Berkooz, *Turbulence, Coherent Structures, Dynamical Systems and Symmetry* (Cambridge University Press, Cambridge, England, 1996).
- [24] J. M. Hyman and B. Nicolaenko, *Physica D (Amsterdam)* **18**, 113 (1986).
- [25] S. M. Cox and P. C. Matthews, *J. Comput. Phys.* **176**, 430 (2002).
- [26] A.-K. Kassam and L. N. Trefethen, *SIAM J. Sci. Comput.* **26**, 1214 (2005).
- [27] X. Ding and P. Cvitanović, *SIAM J. Appl. Dyn. Syst.* (to be published).
- [28] References [8,9,15] include the marginal Galilean symmetry mode in the mode count; here, this mode is absent, as we have set  $\int u(x, t) dx = 0$ . Consequently, the number of entangled modes (the dimension of the physical manifold) differs by one.
- [29] C. Pugh, M. Shub, and A. Starkov, *Bull. Am. Math. Soc.* **41**, 1 (2004).
- [30] J. Bochi and M. Viana, in *Modern Dynamical Systems and Applications*, edited by M. Brin, B. Hasselblatt, and Y. Pesin (Cambridge University Press, Cambridge, England, 2004), pp. 271–297.
- [31] H. Bosetti and H. A. Posch, *Commun. Theor. Phys.* **62**, 451 (2014).
- [32] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon, Oxford, 1959).
- [33] A. Björck and G. H. Golub, *Math. Comput.* **27**, 579 (1973).
- [34] A. V. Knyazev and M. E. Argentati, *SIAM J. Sci. Comput.* **23**, 2008 (2002).
- [35] N. B. Budanur, P. Cvitanović, R. L. Davidchack, and E. Siminos, *Phys. Rev. Lett.* **114**, 084102 (2015).
- [36] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.000.000000> for a discussion of the in-slice Floquet vectors.