

GAN Crash course

Eiríkur Agustsson

March 21, 2018

f -divergences

Given two distributions P and Q with density $p(x)$ and $q(x)$ over a domain \mathcal{X} , we define the f -divergence,

f -divergences

Given two distributions P and Q with density $p(x)$ and $q(x)$ over a domain \mathcal{X} , we define the f -divergence,

$$D_f(P\|Q) = \int_{\mathcal{X}} q(x) f\left(\frac{p(x)}{q(x)}\right) dx, \quad (1)$$

where the function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a convex (lower-semicontinuous) function, with $f(1) = 0$.

f -divergences

Given two distributions P and Q with density $p(x)$ and $q(x)$ over a domain \mathcal{X} , we define the f -divergence,

$$D_f(P\|Q) = \int_{\mathcal{X}} q(x) f\left(\frac{p(x)}{q(x)}\right) dx, \quad (1)$$

where the function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a convex (lower-semicontinuous) function, with $f(1) = 0$.

Example, set $f(u) = -\log(u) = \log(1/u)$, obtaining

$$D_f(P\|Q) = \int_{\mathcal{X}} q(x) \log\left(\frac{p(x)}{q(x)}\right) dx = D_{\text{KL}}(Q\|P), \quad (2)$$

which is the well known Kullback-Leibner divergence.

f -divergences

Other divergences that belong to the f divergence family:

Name	$D_f(P\ Q)$	Generator $f(u)$
Kullback-Leibler	$\int p(x) \log \frac{p(x)}{q(x)} dx$	$u \log u$
Reverse KL	$\int q(x) \log \frac{q(x)}{p(x)} dx$	$-\log u$
Pearson χ^2	$\int \frac{(q(x)-p(x))^2}{p(x)} dx$	$(u-1)^2$
Squared Hellinger	$\int \left(\sqrt{p(x)} - \sqrt{q(x)} \right)^2 dx$	$(\sqrt{u}-1)^2$
Jensen-Shannon	$\frac{1}{2} \int p(x) \log \frac{2p(x)}{p(x)+q(x)} + q(x) \log \frac{2q(x)}{p(x)+q(x)} dx$	$-(u+1) \log \frac{1+u}{2} + u \log u$

f -divergences

Other divergences that belong to the f divergence family:

Name	$D_f(P\ Q)$	Generator $f(u)$
Kullback-Leibler	$\int p(x) \log \frac{p(x)}{q(x)} dx$	$u \log u$
Reverse KL	$\int q(x) \log \frac{q(x)}{p(x)} dx$	$-\log u$
Pearson χ^2	$\int \frac{(q(x)-p(x))^2}{p(x)} dx$	$(u-1)^2$
Squared Hellinger	$\int \left(\sqrt{p(x)} - \sqrt{q(x)} \right)^2 dx$	$(\sqrt{u}-1)^2$
Jensen-Shannon	$\frac{1}{2} \int p(x) \log \frac{2p(x)}{p(x)+q(x)} + q(x) \log \frac{2q(x)}{p(x)+q(x)} dx$	$-(u+1) \log \frac{1+u}{2} + u \log u$

Expressive family, but note that the form is constrained an integral of a pointwise function between p and q .

f -divergences

Other divergences that belong to the f divergence family:

Name	$D_f(P\ Q)$	Generator $f(u)$
Kullback-Leibler	$\int p(x) \log \frac{p(x)}{q(x)} dx$	$u \log u$
Reverse KL	$\int q(x) \log \frac{q(x)}{p(x)} dx$	$-\log u$
Pearson χ^2	$\int \frac{(q(x)-p(x))^2}{p(x)} dx$	$(u-1)^2$
Squared Hellinger	$\int \left(\sqrt{p(x)} - \sqrt{q(x)} \right)^2 dx$	$(\sqrt{u}-1)^2$
Jensen-Shannon	$\frac{1}{2} \int p(x) \log \frac{2p(x)}{p(x)+q(x)} + q(x) \log \frac{2q(x)}{p(x)+q(x)} dx$	$-(u+1) \log \frac{1+u}{2} + u \log u$

Expressive family, but note that the form is constrained an integral of a pointwise function between p and q .

Cannot capture e.g. Wasserstein distance:

$$W(P, Q) = \inf \{ E_{(x,y) \sim R} [c(x, y)] \mid x \sim P, y \sim Q \}, \quad (3)$$

which measure minimal expected cost $c : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}^+$ over couplings R between P and Q .

Standard GAN

It turns out the standard GAN can be related to the JS-divergence.

Standard GAN

It turns out the standard GAN can be related to the JS-divergence. Recall, that we then have a discriminator D and a generator G playing the following two-player minimax game:

$$\min_G \max_D \mathbb{E}_{x \sim P_{\text{data}}} [\log D(x)] + \mathbb{E}_{x \sim G(z)} [\log(1 - D(x))] \quad (4)$$

$$= \min_G \max_D \int_x p_{\text{data}}(x) \log(D(x)) + p_{G(z)}(x) \log(1 - D(x)) dx \quad (5)$$

where z has a fixed prior distribution (e.g. uniform).

Standard GAN

It turns out the standard GAN can be related to the JS-divergence. Recall, that we then have a discriminator D and a generator G playing the following two-player minimax game:

$$\min_G \max_D \mathbb{E}_{x \sim p_{\text{data}}} [\log D(x)] + \mathbb{E}_{x \sim G(z)} [\log(1 - D(x))] \quad (4)$$

$$= \min_G \max_D \int_x p_{\text{data}}(x) \log(D(x)) + p_{G(z)}(x) \log(1 - D(x)) dx \quad (5)$$

where z has a fixed prior distribution (e.g. uniform).

Now, since $a \log(y) + b \log(1 - y)$ is maximized at $y = a/(a + b)$, we know that the optimal D is $D_G^*(x) = \frac{p_{\text{data}}(x)}{p_{\text{data}}(x) + p_{G(z)}(x)}$.

Standard GAN

Now suppose when maximizing D we find D_G^* . Then we can compute

$$\begin{aligned} \min_G \max_D \mathbb{E}_{x \sim p_{\text{data}}} [\log D(x)] + \mathbb{E}_{x \sim G(z)} [\log(1 - D(x))] \\ = \min_G \mathbb{E}_{x \sim p_{\text{data}}} \left[\log \left(\frac{p_{\text{data}}(x)}{p_{\text{data}}(x) + p_{G(z)}(x)} \right) \right] + \mathbb{E}_{x \sim G(z)} \left[\log \left(\frac{p_{G(z)}(x)}{p_{\text{data}}(x) + p_{G(z)}(x)} \right) \right] \\ = \min_G \int_x p_{\text{data}}(x) \log \left(\frac{p_{\text{data}}(x)}{\frac{1}{2}(p_{\text{data}}(x) + p_{G(z)}(x))} \right) \\ + p_{G(z)}(x) \log \left(\frac{p_{G(z)}(x)}{\frac{1}{2}(p_{\text{data}}(x) + p_{G(z)}(x))} \right) dx - \log(4) \\ = \min_G D_{\text{KL}}(p_{\text{data}} \| \tfrac{1}{2}(p_{\text{data}} + p_{G(z)})) + D_{\text{KL}}(p_{G(z)} \| \tfrac{1}{2}(p_{\text{data}} + p_{G(z)})) - \log(4) \\ = \min_G 2D_{\text{Jensen-Shannon}}(p_{\text{data}} \| p_{G(z)}) - \log(4) \end{aligned}$$

So, in this case, G is minimizing the Jensen-Shannon divergence between p_{data} and $p_{G(z)}$!

Standard GAN

- So what is missing?

Standard GAN

- So what is missing?
- First, in practice, we only consider a limited family of functions

$$\min_{\theta \in \Theta} \max_{\omega \in \Omega} \mathbb{E}_{x \sim P_{\text{data}}} [\log D_{\omega}(x)] + \mathbb{E}_{x \sim G_{\theta}(z)} [\log(1 - D_{\omega}(x))], \quad (6)$$

where $\theta \in \Theta$ and $\omega \in \Omega$ parameterize families of functions (e.g. neural networks).

Standard GAN

- So what is missing?
- First, in practice, we only consider a limited family of functions

$$\min_{\theta \in \Theta} \max_{\omega \in \Omega} \mathbb{E}_{x \sim p_{\text{data}}} [\log D_{\omega}(x)] + \mathbb{E}_{x \sim G_{\theta}(z)} [\log(1 - D_{\omega}(x))], \quad (6)$$

where $\theta \in \Theta$ and $\omega \in \Omega$ parameterize families of functions (e.g. neural networks).

- Since $D_G^*(x) = \frac{p_{\text{data}}(x)}{p_{\text{data}}(x) + p_{G(z)}(x)}$ depends on p_{data} , it is unlikely it is an element of the function family $\{D_{\omega} | \omega \in \Omega\}$.

Standard GAN

- So what is missing?
- First, in practice, we only consider a limited family of functions

$$\min_{\theta \in \Theta} \max_{\omega \in \Omega} \mathbb{E}_{x \sim p_{\text{data}}} [\log D_{\omega}(x)] + \mathbb{E}_{x \sim G_{\theta}(z)} [\log(1 - D_{\omega}(x))], \quad (6)$$

where $\theta \in \Theta$ and $\omega \in \Omega$ parameterize families of functions (e.g. neural networks).

- Since $D_G^*(x) = \frac{p_{\text{data}}(x)}{p_{\text{data}}(x) + p_{G(z)}(x)}$ depends on p_{data} , it is unlikely it is an element of the function family $\{D_{\omega} | \omega \in \Omega\}$.
- Second, instead of finding the global mini-maximum, we only find a saddle point via double-loop SGD.
- Thus, we cannot simply plug D_G^* into the GAN objective!

Othe divergences

Can obtain any f -divergence (not just JS) by changing the loss

Othe divergences

Can obtain any f -divergence (not just JS) by changing the loss

$$\min_G \max_D \mathbb{E}_{x \sim P_{\text{data}}} [\log D(x)] + \mathbb{E}_{x \sim G(z)} [\log(1 - D(x))] \quad (7)$$

(8)

Othe divergences

Can obtain any f -divergence (not just JS) by changing the loss

$$\min_G \max_D \mathbb{E}_{x \sim P_{\text{data}}} [\log D(x)] + \mathbb{E}_{x \sim G(z)} [\log(1 - D(x))] \quad (7)$$

(8)

Becomes

$$\min_G \max_D \mathbb{E}_{x \sim P_{\text{data}}} [f(D(x))] + \mathbb{E}_{x \sim G(z)} [g(D(x))] \quad (9)$$

(10)

In particular, a notable alternative is the Least Squares GAN

Thank you for your attention!

