



Bastu - ALT2's Dental Health Report

APR 13TH
2020



Table Of Contents

01

Introduction and Key
Concepts

02

Dental Health Summary

03

General Health Implications

04

Periodontal Disease Risk -
Detailed Assessment

05

Tooth Resorption Risk -
Detailed Assessment

06

Bad Breath Risk - Detailed
Assessment

07

What's Next?

09

Appendix

10

Bibliography


Did you know?



Dental disease affects 50-90% of cats over the age of four.

Fortunately, most dental diseases are preventable with a good dental care routine. They are also mostly treatable, if caught early. The microbes in a cat's mouth can inform us of developing dental issues, before they are advanced enough to be visible by a veterinarian during a routine checkup. The Basepaws Dental Health test looks for microbial signatures associated with three of the most common dental conditions in cats - Periodontal Disease, Tooth Resorption, Halitosis (bad breath).

The oral microbiome



Environmental factors and various food sources make the feline oral cavity a fascinating place, characterized by unique interactions between a cat's mouth and the microbes within it (the oral microbiome). The almost constant exposure to foreign microbial organisms has made the oral microbiome fiercely competitive. Once in a while, pathogenic microbes manage to colonize parts of the oral cavity which can be associated with dental problems.

The feline oral microbiome can reveal information about developing dental issues.


The oral microbiome also has implications for general health.

Can the oral microbiome change?

YES! The oral microbiome is not static.

Different factors such as diet (dry versus wet food), environment (indoor versus outdoor), supplement intake, medications (particularly antibiotics) and dental care routine can all influence the composition of the oral microbiome.

This is why testing early and testing often is key for optimal dental health!



Bastu - ALT2's Dental health summary

How does this test work? We used our oral microbiome database containing healthy cats and cats suffering from periodontal disease, tooth resorption or halitosis to identify a set of predictive microbes whose compositional abundance is associated with each condition. Based on these results, we developed a 0 - 10 risk score system for each condition. The results below show Bastu - ALT2's overall risk for each of the three conditions, as well as a breakdown of the predictive microbes whose compositional abundance is associated with high, medium or low risk for each dental condition. **The purple line and the number next to it indicate your cat's overall risk score for each condition.**


low risk: 0 - 3.3 medium risk: >3.3 - 6.6 high risk: >6.6 - 10

Risk for periodontal disease

Periodontal disease affects the tissues surrounding the teeth. Initial stages are classified as gingivitis, while advanced cases are known as periodontitis.

OVERALL RISK:

- ☒ LOW
☐ MEDIUM
☐ HIGH




Risk for tooth resorption

Tooth resorption is a relatively common condition characterized by progressive dentin erosion.

OVERALL RISK:

- ☒ LOW
☐ MEDIUM
☐ HIGH




Risk for bad breath (halitosis)

When bad breath is a persistent problem for a cat, this could be indicative of more serious general health issues.

OVERALL RISK:

- ☒ LOW
☐ MEDIUM
☐ HIGH



What's next?


- Adopt a daily dental care routine to maintain Bastu - ALT2's dental health
- Check for emerging problems by doing regular 'flip the lip' exams
- Keep up with Bastu - ALT2's routine checkups with the veterinarian

**Next
recommended
dental health
test in:
6-8 months**

What are Bastu - ALT2's health implications?

Great news! Currently, your cat's oral health is unlikely to be having a negative effect on their general health.

Cats with good oral health are less prone to developing chronic kidney disease, diabetes mellitus, cardiovascular problems and some autoimmune diseases.




What can you do?

- It is important to regularly assess a cat's dental health in order to address any emerging issues early on and minimize chances of them negatively impacting general health.
- Adopting a thorough and consistent dental care routine at home can significantly reduce the chance of developing dental diseases. This will in turn reduce your cat's likelihood of developing more serious general health problems.

Periodontal Disease

Periodontal disease (PD) is a group of inflammatory disorders affecting the tissues surrounding the teeth. Periodontal disease is initiated by the build-up of plaque on the tooth surface resulting in the gingiva becoming inflamed (gingivitis). Without an effective oral care regime, inflammation can begin to destroy the structures that support the tooth (periodontitis). Periodontal disease affects up to 80% of the adult feline population. Below you will see how your cat's oral microbiome compares to a healthy population when it comes to microbial signatures of periodontal disease.



We analyzed Bastu - ALT2's oral microbiome to establish the compositional abundance of 108 microbes predictive of periodontal disease. We ranked each microbe's abundance on a scale from 1 to 5, where 1 represents abundance levels close to a healthy control population and 5 represents abundance levels close to cats with periodontal disease. Below are Bastu - ALT2's TOP 3 most significant microbes associated with high, medium, and low risk, respectively.

Currently, Bastu - ALT2's compositional abundance levels for 24 out of 108 microbes are consistent with having periodontal disease (22%).

Top 3 high risk microbes

Moraxella ovis
Corynebacterium xerosis
Actinomyces sp. Chiba101




Top 3 medium risk microbes

Ottowia sp. oral taxon 894
Roseburia hominis
Capnocytophaga canimorsus




Top 3 low risk microbes

Rodentibacter pneumotropicus
Avibacterium paragallinarum
Lautropia mirabilis



Tooth Resorption

Every tooth is composed of a root canal (containing nerves, blood and lymphatic vessels) and bony substances called dentin and enamel. When a cat suffers from tooth resorption, the dentin of the affected tooth starts to progressively erode. Unfortunately, tooth resorption is relatively common, affecting 20-60% of all cats and over 70% of cats over the age of five. Below, you can see how your cat compares to the healthy feline population with regards to abundance of microbes associated with tooth resorption.



We analyzed Bastu - ALT2's oral microbiome to establish the compositional abundance of 74 microbes predictive of tooth resorption. We ranked each microbe's abundance on a scale from 1 to 5, where 1 represents abundance levels close to a healthy control population and 5 represents abundance levels close to cats with tooth resorption. Below are Bastu - ALT2's TOP 3 most significant microbes associated with high, medium, and low risk, respectively.

Currently, Bastu - ALT2's compositional abundance levels for 19 out of 74 microbes are consistent with having tooth resorption (26%).

Top 3 high risk microbes

Moraxella ovis	=====
Corynebacterium xerosis	=====
Wolinella succinogenes	=====

Top 3 medium risk microbes


Parabacteroides distasonis	=====
Capnocytophaga stomatis	=====
Escherichia coli	=====

Top 3 low risk microbes

Avibacterium paragallinarum	=====
Acinetobacter junii	=====
Rodentibacter pneumotropicus	=====

Bad Breath

Occasional bad breath is usually not something you should worry about. When bad breath is a persistent problem, this could be indicative of more serious issues. The most common cause of bad breath is periodontal disease. Different types of bad breath can also indicate general health problems, such as kidney disease, diabetes and some liver disorders. Here is how your cat's oral microbiome compares to the general healthy population when it comes to halitosis.



We analyzed Bastu - ALT2's oral microbiome to establish the compositional abundance of 182 microbes predictive of bad breath. We ranked each microbe's abundance on a scale from 1 to 5, where 1 represents abundance levels close to a healthy control population and 5 represents abundance levels close to cats with bad breath. Below are Bastu - ALT2's TOP 3 most significant microbes associated with high, medium, and low risk, respectively.

Currently, Bastu - ALT2's compositional abundance levels for 12 out of 182 microbes are consistent with having bad breath (7%).

Top 3 high risk microbes

Mycoplasma arginini	=====
Gemella sp. oral taxon 928	=====
Streptococcus anginosus	=====

Top 3 medium risk microbes

Mannheimia pernigra	=====
Frederiksenia canicola	=====
A. sp.,oral,taxon,190	=====

Top 3 low risk microbes

Alysiella filiformis	=====
Avibacterium paragallinarum	=====
Neisseria dentiae	=====


What's next for Bastu - ALT2?

At home care

Congratulations! Your cat seems to be at low risk for periodontal disease, tooth resorption and bad breath. If you have an established dental care routine, continue doing what you are doing. If you don't, consider adopting one, as it is likely that, as your cat ages, their dental health will deteriorate. If done properly, the most effective at home treatment is daily tooth brushing.

We teamed up with some of the world's top veterinary dentistry professionals to provide you with support and innovative solutions on how to approach brushing your cat's teeth and other tips and tricks for optimal dental hygiene.

Watch the video to learn how to tailor your routine to your cat's personality and comfort level.




[Learn More](#)

You can also read about some effective, off-the-beaten-path ways to maximize the effect of your cat's dental care routine in this article.

In addition, you can also learn how to perform routine 'flip the lip' exams at home and recognize signs of dental pain in your cat (dropping food, loss of appetite, favoring one side over the other when chewing, drooling).


Finally, you can consider incorporating some of the products recommended by the [Veterinary Oral Health Council \(VOHC\)](#) to support your preventative dental care routine. These products have demonstrated efficacy in fighting plaque and tartar buildup, which are the root cause of many dental problems.

Veterinary oral exam



Have a question for Basepaws?

If you have a question about any part of this report, you can send us an email at: help@basepaws.com



While your cat's mouth looks good to us, it is much easier to resolve dental issues that are spotted early

Therefore, a proactive approach to feline dentistry is recommended - keep up with your cat's yearly vet visits.

Want to discuss your results with other cat parents? Join our facebook group!



Would you like to tell us more about your cat's dental and general health? Contribute to feline dental health research by filling out this [survey](#).

Next dental health test recommended in:
6 - 8 months or sooner if any of the following change:
diet, medications, supplements, general health

Appendix

Sequencing and analysis methodology

Most direct-to-consumer microbiome tests use a technique called '16S rRNA gene sequencing'. This technique can only provide information about the bacteria present in the microbiome.

However, it is well-known that the microbiome is composed of viruses, protozoa, fungi and archaea species, in addition to bacteria. This means that the 16S approach zooms in on just one part of the microbiome, ignoring the rest. Additionally, 16S sequencing does not provide sufficient resolution to reliably and consistently go beyond the genus level of bacterial classification. Therefore, in most cases, we don't know the exact species of bacteria in the microbiome, making analysis somewhat vague and relying on approximation.

To address these problems, Basepaws uses metagenomic sequencing instead of 16S sequencing. Our method allows us to capture organisms across all domains of life, not restricting us to just bacteria. In addition, we can reliably identify organisms to the species or even strain level, making our analysis more accurate and improving our confidence in the results. These results paint a much richer and unbiased picture of your feline companion's mouth. We used pairwise log ratio transformation to estimate the compositional abundance of microbial species and Gaussian mixture modeling to determine your cat's risk for periodontal disease, tooth resorption and bad breath.

Limitations

The Basepaws oral microbiome report is based on our ability to identify thousands of microbial species with each test.

Our large oral microbiome reference database allows us to identify a multitude of novel associations between microbes found in the mouth and a variety of diseases, as well as confirm previously reported findings. However, the field of feline oral microbiome science is extremely young and understudied, which is why we report only on conditions and microbes where previous knowledge exists and/or we see a particularly strong signal coming through in our data.

As we accumulate more data and conduct more analyses, we will aim to continuously enrich this report, providing even more helpful insights. We want to emphasize that the identification of a certain microbial signature associated with a dental disease does not constitute a diagnosis. Conversely, not detecting a particular microbial signature does not exclude the possibility of an unknown disease-causing pathogen being present or dental disease being caused by something other than pathogenic microbes. This report does not aim to substitute a diagnosis by a professional.

Bibliography

- Amado, P.P.P., Kawamoto, D., Albuquerque-Souza, E., Franco, D.C., Saraiva, L., Casarin, R.C.V., Horliana, A.C.R.T. and Mayer, M.P.A., 2020. Oral and fecal microbiome in molar-incisor pattern periodontitis. *Frontiers in cellular and infection microbiology*, 10. [link](#)
- Anesti, V., McDonald, I.R., Ramaswamy, M., Wade, W.G., Kelly, D.P. and Wood, A.P., 2005. Isolation and molecular detection of methylotrophic bacteria occurring in the human mouth. *Environmental Microbiology*, 7(8), pp.1227-1238. [link](#)
- Arora, N., Mishra, A. and Chugh, S., 2014. Microbial role in periodontitis: Have we reached the top? Some unsung bacteria other than red complex. *Journal of Indian Society of Periodontology*, 18(1), p.9. [link](#)
- Awano, S., Gohara, K., Kurihara, E., Ansai, T. and Takehara, T., 2002. The relationship between the presence of periodontopathogenic bacteria in saliva and halitosis. *International dental journal*, 52(S5P1), pp.212-216. [link](#)
- Binti Badlishah Sham, N.I., Lewin, S.D. and Grant, M.M., 2020. Proteomic Investigations of In Vitro and In Vivo Models of Periodontal Disease. *PROTEOMICS–Clinical Applications*, 14(3), p.1900043. [link](#)
- Blum, S., Elad, D., Zukin, N., Lysnyansky, I., Weisblith, L., Perl, S., Netanel, O. and David, D., 2010. Outbreak of *Streptococcus equi* subsp. *zooeconomicus* infections in cats. *Veterinary microbiology*, 144(1-2), pp.236-239. [link](#)
- Booij-Vrieling, H.E., van der Reijden, W.A., Houwers, D.J., De Wit, W.E.A.J., Bosch-Tijhof, C.J., Penning, L.C., Van Winkelhoff, A.J. and Hazewinkel, H.A.W., 2010. Comparison of periodontal pathogens between cats and their owners. *Veterinary microbiology*, 144(1-2), pp.147-152. [link](#)
- Boyanova, L., Setchanova, L., Gergova, G., Kostyanov, T., Yordanov, D., Popova, C., Kotsilkov, K. and Mitov, I., 2009. Microbiological diagnosis of the severe chronic periodontitis. *Journal of IMAB*, pp.89-94. [link](#)
- Byrne-Bailey, K.G., Weber, K.A., Bose, S., Knox, T., Spanbauer, T.L., Chertkov, O. and Coates, J.D., 2010. Completed genome sequence of the anaerobic iron-oxidizing bacterium *Acidovorax ebreus* strain TSPY. *Journal of bacteriology*, 192(5), pp.1475-1476. [link](#)
- Chan, Y., Huo, Y.B., Yu, X., Zeng, H., Leung, W.K. and Watt, R.M., 2020. Complete Genome Sequence of Human Oral Phylogroup 1 *Treponema* sp. Strain OMZ 804 (ATCC 700766), Originally Isolated from Periodontitis Dental Plaque. *Microbiology Resource Announcements*, 9(22). [link](#)
- Chan, Y., Ma, A.P., Lacap-Bugler, D.C., Huo, Y.B., Leung, W.K., Leung, F.C. and Watt, R.M., 2014. Complete genome sequence for *Treponema* sp. OMZ 838 (ATCC 700772, DSM 16789), isolated from a necrotizing ulcerative gingivitis lesion. *Genome announcements*, 2(6). [link](#)
- Chávez de Paz, L., 2004. Gram-positive organisms in endodontic infections. *Endodontic Topics*, 9(1), pp.79-96. [link](#)
- Cho, E., Park, S.N., Lim, Y.K., Shin, Y., Paek, J., Hwang, C.H., Chang, Y.H. and Kook, J.K., 2015. *Fusobacterium hwasookii* sp. nov., isolated from a human periodontitis lesion. *Current microbiology*, 70(2), pp.169-175. [link](#)
- Colombo, A.P.V., Boches, S.K., Cotton, S.L., Goodson, J.M., Kent, R., Haffajee, A.D., Socransky, S.S., Hasturk, H., Van Dyke, T.E., Dewhirst, F. and Paster, B.J., 2009. Comparisons of subgingival microbial profiles of refractory periodontitis, severe periodontitis, and periodontal health using the human oral microbe identification microarray. *Journal of periodontology*, 80(9), pp.1421-1432. [link](#)
- Colombo, A.P.V., Magalhães, C.B., Hartenbach, F.A.R.R., do Souto, R.M. and da Silva-Boghossian, C.M., 2016. Periodontal-disease-associated biofilm: A reservoir for pathogens of medical importance. *Microbial pathogenesis*, 94, pp.27-34. [link](#)
- Cortelli, J.R., Barbosa, M.D.S. and Westphal, M.A., 2008. Halitosis: a review of associated factors and therapeutic approach. *Brazilian oral research*, 22, pp.44-54. [link](#)
- Davis, I.J., Bull, C., Horsfall, A., Morley, I. and Harris, S., 2014. The Unculturable: targeted isolation of bacterial species associated with canine periodontal health or disease from dental plaque. *BMC microbiology*, 14(1), p.196. [link](#)

Bibliography

Davis, I.J., Bull, C., Horsfall, A., Morley, I. and Harris, S., 2014. The Unculturables: targeted isolation of bacterial species associated with canine periodontal health or disease from dental plaque. BMC microbiology, 14(1), p.196. [link](#)

Dilegge, S.K., Edgcomb, V.P. and Leadbetter, E.R., 2011. Presence of the oral bacterium *Capnocytophaga canimorsus* in the tooth plaque of canines. Veterinary microbiology, 149(3-4), pp.437-445. [link](#)

Donati, C., Zolfo, M., Albanese, D., Truong, D.T., Asnicar, F., Iebba, V., Cavalieri, D., Jousson, O., De Filippo, C., Huttenhower, C. and Segata, N., 2016. Uncovering oral *Neisseria* tropism and persistence using metagenomic sequencing. Nature microbiology, 1(7), p.16070. [link](#)

Ebrahim-Saraie, H.S., Motamedifar, M., Mansury, D., Ebrahimi, H., Pourshahidi, S., Halaji, M. and Shahraki, H.R., 2015. Association of *Fusobacterium* Isolation From Periodontal Pockets With Halitosis and the Related Risk Factors in Shiraz, Iran. Archives of Clinical Infectious Diseases, 10(4). [link](#)

Forsblom, B., Sarkiala-Kessel, E., Kanervo, A., Vaeisaenen, M.L., Helander, I.M. and Jousimies-Somer, H., 2002. Characterisation of aerobic gram-negative bacteria from subgingival sites of dogs—potential bite wound pathogens. Journal of medical microbiology, 51(3), pp.207-220. [link](#)

Galperin, M.Y., 2007. Dark matter in a deep-sea vent and in human mouth. Environmental microbiology, 9(10), p.2385. [link](#)

Gerner-Smidt, P., Keiser-Nielsen, H., Dorsch, M., Stackebrandt, E., Ursing, J., Blom, J., Christensen, A.C., Christensen, J.J., Frederiksen, W., Hoffmann, S. and Holten-Andersen, W., 1994. *Lautropia mirabilis* gen. nov., sp. nov., a Gram-negative motile coccus with unusual morphology isolated from the human mouth. Microbiology, 140(7), pp.1787-1797. [link](#)

Hampelska, K., Jaworska, M.M., Babalska, Z.Ł. and Karpiński, T.M., 2020. The role of oral microbiota in intra-oral halitosis. Journal of Clinical Medicine, 9(8), p.2484. [link](#)

Han, X., Lin, X., Yu, X., Lin, J., Kawai, T., LaRosa, K.B. and Taubman, M.A., 2013. *Porphyromonas gingivalis* infection-associated periodontal bone resorption is dependent on receptor activator of NF-κB ligand. Infection and immunity, 81(5), pp.1502-1509. [link](#)

Han, Y.W., Shi, W., Huang, G.T.J., Haake, S.K., Park, N.H., Kuramitsu, H. and Genco, R.J., 2000. Interactions between periodontal bacteria and human oral epithelial cells: *Fusobacterium nucleatum* adheres to and invades epithelial cells. Infection and immunity, 68(6), pp.3140-3146. [link](#)

Harris, S., Croft, J., O'Flynn, C., Deusch, O., Colyer, A., Allsopp, J., Milella, L. and Davis, I.J., 2015. A pyrosequencing investigation of differences in the feline subgingival microbiota in health, gingivitis and mild periodontitis. PLoS One, 10(11), p.e0136986. [link](#)

Holcombe, L.J., Patel, N., Colyer, A., Deusch, O., O'Flynn, C. and Harris, S., 2014. Early canine plaque biofilms: characterization of key bacterial interactions involved in initial colonization of enamel. PLoS One, 9(12), p.e113744. [link](#)

Iwamoto, T., Suzuki, N., Tanabe, K., Takeshita, T. and Hirofuji, T., 2010. Effects of probiotic *Lactobacillus salivarius* WB21 on halitosis and oral health: an open-label pilot trial. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology, 110(2), pp.201-208. [link](#)

Johnson, L., Almeida-da-Silva, C.L.C., Takiya, C.M., Figliuolo, V., Rocha, G.M., Weissmüller, G., Scharfstein, J., Coutinho-Silva, R. and Ojcius, D.M., 2018. Oral infection of mice with *Fusobacterium nucleatum* results in macrophage recruitment to the dental pulp and bone resorption. biomedical journal, 41(3), pp.184-193. [link](#)

Kačirová, J., Maďar, M., Štrkolcová, G., Maďari, A. and Nemcová, R., 2019. Dental Biofilm as Etiological Agent of Canine Periodontal Disease. In Bacterial Biofilms. IntechOpen. [link](#)

Khazandi, M., Bird, P.S., Owens, J., Wilson, G., Meyer, J.N. and Trott, D.J., 2014. In vitro efficacy of cefovecin against anaerobic bacteria isolated from subgingival plaque of dogs and cats with periodontal disease. Anaerobe, 28, pp.104-108. [link](#)

Bibliography

- Khazandi, M., Bird, P.S., Owens, J., Wilson, G., Meyer, J.N. and Trott, D.J., 2014. In vitro efficacy of cefovecin against anaerobic bacteria isolated from subgingival plaque of dogs and cats with periodontal disease. *Anaerobe*, 28, pp.104-108. [link](#)
- Kushkevych, I., Coufalová, M., Vítězová, M. and Rittmann, S.K.M., 2020. Sulfate-Reducing Bacteria of the Oral Cavity and Their Relation with Periodontitis—Recent Advances. *Journal of Clinical Medicine*, 9(8), p.2347. [link](#)
- Langendijk, P.S., Kulik, E.M., Sandmeier, H., Meyer, J. and Van Der Hoeven, J.S., 2001. Isolation of *Desulfomicrobium orale* sp. nov. and *Desulfovibrio* strain NY682, oral sulfate-reducing bacteria involved in human periodontal disease. *International Journal of Systematic and Evolutionary Microbiology*, 51(3), pp.1035-1044. [link](#)
- Liu, G., Tang, C.M. and Exley, R.M., 2015. Non-pathogenic *Neisseria*: members of an abundant, multi-habitat, diverse genus. *Microbiology*, 161(7), pp.1297-1312. [link](#)
- Loubinoux, J., Bisson-Boutelliez, C., Miller, N. and Le Faou, A.E., 2002. Isolation of the provisionally named *Desulfovibrio fairfieldensis* from human periodontal pockets. *Oral microbiology and immunology*, 17(5), pp.321-323. [link](#)
- Love, D.N., Johnson, J.L., Jones, R.F. and Bailey, M.A.R.I.L.Y.N., 1985. Comparison of *Bacteroides* zoogloeiformans strains isolated from soft tissue infections in cats with strains from periodontal disease in humans. *Infection and immunity*, 47(1), pp.166-168. [link](#)
- Love, D.N., Johnson, J.L. and Moore, L.V.H., 1989. *Bacteroides* species from the oral cavity and oral-associated diseases of cats. *Veterinary microbiology*, 19(3), pp.275-281. [link](#)
- Mallonee, D.H., Harvey, C.E., Venner, M. and Hammond, B.F., 1988. Bacteriology of periodontal disease in the cat. *Archives of oral biology*, 33(9), pp.677-683. [link](#)
- Mineoka, T., Awano, S., Rikimaru, T., Kurata, H., Yoshida, A., Ansai, T. and Takehara, T., 2008. Site-specific development of periodontal disease is associated with increased levels of *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia* in subgingival plaque. *Journal of periodontology*, 79(4), pp.670-676. [link](#)
- Nises, J., Rosander, A., Pettersson, A. and Backhans, A., 2018. The occurrence of *Treponema* spp. in gingival plaque from dogs with varying degree of periodontal disease. *PloS one*, 13(8), p.e0201888. [link](#)
- Nordhoff, M., Rühle, B., Kellermeier, C., Moter, A., Schmitz, R., Brunnberg, L. and Wieler, L.H., 2008. Association of *Treponema* spp. with canine periodontitis. *Veterinary microbiology*, 127(3-4), pp.334-342. [link](#)
- Norris, J.M. and Love, D.N., 1999. Associations amongst three feline *Porphyromonas* species from the gingival margin of cats during periodontal health and disease. *Veterinary microbiology*, 65(3), pp.195-207. [link](#)
- Özavci, V., Erbas, G., Parin, U., Yüksel, H.T. and Kirkan, Ş., 2019. Molecular detection of feline and canine periodontal pathogens. *Veterinary and Animal Science*, 8, p.100069. [link](#)
- Park, S.N., Lim, Y.K., Shin, J.H., Kim, H.S., Jo, E., Lee, W.P., Shin, Y., Paek, J., Chang, Y.H., Kim, H. and Kook, J.K., 2019. *Fusobacterium pseudoperiodonticum* sp. nov., Isolated from the Human Oral Cavity. *Current microbiology*, 76(6), pp.659-665. [link](#)
- Prates, M., Fernandes, F., Proença, F., Mussá, Y., Tavares, A. and Pereira, A., 2020. Oral Infection Caused by *Stenotrophomonas maltophilia*: A Rare Presentation of an Emerging Opportunistic Pathogen. *Case Reports in Infectious Diseases*, 2020. [link](#)
- Richards, A.M., Abu Kwaik, Y. and Lamont, R.J., 2015. Code blue: A *Cinetobacter baumannii*, a nosocomial pathogen with a role in the oral cavity. *Molecular oral microbiology*, 30(1), pp.2-15. [link](#)
- Riggio, M.P., Lennon, A., Taylor, D.J. and Bennett, D., 2011. Molecular identification of bacteria associated with canine periodontal disease. *Veterinary microbiology*, 150(3-4), pp.394-400. [link](#)

Bibliography

Riggio, M.P., Lennon, A., Taylor, D.J. and Bennett, D., 2011. Molecular identification of bacteria associated with canine periodontal disease. *Veterinary microbiology*, 150(3-4), pp.394-400. [link](#)

Rodrigues, M.X., Bicalho, R.C., Fiani, N., Lima, S.F. and Peralta, S., 2019. The subgingival microbial community of feline periodontitis and gingivostomatitis: characterization and comparison between diseased and healthy cats. *Scientific reports*, 9(1), pp.1-10. [link](#)

Ruparell, A., Inui, T., Staunton, R., Wallis, C., Deusch, O. and Holcombe, L.J., 2020. The canine oral microbiome: variation in bacterial populations across different niches. *BMC microbiology*, 20(1), pp.1-13. [link](#)

Sanguansermsri, P., Nobbs, A.H., Jenkinson, H.F. and Surarit, R., 2018. Interspecies dynamics among bacteria associated with canine periodontal disease. *Molecular oral microbiology*, 33(1), pp.59-67. [link](#)

Sela, M.N., 2001. Role of *Treponema denticola* in periodontal diseases. *Critical Reviews in Oral Biology & Medicine*, 12(5), pp.399-413. [link](#)

Sundqvist, G., 1990. Endodontic microbiology. *Experimental endodontics*, 6, pp.131-153.

Suzuki, M., Kimura, M., Imaoka, K. and Yamada, A., 2010. Prevalence of *Capnocytophaga canimorsus* and *Capnocytophaga cynodegmi* in dogs and cats determined by using a newly established species-specific PCR. *Veterinary microbiology*, 144(1-2), pp.172-176. [link](#)

Svartström, O., Mushtaq, M., Pringle, M. and Segerman, B., 2013. Genome-wide relatedness of *Treponema pedis*, from gingiva and necrotic skin lesions of pigs, with the human oral pathogen *Treponema denticola*. *PloS one*, 8(8), p.e71281. [link](#)

Sykora, S., Pieber, K., Simhofer, H., Hackl, V., Brodesser, D. and Brandt, S., 2014. Isolation of *Treponema* and *Tannerella* spp. from equine odontoclastic tooth resorption and hypercementosis related periodontal disease. *Equine veterinary journal*, 46(3), pp.358-363. [link](#)

Umeda, K., Hatakeyama, R., Abe, T., Takakura, K.I., Wada, T., Ogasawara, J., Sanada, S.I. and Hase, A., 2014. Distribution of *Capnocytophaga canimorsus* in dogs and cats with genetic characterization of isolates. *Veterinary microbiology*, 171(1-2), pp.153-159. [link](#)

Van Dyke, T.E., Offenbacher, S., Place, D., Dowell, V.R. and Jones, J., 1988. Refractory Periodontitis: Mixed Infection with *Bacteroides gingivalis* and Other Unusual *Bacteroides* Species: A Case Report. *Journal of periodontology*, 59(3), pp.184-189. [link](#)

Wallis, C.V., Marshall-Jones, Z.V., Deusch, O. and Hughes, K.R., 2017. 17 Canine and Feline Microbiomes. Understanding Host-Microbiome Interactions-An Omics Approach: Omics of Host-Microbiome Association, p.279. [link](#)

Wyss, C., Moter, A., Choi, B.K., Dewhirst, F.E., Xue, Y., Schüpbach, P., Göbel, U.B., Paster, B.J. and Guggenheim, B., 2004. *Treponema putidum* sp. nov., a medium-sized proteolytic spirochaete isolated from lesions of human periodontitis and acute necrotizing ulcerative gingivitis. *International journal of systematic and evolutionary microbiology*, 54(4), pp.1117-1122. [link](#)

Yasukawa, T., Ohmori, M. and Sato, S., 2010. The relationship between physiologic halitosis and periodontopathic bacteria of the tongue and gingival sulcus. *Odontology*, 98(1), pp.44-51. [link](#)

Ye, W., Zhang, Y., He, M., Zhu, C. and Feng, X.P., 2019. Relationship of tongue coating microbiome on volatile sulfur compounds in healthy and halitosis adults. *Journal of breath research*, 14(1), p.016005. [link](#)

Yoshino, Y., Kitazawa, T., Kamimura, M., Tatsuno, K., Ota, Y. and Yotsuyanagi, H., 2011. *Pseudomonas putida* bacteremia in adult patients: five case reports and a review of the literature. *Journal of Infection and Chemotherapy*, 17(2), pp.278-282. [link](#)

Zambori, C., Tirziu, E., Nichita, I., Cumpanasoiu, C., Gros, R.V., Seres, M., Mladin, B. and Mot, D., 2012. Biofilm implication in oral diseases of dogs and cats. *Scientific Papers Animal Science and Biotechnologies*, 45(2), pp.208-212.

Bibliography

Zambori, C., Tirziu, E., Nichita, I., Cumpanasoiu, C., Gros, R.V., Seres, M., Mladin, B. and Mot, D., 2012. Biofilm implication in oral diseases of dogs and cats. Scientific Papers Animal Science and Biotechnologies, 45(2), pp.208-212.

Zangenah, S., Abbasi, N., Andersson, A.F. and Bergman, P., 2016. Whole genome sequencing identifies a novel species of the genus Capnocytophaga isolated from dog and cat bite wounds in humans. Scientific reports, 6, p.22919. [link](#)

Zhang, Y., Zhen, M., Zhan, Y., Song, Y., Zhang, Q. and Wang, J., 2017. Population-genomic insights into variation in Prevotella intermedia and Prevotella nigrescens isolates and its association with periodontal disease. Frontiers in cellular and infection microbiology, 7, p.409. [link](#)