

## ENERGY LEVELS OF LIGHT NUCLEI $A = 16-17^{\dagger}$

F. AJZENBERG-SELOVE

*University of Pennsylvania, Philadelphia, Pennsylvania 19104*

Received 24 August 1981

**Abstract:** Compilation of energy levels of  $A = 16$  and  $17$  nuclei, with emphasis on the review of material leading to information about the structure of the  $A = 16$  and  $17$  systems.

### Introduction

The first section <sup>1)</sup> in the present generation of review papers summarizing the information on the light nuclei covered  $A = 5-10$ . The reader is referred to that article for an exposition of the style, format and conventions used here. We list below some other recent compilations of similar character:

$A = 3$  and  $4$ : Fiarman, Hanna and Meyerhof <sup>2,3)</sup>,

$A = 11-12, 13-15, 18-20$ : Ajzenberg-Selove <sup>4-6)</sup>,

$A = 21-44$ : Endt and Van der Leun <sup>7)</sup>.

Information about mass chains with  $A > 44$  can be found in *Nuclear Data Sheets*.

### Acknowledgments

The author is grateful to her many colleagues who have pointed out errors in the preliminary version of this paper and who have supplied corrections and new information. She is particularly indebted to Drs. F. C. Barker, F. D. Becchetti, P. M. Endt, F. Hintenberger and H. T. Richards for their help.

The author wishes to acknowledge the intelligence and skill of Glenda C. Marshall who prepared the manuscript. She is very grateful also to Mary Jane Potter who prepared the figures, to Michael Kearney, Roy Marshall and William Schriver of the Physics Computer and to Marion Kreiter and Revathy Mittal of the Physics Library. We thank the US Department of Energy and Drs. W. Hass, P. D. Greaves and W. H. Wimmers and the staff of *Nuclear Physics* and of the North-Holland Publishing Company for their support.

<sup>†</sup> This work has been supported by the US Department of Energy [EY-76-S-02-2785.\*000].

### References to Introduction

- 1) F. Ajzenberg-Selove, Nucl. Phys. **A320** (1979) 1
- 2) S. Fiarman and S. S. Hanna, Nucl. Phys. **A251** (1975) 1
- 3) S. Fiarman and W. E. Meyerhof, Nucl. Phys. **A206** (1973) 1
- 4) F. Ajzenberg-Selove and C. L. Busch, Nucl. Phys. **A336** (1980) 1
- 5) F. Ajzenberg-Selove, Nucl. Phys. **A360** (1981) 1
- 6) F. Ajzenberg-Selove, Nucl. Phys. **A300** (1978) 1
- 7) P. M. Endt and C. Van der Leun, Nucl. Phys. **A310** (1978) 1

**TABLE 1**  
Parameters of the ground states of the light nuclei

	Atomic mass excess (keV) <sup>a)</sup>	$\tau_{1/2}^{\text{b})}$ or $\Gamma_{\text{c.m.}}$	Decay	$J^{\pi \text{b,c)}}; T$
<sup>16</sup> He	see text		n	(0 <sup>+</sup> ); 6
<sup>16</sup> Be	see text		n	(0 <sup>+</sup> ); 4
<sup>16</sup> B	see text		n	$T = 3$
<sup>16</sup> C	$13\ 694 \pm 4^{\text{b})}$	$0.747 \pm 0.008$ sec	$\beta^-$	$0^+; 2$
<sup>16</sup> N	$5681.6 \pm 2.3^{\text{d})}$	$7.13 \pm 0.02$ sec	$\beta^-$	$2^-; 1$
<sup>16</sup> O	$-4737.02 \pm 0.04^{\text{d})}$	stable		$0^+; 0$
<sup>16</sup> F	$10\ 692 \pm 14^{\text{d})}$	$40 \pm 20$ keV	$\beta^+$	(0 <sup>-</sup> ); 1
<sup>16</sup> Ne	$24\ 020 \pm 40^{\text{b})}$	$50^{+50}_{-45}$ keV	p	$0^+; 2$
<sup>17</sup> Be	see text		n	$T = \frac{9}{2}$
<sup>17</sup> B	see text		$(\beta^-)$	$(\frac{1}{2}^-); \frac{7}{2}$
<sup>17</sup> C	$21\ 023 \pm 35^{\text{b})}$		$(\beta^-)$	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+; \frac{5}{2}$
<sup>17</sup> N	$7870 \pm 15^{\text{d})}$	$4.173 \pm 0.004$ sec	$\beta^-$	$\frac{1}{2}^-; \frac{3}{2}$
<sup>17</sup> O	$-809.9 \pm 0.8^{\text{d})}$	stable <sup>e)</sup>		$\frac{3}{2}^+; \frac{1}{2}$
<sup>17</sup> F	$1951.66 \pm 0.18^{\text{d})}$	$64.49 \pm 0.16$ sec <sup>f)</sup>	$\beta^+$	$\frac{5}{2}^+; \frac{1}{2}$
<sup>17</sup> Ne	$16\ 480 \pm 50^{\text{b})}$	$109.0 \pm 1.0$ msec	$\beta^+$	$\frac{1}{2}^-; \frac{5}{2}$
<sup>17</sup> Na	see text		p	$T = \frac{5}{2}$

<sup>a)</sup> The values of the mass excess shown here were used to calculate  $Q_m$ .

<sup>b)</sup> From data reviewed in the present article.

<sup>c)</sup>  $J^\pi$  values in parentheses are derived from systematics.

<sup>d)</sup> (Wa 77).

<sup>e)</sup>  $\mu = -1.89379 \pm 0.00009$  n.m. [see (Le 78n)] and  $Q = -25.78$  mb (Sc 69u).

<sup>f)</sup>  $\mu = 4.7223 \pm 0.0012$  n.m. [see (Le 78n)] and  $Q = 100 \pm 20$  mb (Mi 741).

TABLE 2  
Electromagnetic transitions in  $A = 16-17^a$ )

Nucleus	$E_u \rightarrow E_u$ (MeV)	$J_i^\pi(T_i) \rightarrow J_f^\pi(T_f)$	$\Gamma_\gamma$ (eV)	Branching ratio (percent)	Multipol.	$\Gamma_\gamma/\Gamma_W$ (W.u.)
$^{16}\text{N}^b)$	0.12 $\rightarrow$ 0	$0^-(1) \rightarrow 2^-(1)$	$(8.7 \pm 0.1) \times 10^{-11}$	100	E2	$1.7 \pm 0.1$
	0.40 $\rightarrow$ 0	$1^- \rightarrow 2^-$	$(3.1 \pm 0.2) \times 10^{-5}$	$26.6 \pm 0.6$	M1	$0.023 \pm 0.001$
	$\rightarrow 0.12$	$\rightarrow 0^-$	$(8.6 \pm 0.5) \times 10^{-5}$	$73.4 \pm 1.6$	M1	$0.19 \pm 0.01$
$^{16}\text{O}^g)$	6.13 $\rightarrow$ 0	$3^-(0) \rightarrow 0^+(0)$	$(2.60 \pm 0.13) \times 10^{-5}$	100	E3	$13.5 \pm 0.7$
	6.92 $\rightarrow$ 0	$2^+ \rightarrow 0^+$	$0.100 \pm 0.004$	>99	E2	$3.1 \pm 0.1$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$(2.7 \pm 0.3) \times 10^{-5}$	$(2.7 \pm 0.3) \times 10^{-2}$	E2	$27 \pm 3$
	$\rightarrow 6.13$	$\rightarrow 3^-$	$\leq 9 \times 10^{-6}$	$\leq 8 \times 10^{-3}$	E1	$\leq 4 \times 10^{-5}$
	7.12 $\rightarrow$ 0	$1^- \rightarrow 0^+$	$(5.7 \pm 0.6) \times 10^{-2}$	>99	E1	$(3.6 \pm 0.4) \times 10^{-4}$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$< 3 \times 10^{-7}$	$\leq 6 \times 10^{-4}$	E1	$< 6 \times 10^{-7}$
	$\rightarrow 6.13$	$\rightarrow 3^-$	$(4 \pm 1) \times 10^{-5}$	$(7.0 \pm 1.4) \times 10^{-2}$	E2	$21 \pm 5$
	8.87 $\rightarrow$ 0	$2^- \rightarrow 0^+$	$(2.4 \pm 0.4) \times 10^{-4}$	$7.2 \pm 0.8$	M2	$0.046 \pm 0.008$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$(4.0 \pm 1.0) \times 10^{-6}^a)$	$0.122 \pm 0.033$	M2	$0.24 \pm 0.06$
	$\rightarrow 6.13$	$\rightarrow 3^-$	$(8.5_{-2.5}^{+4.5}) \times 10^{-4}$	$76.0 \pm 3.0$	M1	$(2.0_{-0.6}^{+1.0}) \times 10^{-3}$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$(1.7 \pm 0.5) \times 10^{-3}$		E2	$5.6 \pm 1.6$
$^{16}\text{O}^g)$	$\rightarrow 7.12$	$\rightarrow 1^-$	$(1.4 \pm 0.4) \times 10^{-4}^a)$	$4.2 \pm 0.8$	E1	$(4.4 \pm 1.2) \times 10^{-5}$
	9.63 $\rightarrow$ 0	$1^- \rightarrow 0^+$	$(4.2 \pm 0.8) \times 10^{-4}^a)$	$12.6 \pm 2.0$	M1 <sup>k)</sup>	$(3.7 \pm 0.7) \times 10^{-3}$
	9.85 $\rightarrow$ 0	$2^+ \rightarrow 0^+$	$(2.3 \pm 0.3) \times 10^{-2}$	$\approx 100$	E1	$(6.0 \pm 0.9) \times 10^{-5}$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$(6.0 \pm 0.5) \times 10^{-3}$	$61 \pm 4$	E2	$0.032 \pm 0.003$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$(1.8 \pm 0.4) \times 10^{-3}^a)$	$18 \pm 4$	E2	$1.1 \pm 0.2$
	10.36 $\rightarrow$ 0	$4^+ \rightarrow 0^+$	$(2.1 \pm 0.4) \times 10^{-3}^a)$	$21 \pm 4$	M1 <sup>k)</sup>	$(4.0 \pm 0.8) \times 10^{-3}$
	$\rightarrow 6.13$	$\rightarrow 3^-$	$(5.6 \pm 2.0) \times 10^{-8}$		E4	$3.7 \pm 1.3$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$< 1 \times 10^{-3}$		E1	$< 3 \times 10^{-5}$
	10.96 $\rightarrow$ 7.12	$0^- \rightarrow 1^-$	$(5.8 \pm 0.7) \times 10^{-2}$	$\approx 100$	E2	$61 \pm 7$
	11.096 $\rightarrow$ 6.13	$4^+ \rightarrow 3^-$	$(8 \pm 5) \times 10^{-4}$	>99	M1	$(7 \pm 4) \times 10^{-4}$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$(3.1 \pm 1.3) \times 10^{-3}$		E1	$(5.9 \pm 2.5) \times 10^{-5}$
$^{16}\text{O}^g)$	11.52 $\rightarrow$ 0	$2^+ \rightarrow 0^+$	$(2.5 \pm 0.6) \times 10^{-3}$		E2	$1.0 \pm 0.3$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$0.61 \pm 0.02$	$91.7$	E2	$1.5 \pm 0.5$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$(2.8 \pm 0.6) \times 10^{-2}^a)$	$4.2 \pm 0.7$	E2	$2.9 \pm 0.6$
			$(2.7 \pm 0.7) \times 10^{-2}^a)$	$4.0 \pm 1.0$	M1 <sup>k)</sup>	$0.013 \pm 0.003$

TABLE 2—continued

Nucleus	$E_{\text{in}} \rightarrow E_{\text{ex}}$ (MeV)	$J_i^{\pi}(T_i) \rightarrow J_f^{\pi}(T_f)$	$\Gamma_{\gamma}$ (eV)	Branching ratio (percent)	Multipol.	$\Gamma_{\gamma}/\Gamma_w$ (W.u.)
$^{17}\text{N}^d)$	$\rightarrow 7.12$	$\rightarrow 1^-$	$\leq 5 \times 10^{-3}$	$\leq 0.8$	E1	$\leq 1 \times 10^{-4}$
	$12.44 \rightarrow 0$	$1^- \rightarrow 0^+$	$12 \pm 2$	$\simeq 100$	E1	$0.014 \pm 0.002$
	$\rightarrow 6.05$	$\rightarrow 0^+$	$0.12 \pm 0.04$	$1.2 \pm 0.4$	E1	$(1.1 \pm 0.4) \times 10^{-3}$
	$12.53 \rightarrow 0$	$2^- \rightarrow 0^+$	<sup>b)</sup>		M2	
	$\rightarrow 6.13$	$\rightarrow 3^-$	$2.1 \pm 0.2$	$60 \pm 6$	M1 <sup>b)</sup>	$0.38 \pm 0.04$
	$\rightarrow 6.92$	$\rightarrow 2^+$	$\leq 0.34$	$\leq 10$	E1	$\leq 4.5 \times 10^{-3}$
	$\rightarrow 7.12$	$\rightarrow 1^-$	$0.5 \pm 0.1$	$15 \pm 3$	M1 <sup>b)</sup>	$0.15 \pm 0.03$
	$\rightarrow 8.87$	$\rightarrow 2^-$	$0.9 \pm 0.1$	$25 \pm 3$	M1	$0.9 \pm 0.1$
	$12.80 \rightarrow 7.12$	$0^-(1) \rightarrow 1^-(0)$	$2.5 \pm 0.2$	$\simeq 100$	M1	$0.65 \pm 0.06$
	$12.97 \rightarrow 0$	$2^-(1) \rightarrow 0^+(0)$	$(7.1 \pm 0.2) \times 10^{-2}$		M2	$2.0 \pm 0.1$
	$\rightarrow 6.13$	$\rightarrow 3^-(0)$	$2.3 \pm 0.2$	$63 \pm 6$	M1	$0.34 \pm 0.03$
	$\rightarrow 7.12$	$\rightarrow 1^-(0)$	$0.44 \pm 0.10$	$12 \pm 3$	M1	$0.10 \pm 0.02$
	$\rightarrow 8.87$	$\rightarrow 2^-(0)$	$0.90 \pm 0.10$	$25 \pm 3$	M1	$0.62 \pm 0.07$
	$13.09 \rightarrow 0$	$1^-(1) \rightarrow 0^+(0)$	$35 \pm 5$	$\simeq 100$	E1	$0.036 \pm 0.005$
	$\rightarrow 6.05$	$\rightarrow 0^+(0)$	$0.21 \pm 0.07$ <sup>b)</sup>	$0.58 \pm 0.12$	E1	$(1.4 \pm 0.5) \times 10^{-3}$
	$\rightarrow 7.12$	$\rightarrow 1^-(0)$	$1.1 \pm 0.3$ <sup>b)</sup>	$3.1 \pm 0.8$	M1	$0.25 \pm 0.07$
	$13.26 \rightarrow 6.13$	$3^-(1) \rightarrow 3^-(0)$	$9.2 \pm 1.5$	$> 85$	M1	$1.2 \pm 0.2$
	$1.37 \rightarrow 0$	$\frac{3}{2}^-(\frac{5}{2}) \rightarrow \frac{1}{2}^+(\frac{3}{2})$	$(7 \pm 3) \times 10^{-3}$	100	M1	$0.13 \pm 0.06$
	$1.85 \rightarrow 0$	$\frac{1}{2}^+ \rightarrow \frac{1}{2}^+$	$(1.4 \pm 0.7) \times 10^{-5}$	$86.5 \pm 2.5$	E1	$(5.0 \pm 2.5) \times 10^{-6}$
	$\rightarrow 1.37$	$\rightarrow \frac{3}{2}^-$	$(2.2 \pm 1.1) \times 10^{-6}$	$13.5 \pm 2.5$	E1	$(4.4 \pm 2.2) \times 10^{-5}$
	$1.91 \rightarrow 0$	$\frac{5}{2}^- \rightarrow \frac{3}{2}^-$	$(4.6 \pm 0.9) \times 10^{-5}$	$77.0 \pm 2.5$	E2	$0.8 \pm 0.2$
	$\rightarrow 1.37$	$\rightarrow \frac{3}{2}^+$	$(1.4 \pm 0.3) \times 10^{-5}$	$23.0 \pm 2.5$	M1	$(4.2 \pm 0.9) \times 10^{-3}$
	$2.53 \rightarrow 0$	$\frac{5}{2}^+ \rightarrow \frac{1}{2}^-$	$(2.2 \pm 0.3) \times 10^{-6}$ <sup>b)</sup>	$11 \pm 1$	M2	$0.21 \pm 0.03$
	$\rightarrow 1.37$	$\rightarrow \frac{3}{2}^+$	$(6.9 \pm 1.0) \times 10^{-6}$ <sup>b)</sup>	$34 \pm 3$	E1	$(9.8 \pm 1.5) \times 10^{-6}$
	$\rightarrow 1.85$	$\rightarrow \frac{1}{2}^+$	$(2.4 \pm 0.5) \times 10^{-6}$ <sup>b)</sup>	$12.0 \pm 1.5$	E2	$7.7 \pm 1.5$
	$\rightarrow 1.91$	$\rightarrow \frac{3}{2}^-$	$(8.4 \pm 1.0) \times 10^{-6}$ <sup>b)</sup>	$41.0 \pm 2.5$	E1	$(7.8 \pm 0.9) \times 10^{-5}$
	$3.13 \rightarrow 1.91$	$\frac{7}{2}^- \rightarrow \frac{3}{2}^-$	$(2.4 \pm 0.7) \times 10^{-3}$	100	M1	$0.063 \pm 0.018$
	$3.63 \rightarrow 1.91$	$(\frac{7}{2}, \frac{5}{2})^- \rightarrow \frac{3}{2}^-$	$(2.6 \pm 0.8) \times 10^{-5}$	$47 \pm 10$	(E2) <sup>b)</sup>	$0.8 \pm 0.3$
	$\rightarrow 3.13$	$\rightarrow \frac{1}{2}^-$	$(2.9 \pm 1.0) \times 10^{-5}$	$53 \pm 10$	M1 <sup>b)</sup>	$0.011 \pm 0.004$

$^{17}\text{O}^*$	0.87 → 0 3.06 → 0.87	$\frac{1}{2}^+(\frac{1}{2}) \rightarrow \frac{5}{2}^+(\frac{1}{2})$ $\frac{1}{2}^- \rightarrow \frac{5}{2}^+$ $\frac{5}{2}^- \rightarrow \frac{5}{2}^+$ $\frac{5}{2}^- \rightarrow \frac{1}{2}$	$(2.55 \pm 0.03) \times 10^{-6}$ $(5.5 \pm 3.7) \times 10^{-3}$ $>0.026$	100 100 100	E2 E1 E1	$2.39 \pm 0.03$ $(1.2 \pm 0.5) \times 10^{-3}$ $>1 \times 10^{-3}$
$^{17}\text{F}^t$	3.84 → 0 0.50 → 0 3.10 → 0.50 3.86 → 0 11.19 → 0.50	$\frac{1}{2}^+(\frac{1}{2}) \rightarrow \frac{5}{2}^+(\frac{1}{2})$ $\frac{1}{2}^- \rightarrow \frac{1}{2}^+$ $\frac{1}{2}^- \rightarrow \frac{5}{2}^+$ $\frac{1}{2}^- \rightarrow \frac{1}{2}$ $\frac{5}{2}^- \rightarrow \frac{1}{2}^+$	$(1.60 \pm 0.03) \times 10^{-6}$ $(12 \pm 2) \times 10^{-3}$ $0.11 \pm 0.02$ $6.0 \pm 2.5$	100 100 100 100	E2 E1 E1 E1	$25.0 \pm 0.5$ $(1.5 \pm 0.3) \times 10^{-3}$ $(4.3 \pm 0.8) \times 10^{-3}$ $0.011 \pm 0.005$

<sup>a</sup>) See also (En 79a). The last columns give the  $\gamma$ -ray strengths expressed in Weisskopf units [see D. H. Wilkinson, in Nuclear spectroscopy B, ed. F. Ajzenberg-Selove, Academic Press, NY, 1960]. The Weisskopf estimates ( $\Gamma_W$  in eV,  $E_\gamma$  in MeV) are

$$\Gamma_W(E1) = 6.8 \times 10^{-2} A^{2/3} E_\gamma^3$$

$$\Gamma_W(E2) = 4.9 \times 10^{-8} A^{4/3} E_\gamma^5$$

$$\Gamma_W(E3) = 2.3 \times 10^{-14} A^2 E_\gamma^7$$

$$\Gamma_W(E4) = 6.8 \times 10^{-21} A^{8/3} E_\gamma^9$$

$$\Gamma_W(M1) = 2.1 \times 10^{-2} E_\gamma^3$$

$$\Gamma_W(M2) = 1.5 \times 10^{-8} A^{2/3} E_\gamma^5$$

The values for these  $\gamma$ -ray strengths are occasionally different from those listed in other tables of this paper because different values of  $r$  were used. In this table  $r_0 = 1.2$  fm is used consistently. The multipolarities in the next to the last column were used to calculate the  $\Gamma_W$ .

<sup>b</sup>) See also table 16.4.

<sup>c</sup>) See also table 16.12.

<sup>d</sup>) See also table 17.5.  $\Gamma_\gamma$  based on  $\tau_m$  and mean value of branching ratio if more than one measurement is reported. See table 17.5 also for the  $\gamma$ -decay of other  $^{17}\text{N}$  states.

<sup>e</sup>) See also table 17.7.

<sup>f</sup>) See reaction 10 in  $^{17}\text{F}$  and table 17.11. For other resonances in  $^{16}\text{O}(p, \gamma)^{17}\text{F}$  see table 17.20.

<sup>g</sup>) Based on branching ratio in table 16.12 and on  $\Gamma_\gamma$  shown here for ground state transition.

<sup>h</sup>) See table 16.12.

<sup>i</sup>) (Gu 76d).

<sup>j</sup>)  $\frac{5}{2}^-$  assumed for  $^{17}\text{N}^*(3.63)$ .

<sup>k</sup>) Assumed, but  $\delta$  is not known. I am indebted to Professor P. M. Endt for his comments.

<sup>l</sup>) Values corrected for sum of branching ratios = 100%.

# $^{16}\text{He}$ , $^{16}\text{Be}$ , $^{16}\text{B}$ , $^{16}\text{C}$ MASTER TABLE

6

F. AJZENBERG-SELOVE

## $^{16}\text{He}$

(Not illustrated)

This nucleus has not been observed. See also (Na 78e; theor.).

## $^{16}\text{Be}$

(Not illustrated)

This nucleus has not been observed. Its atomic mass excess is calculated to be 59.22 MeV. It is then unstable with respect to breakup into  $^{14}\text{Be} + 2\text{n}$  by 2.4 MeV (Th 74). See also (Se 81c; theor.).

## $^{16}\text{B}$

(Not illustrated)

This nucleus has not been observed in the 4.8 GeV proton bombardment of a uranium target: it is particle unstable (Bo 74a). Its mass excess is predicted to be 37.97 MeV: it would then be unstable with respect to decay into  $^{15}\text{B} + \text{n}$  by 0.9 MeV [mass excess calculated using the transverse form of the mass equation] (Th 74, Je 75). See also (Se 81c; theor.).

## $^{16}\text{C}$

(Figs. 1 and 5)

### GENERAL<sup>†</sup>


*Experimental work:* (Ar 77b, Ch 81e).

*Theory:* (A1 79n).

TABLE 16.1  
Energy levels of  $^{16}\text{C}$

$E_x$ (MeV $\pm$ keV)	$J''; T$	$\tau_{1/2}$ (sec) or $\Gamma$ (keV)	Decay	Reactions
0 $1.766 \pm 10$	$0^+; 2$ $2^+$	$\tau_{1/2} = 0.747 \pm 0.008$	$\beta^-$	1, 2, 3
$3.027 \pm 12$	$(0^+)$		$\gamma$	2
$3.986 \pm 7$	2		$(\gamma)$	2
$4.088 \pm 7$	$3(^+)$		$\gamma$	2
$4.142 \pm 7$	$4^+$		$\gamma$	2
$6.109 \pm 15$	$(2^+, 3^-, 4^+)$	$\Gamma \leq 25$	$\gamma$	2

<sup>†</sup> See also (Aj 77).

Fig. 1. Energy levels of  $^{16}\text{C}$ . For notation see fig. 2.

**Mass of  $^{16}\text{C}$ :** The atomic mass excess of  $^{16}\text{C}$  based on the  $^{14}\text{C}(t, p)Q_0$  measurements of (Fo 77d, Se 78a) and on the (Wa 77) masses for  $^{14}\text{C}$ ,  $t$  and  $p$  is  $13\,694 \pm 4$  keV.

$$1. \quad ^{16}\text{C}(\beta^-)^{16}\text{N} \qquad Q_m = 8.012$$

The half-life of  $^{16}\text{C}$  is  $0.747 \pm 0.008$  sec; it decays to  $^{16}\text{N}^*(3.34, 4.32)$  [both  $J^\pi = 1^+$ ] with branchings of 84% and 16% respectively [ $\log ft = 3.55, 3.83$ ]; see table 16.2 (Al 76). See also (Fo 78g).

TABLE 16.2  
The  $\beta^-$  decay of  $^{16}\text{C}^*$ )

Decay to $^{16}\text{N}^*$ (MeV)	$J^\pi$	Branch (%)	$\log ft$
0.120	$0^-$	<0.5 <sup>b)</sup>	>6.85
0.297	$3^-$	<0.5 <sup>b)</sup>	>6.83
0.398	$1^-$	<0.7 <sup>b)</sup>	>6.64
3.36	$1^+$	$84.4 \pm 1.7$	$3.551 \pm 0.012$
4.32	$1^+$	$15.6 \pm 1.7$	$3.83 \pm 0.05$

<sup>a)</sup> (Al 76).

<sup>b)</sup> The combined branching to  $^{16}\text{N}^*(0.120, 0.297, 0.398)$  is <1.2% (Al 76).

TABLE 16.3  
States of <sup>16</sup>C from <sup>14</sup>C (t, p)

$E_x$ (keV) (Fo 77d)	$E_x$ (keV) (Se 78a)	$L$ <sup>a)</sup>	$J$ <sup>b)</sup> (Ba 77qq)
0	0	0	
$1766 \pm 10$	$1.77^c)$	2	2
$3020 \pm 15$	$3039 \pm 20$	$(0)^c)$	
$3983 \pm 10$	$3990 \pm 10$	$(2)^c)$	$2^f)$
$4083 \pm 10$	$4094 \pm 10$	$(3)^c)$	$3^g)$
$4136 \pm 10$	$4194 \pm 10$	4	4
$6109 \pm 15^d)$		2, 3, 4	

<sup>a)</sup> From angular distribution measurements analyzed by DWBA  
 $E_t = 18$  MeV (Fo 77d) and 23 MeV (Se 78a).

<sup>b)</sup> From  $\gamma\gamma$  correlation measurements at  $E_t = 12$  MeV (Ba 77qq).

<sup>c)</sup> This state is weakly excited.

<sup>d)</sup>  $\Gamma \leq 25$  keV.

<sup>e)</sup> State observed but  $E_x$  not determined.

<sup>f)</sup> (Se 78a) suggest that the state is  $2^-$  by comparison with the transition to <sup>14</sup>C\*(7.38).

<sup>g)</sup> The very low intensity of the proton group to <sup>16</sup>C\*(4.14) suggests  $J^\pi = 3^+$ : see e.g. (Se 78a).

## 2. <sup>14</sup>C(t, p)<sup>16</sup>C

$$Q_m = -3.013$$

$$Q_0 = -3015 \pm 8 \text{ keV (Fo 77d)}$$

$$Q_0 = -3013 \pm 4 \text{ keV (Se 78a)}$$

Observed proton groups are displayed in table 16.3 (Fo 77d, Se 78a), together with the angular correlation results of (Ba 77qq). The similarities in the spectra of <sup>14</sup>C and <sup>16</sup>C are discussed by (Fo 78f, Se 78a). See also (Fo 77g, Fo 78g).

## 3. <sup>14</sup>C(<sup>18</sup>O, <sup>16</sup>O)<sup>16</sup>C

$$Q_m = 6.720$$

See (Aj 77).

## <sup>16</sup>N

(Figs. 2 and 5)

## GENERAL<sup>†</sup>

*Model calculations:* (Ro 79c, Ha 80p).

*Reactions involving muons:* (Bo 77o, Er 77d, Na 77a, Gu 78c, Pa 78h, Ro 78l, Er 79c, Gu 79, Gu 79c, Ho 79d, Ki 79m, Rh 79, St 79l, Wu 79d, Br 80, Ch 80b, Gr 80, Ei 81).

<sup>†</sup> See also (Aj 77).




Fig. 2. Energy levels of  $^{16}\text{N}$ . In these diagrams, energy values are plotted vertically in MeV, based on the ground state as zero. Uncertain levels or transitions are indicated by dashed lines; levels which are known to be particularly broad are cross-hatched. Values of total angular momentum  $J$ , parity, and isobaric spin  $T$  which appear to be reasonably well established are indicated on the levels; less certain assignments are enclosed in parentheses. For reactions in which  $^{16}\text{N}$  is the compound nucleus, some typical thin-target excitation functions are shown schematically, with the yield plotted horizontally and the bombarding energy vertically. Bombarding energies are indicated in laboratory coordinates and plotted to scale in c.m. coordinates. Excited states of the residual nuclei involved in these reactions have generally not been shown; where transitions to such excited states are known to occur, a brace is sometimes used to suggest reference to another diagram. For reactions in which the present nucleus occurs as a residual product, excitation functions have not been shown; a vertical arrow with a number indicating some bombarding energy, usually the highest, at which the reaction has been studied, is used instead. Further information on the levels illustrated, including a listing of the reactions in which each has been observed, is contained in the master table, entitled "Energy levels of  $^{16}\text{N}$ ".

# <sup>16</sup>N MASTER TABLE

10

F. AJZENBERG-SELOVE

*Reactions involving pions:* (Al 77aa, Ba 77xx, Be 77aa, Er 77d, Ho 77f, Sh 77n, Su 77h, Fu 78g, Na 78i, Oh 78, Bo 79ee, De 79k, Er 79c, Kn 79c, Na 79q, Pe 79a, Sr 79a, Tr 79h, Wi 79d, Wi 79e, Bo 80a, Bo 80q, De 80bb, Gr 80, St 80d, Wu 80, Ya 81).

*Other topics:* (Le 76t, Ar 77b, Ba 77t, Br 77, Ha 77i, Bu 78k, Fo 78g, Ga 78c, Ge 78b, He 78h, Ko 78a, Le 78r, Ab 79a, Al 79n, Gr 81b).

TABLE 16.4  
Energy levels of <sup>16</sup>N

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$2^-; 1$	$\tau_{1/2} = 7.13 \pm 0.02$ sec	$\beta^-$	1, 3, 10, 12 $\rightarrow$ 15, 18, 19, 24 $\rightarrow$ 26, 28, 29
$0.1201 \pm 0.5$	$0^-$	$\tau_m = 7.58 \pm 0.09$ $\mu$ sec	$\gamma, \beta^-$	10, 12, 13, 14, 18, 21, 24, 28, 29
$0.2970 \pm 0.7$	$3^-$	$^*)$	$\gamma$	5, 10, 12, 13, 14, 18, 19, 24, 26, 28, 29
$0.3975 \pm 0.7$	$1^-$	$\tau_m = 5.6 \pm 0.3$ psec $ g  = 1.5 \pm 0.1$	$\gamma$	5, 10 $\rightarrow$ 14, 18, 21, 24, 28, 29
$3.355 \pm 5$	$1^+$	$\Gamma = 15 \pm 5$	n	3, 10, 12, 16, 18, 20, 28
$3.519 \pm 5$	$2^+$	3	n	3, 10, 12, 16, 18, 28
$3.960 \pm 5$	$3^+$	$\leq 2$	n	3, 7, 10, 11, 12, 16, 18, 28
$4.319 \pm 5$	$1^+$	$20 \pm 5$	n	3, 10, 12, 16, 18, 19, 28
$4.387 \pm 6$	$1^-$	$82 \pm 20$	n	3, 10, 12, 16, 18, 28
$4.76 \pm 50$	$1^-$	$250 \pm 50$	n	12, 16, 18
$4.776 \pm 10$	$2^+$	$59 \pm 8$	n	10, 12, 16, 18, 28
( $4.90 \pm 10$ )				18
$5.050 \pm 6$	$2^-$	$19 \pm 6$	n	10, 12, 16, 18, 28
$5.130 \pm 7$	$\geq 2$	$\leq 7 \pm 4$	n	10, 12, 16, 18
$5.150 \pm 7$	( $2, 3$ ) <sup>-</sup>	$\leq 7 \pm 4$	n	12, 16, 18, 28
$5.232 \pm 5$	$3^+$	$\leq 4$	n	10, 12, 16, 18, 28
5.24	$1^+$	260	n	16
$5.25 \pm 70$	$2^-$	$320 \pm 80$	n	12, 18
$5.518 \pm 6$	$3^+$	$\leq 7 \pm 4$	n	10, 12, 16, 18, 28
$5.730 \pm 6$	( $5^+$ )	$\leq 7 \pm 4$	n	7, 10, 11, 12, 16, 18, 28
$6.009 \pm 10$	$1^-$	$270 \pm 30$	n	12, 16, 28
$6.168 \pm 4$	( $4^-$ )	$\leq 7 \pm 4$	n	10, 12, 18, 26, 28
$6.373 \pm 6$	( $3^-$ )	$30 \pm 6$	n	12, 16, 18, 26, 28
$6.426 \pm 7$		$300 \pm 30$		12, 18
$6.513 \pm 6$	$1^+$	$34 \pm 6$	n	12, 16, 18, 28
$6.613 \pm 6$		$\leq 7 \pm 4$		12, 13, 14, 18, 28
$6.848 \pm 6$		$\leq 7 \pm 4$		10, 12, 18, 28
( $6.84$ )	$\geq 2$	$> 140$	n	16
$7.02 \pm 20$	$1^+$	$22 \pm 5$	n	12, 16, 18, 28
$7.134 \pm 7$		$\leq 7 \pm 4$		10, 12, 18, 28
$7.250 \pm 7$	$\geq 2$	$17 \pm 5$	n	12, 16, 18, 28
$7.573 \pm 6$	$\geq 3$	$\leq 7 \pm 4$	n	10, 12, 16, 18, 28
$7.637 \pm 5$	( $3, 4, 5$ ) <sup>+</sup>	$\leq 7 \pm 4$	n	7, 10, 11, 12, 18, 28
$7.675 \pm 5$		$\leq 7 \pm 4$	n	7, 12, 13, 16, 18, 28
$7.877 \pm 9$	$\geq 4$	$100 \pm 15$	n	12, 16, 18, 28

TABLE 16.4—continued

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$E_{\text{c.m.}}$ (keV)	Decay	Reactions
8.048 $\pm$ 9		85 $\pm$ 15	n	12, 16, 28
8.182 $\pm$ 9	(3, 2) <sup>+</sup>	28 $\pm$ 8		10, 12, 17
8.282 $\pm$ 8		24 $\pm$ 8		12, 28
8.365 $\pm$ 8	$\geq 1$	18 $\pm$ 8	n	12, 16, 28
8.49 $\pm$ 30	$\geq 1$	$\leq 50$	n	16, 28
8.72	$\geq 1$	40	n	16
8.819 $\pm$ 15		$\leq 50$	n	16, 28
9.035 $\pm$ 15		$\leq 50$		28
9.16 $\pm$ 30	$\geq 2$	100	n	16, 28
9.34 $\pm$ 30		$\leq 50$	n	16, 28
9.459 $\pm$ 15	$\geq 2$	100	n	16, 28
9.760 $\pm$ 10	$T = 1$	15 $\pm$ 8		10
9.813 $\pm$ 10	$T = 1$			10
9.928 $\pm$ 7	$0^+; T = 2$	<12		10, 27
10.055 $\pm$ 15	$\geq 3$	30	n	16
10.27	$\geq 2$	165	n	16
10.71	$\geq 2$	120	n	16
11.49	$\geq 3$		n	6, 16
11.62	$\geq 3$	220	n, d	9
11.701 $\pm$ 7	$1^-, 2^+;$ $T = 2$	<12		10
(11.92)		390	n, d	9
(12.09)			n	6, 16
12.26		290	n, p, d	9
(12.46)			n	16
12.61		180	n, p, d	9
12.88		155	n, p, d	9
(12.97)		175	n, d	9
13.12			n, (d)	9, 16
13.83			n	16
14.41 $\pm$ 50	(3) <sup>+</sup>	180	d	9

<sup>a</sup>) The previously reported  $\tau_m$  needs, in the opinion of the reviewer, to be remeasured: see (Aj 71) for the previously reported value.

$$1. \ ^{16}\text{N}(\beta^-)^{16}\text{O} \quad Q_m = 10.418$$

The half-life of  $^{16}\text{N}$  is  $7.13 \pm 0.02$  sec: see table 16.3 in (Aj 71). See also (Sa 75i). From the character of the beta decay [see table 16.21] it is concluded that  $^{16}\text{N}_{\text{g.s.}}$  has  $J^\pi = 2^-$ : see  $^{16}\text{O}$ . The beta decay of  $^{16}\text{N}^*(0.12)[J^\pi = 0^-, \tau_{1/2} = 5.26 \pm 0.06 \mu\text{sec}]$  to  $^{16}\text{O}_{\text{g.s.}}$  has been studied: the  $\beta$ -decay rate  $\lambda_\beta = 0.43 \pm 0.10 \text{ sec}^{-1}$  (Pa 75a). See also (Ak 77a; applications) and (Bo 77o, Gu 78c, Gu 79c, Ko 79bb, Mu 79b, Rh 79, Ch 80b, Ch 81, To 81e; theor.).

2. (a)  $^9\text{Be}(^7\text{Li}, t)^{13}\text{C}$   $Q_m = 8.181$   $E_b = 20.575$
- (b)  $^9\text{Be}(^7\text{Li}, \alpha)^{12}\text{B}$   $Q_m = 10.462$
- (c)  $^9\text{Be}(^7\text{Li}, {^8\text{Li}})^8\text{B}$   $Q_m = -17.613$

The yields of  $t_0$  and of  $\alpha_0$ ,  $\alpha_1$  and  $\alpha_2$  have been measured for  $E(^7\text{Li}) = 4$  to 14 MeV at  $0^\circ$ : several broad peaks are observed (Wy 71). The yields of  $\alpha_0$  and  $\alpha_2$  have also been studied for  $E(^7\text{Li}) = 3.3$  and 5.0 to 6.2 MeV (Sn 69b). The cross section for reaction (c) rises monotonically for  $E(^7\text{Li}) = 1.1$  to 4 MeV (No 57a, No 59a).

$$3. \ ^{10}\text{B}(^7\text{Li}, p)^{16}\text{N} \quad Q_m = 13.989$$

Angular distributions have been obtained at  $E(^7\text{Li}) = 16$  MeV to  $^{16}\text{N}^*(3.36, 3.52, 3.96, 4.32, 4.39)$ . The  $\sigma_t$  for  $^{16}\text{N}^*(3.36, 3.96, 4.32, 4.39)$ , whose  $J''$  are known, follow the  $2J_f + 1$  relation. From the  $\sigma_t$  for  $^{16}\text{N}^*(3.52)$  it follows then that  $J = 2$  if it is a single state or  $J = 0$  and 1 (with one of the two states having odd parity) if the group corresponds to two unresolved states (Fo 75i). See also (Aj 71).

$$4. \ ^{11}\text{B}(^6\text{Li}, p)^{16}\text{N} \quad Q_m = 9.785$$

See (Mc 66a).

$$5. \ ^{11}\text{B}(^7\text{Li}, d)^{16}\text{N} \quad Q_m = 4.759$$

See (Mc 66a) and (Th 69a:  $E_x = 297.6 \pm 0.9$  and  $397.8 \pm 1.0$  keV).

$$6. \ ^{13}\text{C}(\alpha, p)^{16}\text{N} \quad Q_m = -7.421$$

At  $E_\alpha = 50$  MeV proton groups are reported to  $^{16}\text{N}^*(5.1, 5.6, 7.6, 11.4, 12.1)$  (Bu 76k). See also (Fo 77g, Ha 81d).

$$7. \ ^{13}\text{C}(^6\text{Li}, ^3\text{He})^{16}\text{N} \quad Q_m = 3.401$$

At  $E(^6\text{Li}) = 44$  MeV angular distributions to  $^{16}\text{N}^*$  (3.96, 5.73, 7.65, 11.21, 11.81) are strongly forward peaked (Ma 771; prelim.).

$$8. \ ^{14}\text{C}(d, \gamma)^{16}\text{N} \quad Q_m = 10.474$$

TABLE 16.5  
Resonances in  $^{14}\text{C} + d$

$E_d$ (MeV)	Resonant for	$\Gamma_{c.m.}$ (keV)	$E_x$ (MeV)	Refs.
1.30	$n_0$	220	11.61	(Ch 61a, Im 63)
1.65	$n_0$	390	11.92	(Ch 61a)
2.04	$n_0, p$	290	12.26	(Ch 61a, Im 63, Do 56b)
2.44	$n_0, p$	180	12.61	(Ch 61a, Do 56b)
2.75	$n_0, p$	155	12.88	(Ch 61a, Im 63, Do 56b)
2.86	$n_0$	175	12.97	(Ch 61a)
(3.10)	$n_0$	(175)	(13.18)	(Ch 61a)
$4.50 \pm 0.05$	$d_0$	180 *)	14.41	(Co 73u)

\*)  $\Gamma_d = 45$  keV;  $I_d = 4$ ;  $J'' = 3^+, (4^+, 5^+)$  (Co 73u:  $E_d = 4$  to 10 MeV).

The cross section has been measured for  $1.2 < E_d < 2.6$  MeV. It shows some evidence of structure (Ne 64f). See also (Aj 71).

9. (a) $^{14}\text{C}(\text{d}, \text{n})^{15}\text{N}$	$Q_m = 7.9828$	$E_b = 10.474$
(b) $^{14}\text{C}(\text{d}, \text{p})^{15}\text{C}$	$Q_m = -1.0065$	
(c) $^{14}\text{C}(\text{d}, \text{d})^{14}\text{C}$		
(d) $^{14}\text{C}(\text{d}, \text{t})^{13}\text{C}$	$Q_m = -1.9192$	
(e) $^{14}\text{C}(\text{d}, \alpha)^{12}\text{B}$	$Q_m = 0.361$	

Observed resonances for  $n_0$ ,  $p$  and  $d_0$  are displayed in table 16.5. For polarization measurements in reaction (a) see (Aj 77). For reaction (b) see also (Aj 77). See also  $^{12}\text{B}$  in (Aj 68) and  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{C}$  and  $^{15}\text{N}$  in (Aj 81).

10. $^{14}\text{C}(^3\text{He}, \text{p})^{16}\text{N}$	$Q_m = 4.981$
---	---------------

Proton groups have been observed to  $^{16}\text{N}$  states with  $E_x < 12$  MeV and angular distributions (with  $E(^3\text{He}) \leq 15$  MeV) lead to the  $J^\pi$  assignments shown in table 16.6.

TABLE 16.6  
Excited states in  $^{16}\text{N}$  from  $^{14}\text{C}(^3\text{He}, \text{p})^{16}\text{N}$

$E_x$ (MeV $\pm$ keV) (Ga 66d)	$\Gamma$ (keV)	$J^\pi; T$
$0.121 \pm 6$		$0^-$
$0.298 \pm 6$		$3^-$
$0.396 \pm 7$		
$3.348 \pm 7$		$1^+$
$3.517 \pm 7$		$2^+, (3)^+$
$3.958 \pm 7$		$(2)^+, 3^+$
$4.313 \pm 9$		$1^+$
$4.386 \pm 9$		
$4.768 \pm 11$		
$5.052 \pm 9$		
$5.137 \pm 9$		
$5.234 \pm 9$		
$5.512 \pm 5$		$(1, 2, 3)^+$
$5.724 \pm 5$		$(1, 2, 3)^+$
$6.168 \pm 5$		$5^+$
$6.843 \pm 5$		
$7.113 \pm 5$		
$7.570 \pm 5$		
$7.636 \pm 5$		
$7.673 \pm 5$		
$8.205 \pm 5$		
$9.760 \pm 10$	$15 \pm 8$	$T = 1$
$9.813 \pm 10$		$T = 1$
$9.928 \pm 7$	$< 12$	$0^+; 2$
$11.701 \pm 7$	$< 12$	$1^-, 2^+; 2$

TABLE 16.7  
States in <sup>16</sup>N from <sup>14</sup>N(t, p)<sup>16</sup>N

(He 66a)		<i>L</i> <sup>a)</sup>	<i>J</i> <sup>"</sup> <sup>a)</sup>
<i>E</i> <sub>x</sub> (MeV ± keV)	<i>Γ</i> (keV)		
0		3	2 <sup>- b)</sup>
0.120 ± 10		1	0 <sup>- b)</sup>
0.300 ± 10		3	3 <sup>- b)</sup>
0.399 ± 10		1	1 <sup>- b)</sup>
3.359 ± 10	15 ± 5	0	1 <sup>+ b)</sup>
3.519 ± 10	≤7 ± 4	°)	
3.957 ± 10	≤7 ± 4	2	3 <sup>+ b)</sup>
4.318 ± 10	20 ± 5	0	1 <sup>+ b)</sup>
4.391 ± 10	82 ± 20	1	1 <sup>- b)</sup>
4.725 ± 10 <sup>d)</sup>	290 ± 30	1	1 <sup>-</sup>
4.774 ± 10	59 ± 8	2	2 <sup>- b)</sup>
5.053 ± 10	19 ± 6	(1 + 3)	2 <sup>-</sup>
5.130 ± 10	≤7 ± 4	°)	
5.150 ± 10	≤7 ± 4		
5.226 ± 10	≤7 ± 4	2	(1, 2, 3) <sup>+</sup>
5.305 ± 10 <sup>d)</sup>	260 ± 30	°)	
5.520 ± 10	≤7 ± 4	(0, 1) + 2 + 4 <sup>e)</sup>	
5.730 ± 10	≤7 ± 4	(1, 3) + 4 <sup>e)</sup>	
6.009 ± 10	270 ± 30	1	1 <sup>-</sup>
6.167 ± 10	≤7 ± 4	(3)	(4 <sup>-</sup> )
6.371 ± 10	30 ± 6	(3)	(3 <sup>-</sup> )
6.422 ± 10	300 ± 30	0 <sup>+(2, 4)<sup>e)</sup></sup>	
6.512 ± 10	34 ± 6	0 <sup>+(2, 3)</sup>	1 <sup>+</sup>
6.613 ± 10	≤7 ± 4	(2 + 4) or 3	
6.854 ± 10	≤7 ± 4	3 or (2 + 4)	
7.006 ± 10	22 ± 5	0 <sup>+(+2)</sup>	1 <sup>+</sup>
7.133 ± 10	≤7 ± 4	(3, 2)	
7.250 ± 10	17 ± 5	(2 + 4) or 3	
7.573 ± 10	≤7 ± 4	3 or (2 + 4)	
7.640 ± 10	≤7 ± 4	4	3, 4 <sup>-</sup>
7.675 ± 10	≤7 ± 4	(1 + 4)	(3, 4, 5) <sup>+</sup>
7.876 ± 10	100 ± 15	1 + 4 <sup>e)</sup>	
8.043 ± 10	85 ± 15	(2 + 4) or 3	
8.183 ± 10	28 ± 8	2 (+4)	(3, 2) <sup>+</sup>
8.280 ± 10	24 ± 8	(1)	((0, 1, 2) <sup>-</sup> )
8.361 ± 10	18 ± 8	(1 + 4) <sup>e)</sup>	

<sup>a)</sup> From reanalysis of data of (He 66a): see (Cr 75a, Fo 78c).

<sup>b)</sup> Identified with shell-model counterparts (Cr 75a).

<sup>c)</sup> Results are ambiguous (Cr 75a).

<sup>d)</sup> The errors listed here for the *E*<sub>x</sub> to these two broad peaks are probably underestimates: I am indebted to Dr. H. Fuchs for his comments.

<sup>e)</sup> May be a doublet (Fo 78c).



At  $E_\alpha = 46$  MeV the angular distributions of the groups to  $^{16}\text{N}^*(0.30, 3.96, 5.73, 7.60)$  have been determined: the most strongly populated state is the ( $5^+$ ) state  $^{16}\text{N}^*(5.73)$  (Lu 69a).



Observed proton groups are displayed in table 16.7. Angular distributions have also been measured at  $E_t = 15$  MeV to the first four states of  $^{16}\text{N}$ : cross sections obtained by DWBA analysis using microscopic wave functions are in poor agreement with the data (Fo 79). These distributions have also been compared with those in the ( $^3\text{He}, \text{p}$ ) reaction to analog states in  $^{16}\text{O}$ : see reaction 38 in  $^{16}\text{O}$  (Fo 78r).  $\tau_m$  for  $^{16}\text{N}^*(0.40) = 5.1 \pm 0.3$  psec (He 77e, He 77b). See also (Aj 77).



At  $E_\alpha = 65$  MeV strong transitions are observed to the unresolved ground state quartet and to  $^{16}\text{N}^*(5.25, 6.61, 7.68)$  (Ja 78i).




At  $E({}^{10}\text{B}) = 100$  MeV strong transitions are observed to the unresolved ground state quartet and to  $^{16}\text{N}^*(5.14, 6.59, 7.65)$  (Ha 78a). For reaction (b) see (Pr 80c).



The thermal cross section is  $24 \pm 8$   $\mu\text{b}$  (Mu 73i).



The scattering amplitude (bound)  $a = 6.44 \pm 0.03$  fm,  $\sigma_{\text{free}} = 4.59 \pm 0.05$  b,  $\sigma_{\text{inc}}^{\text{spin}}$  (bound nucleus)  $< 1$  mb (Ko 79w). The total cross section has been measured for  $E_n = 0.4$  to 32 MeV: see (Aj 71, Ga 76f) and (Ze 71). Observed resonances and parameters derived from  $R$ -matrix and phase-shift analyses of these data, angular distributions and polarization measurements are displayed in table 16.8 (Ze 71, Do 71a). See also (Aj 77) and (Ro 76u, Ro 77k, Ro 78j, Ab 79e, Ro 79d, Ha 80p; theor.).



For reaction (b) see (Aj 76, Ba 78e). For reaction (c) see (Ba 78e). For reactions (a) and (d) see (Aj 77).

## F. AJZENBERG-SELOVE

 TABLE 16.8  
 Resonances in <sup>15</sup>N(n, n)<sup>15</sup>N<sup>a,b)</sup>

<i>E<sub>n</sub></i> (MeV ± keV)	<i>E<sub>lab</sub></i> (keV)	<i>E<sub>x</sub></i> (MeV)	<i>J<sup>π</sup></i>
0.921	14	3.354	1 <sup>+ c)</sup>
1.095	3	3.517	1
1.563	≤2	3.955	1
1.944	29	4.312	1 <sup>+ d)</sup>
2.038	56	4.400	1 <sup>- d)</sup>
2.30 ± 70 <sup>e)</sup>	410 ± 100 <sup>e)</sup>	4.65	1 <sup>- d)</sup>
2.399	107	4.738	2 <sup>+ d)</sup>
2.732	35	5.050	1 <sup>- e)</sup>
2.830	12	5.142	3 <sup>(-)</sup>
2.84 ± 70 <sup>f)</sup>	710 ± 100 <sup>f)</sup>	5.15	2 <sup>- d)</sup>
2.915	4	5.222	≥2
2.93	260	5.24	1 <sup>+</sup>
3.225		5.512	
3.454	24	5.727	1 <sup>+</sup>
3.69	297	5.95	1 <sup>-</sup>
3.987	88	6.226	(1 <sup>+</sup> )
4.126	78	6.356	(3 <sup>-</sup> )
4.252	113	6.474	(2 <sup>+</sup> )
4.64	>150	6.84	≥2
4.80	37	6.99	≥1
5.055	25	7.227	≥2
5.43	30	7.58	≥3
5.56		7.70	
5.73	165	7.86	≥4
5.90		8.02	
6.28		8.37	≥1
6.42		8.51	≥1
6.65	45	8.72	≥1
6.76		8.82	
7.10	110	9.14	≥2
7.31		9.34	
7.44	105	9.46	≥2
7.71	150	9.71	≥2
8.07	30	10.05	≥3
8.30	175	10.27	≥2
8.77	130	10.71	≥2
9.61		11.49	≥3
9.77		11.64	≥3
10.25		12.09	
10.64		12.46	
11.09		12.88	
11.41		13.12	
12.10		13.83	

18.  $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$   $Q_m = 0.267$

Levels derived from observed proton groups and  $\gamma$ -rays are listed in table 16.9. Gamma transitions are shown in the inset of fig. 2: (Pa 71f) report that the branchings of  $^{16}\text{N}^*(0.40)$  to  $^{16}\text{N}^*(0, 0.12, 0.30)$  are, respectively,  $26.6 \pm 0.6\%$ ,  $73.4 \pm 1.6\%$  and  $\leq 0.15\%$ . The  $0.30 \rightarrow 0.12$  transition is  $\leq 1\%$  (Pa 71f).

The mean life of  $^{16}\text{N}^*(0.12)$  is  $7.58 \pm 0.09$   $\mu\text{sec}$  (Be 67j); together with the angular distribution analyses this leads to  $J^\pi = 0^-$  for this state. The very strong evidence for  $J^\pi = 2^-, 3^-$  and  $1^-$ , respectively, for  $^{16}\text{N}^*(0, 0.30, 0.40)$  is reviewed in (Aj 71). See also (Ba 77t, Pa 78b, Pa 79a; theor.).

19. (a)  $^{15}\text{N}({}^{11}\text{B}, {}^{10}\text{B})^{16}\text{N}$   $Q_m = -8.964$   
 (b)  $^{15}\text{N}({}^{13}\text{C}, {}^{12}\text{C})^{16}\text{N}$   $Q_m = -2.455$

See (Pr 80c).

20.  $^{16}\text{C}(\beta^-)^{16}\text{N}$   $Q_m = 8.012$

See  $^{16}\text{C}$ .

21.  $^{16}\text{O}(\mu^-, \nu)^{16}\text{N}$   $Q_m = 95.241$

Partial  $\mu^-$ -capture rates leading to  $^{16}\text{N}^*(0.12, 0.40)$  [ $J^\pi = 0^-, 1^-$ ] are consistent with the assumption of a large mesonic exchange effect in the time part of the weak axial current (Gu 79). See also (Gi 81c), (Gu 78c; theor.) and p. 8.

22.  $^{16}\text{O}(\pi^-, \gamma)^{16}\text{N}$   $Q_m = 129.148$

See (Tr 79h).

23.  $^{16}\text{O}(\text{n}, \text{p})^{16}\text{N}$   $Q_m = -9.636$

See (Ne 78, Ne 78b;  $E_n = 60$  MeV). See also (Aj 77).

24.  $^{16}\text{O}(\text{t}, {}^3\text{He})^{16}\text{N}$   $Q_m = -10.400$

At  $E_t = 23.5$  MeV  $^{16}\text{N}^*(0, 0.30)$  [ $J^\pi = 2^-, 3^-$ ] are strongly populated relative to  $^{16}\text{N}^*(0.12, 0.40)$  [ $0^-, 1^-$ ]. This suggests that the  $2^-$  and  $3^-$  states in  $^{16}\text{F}$  are those

#### Footnotes to table 16.8

<sup>a)</sup> (Ze 71, Do 71a). See table 16.7 in (Aj 71) for the earlier work.

<sup>b)</sup> Below  $E_n = 4.5$  MeV, the multilevel  $R$ -matrix formalism was used to determine  $E_\lambda$ ,  $\Gamma_\lambda$  and whenever possible  $J^\pi$  by a  $\chi^2$  fitting and minimization technique. Above this energy the  $2J+1$  dependence was used; the parity cannot be determined because no marked interference effects are observed between resonance and potential scattering. Above 5.65 MeV all  $J$ -values are lower limits because the inelastic channel is open. [A channel radius  $a = 4.69$  fm was used.] (Ze 71)

<sup>c)</sup> Parity determined from angular distribution.

<sup>d)</sup>  $J^\pi$  also obtained by phase-shift analysis.

<sup>e)</sup> The phase-shift analysis indicates that the resonance is at  $E_n = 2.42 \pm 0.08$  MeV with  $\Gamma = 250 \pm 50$  keV. This is one of two ( $d_{3/2}p_{1/2}^{-1}$ ) single-particle resonances (Ze 71).

<sup>f)</sup> The phase-shift analysis finds  $E_\lambda = 2.94 \pm 0.1$  MeV,  $\Gamma = 320 \pm 80$  keV. This is the other ( $d_{3/2}p_{1/2}^{-1}$ ) single-particle resonance (Ze 71).

<sup>g)</sup> See also (Pa 79a).

that are strongly populated in the  $^{16}\text{O}(^3\text{He}, \text{t})^{16}\text{F}$  reaction [ $^{16}\text{F}^*(0.42, 0.72)$ ] and that the other two states in  $^{16}\text{F}$  [ $^{16}\text{F}^*(0, 0.20)$ ] are  $0^-$  and  $1^-$  [the ordering within the  $2^-$  and  $3^-$ , and the  $0^-$  and  $1^-$  states, in  $^{16}\text{F}$  is ambiguous] (Fl 74a). See also  $^{16}\text{O}(^3\text{He}, \text{t})$  in  $^{16}\text{F}$ .

25.  $^{16}\text{O}(^{11}\text{B}, ^{11}\text{C})^{16}\text{N}$  $Q_m = -12.401$ 

At  $E(^{11}\text{B}) = 115$  MeV unresolved groups near  $E_x = 0$  and 6 MeV are relatively strongly populated (Ra 79c).

TABLE 16.9  
Levels of  $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$  and  $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$

$(\text{He 66a})^a$ $E_x$ (MeV $\pm$ keV)	$I_n$ <sup>a,b)</sup>	$(\text{He 66a})^c$	$(\text{Bo 70b})^c$	$J^\pi$ <sup>d)</sup>
0	k)	0		$2^{-1})$
$0.1201 \pm 0.5$ <sup>e)</sup>	k)	$0.119 \pm 15$		$0^-$
$0.2962 \pm 1.0$ <sup>e)</sup>	k)	$0.301 \pm 15$		$3^{-1})$
$0.3973 \pm 1.0$ <sup>e)</sup>	k)	$0.400 \pm 15$		$1^-$
$3.365 \pm 10$		$3.358 \pm 15$		$1^{+1})$
$3.523 \pm 10$	2 or 1+3	$3.524 \pm 15$	f)	$2^{+m})$
$3.964 \pm 10$	3	$3.964 \pm 15$		$3^{+l,m})$
$4.325 \pm 10$	1	$4.324 \pm 15$		$1^{+1})$
$4.40$ <sup>b)</sup>	0	$4.383 \pm 15$		$(0, 1)^-$
$4.715 \pm 10$	1			$(1, 2, 3)^+$
$4.780 \pm 10$		$4.787 \pm 15$	f)	
$(4.90 \pm 10)$				
$5.032 \pm 10$	2	$5.065 \pm 15$		$2^{-m})$
$5.128 \pm 10$	$\geq 2$	$5.139 \pm 15$		$\geq 2$
$5.150 \pm 10$	2			$(2, 3)^-$
$5.231 \pm 10$	3	$5.240 \pm 15$		$3^{+m})$
$5.310 \pm 10$				
$5.523 \pm 10$	3	$5.528 \pm 15$	f)	$3^{+m})$
$5.739 \pm 10$	2	$5.740 \pm 15$	f)	$(1, 2)^{-n})$
			$6.01 \pm 15$ <sup>i)</sup>	
$6.170 \pm 10$	$\geq 3$	$6.168 \pm 15$	<sup>a)</sup>	$4^{-1})$
$(6.28 \pm 10)$	1			$(0, 1, 2)^+$
$6.376 \pm 10$	2		$6.37 \pm 15$ <sup>j)</sup>	$(1, 2, 3)^-$
$6.431 \pm 10$				
$6.514 \pm 10$	1	$6.512 \pm 15$	<sup>a)</sup>	$(0, 1, 2)^+$
$6.609 \pm 10$		$6.620 \pm 15$	<sup>a)</sup>	
$(6.79 \pm 10)$				
$6.847 \pm 10$		$6.852 \pm 15$	<sup>a)</sup>	
$7.034 \pm 10$			$7.01 \pm 15$ <sup>j)</sup>	
$7.135 \pm 10$		$7.141 \pm 15$	<sup>a)</sup>	
$7.250 \pm 10$		$7.247 \pm 15$	<sup>a)</sup>	
$7.577 \pm 10$		$7.596 \pm 15$	<sup>a)</sup>	
$7.638 \pm 10$			$7.64 \pm 15$ <sup>j)</sup>	
$7.676 \pm 10$		$7.683 \pm 15$		

TABLE 16.9—continued

(He 66a) <sup>a)</sup> <i>E<sub>x</sub></i> (MeV ± keV)	<i>I<sub>n</sub></i> <sup>a,b)</sup>	(He 66a) <sup>c)</sup>	(Bo 70b) <sup>c)</sup>	<i>J"</i> <sup>d)</sup>
7.840 ± 10		8.286 ± 15 8.374 ± 15	7.88 ± 15 <sup>j)</sup> 8.06 ± 15 <sup>j)</sup> 8.18 ± 15 <sup>j)</sup> 8.49 ± 30 <sup>h)</sup> 8.819 ± 15 <sup>i)</sup> 9.035 ± 15 (9.16 ± 30) (9.34 ± 30) 9.459 ± 15 (9.66 ± 40) 9.794 ± 15 <sup>i)</sup> 9.90 ± 30 10.055 ± 15 <sup>i)</sup> (10.17 ± 30) (10.26 ± 30)	

<sup>a)</sup>  $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$ .<sup>b)</sup> (Fu 72c, Fu 71a;  $E_{\text{d}} = 12$  MeV). I am very much indebted to H. Fuchs for his comments.<sup>c)</sup>  $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$ .<sup>d)</sup>  $J''$  assignment from angular distribution analyses and  $\gamma$ -decay; see (Aj 77) for references.<sup>e)</sup> From  $\gamma$ -decay studies (Gi 63e).<sup>f)</sup> Angular distribution reported in  $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$  at  $E_{\text{d}} = 10.0 \rightarrow 11.2$  MeV but  $L$  not determined (Bo 70b).<sup>g)</sup> Alpha group seen but  $E_{\text{x}}$  not determined.<sup>h)</sup>  $\Gamma$  for this level and the ones listed below  $\leq 40-50$  keV (Bo 70b).<sup>i)</sup> These levels appear to be correlated with thresholds for neutron emission to excited states of  $^{15}\text{N}$  (Bo 70b, Bo 70c).<sup>j)</sup> T. I. Bonner, private communication.<sup>k)</sup> Absolute spectroscopic factors for  $^{16}\text{N}^*(0, 0.12, 0.30, 0.40)$  are 0.55, 0.46, 0.54, 0.52 (Bo 73o; average values in the range  $E_{\text{d}} = 5$  to 6 MeV).<sup>l)</sup> Polarization measurements at  $E_{\text{d}} = 52$  MeV are consistent with  $J'' = 2^-, 3^-, 3^+$  and  $4^-$  for  $^{16}\text{N}^*(0, 0.30, 3.96, 6.17)$  and with  $1^+$  for  $^{16}\text{N}^*(3.36, 4.32)$  [admixture of  $L = 0+2$ ] (Ma 81c).<sup>m)</sup> Polarization measurements at  $E_{\text{d}} = 8.5$  to 11.3 MeV are in agreement with the previously assigned  $J''$  values for  $^{16}\text{N}^*(0, 0.12, 0.30, 0.40, 3.36, 4.32, 5.05)$  and lead to the unique  $J''$  values for  $^{16}\text{N}^*(3.52, 3.96, 5.05, 5.23, 5.52)$  shown here (Ba 78r).<sup>n)</sup> A closely spaced doublet appears to be present. At least one of the states has unnatural parity (Ba 78r).26.  $^{17}\text{O}(\text{d}, ^3\text{He})^{16}\text{N}$ 

$$Q_m = -8.287$$

$^3\text{He}$  groups observed in this reaction are displayed in table 16.10 (Ma 78p). Polarization data is consistent with  $1p_{1/2}$  for  $^{16}\text{N}^*(0, 0.30)$ , with perhaps some small  $1p_{3/2}$  admixture for  $^{16}\text{N}_{\text{g.s.}}$ . There is no significant deviation for a pure  $1p_{3/2}$  VAP for the groups to  $^{16}\text{N}^*(6.17, 6.36)$  [ $J'' = 4^-, 3^-$ ] (Ma 81d). Comparisons of the  $^{16}\text{N}$

states with  $T = 1$  states in  $^{16}\text{O}$  are discussed by (Wa 771, Ma 78p): see reaction 81 in  $^{16}\text{O}$  and table 16.20. See also (Aj 77).



At  $E_p = 43.7$  MeV, the angular distribution of the  $^3\text{He}$  nuclei corresponding to a state at  $E_x = 9.9$  MeV fixes  $L = 0$  and therefore  $J'' = 0^+$  for  $^{16}\text{N}^*(9.9)$ : it is presumably the  $T = 2$  analog of the ground state of  $^{16}\text{C}$ . Some lower-lying  $T = 1$  states were also observed (Ce 64b). See also (Fo 78g).

TABLE 16.10  
States of  $^{16}\text{N}$  from  $^{17}\text{O}(\text{d}, ^3\text{He})^*$ )

$E_x$ (MeV) <sup>b</sup>	$l$	$C^2 S$	$J''$
0	1	0.94	$2^-$
=0.30	1	1.33	$3^-$
3.35	2	0.02	$1^+$
3.52	2	0.06	$(2)^+$
=3.96	0	0.10	$(3)^+$
5.14 <sup>c</sup> )	$1^+(2)$	$0.2 + 0.03$	$3^- + \geq 2$
5.53	0	0.08	$(2, 3)^+$
5.74	1	0.40	$(1, 2, 3, 4)^-$
6.17	1	1.20	$(4)^- \text{d})$
6.36	1	0.80	$3^-$
7.66	1	0.30	$(2, 4)^- \text{d})$
9.48	1	0.25	$(1, 2, 3, 4)^-$

<sup>a</sup>) (Ma 78p;  $E_d = 52$  MeV.)

<sup>b</sup>) Resolution of  $^3\text{He}$  groups was 120 keV FWHM.

<sup>c</sup>) Unresolved doublet; angular distribution dominated by  $l = 1$  proton pickup.

<sup>d</sup>) Based on analog relation with the  $T = 1$  states in  $^{16}\text{O}$ : see reaction 81 in  $^{16}\text{O}$  and table 16.20.



Alpha-particle groups observed in this reaction are displayed in table 16.9. Polarization measurements are reported at  $E_\alpha = 8.5$  to 11.3 MeV (Ba 78r) and at 52 MeV (Ma 81c): see footnotes j), k) and l) to table 16.9.  $^{16}\text{N}^*(8.82, 9.8, 10.06)$  may be related to nearly bound virtual states of a  $2s_{1/2}$  neutron with  $^{15}\text{N}^*(6.32, 7.30, 7.57)$  (Bo 70c, Bo 70b). (As 75a) find  $\tau_m = 6.5 \pm 0.5$  psec for  $^{16}\text{N}^*(0.40)$  and  $|g| = 1.83 \pm 0.13$ ;  $|M|^2$  for the M1 transition to  $^{16}\text{N}^*(0.12)$  is  $0.17 \pm 0.02$  W.u. [Based on  $\tau_m = 5.6 \pm 0.3$  psec,  $|g| = 1.5 \pm 0.1$ .]



Angular distributions have been reported for  $E_n = 4.7$  to 14.4 MeV [see (Aj 71, Aj 77)] and at  $E_n = 13.9$  and 15.6 MeV (Ro 77i;  $\alpha_{0+1+2+3}$ ). See also  $^{20}\text{F}$  in (Aj 83).

<sup>16</sup>O

(Figs. 3 and 5)

GENERAL<sup>†</sup>

*Shell model:* (Ap 76a, Be 76v, Na 76p, Ap 77, Br 77q, Ca 77, Ca 77c, Ca 77u, Go 77p, He 77i, He 77k, Li 77l, Sa 77dd, Wa 77k, Zh 77, Ap 78, Ba 78aa, Ch 78z, Da 78n, He 78a, Ho 78f, Ay 79, Ha 79v, Jo 79i, Ka 79a, Pe 79b, Ro 79c, Sa 79u, St 79q, Va 79a, An 80k, Gi 80a, Ha 80p, Mc 80a, Pe 80j, Wa 80j, Ka 81e).

*Collective, deformed and rotational models:* (Ho 75m, Ca 77c, Vo 77b, Ra 78i, Ma 79f, Vl 79, Ba 80o, Be 80t, Fu 80i, Ku 80d, Ok 80d, Pe 80j, Ka 81e).

*Cluster and  $\alpha$ -particle models:* (Fa 75i, Ho 75m, Ap 76, Ap 76a, Ap 76b, Fl 76b, Ho 76j, Su 76e, Ba 77rr, Be 77n, He 77i, Ho 77l, Ho 77m, Sa 77v, Sa 77cc, Su 77f, Ap 78, Ch 78z, He 78a, Is 78a, Kh 78b, Ad 79, Go 79t, Gr 79q, In 79b, Ro 79p, Wi 79o, Za 79b, Ag 80, Du 80b, Fu 80h, Fu 80i, Gu 80b, Ik 80a, Su 80d, El 81c, Em 81, Em 81a, Ma 81g, Wi 81b).

*Special states:* (Fa 75i, Br 75s, Si 75f, Ap 76, Ap 76a, Ap 76b, Ba 76cc, Bo 76z, Fa 76f, He 76k, Li 76s, St 76l, Su 76e, Ap 77, Bl 77b, Bu 77s, Ca 77c, Fa 77a, Go 77p, Gr 77k, He 77h, Ka 77ee, Kn 77, Kr 77, Mo 77v, Pe 77i, Su 77f, Su 77g, Wa 77k, Al 78f, Ap 78, Ba 78aa, Ba 78f, Be 78cc, Ch 78b, Co 78p, Da 78n, De 78n, En 78c, Ha 78w, Ka 78w, Mc 78d, Ra 78i, Sa 78n, Sc 78i, Sh 78d, Ab 79a, Ch 79j, Gm 79, In 79d, Ka 79n, Ki 79k, Ku 79l, Pe 79b, Pr 79c, Ro 79c, Ro 79l, Ro 79p, Sh 79t, Vl 79, Be 80t, Br 80n, Ca 80e, Fu 80i, Go 80n, Ha 80p, Hi 80c, Ka 80n, Me 80b, Ok 80d, Ov 80, Sh 80n, Sp 80c, Su 80d, Ta 80p, Wa 80e, Wa 80j, Ch 81, Cu 81f, Em 81a, Ka 81e, Ro 81l, Sp 81a, Wi 81b).

*Electromagnetic transitions:* (Ba 76cc, He 76k, Ho 76h, St 76l, Su 76e, An 77a, Be 77n, Bi 77d, Bl 77b, Br 77, De 77d, Do 77a, Go 77, Gr 77k, He 77l, Ko 77g, Kr 77, Pe 77i, Wa 77k, Ad 78b, Ba 78aa, Sc 78i, Tr 78d, Ka 79n, Ku 79l, Mo 79g, Mo 79p, Ar 80d, Du 80b, Mu 80, Pe 80j, To 80e, Sp 81a).

*Giant resonances<sup>††</sup>:* (Ge 75o, Kr 75f, Kr 75g, Am 76b, Be 76bb, Mu 76b, Ab 77a, Br 77, De 77d, Gm 77, Go 77, He 77i, Kn 77, Kr 77, Li 77g, Sa 77ee, Ad 78b, He 78a, Ro 78j, St 78j, Yu 78a, Bl 79c, Bo 79g, Do 79o, Fa 79b, Gm 79, Go 79c, Ha 79k, Im 79b, Iz 79, Ki 79e, Kn 79j, Ko 79o, Kr 79a, Sh 79t, Wa 79c, Ar 80d, Ba 80o, Bo 80l, Go 80f, Mu 80, Sh 80n, Sp 80d, Kh 81b, Li 81d).

*Astrophysical questions:* (Bo 76cc, No 76l, Qu 76a, Si 76g, Va 76m, Ar 77a, Au 77f, Au 77h, Au 77i, Au 77j, Bu 77l, Ca 77n, Ca 77q, Cl 77i, En 77a, Fr 77g, Ib 77a,

<sup>†</sup> See also (Aj 77).<sup>††</sup> See also reactions 54 and 58.




Fig. 3. Energy levels of  $^{16}\text{O}$ . For notation see fig. 2.

Jo 77c, Ki 77, Pa 77c, Pr 77h, Si 77e, St 77e, To 77g, Vo 77, Wa 77t, Bu 78g, Bu 78k, By 78a, Cl 78g, Di 78d, Fe 78e, Gl 78b, Ib 78, Ib 78a, Ka 78s, La 78a, Le 78t, Lu 78, Mc 78a, Me 78f, Or 78, Pe 78i, Po 78b, St 78i, To 78d, Tr 78, Wo 78f, Ch 79i, De 79n, Gl 79a, Ka 79c, La 79c, La 79f, Le 79m, Ma 79c, Mc 79b, Pe 79e, Ra 79h, Sa 79c, Sw 79, Ti 79a, We 79, Wi 79m, Ca 80k, Cl 80c, Co 80s, Ka 80p, Mc 80d, Me 80, Mo 80n, Re 80, Du 81c, Ga 81b, Gu 81b, Wi 81c, Wi 81i).

*Applications:* (Ch 76r, Ch 76u, Le 76q, Ep 77a, Gr 77j, Lo 77h, Ma 77ww, Mo 77a, Th 77c, Em 78, He 78d, Tr 78h, Gr 79c, Ja 79h, Le 79e, Sp 79a, He 80d, La 80i, Mc 80f).

*Special reactions involving <sup>16</sup>O:* (Ch 76n, Da 76p, Ho 76f, Ja 76q, Le 76t, Ar 77b, Ce 77, Ch 77e, Co 77m, Fe 77e, Ga 77l, Ha 77h, Ha 77aa, Ho 77o, Ja 77, Ka 77cc, Li 77k, Ma 77e, Ma 77f, Mo 77p, Na 77, Ob 77, Pr 77a, Sh 77o, St 77e, To 77j, Tr 77g, Va 77, Ar 78l, Ba 78p, Bh 78b, Bi 78a, Co 78e, Fr 78j, Go 78n, Gy 78, He 78b, He 78c, He 78m, Ja 78o, Ko 78a, Ku 78e, Ku 78h, Ob 78a, Vo 78, Vo 78a, Ba 79mm, Be 79u, Bl 79f, De 79f, Dy 79, Ga 79, Ge 79b, Go 79j, He 79i, Ko 79s, Mc 79d, Me 79q, Mo 79v, Po 79a, Sa 79l, Sa 79m, Sc 79c, St 79j, St 79m, Va 79h, Ak 80b, Ba 80a, Ce 80a, Gr 80d, Ko 80e, Mi 80a, Ol 80, Ov 80, Sa 80i, Sc 80f, Bh 81, Bo 81k, Ci 81, Eg 81b, Ma 81g, Si 81, Ta 81, Tr 81a, Uc 81).

*Muon and neutrino capture and reactions:* (Ko 74v, Da 76q, Ba 77ww, Bo 77o, Er 77d, Gr 77m, La 77a, Mu 77b, Wa 77o, Ba 78ff, Gu 78c, Gu 78f, It 78, Pa 78h, Ro 78l, Bu 79e, Do 79o, Er 79c, Fr 79g, Gm 79a, Gu 79, Gu 79c, Ho 79d, Ki 79m, Ko 79bb, Me 79r, Mu 79b, Rh 79, St 79l, Su 79d, Ve 79a, Wu 79d, Ba 80aa, Br 80, Ch 80b, Ch 80m, Ch 80n, Er 80b, Gr 80, Ko 80s, Su 80c, Ch 81, Ei 81, Er 81a, Gi 81c, To 81e, Va 81e, Va 81g).

*Pion capture and reactions:* (Bu 76i, De 76l, Li 76r, Li 76t, Ro 76r, Al 77f, Al 77aa, Ba 77m, Ba 77kk, Ba 77xx, Ba 77ab, Ba 77aj, Be 77gg, Be 77hh, Bo 77d, Bo 77s, Br 77a, Bu 77c, Ch 77q, Co 77v, Do 77a, Er 77d, Gi 77a, Gi 77c, Hi 77f, Ho 77f, Ho 77p, Ka 77c, Ko 77cc, Le 77p, Ma 77h, Ma 77gg, Ma 77ww, Mc 77, Mi 77c, Mi 77e, Na 77a, Na 77d, Na 77r, Pr 77g, Ro 77u, Sa 77dd, Sc 77d, Si 77, Sm 77c, Sp 77d, St 77o, Su 77h, Vi 77a, Wa 77p, Wa 77r, Al 78a, Ba 78l, Ba 78w, Be 78a, Be 78p, Be 78z, Bo 78, Bo 78r, Bo 78s, Br 78a, Br 78c, Er 78b, Er 78c, Fu 78c, Fu 78g, Ga 78, Gi 78b, Gi 78c, Gr 78g, In 78, Ja 78f, Ja 78j, Ja 78k, Ko 78n, Kw 78, Li 78d, Ma 78g, Me 78j, Mo 78b, Na 78, Na 78i, Oh 78, Ro 78h, Sc 78b, Sh 78e, Sh 78g, Sh 78j, Sp 78, We 78d, We 78h, Yo 78c, Ab 79c, Al 79k, Am 79e, An 79d, An 79l, Ar 79k, Ba 79i, Bl 79e, Bo 79j, Bo 79k, Bo 79r, Bo 79ee, Br 79h, Bu 79, Ch 79a, De 79k, Di 79b, Do 79o, Dy 79a, Ep 79a, Er 79c, Gi 79c, Hi 79c, Hu 79b, Hu 79c, In 79c, Ja 79f, Ja 79j, Jo 79g, Kl 79e, Kl 79f, Kn 79, Ko 79p, Li 79a, Li 79g, Ma 79j, Ma 79q, Na 79c, Na 79q, Oh 79, Os 79c, Os 79i, Pi 79f, Re 79a, Ro 79e, Sc 79b, Sh 79g, Sh 79j, Sr 79a, St 79, Tr 79d, Tr 79h, Ul 79, Wa 79b, Wa 79g, Wi 79d, Wi 79e, Al 80e, Al 80g, Ar 80, As 80, Ba 80f, Ba 80y, Be 80q, Be 80s,

Be 80u, Bh 80, Bh 80b, Br 80a, Bu 80b, Ch 80n, Co 80a, Cr 80a, De 80m, De 80aa, Er 80b, Fr 80h, Gr 80, Gr 80b, Gr 80f, Ho 80k, Ho 80l, Je 80c, Ka 80h, La 80a, Le 80c, Li 80j, Li 80k, Li 80n, Mi 80f, Na 80a, Na 80i, Ob 80a, Sc 80, Sc 80j, Se 80e, Si 80c, Sp 80, St 80d, St 80g, Th 80, Th 80a, Tr 80h, Wu 80, As 81a, Ba 81l, Bo 81, Bu 81e, Cu 81b, Do 81, Do 81a, Fr 81e, Fr 81f, Gi 81b, Gr 81c, Io 81, Ki 81a, Le 81e, Li 81b, Li 81h, Ma 81h, Mo 81e, Mo 81f, Os 81a, Sa 81g, Th 81b, Th 81d, We 81e, Wh 81, Wh 81b).

*K-mesons and other meson interactions:* (Bo 76y, Br 76q, Da 76o, Ba 77aj, Bo 77u, Ju 77c, Ki 77j, La 77j, Po 77e, Th 77f, At 78, Co 78o, Gr 78g, Po 78a, Sc 78b, So 78d, Ba 79hh, Bo 79k, Ch 79u, Ga 79e, Ga 79k, Gi 79, Ki 79g, Po 79, Ra 79f, Ba 80v, Do 80g, Ga 80, Po 80, Be 81u, Bo 81j, Hu 81a, Hu 81b).

*Antiproton interactions:* (Ba 77yy, We 77g, Gr 78g, Po 78, Au 81).

*Other topics:* (Er 75b, Go 75cc, Bo 76y, Bo 76z, Da 76o, Fu 76g, Go 76k, Kn 76g, Li 76s, Lo 76g, Na 76p, Pr 76a, St 76g, Ak 77, Au 77d, Bo 77u, Br 77k, Bu 77s, Ca 77, Ca 77u, De 77f, Go 77, Go 77p, Ho 77b, Ho 77h, Je 77a, Ka 77ff, Ki 77h, Ki 77j, Kr 77, Li 77e, Li 77l, Mo 77v, Ne 77c, Os 77b, Po 77e, Pr 77, Sa 77aa, Sm 77, So 77a, Su 77g, Tr 77a, Va 77d, Wa 77k, Za 77, An 78f, Au 78b, Ba 78f, Be 78v, Bi 78l, Bi 78n, Br 78m, Da 78s, De 78h, De 78n, En 78c, Fa 78d, Ga 78d, Ha 78o, Ha 78w, Ho 78f, Ka 78r, Ka 78w, Kh 78a, Ko 78h, Kw 78, Le 78c, Mc 78d, Mi 78e, Mu 78e, On 78, Po 78a, Ro 78p, Sc 78i, Sh 78d, Sh 78q, Si 78b, Si 78d, So 78c, So 78d, St 78j, Ul 78, Ab 79a, Ay 79, Be 79k, Bi 79b, Bo 79d, Bo 79g, Bo 79y, Br 79j, Ch 79u, De 79o, Du 79a, Fi 79, Ga 79e, Ga 79k, Go 79c, Go 79t, Go 79u, Gu 79g, Ha 79v, He 79, Ja 79m, Ja 79u, Ka 79a, Ka 79d, Ka 79n, Ka 79s, Ki 79g, Ki 79n, Ko 79r, Ko 79gg, Ko 79ll, Le 79g, Mu 79c, No 79c, Pe 79b, Sa 79u, St 79q, Ta 79h, To 79d, Va 79a, Va 79l, Za 79a, Za 79b, An 80k, Bo 80f, Bo 80n, Br 80n, Ca 80e, Da 80b, De 80, Dw 80, Fa 80b, Fa 80e, Gi 80a, Go 80n, He 80o, Hi 80c, Hy 80a, Ja 80e, Ka 80m, Me 80c, Mu 80, Mu 80g, Ng 80, Po 80, Qu 80, Sh 80n, Ta 80p, To 80e, Va 80b, Zo 80, Bl 81, Bo 81r, Br 81d, Cu 81f, Da 81c, Du 81b, Li 81d, Ro 81l, Sh 81h, Za 81a).

*Ground state of <sup>16</sup>O:* (Fu 76h, Ga 76k, Gi 76e, Sa 76i, An 77b, An 77f, Br 77, Bu 77s, Dy 77c, Go 77n, Gr 77b, Je 77a, Ma 77gg, No 77d, Pi 77d, Va 77e, Al 78f, Al 78k, An 78b, An 78f, Be 78v, Be 78cc, Bi 78l, Bi 78m, Br 78j, Ch 78y, Ch 78z, Fa 78d, He 78k, Ka 78r, Ko 78h, Mu 78e, Ne 78a, On 78, Ro 78p, Sh 78q, Si 78d, Sm 78d, So 78c, St 78g, Ul 78, Za 78a, Za 78b, Ay 79, Be 79q, Bi 79b, Br 79i, Br 79j, Du 79a, Fi 79d, Go 79u, Gu 79c, Gu 79g, Ha 79v, In 79d, Ja 79m, Ka 79n, Ko 79ll, Ma 79ll, Sa 79l, Sa 79u, St 79q, Va 79l, An 80l, Ar 80d, Bo 80n, Br 80i, De 80m, Fa 80b, Gi 80a, Go 80f, Hi 80c, Ho 80d, Hy 80a, Me 80b, My 80, Va 80a, Wa 80e, Ar 81, At 81, Br 81d, Du 81a, Si 81b, Za 81a).

$$\langle r^2 \rangle^{1/2} = 2.710 \pm 0.015 \text{ fm (Ki 78a). See also (Ba 80aa).}$$

$$g = +0.55 \pm 0.03 \text{ for } ^{16}\text{O}^*(6.13) (\text{Br 73m, Ra 73c, Ka 77}).$$

**<sup>16</sup>O MASTER TABLE**

 ENERGY LEVELS OF LIGHT NUCLEI  $A = 16 - 17$ 

25

 TABLE 16.11  
 Energy levels of <sup>16</sup>O<sup>a)</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\Gamma_{c.m.}$ or $\tau_m$ (keV)	Decay	Reactions
0	$0^+; 0$		stable		
$6.0494 \pm 1.0$	$0^+; 0$	$0^+$	$\tau_m = 96 \pm 7$ psec	$\pi$	2, 3, 9 $\rightarrow$ 18, 25 $\rightarrow$ 28, 30, 31, 38, 40, 42, 48 $\rightarrow$ 98
$6.13043 \pm 0.05$	$3^-; 0$		$\tau_m = 26.6 \pm 0.7$ psec $ g  = 0.55 \pm 0.03$	$\gamma$	2, 3, 9, 10, 14, 18, 26, 38, 39, 40, 48, 49, 52, 57, 59, 60, 61, 63, 64, 68, 80, 81, 82, 84, 88, 89
$6.9171 \pm 0.6$	$2^+; 0$	$0^+$	$\tau_m = 6.6 \pm 0.4$ fsec	$\gamma$	2, 3, 9, 10, 14, 38, 39, 40, 48, 49, 56, 57, 60, 61, 63, 64, 81, 82, 84, 85, 88, 89
$7.11685 \pm 0.14$	$1^-; 0$		$\tau_m = 11.6 \pm 1.0$ fsec	$\gamma$	2, 3, 9, 10, 38, 39, 40, 48, 49, 52, 56, 57, 60, 61, 64, 80, 81, 82, 84, 85, 88, 89
$8.8719 \pm 0.5$	$2^-; 0$		$\tau_m = 180 \pm 16$ fsec	$\gamma, \alpha$	2, 3, 7, 9, 10, 11, 17, 26, 38, 39, 42, 48, 49, 52, 60, 61, 63, 64, 80, 81, 82, 84, 88, 89, 92, 96
$9.632 \pm 21$	$1^-; 0$	$0^-$	$\Gamma_{c.m.} = 400 \pm 10$	$\gamma, \alpha$	3, 7, 9, 10, 38, 49, 52, 64
$9.847 \pm 3$	$2^+; 0$		$0.625 \pm 0.100$	$\gamma, \alpha$	2, 3, 7, 9, 10, 11, 26, 38, 39, 42, 48, 49, 52, 57, 60, 61, 63, 64, 82, 84, 88, 92, 96
$10.355 \pm 3$	$4^+; 0$	$0^+$	$25 \pm 4$	$\gamma, \alpha$	2, 3, 7, 9, 10, 11, 13, 16, 17, 26, 27, 38, 39, 49, 57, 60, 61, 63, 64, 80, 82, 84, 85, 88, 92, 96
$10.957 \pm 1$	$0^-; 0$		$\tau_m = 8 \pm 5$ fsec		2, 38, 48, 49, 60, 61, 82
$11.080 \pm 3$	$3^+; 0$		$\Gamma < 12$	$\gamma$	2, 38, 48, 49, 82, 88
$11.096 \pm 2$	$4^+; 0$		$0.28 \pm 0.05$	$\gamma, \alpha$	2, 3, 7, 9, 11, 13, 17, 26, 38, 39, 60, 61, 63, 64
(11.26) <sup>b)</sup>	( $0^+; 0$ )		(2500)	( $\alpha$ )	7, 49, 82, 84
$11.520 \pm 4$	$2^+; 0$		$74 \pm 4$	$\gamma, \alpha$	2, 3, 7, 38, 57, 60, 61, 63, 64
$11.60 \pm 20$	$3^-; 0$	$0^-$	$800 \pm 100$	$\alpha$	7, 10
$12.049 \pm 2$	$0^+; 0$		$1.5 \pm 0.5$	$\gamma, \alpha$	7, 38, 57, 60, 61, 63, 64, 84
$12.438 \pm 3$	$1^-; 0$		$90 \pm 10$	$\gamma, p, \alpha$	3, 5, 7, 38, 43, 44, 47, 48, 49, 60, 61, 64
$12.530 \pm 1$	$2^-; 0$		0.8	$\gamma, p, \alpha$	2, 38, 43, 44, 47, 48, 49, 57, 81
$12.797 \pm 4$	$0^-; 1$		$38 \pm 4$	$\gamma, p$	38, 44, 48, 49
$12.9685 \pm 0.4$	$2^-; 1$		$1.9 \pm 0.2$	$\gamma, p, \alpha$	38, 43, 44, 47, 48, 49, 57, 80, 81, 82

TABLE 16.11—continued

$E_x$ (MeV ± keV)	$J^\pi; T$	$K^\pi$	$\Gamma_{c.m.}$ (keV)	Decay	Reactions
13.02 ± 10	2 <sup>+</sup> ; 0		150 ± 10	$\gamma, p, \alpha$	3, 7, 44, 47, 57, 60, 61, 63, 64
13.090 ± 5	1 <sup>-</sup> ; 1		130 ± 5	$\gamma, p, \alpha$	3, 5, 7, 9, 38, 43, 44, 47, 48, 57, 82
13.120 ± 10	3 <sup>-</sup> ; 0		130 ± 30	$\gamma, p, \alpha$	2, 3, 5, 7, 38, 48
13.258 ± 2	3 <sup>-</sup> ; 1		21 ± 1	$\gamma, p, \alpha$	3, 5, 7, 38, 44, 47, 48, 49, 60, 80, 81, 82, 85
13.664 ± 3	1 <sup>+</sup> ; 0		63 ± 3	$\gamma, p, \alpha$	38, 43, 44, 47, 61
13.869 ± 2	4 <sup>+</sup> ; 0		84 ± 2	$p, \alpha$	2, 7, 38, 44, 47, 60, 63
13.980 ± 2	2 <sup>-</sup>		20 ± 2	$p, \alpha$	2, 38, 39, 44, 47
14.032 ± 15	0 <sup>+</sup>		200 ± 50	$\alpha$	7
14.1 ± 100	3 <sup>-</sup>		750 ± 200	$\alpha$	7
14.302 ± 3			34 ± 12		26, 38, 39
14.399 ± 2	≥ 5		27 ± 5		2, 10, 26, 38, 39
14.620 ± 11	(4 <sup>+</sup> )		490 ± 20	$\alpha$	7, 9
14.660 ± 11	5 <sup>-</sup>		650 ± 50	$\alpha$	7, 9, 10, 11, 13
14.816 ± 2	6 <sup>+</sup> ; 0		70 ± 8	$\alpha$	2, 3, 7, 9, 26, 38, 39, 64
14.922 ± 6	2 <sup>+</sup>		65 ± 5	$p, \alpha$	2, 37, 38, 44, 47
15.100 ± 3	0 <sup>+</sup>		166 ± 30	$p, \alpha$	5, 7, 38
15.196 ± 3	2 <sup>-</sup>		58 ± 10	$p, \alpha$	38, 39, 44, 47, 81
15.26 ± 20.	2 <sup>+</sup> ; (0)		300 ± 100	$p, \alpha$	44, 47, 60, 63
15.408 ± 2	3 <sup>-</sup> ; 0		133 ± 7	$p, \alpha$	5, 7, 37, 38, 39, 44, 47, 60, 64, 80, 81
15.785 ± 5	(3 <sup>+</sup> )		40 ± 10		38, 39
15.828 ± 30	3 <sup>-</sup>		700 ± 120	$\alpha$	7
(15.9)	(2 <sup>+</sup> )		~600	$\gamma, \alpha$	3
16.209 ± 2	(4 <sup>+</sup> )		40 ± 10		38, 39
16.22 ± 20	1 <sup>+</sup> ; 1		18 ± 3	$\gamma, n, p$	37, 44, 45, 57
16.275 ± 7	6 <sup>+</sup> ; 0	0 <sup>+</sup>	420 ± 20	$\alpha$	2, 7, 9, 10, 11, 13, 27, 39
16.350 ± 13	(0 <sup>+</sup> , 1 <sup>-</sup> )		65 ± 45	$p, \alpha$	5, 7, 38, 84
16.442 ± 2	2 <sup>+</sup> ; (1)		22 ± 3	$\gamma, n, p, \alpha$	3, 4, 5, 7, 38, 44, 47, 57, 60, 63, 64
16.817 ± 2	3 <sup>+</sup>		70 ± 10	$\gamma, p, \alpha$	38, 44, 47, 57
16.844 ± 21	4 <sup>+</sup>		570 ± 60	$\alpha$	7
16.93 ± 50	2 <sup>+</sup>		~280	$\alpha$	7, 8, 60
17.0	1 <sup>-</sup> ; 1		~1500	$\gamma, p$	43
17.129 ± 5	2 <sup>+</sup>		107 ± 14	$n, p, \alpha$	4, 5, 7
17.14 ± 20	1 <sup>-</sup> ; 1		36 ± 5	$\gamma, n, p, \alpha$	7, 39, 43, 44, 45, 49, 57
17.20 ± 20	2 <sup>+</sup>		160 ± 60	$\alpha$	2, 7, 8, 49, 60, 63, 64
17.27 ± 20	1 <sup>-</sup> ; 1		90 ± 10	$\gamma, n, p, \alpha$	4, 43, 44, 45
17.510 ± 26	1 <sup>-</sup>		180 ± 60	$\alpha$	7
17.555 ± 21	(6 <sup>+</sup> )		180 ± 70	$n, \alpha$	4, 7
17.618 ± 20	(0 <sup>+</sup> , 1 <sup>-</sup> )		175 ± 60	$p, \alpha$	5, 7
17.72	(0 <sup>+</sup> , 2 <sup>+</sup> )		~75	(p), $\alpha$	5, 8
17.784 ± 15	4 <sup>+</sup>		400 ± 40	$n, \alpha$	4, 7, 8
17.788 ± 16	4 <sup>-</sup> ; 0		150 ± 60		57, 58, 60, 63, 64, 81, 82
18.016 ± 1	4 <sup>+</sup> ; (0)		14 ± 2	(n), $p, \alpha$	4, 5, 7, 8
18.033 ± 10	3 <sup>+</sup> ; 1		26 ± 5	$\gamma, n, p$	43, 44, 45, 81, 82
18.11 ± 30	(0, 2) <sup>+</sup> ; 0		300 ± 50	(γ), $n, p, \alpha$	3, 5, 7, 45, 60, 64

TABLE 16.11—continued

$E_x$ (MeV ± keV)	$J^\pi; T$	$K^\pi$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
18.29			~300	$\gamma, p, \alpha$	3, 5, 7
18.404 ± 12	5 <sup>-</sup>		550 ± 40	$\alpha$	7
18.46 ± 25	2 <sup>+</sup> ; 0		60 ± 10	n, p	45, 57, 60, 63, 64
18.6	(1 <sup>-</sup> , 5 <sup>-</sup> )		~150	$\alpha$	7
18.6	(4) <sup>+</sup>		~300	$\alpha$	7, 8
18.69 ± 30			260 ± 30	n, p	2, 45, 60, 64
18.773 ± 22	1 <sup>-</sup>		215 ± 45	p, $\alpha$	5, 7
18.785 ± 6	4 <sup>+</sup>		260 ± 20	n, p, $\alpha$	4, 5, 7, 8
18.975 ± 10	4 <sup>-</sup> ; 1		≤ 40	p	44, 57, 58, 60, 63, 81, 82
18.99 ± 30	1 <sup>+</sup>		~250	$\gamma, p$	43
19.0	(5 <sup>-</sup> )		~550	$\alpha$	7
19.08 ± 30	2 <sup>+</sup> ; (1)		~120	$\gamma, p$	43, 44
19.206 ± 12	3 <sup>-</sup> ; 1		68 ± 10		81, 82
19.25 ± 20	(2 <sup>-</sup> ; 1)		90 ± 10	n, p, ( $\alpha$ )	7, 45
19.257 ± 9	2 <sup>+</sup>		155 ± 25	$\gamma, p, \alpha$	5, 7, 43
19.319 ± 14	6 <sup>+</sup>		65 ± 35	n, $\alpha$	5, 7, 8
19.375 ± 2	4 <sup>+</sup>		23 ± 4	p, $\alpha$	5, 7
19.48 ± 25	1 <sup>-</sup> ; 1		250 ± 50	$\gamma, n, p$	43, 44, 45, 57
19.53 ± 30	2 <sup>+</sup> ; 0		255 ± 75	n, p, $\alpha$	2, 4, 7, 45, 60, 64
19.754 ± 16	2 <sup>+</sup>		290 ± 50	p, $\alpha$	5, 7
19.802 ± 16	4 <sup>-</sup> ; 0		36 ± 5		57, 58, 81, 82
19.90 ± 20	3; 0		100 ± 30	$\gamma, n, p, \alpha$	2, 7, 43, 44, 45, 58, 60
20.055 ± 13	2 <sup>+</sup> ; 0		350 ± 50	$\gamma, n, (p), \alpha$	3, 4, 5, 7, 64
20.43 ± 30	2 <sup>-</sup> ; 1 (20.5)		190 ± 40 (~300)	$\gamma, n, p$ $\alpha$	43, 44, 45, 57, 81 7
20.541 ± 2	5 <sup>-</sup>		11 ± 2	p, $\alpha$	2, 5, 7
20.560 ± 2			< 5	$\alpha$	7
20.615 ± 3 (20.8)			< 10 (~60)	$\alpha$ n, (p), $\alpha$	7, 27 4, 5
20.857 ± 14	7 <sup>-</sup>	0 <sup>-</sup>	900 ± 100	$\alpha$	7, 9, 10, 11, 13
20.945 ± 20	1 <sup>-</sup> ; 1		300 ± 10	$\gamma, n, p$	3, 43, 44, 45, 57
21.05 ± 50	(2 <sup>+</sup> ; 0)		320 ± 50		60, 64
21.052 ± 6	6 <sup>+</sup>		205 ± 20	$\alpha$	7
21.175 ± 15					2
21.50	(1 → 4)		120	p	44
21.52	7 <sup>-</sup>		61 ± 32	(n), $\alpha$	4, 7
21.648 ± 3	6 <sup>+</sup>		115 ± 8	n, $\alpha$	4, 7, 9
21.776 ± 9	3 <sup>-</sup>		43 ± 20	n, p, $\alpha$	2, 4, 5, 7
22.04			60	n, d, $\alpha$	4, 32
22.150 ± 10	1 <sup>-</sup> ; 1		730 ± 10	$\gamma, n, p, d$	31, 36, 43, 44, 45
22.44 ± 100	(1 <sup>-</sup> ; 1)		300 ± 100	n, p, d, $\alpha$	32, 36, 45, 60
22.5 ± 500	(8) <sup>+</sup>	(0 <sup>+</sup> )			13
22.65 ± 30				n, $\alpha$	2, 4, 8
22.720 ± 5	0 <sup>+</sup> ; T = 2		12.5 ± 2.5	n, p, d, $\alpha$	4, 5, 7, 29, 33, 36, 84
22.89 ± 10	1 <sup>-</sup> ; 0		300 ± 10	$\gamma, p, d$	31, 43, 44
23.0 ± 100	6 <sup>+</sup>		≤ 500	$\alpha$	8, 9
23.11			~20	$\alpha$	7, 8
23.2 ± 80	(1 <sup>-</sup> ; 1)		550 ± 150	n, p	45, 60
23.51 ± 30			300	p, d, $\alpha$	2, 7, 33, 34, 36

TABLE 16.11—continued

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\Gamma_{c.m.}$ (keV)	Decay	Reactions
23.879 $\pm$ 6	6 <sup>+</sup>		26 $\pm$ 4	p, $\alpha$	5, 7, 8, 9
24.07 $\pm$ 30	1 <sup>-</sup> ; 1		550 $\pm$ 50	$\gamma$ , (n), p, <sup>3</sup> He	18, 19, 43, 44, 60
24.35 $\pm$ 70	(2 <sup>+</sup> , 3 <sup>-</sup> ); 0		400 $\pm$ 50	n, p	45, 64
24.522 $\pm$ 11	2 <sup>+</sup> ; 2		< 50		29, 84
24.76 $\pm$ 60	(2, 4) <sup>+</sup> ; 1		340 $\pm$ 60	$\gamma$ , n, p, <sup>3</sup> He, $\alpha$	23, 43, 44, 45
25.12 $\pm$ 50	1 <sup>-</sup> ; 1		3000 $\pm$ 300	$\gamma$ , p, <sup>3</sup> He	18, 43, 44, 47, 63
25.5 $\pm$ 150	1 <sup>-</sup> ; 1		1300 $\pm$ 300		60
25.6	(3 <sup>-</sup> ); 1		450	$\alpha$	7
26.0 $\pm$ 100	1 <sup>-</sup> ; (1)		500–1000	$\gamma$ , <sup>3</sup> He, $\alpha$	18, 23
26.3	2 <sup>+</sup>		1200	$\alpha$	7
26.4 $\pm$ 100	(2, 4) <sup>+</sup> ; 1		550 $\pm$ 100	$\gamma$ , n, p, <sup>3</sup> He, $\alpha$	18, 20, 43, 44, 45, 47
27.0 $\pm$ 100	(T = 1)		broad	<sup>3</sup> He, $\alpha$	23
27.3 $\pm$ 70	(2, 4) <sup>+</sup> ; 1		830 $\pm$ 110	$\gamma$ , p, <sup>3</sup> He, $\alpha$	18, 23, 24, 43, 44, 47
27.5	(3 <sup>-</sup> ; 0)		$\sim$ 2500	$\gamma$ , <sup>3</sup> He	18
28.6 $\pm$ 200				$\gamma$ , <sup>3</sup> He	18
29.7 $\pm$ 100	(T = 1)		470 $\pm$ 150	n, p, d, <sup>3</sup> He, $\alpha$	23, 34, 45
31.8 $\pm$ 600				$\gamma$	56

<sup>a</sup>) See also table 16.12.<sup>b</sup>) I am indebted to Professor H. T. Richards for his remarks concerning the existence of this level.TABLE 16.12  
Radiative decays in <sup>16</sup>O <sup>a</sup>)

$E_t$ (MeV)	$J_t^\pi; T$	$E_t$ (MeV)	$J_t^\pi; T$	Branch (%)	$\Gamma_{rad}$ (eV)
6.05	0 <sup>+</sup> ; 0	0	0 <sup>+</sup> ; 0	100	$3.55 \pm 0.21$ <sup>b</sup> )
6.13	3 <sup>-</sup> ; 0	0	0 <sup>+</sup> ; 0	100	$(2.60 \pm 0.13) \times 10^{-5}$
6.92	2 <sup>+</sup> ; 0	0	0 <sup>+</sup> ; 0	> 99	$0.100 \pm 0.004$ <sup>c</sup> )
		6.05	0 <sup>+</sup> ; 0	$(2.7 \pm 0.3) \times 10^{-2}$	$(2.7 \pm 0.3) \times 10^{-5}$
		6.13	3 <sup>-</sup> ; 0	$\leq 8 \times 10^{-3}$	
7.12	1 <sup>-</sup> ; 0	0	0 <sup>+</sup> ; 0	> 99	$0.057 \pm 0.006$ <sup>d</sup> )
		6.05	0 <sup>+</sup> ; 0	$< 6 \times 10^{-4}$	
		6.13	3 <sup>-</sup> ; 0	$(7.0 \pm 1.4) \times 10^{-2}$	
8.87	2 <sup>-</sup> ; 0	0	0 <sup>+</sup> ; 0	7.2 $\pm$ 0.8	$(2.4 \pm 0.4) \times 10^{-4}$
		6.05	0 <sup>+</sup> ; 0	$0.122 \pm 0.033$	$(2.9 \pm 1.0) \times 10^{-6}$
		6.13 <sup>b</sup> )	3 <sup>-</sup> ; 0	76.0 $\pm$ 3.0	<sup>e</sup> )
		6.92	2 <sup>+</sup> ; 0	4.2 $\pm$ 0.8	$(1.72 \pm 0.25) \times 10^{-4}$
		7.12	1 <sup>-</sup> ; 0	12.6 $\pm$ 2.0	
9.63	1 <sup>-</sup> ; 0	0	0 <sup>+</sup> ; 0	$\sim$ 100	$(23 \pm 3) \times 10^{-3}$
9.85	2 <sup>+</sup> ; 0	0	0 <sup>+</sup> ; 0	61 $\pm$ 4	$(6.0 \pm 0.4) \times 10^{-3}$
		6.05	0 <sup>+</sup> ; 0	18 $\pm$ 4	$(1.9 \pm 0.4) \times 10^{-3}$
		6.92	2 <sup>+</sup> ; 0	21 $\pm$ 4	$(2.2 \pm 0.4) \times 10^{-3}$

TABLE 16.12—continued

$E_i$ (MeV)	$J_i^{\pi}; T$	$E_f$ (MeV)	$J_f^{\pi}; T$	Branch (%)	$\Gamma_{\text{rad}}$ (eV)
10.35	$4^+; 0$	0	$0^+; 0$		$(5.6 \pm 2.0) \times 10^{-8}$
		6.13	$3^-; 0$		$<1.0 \times 10^{-3}$
		6.92	$2^+; 0$	~100	$(5.8 \pm 0.7) \times 10^{-2}$
10.96	$0^-; 0^c$	7.12	$1^-; 0$	>99	$(0.08 \pm 0.05) \times 10^{-3}$
11.10	$4^+; 0$	6.13	$3^-; 0$	"	$(3.1 \pm 1.3) \times 10^{-3}$
		6.92	$2^+; 0$	"	$(2.5 \pm 0.6) \times 10^{-3}$
11.52	$2^+; 0$	0	$0^+; 0$	91.7	$0.61 \pm 0.02$
		6.05	$0^+; 0$	$4.2 \pm 0.7$	$(2.8 \pm 0.6) \times 10^{-2}$
		6.92	$2^+; 0$	$4.0 \pm 1.0$	$(29 \pm 7) \times 10^{-3}$
		7.12	$1^-; 0$	≤0.8	
12.05	$0^+; 0$	0	$0^+; 0$		$4.03 \pm 0.09$
12.44	$1^-; 0$	0	$0^+; 0$		$12 \pm 2$
		6.05	$0^+; 0$	~100	$0.12 \pm 0.04$
12.53	$2^-; 0$	0	$0^+; 0$	$1.2 \pm 0.4$	$(108 \pm 15) \times 10^{-3}$
		6.13	$3^-; 0$	60±6	$2.1 \pm 0.2$
		6.92	$2^+; 0$	<10	≤0.34
		7.12	$1^-; 0$	15±3	$0.5 \pm 0.1$
		8.87	$2^-; 0$	25±3	$0.9 \pm 0.1$
12.80	$0^-; 1$	7.12	$1^-; 0$	~100	$2.5 \pm 0.2$
12.97	$2^-; 1$	0	$0^+; 0$		$(71 \pm 2) \times 10^{-3}$
		6.13	$3^-; 0$	63±6	$2.3 \pm 0.2$
		7.12	$1^-; 0$	12±3	$0.44 \pm 0.10$
		8.87	$2^-; 0$	25±3	$0.90 \pm 0.10$
13.09	$1^-; 1$	0	$0^+; 0$	~100	$32 \pm 5$
		6.05	$0^+; 0$	$0.58 \pm 0.12$	
		7.12	$1^-; 0$	$3.1 \pm 0.8$	$1.4 \pm 0.04$
13.26 <sup>d</sup>	$3^-; 1$	6.13	$3^-; 0$	>85	$9.2 \pm 1.5$

<sup>a</sup>) See tables 16.12 in (Aj 71) and 16.15 in (Aj 77) for the earlier work. See the latter table for the references for the values displayed here.

<sup>b</sup>) Monopole matrix element in fm<sup>2</sup>.

<sup>c</sup>) Pairs due to this transition are not observed (Al 78d).

<sup>d</sup>) For the radiative decay of higher states see tables 16.13, 16.18 and 16.22.

<sup>e</sup>) See also (La 77f: 94±10 meV).

<sup>f</sup>) "Best" value based on (La 77f: 60±10 meV) and earlier values displayed in (Aj 77).

<sup>g</sup>)  $(1.70^{+0.35}_{-0.50}) \times 10^{-3}$  (E2),  $(8.5^{+4.5}_{-2.5}) \times 10^{-4}$  (M1).

<sup>h</sup>)  $E_{\gamma} = 2471.5 \pm 0.5$  keV for (8.87→6.13) transition (Ga 70).

1. (a)  ${}^{10}\text{B}({}^6\text{Li}, p){}^{15}\text{N}$   $Q_m = 18.748$   $E_b = 30.876$
- (b)  ${}^{10}\text{B}({}^6\text{Li}, d){}^{14}\text{N}$   $Q_m = 10.140$
- (c)  ${}^{10}\text{B}({}^6\text{Li}, t){}^{13}\text{N}$   $Q_m = 5.843$
- (d)  ${}^{10}\text{B}({}^6\text{Li}, {}^3\text{He}){}^{13}\text{C}$   $Q_m = 8.083$
- (e)  ${}^{10}\text{B}({}^6\text{Li}, \alpha){}^{12}\text{C}$   $Q_m = 23.714$
- (f)  ${}^{10}\text{B}({}^6\text{Li}, {}^6\text{Li}){}^{10}\text{B}$

At  $E(^6\text{Li})=4.9$  MeV, the cross sections for reactions (a) to (e) leading to low-lying states in the residual nuclei are proportional to  $2J_t+1$ : this is interpreted as indicating that the reactions proceed via a statistical compound nucleus mechanism. For highly excited states, the cross section is higher than would be predicted by a  $2J_t+1$  dependence (Mc 66a). For reaction (f) see (Po 76a). See also (Aj 77), and <sup>12</sup>C in (Aj 80), <sup>13</sup>C, <sup>13</sup>N, <sup>14</sup>N and <sup>15</sup>N in (Aj 81).

2.  $^{10}\text{B}(^{10}\text{B}, \alpha)^{16}\text{O}$   $Q_m = 26.4155$

States of <sup>16</sup>O observed at  $E(^{10}\text{B}) = 20$  MeV are displayed in table 16.10 of (Aj 77). At the higher excitation energies, states are reported at  $E_x = 17.200 \pm 0.020$ ,  $17.825 \pm 0.025$ ,  $18.531 \pm 0.025$ ,  $18.69 \pm 0.03$ ,  $18.90 \pm 0.035$ ,  $19.55 \pm 0.035$ ,  $19.91 \pm 0.02$ ,  $20.538 \pm 0.015$ ,  $21.175 \pm 0.015$ ,  $21.84 \pm 0.025$ ,  $22.65 \pm 0.03$  and  $23.51 \pm 0.03$  MeV. The reaction excites known  $T=0$  states:  $\sigma_t$  follows  $2J_t+1$  for 11 of 12 groups leading to states of known  $J$ . The angular distributions show little structure (Aj 76b). See also (Ma 78a) and <sup>20</sup>Ne.

3.  $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$   $Q_m = 7.1620$

The yield of capture  $\gamma$ -rays has been studied for  $E_\alpha < 42$  MeV; see table 16.11 of (Aj 77) and (Sa 81e). Observed resonances are displayed in table 16.13 here.

This reaction plays an important role in astrophysical processes. (Dy 74) has determined  $\sigma_{E1}$  for  $E_\alpha = 1.88$  to 3.92 MeV. From these data  $S(E_\alpha(\text{lab}) = 400 \text{ keV}) = 0.08^{+0.05}_{-0.04} \text{ MeV} \cdot \text{b}$  (Ko 74) [used a hybrid  $R$ -matrix-optical-model analysis of ( $\alpha, \gamma$ ) and ( $\alpha, \alpha$ )],  $0.08^{+0.14}_{-0.07} \text{ MeV} \cdot \text{b}$  (Hu 76h) [used a two channel, two level approximation of a modified  $K$ -matrix]. (Dy 74) state that  $S$  depends quite strongly on  $\theta_\alpha^2(6.92)/\theta_\alpha^2(7.12)$ :  $S$  may have to be increased to allow for the tail of <sup>16</sup>O\*(6.92). For astrophysical considerations see the above references, (Aj 71, Aj 77) and (Fo 76j, Ba 77a, Sa 77b, Ro 78b, Ro 78o, Ta 78s, Ba 80k, Kh 81).

At higher energies the E2 cross section shows resonances at  $E_x = 13.2$ , 15.9, 16.5, 18.3, 20.0 and 26.5 MeV [see table 16.13]. Some E2 strength is also observed for  $E_x = 14$  to 15.5 and 20.5 to 23 MeV. In the range  $E_\alpha = 7$  to 27.5 MeV the  $T=0$  E2 strength is  $\sim 17\%$  of the sum rule. It appears from this and other experiments that the E2 centroid is at  $E_x \sim 15$  MeV, with a 15 MeV spread (Sn 74). Structures are observed in the yield of  $\gamma$ -rays from the decay of <sup>16</sup>O\*( $14.8 \pm 0.1$ ) for  $E_x = 34$ –39 MeV. It is suggested that these correspond to a giant quadrupole excitation with  $J^\pi = 8^+$  built on the  $6_1^+$  state at  $E_x = 14.816$  MeV (Sa 81e). See also (Ha 75ee, We 80d).

4.  $^{12}\text{C}(\alpha, n)^{15}\text{O}$   $Q_m = -8.5019$   $E_b = 7.1620$

For cross-section measurements from threshold to  $E_\alpha = 24.7$  MeV see (Aj 71) and table 16.11 in (Aj 77). Observed resonances are displayed in table 16.13 here. The production of neutrons at  $E_\alpha = 710$  MeV has been studied by (Ce 80). See also (Aj 77, Gr 77i, Li 77j) and (We 78i; applications).

ENERGY LEVELS OF LIGHT NUCLEI  $A=16-17$ 

TABLE 16.13  
Resonances in  $^{12}\text{C} + \alpha$

No.	$E_\alpha$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles <sup>a</sup> ) (x)	$\Gamma_x$	$\Gamma_\alpha/\Gamma$	$^{16}\text{O}^*$ (MeV)	$J^*; T$	Refs. <sup>b</sup>
1	$3.322 \pm 30$	400	$\gamma_0$ $\alpha_0$	$23 \pm 3$ meV $5.9 \pm 0.6$ meV $2.2 \pm 0.4$ meV	$\sim 1$	8.87 9.65 <sup>b</sup> )	$1^-$	(Mc 77h) <sup>b</sup>
2	$3.575 \pm 10$	$0.625 \pm 0.100$	$\gamma_0$ $\gamma_3$ $\alpha_0$	$\gamma_3$ $\gamma_0$ $\gamma_3$		9.842	$2^+$	(Mc 77h)
3	$4.256 \pm 11$	$27 \pm 4$	$\gamma_0$ $\alpha_0$	$58 \pm 7$ meV $\gamma_3$ $\alpha_0$ $\gamma_2$ $\gamma_3$	$10.353$ $1$		$4^+$	
4	$5.245 \pm 8$	$0.28 \pm 0.05$	$\alpha_0$ $\alpha_0$ $\alpha_0$ $\alpha_0$	$3.1 \pm 1.3$ meV $2.5 \pm 0.6$ meV $0.65 \pm 0.08$ eV $29 \pm 7$ meV		11.094	$4^+$	
5	$5.47$	2500	$\alpha_0$ $\gamma_0$ $\gamma_3$ $\alpha_0$	$11.26$ $11.52$			$0^+$	
6	$5.809 \pm 18$	$73 \pm 5$	$\alpha_0$ $\alpha_0$ $\alpha_0$ $\alpha_0$	$9.5 \pm 1.7$ eV <sup>c</sup> $0.12 \pm 0.06$ eV <sup>c</sup>			$2^+$	
7	$5.92 \pm 20$	$800 \pm 100$	$\alpha_0$ $\alpha_0$ $\gamma_0$ $p$	$11$ $1$ $1$ $1.1$ keV			$3^-$	
8	$6.518 \pm 10$	$1.5 \pm 0.5$	$\alpha_0$ $\gamma_1$ $p$	$92 \pm 8$ keV $0.025$ keV <sup>d</sup> $1.1$ keV			$0^+$	
9	$7.045 \pm 5^e$	$99 \pm 7$	$\alpha_0$ $\alpha_1$ $\gamma_0$ $\gamma_1$ $p$	$150 \pm 11$ keV $1.35 \pm 0.4$ eV $100$ keV $45 \pm 18$ keV $1$ keV			$1^-; 0$	(Op 76)
10	$7.82 \pm 10$	$150 \pm 11$	$\alpha_0$ $\alpha_1$ $\gamma_0$ $\gamma_4$ $p$	$0.8$ $1.302$ $13.085^f$ )			$2^+$	
11	$7.915 \pm 10^d$	$130 \pm 5$	$\alpha_0$ $\alpha_1$	$0.3$			$1^-; 1$	(Op 76)

TABLE 16.13—continued

No.	$E_\alpha$ (MeV±keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles <sup>a</sup> (x)	$\Gamma_x$	$\Gamma_a/\Gamma$	<sup>16</sup> O* (MeV)	$J^*; T$	Refs. <sup>b</sup>
12	7.960±10	110±30	$\gamma_0$ p $\alpha_0$ $\alpha_1$	>0.01 eV 1 keV 90±14 keV ~20 keV	0.7	13.129	3 <sup>-</sup> ; 0	
13	8.130±15	26±7	$\gamma$ p $\alpha_0$ $\alpha_1$	4.5 keV 9±4 keV 7.5 keV		13.257	3 <sup>-</sup> ; 1	
14	8.960±10	75±7	$\alpha_0$ $\alpha_1$	49 keV 23 keV	0.65±0.05	13.879±8	4 <sup>+</sup>	
15	9.1	4800	$\alpha_0$	~200 keV	>0.9	(14.0)	(0 <sup>+</sup> )	
16	9.164±15	200±50	$\alpha_0$	0.2±0.1	14.032	14.032	0 <sup>+</sup>	
17	9.3±100	750±200	$\alpha_0$	0.2±0.1	14.1	14.1	3 <sup>-</sup>	
18	9.948	487±12	$\alpha_1$	0.8±0.1	14.620±11	14.620	(4 <sup>+</sup> )	(Am 79a)
19	10.002	672±11	$\alpha_0$	~0.95	14.660±11	14.660	5 <sup>-</sup>	(Am 79a)
20	10.195±7	70±8	$\alpha_0$ $\alpha_1$	22 keV 48 keV	0.45±0.05	14.805	6 <sup>+</sup>	
21	10.544	166±30	$\alpha_0, \alpha_1, p_0$	0.35	15.066±11	15.066	0 <sup>+</sup>	(Am 79a)
22	10.999	133±7	$\alpha_0, \alpha_1, p_0$	0.58	15.408±2	15.408	3 <sup>-</sup>	(Am 79a)
23	11.560	703±113	$\alpha_0, \alpha_1, \gamma_{4.4}$	0.21	15.828±30	15.828	3 <sup>-</sup>	(Am 79a)
24	11.6	~600	$\gamma_0$	$\Gamma_a \Gamma_\gamma / \Gamma \sim 0.4$ eV	15.9	15.9	2 <sup>+</sup>	
25	12.156	422±14	$\alpha_0$	0.93	16.275±7	16.275	6 <sup>+</sup>	(Am 79a)
26	12.272	65±45	$\alpha_0, \alpha_1, \alpha_2, p_0$	0.07	16.362±20	16.362	(0 <sup>+, 1<sup>-</sup>)</sup>	(Am 79a)
27	12.380	22±3	$\gamma_0, n, p_0, \alpha_0,$ $\alpha_1, \alpha_2, \gamma_{4.4}$	0.28	16.443±21 <sup>b</sup>	16.443	2 <sup>+, 1<sup>-</sup>)</sup>	(Am 79a)
28	12.5	730	$p_0, \alpha_0$		(16.5)			

## ENERGY LEVELS OF LIGHT NUCLEI A = 16-17

29	12.915	$567 \pm 60$	$\alpha_0$	0.28	$16.844 \pm 21$	$4^+$
30	13.0	700	$\alpha_0, {}^8\text{Be}$	(16.9)	(Am 79a)	5 <sup>-</sup>
31	13.05	$\sim 280$	$\alpha_2, {}^8\text{Be}$	0.37	$17.129 \pm 5$	$2^+$
32	13.296	$107 \pm 14$	$n, p_0, \alpha_0, \alpha_1, \gamma_{4.4}$		(Am 79a)	$2^+$
33	13.32	$36 \pm 5$	$\alpha_0, \alpha_1$		(Am 79a)	$17.15$
34	13.35	$160 \pm 60$	$\alpha_2, {}^8\text{Be}$		(Am 79a)	$2^+$
35	13.50	$<100$	n		(Am 79a)	
36	13.805	$182 \pm 56$	$\alpha_0, (\alpha_1), \alpha_2$	0.16	$17.510 \pm 26$	$1^-$
37	13.865	$178 \pm 66$	$n, (\alpha_0, \alpha_1)$	0.07	$17.555 \pm 21$	$(6^+)$
38	13.948	$175 \pm 55$	$p_0, \alpha_0$	0.32	$17.618 \pm 20$	$0^+, 1^-$
39	14.08	(~75)	${}^8\text{Be}$		17.72	$(0^+, 2^+)$
40	14.170	$396 \pm 41$	$n, \alpha_0, \alpha_1, \gamma_{4.4}, {}^8\text{Be}$	0.34	$17.784 \pm 15$	$4^+$
41	14.480	$14 \pm 2$	$(n), p_0, \alpha_0, \alpha_1, \gamma_{4.4}, {}^8\text{Be}$	0.36	$18.016 \pm 1$	$4^+; (0)$
42	14.577	$248 \pm 90$	$(\gamma_0), n_0, p_0, \alpha_0$	0.31	$18.089 \pm 25$	$(0^+)$
43	(14.62)	(~45)			(18.12)	$(\neq 4^+)$
44	14.85	$\sim 300$	$\gamma_0, p_0, (\alpha_1, \gamma_{4.4})$		18.29	(Am 79a)
45	14.997	$544 \pm 39$	$\alpha_0$	0.40	$18.404 \pm 12$	$5^-$
46	15.2	$\sim 150$	$\alpha_0, (\alpha_1, \alpha_2, \gamma_{4.4})$		(1 <sup>-</sup> , 5 <sup>-</sup> )	(Am 79a)
47	15.2	$\sim 300$	$\alpha_2, {}^8\text{Be}$		18.6	$(4^+)$
48	15.490	$215 \pm 45$	$p_0, \alpha_0$	0.26	$18.773 \pm 22$	$1^-$
49	15.506	$260 \pm 16$	$n, p_0, \alpha_0, (\alpha_1), {}^8\text{Be}$	0.48	$18.785 \pm 6$	$4^+$
50	15.8	$\sim 550$	$(\alpha_0), \alpha_1, \gamma_{4.4}$		19.0	$(5^-)$
51	15.96	41	$(n), \alpha_0$		(19.12)	$(2^+, 4^+)$
52	16.130	$50 \pm 45$	$(n), (\alpha_0)$	0.04	$19.253 \pm 30$	$(5^-)$
53	16.137	$155 \pm 23$	$p_0, \alpha_0, (\alpha_1)$	0.34	$19.257 \pm 9$	$2^+$
54	16.219	$63 \pm 33$	$p_0, (\alpha_0), \alpha_1, {}^8\text{Be}$	0.07	$19.319 \pm 14$	$(6^+)$
55	16.293	$23 \pm 4$	$p_0, \alpha_0, \alpha_1, \alpha_2$	0.23	$19.375 \pm 2$	$4^+$
56	16.496	$255 \pm 75$	$(n), \alpha_0, (\alpha_1, \alpha_2)$	0.20	$19.527 \pm 26$	$2^+$
57	16.799	$286 \pm 44$	$p_0, \alpha_0, \alpha_1$	0.29	$19.754 \pm 16$	$2^+$
58	(16.92)	(~175)	$a_2$		(19.85)	(Am 79a)
59	(17.05)	(~30)	$(\alpha_0)$		(19.94)	$(\neq 3^-)$
60	17.201	$432 \pm 40$	$\gamma_0, n, (p_0), \alpha_0, (\alpha_1)$	0.43	$20.055 \pm 13$	$2^+$
61	(17.27)	(~45)	$(\alpha_0)$		(20.11)	(Am 79a)
62	17.5	$\sim 1500$	$p_0$		(20.3)	$(\pm 3^-)$
63	(17.66)	(~150)	$n, (p_0), \alpha_2$		(20.40)	$(4^+)$

TABLE 16.13—continued

No.	$E_\alpha$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles <sup>a</sup> (x)	$\Gamma_x$	$\Gamma_\alpha/\Gamma$	$^{16}\text{O}^*$ (MeV)	$J^\pi; T$	Refs. <sup>b</sup>
64	(17.8)	(~300)	( $\alpha_0$ ), $\alpha_1$		0.14 $\pm$ 0.02	(20.5)		(Am 79a)
65	17.849	11 $\pm$ 2	$p_0, \alpha_0, \alpha_1, \alpha_2$			20.51 $\pm$ 2	5 <sup>-</sup>	(Am 79a)
66	17.875	<5	$\alpha_0$			20.560 $\pm$ 2	even	(Am 79a)
67	17.948	<10	$\alpha_0$			20.615 $\pm$ 3	even	(Am 79a)
68	(18.2)	(~60)	n, ( $p_0$ )			(20.8)		(Am 79a)
69	18.271	904 $\pm$ 55	$\alpha_0$	0.60	20.857 $\pm$ 14	7 <sup>-</sup>		(Am 79a)
70	(18.3)	(~50)	$\alpha_0$		(20.9)	(2 <sup>+</sup> )		(Am 79a)
71	(18.48)	240 $\pm$ 80	n, $p_0, (\alpha_0)$	0.20	(21.01)	(Am 79a)		(Am 79a)
72	18.50 $\pm$ 25	900	$\gamma_0, (\alpha_0, \alpha_1)$		21.03	(1 <sup>-</sup> )		(Am 79a)
73	18.5	205 $\pm$ 14	$\alpha_0$	0.50	(21.0)	(5 <sup>-</sup> )		(Am 79a)
74	18.531	306 $\pm$ 46	$\alpha_0$	0.21	21.052 $\pm$ 6	6 <sup>+</sup>		(Am 79a)
75	18.593	61 $\pm$ 32	( $\alpha_0$ )		(21.098)	4 <sup>+</sup>		(Am 79a)
76	19.14		(n), $\alpha_0, \alpha_2$	21.52	21.52	7 <sup>-</sup>		(Am 79a, Fr 81h)
77	19.327	115 $\pm$ 8	n, $\alpha_0, \alpha_1, \alpha_2$	0.41	21.648 $\pm$ 3	6 <sup>+</sup>		(Am 79a)
78	19.498	43 $\pm$ 20	n, $p_0, \alpha_0, \alpha_1, \alpha_2$	0.07	21.776 $\pm$ 9	3 <sup>-</sup>		(Am 79a)
79	19.85	60	n					
80	19.89	340	n		22.04			
81	19.95	<150	n, <sup>8</sup> Be					
82	20.49	375	n		22.07			
83	20.71	60	n, <sup>8</sup> Be					
84	20.760 $\pm$ 5	12.5 $\pm$ 2.5	n, $p_0, \alpha_0, \alpha_2$		22.11			
85	21.28	~20	$\alpha_0, \alpha_1, ^8\text{Be}$		22.52			
86	21.3	$\leq 50$	<sup>8</sup> Be		22.52			
87	21.67	<40	n		22.68			
88	21.85	300	$\alpha_0, \alpha_1$		22.721	0 <sup>+</sup> ; T=2		
89	22.0	1500	$\gamma_{12.71}$		23.11			
90	22.14	120	n		23.1	6 <sup>+</sup>		
91	22.306 $\pm$ 6	26 $\pm$ 4	$p_0, \alpha_0, \alpha_1, \alpha_2, ^8\text{Be}$	j)	23.40			
92	22.37	165	n	0.06 $\pm$ 0.02	23.54			
					23.6			
					23.75			
					23.879			
					23.93			

93	22.75	$\leq 500$	<sup>8</sup> Be			24.21		
94	23.2	750	$\gamma_{12.71}, \gamma_{15.11}$			24.5	$T=1$	
95	24.1	450	$\gamma_{15.11}$			25.2	$T=1$	
96	24.6	450	$\gamma_{15.11}$			25.6	$T=1$	
97	25.5	450	$\gamma_{15.11}$			26.3	$T=1$	
98	25.6	1200	$\alpha_0, \gamma_{12.71}$	$\Gamma_\alpha \Gamma_\gamma / \Gamma = 1.2 \text{ eV}$		26.3	$2^+$	
99	29	4 MeV	$\alpha_0, \alpha_1, p_3$			29		(Bu 78i)

<sup>a</sup>) Po corresponds to <sup>15</sup>N(0).  $\alpha_0, \alpha_1$  correspond to <sup>12</sup>C\*(0, 4.4) and  $\gamma_{1.4}$  corresponds to the  $\gamma$ -ray from the decay of <sup>12</sup>C\*(4.4);  $\gamma_0, \gamma_1, \gamma_2, \gamma_3, \gamma_4$  correspond to the transitions to <sup>16</sup>O\*(0, 6.05, 6.13, 6.92, 7.12).

<sup>b</sup>) Previous ratios are listed in tables 16.11 (Aj 77) and 16.12 (Aj 77). Please note that (Am 79a) is an unpublished thesis. (Am 81b) has been submitted for publication.

<sup>c</sup>)  $7040 \pm 5$  keV (Op 76).

<sup>d</sup>)  $7880 \pm 15$  keV (Op 76).

<sup>e</sup>) Branching ratios to <sup>16</sup>O\*(0, 6.05) = 98.8% and 1.2%.

<sup>f</sup>)  $\Gamma_{\alpha_0} = 0.7 \pm 0.2$  eV (Ke 71e), based on  $\Gamma_{\alpha_0} / \Gamma = 1.0$  (Mo 68h) and  $\Gamma_{\text{c.m.}} = 190 \pm 40$  keV (Ke 71e).

<sup>g</sup>)  $\Gamma_{\alpha_0} \Gamma_{\gamma} / \Gamma^2 = (1.49 \pm 0.17) \times 10^{-4}$  (Ke 71e).

<sup>h</sup>)  $\theta_\alpha^2(7.12) / \theta_\alpha^2(9.63) = 0.19^{+0.16}_{-0.11}$  (Ko 74). See also reactions 9 and 12.

<sup>i</sup>) See column 2 and footnote (d).

<sup>j</sup>)  $\Gamma_{\text{Be}} / \Gamma_{\alpha_0}$  and  $\Gamma_{\alpha_2} \sim 3.5, 1.5 \pm 0.5$  and  $\sim 6$  keV, respectively (Br 76d).

<sup>k</sup>) An attempt is reported by (Mc 77h) to observe a  $0^+$  state in the vicinity of the known  $2^-$  state at 8.87 MeV. No such state is seen:  $\theta_\alpha^2 \leq 2 \times 10^{-4}$ .

<sup>l</sup>) See (Sa 81e) for ( $\alpha, \gamma_{14.8}$ ) measurements which indicate an  $8^+$  GQR built on the  $6_1^+$  state <sup>16</sup>O\*(14.82).

**<sup>16</sup>O**

36

F. AJZENBERG-SELOVE

$$5. \ ^{12}\text{C}(\alpha, \text{p})^{15}\text{N} \quad Q_{\text{m}} = -4.9656 \quad E_{\text{b}} = 7.1620$$

The yield of  $\text{p}_0$  has been studied for  $E_{\alpha} = 7.7$  to 23 MeV: see (Aj 77), (Am 79a, Am 81b;  $15 \leq E_x \leq 22$  MeV) and (Mo 79e;  $17.5 \leq E_x \leq 18.7$  MeV). The excitation curve for  $\text{p}_3$  (to  $^{15}\text{N}^*(6.32)$ ), measured for  $E_{\alpha} = 24$  to 33 MeV, shows a large peak at  $E_x \sim 29$  MeV,  $\Gamma \sim 4$  MeV. It is suggested that it is related to the GQR in  $^{16}\text{O}$  (Bu 78j). For other resonances see table 16.13. See also  $^{15}\text{N}$  in (Aj 81).

$$6. \begin{array}{l} (\text{a}) \ ^{12}\text{C}(\alpha, \text{d})^{14}\text{N} \\ (\text{b}) \ ^{12}\text{C}(\alpha, \text{t})^{13}\text{N} \\ (\text{c}) \ ^{12}\text{C}(\alpha, {^3\text{He}})^{13}\text{C} \end{array} \quad Q_{\text{m}} = -13.57434 \quad E_{\text{b}} = 7.1620$$

$$Q_{\text{m}} = -17.8706$$

$$Q_{\text{m}} = -15.6314$$

See (Aj 81).

$$7. \ ^{12}\text{C}(\alpha, \alpha)^{12}\text{C} \quad E_{\text{b}} = 7.1620$$

The yield of  $\alpha$ -particles corresponding to  $^{12}\text{C}^*(0, 4.4, 7.7)$  and of 4.4, 12.7 and 15.1 MeV  $\gamma$ -rays has been studied at many energies in the range  $E_{\alpha} = 2.5$  to 35.5 MeV: see table 16.11 in (Aj 77), (Am 79a, Am 78a, Am 79, Am 81b;  $E_{\alpha} = 15$  to 20 MeV), and (Ar 79e; 17 to 23 MeV) and (Fr 81h; 17.4 to 20.5 MeV;  $\alpha_2$ ). Observed resonances are displayed in table 16.13. No evidence is observed for a narrow (100 eV)  $0^+$  state in the vicinity of  $^{16}\text{O}^*(8.87)$ ; thus the  $\alpha$ -decay reported in  $^{16}\text{N}(\beta^-)$  [reaction 52] appears to be due to  $^{16}\text{O}^*(8.87)$  ( $J^\pi = 2^-$ ) and is thus parity forbidden (Mc 77h). No evidence is found by (Fr 81h) for  $8^+$  strength.

Total reaction cross-section measurements are reported at 1.55 and 2.89 GeV/c per nucleon (Ja 78m). See also (De 79w, De 79z, De 80b, De 80l, Ba81r, Pe 81). For  $\pi^0$  measurements see (Wa 76k). For spallation measurements see (Fo 77b, Du 78h, Gl 79e, Vi 79, Ab 80b, Go 81b) and (Aj 77).

See also (Fr 78c, Wa79j), (Si 76i, De 80w), (Cu 76f, Ba77l, Ba 77oo, Ba 77rr, Br 77gg, Da 77h, Fr 77b, Hu 77b, Ik 77b, Sa 77p, Ze 77b, Fr 78g, Su 78d, Su 78f, Ve 79e, Ba 80kk, Dm 80, Fu 80e, Fu 80g, Li 80f, St 80i, Su 81a, Wi 81b; theor.) and  $^{12}\text{C}$  in (Aj 80).

$$8. \begin{array}{l} (\text{a}) \ ^{12}\text{C}(\alpha, {^8\text{Be}}){^8\text{Be}} \\ (\text{b}) \ ^{12}\text{C}(\alpha, 2\alpha){^8\text{Be}} \\ (\text{c}) \ ^{12}\text{C}(\alpha, 4\alpha) \end{array} \quad Q_{\text{m}} = -7.4586 \quad E_{\text{b}} = 7.1620$$

$$Q_{\text{m}} = -7.3667$$

$$Q_{\text{m}} = -7.2748$$

The yield of  ${^8\text{Be}}$  (reaction a) shows a number of resonances: see table 16.13. There is no evidence below  $E_x \sim 24$  MeV for  $J^\pi = 8^+$  states although the existence of such states below this energy cannot be ruled out since it is possible that the  $L$  of the entrance channel inhibits the formation of such states. Above 26 MeV  $L = 8$  becomes dominant (Br 76d). For reactions (b) and (c) see (Aj 77).

$$9. \ ^{12}\text{C}({^6\text{Li}}, \text{d})^{16}\text{O} \quad Q_{\text{m}} = 5.6885$$

This reaction has been studied at many energies: see the references in (Aj 77) and table 16.14. At the higher energies the spectra are dominated by states with  $J \geq 4$  (and natural parity). An attempt to locate an  $8^+$  state with  $E_x = 17$  to 24 MeV

TABLE 16.14  
States of <sup>16</sup>O from <sup>12</sup>C(<sup>6</sup>Li, d) and <sup>12</sup>C(<sup>7</sup>Li, t)

$E_x^a)$ (MeV ± keV)	$\Gamma_{c.m.}^b)$ (keV)	$\theta_a^2/\theta_a^2(2^+)^c)$	$\Gamma_{\alpha}/\Gamma^d)$	$J^\pi; K^\pi$
0		0.93, 0.18		$0^+$
6.05		0.38, 1.10		$0^+; 0^+$
6.13		0.23, 0.22		$3^-$
6.92		=1.0		$2^+; 0^+$
7.12		0.53, 0.39		$1^-$
8.87	<20			$2^-$
$9.63 \pm 30^i)$	$409 \pm 10^i)$	0.30, 0.60		$1^-; 0^-$
9.85	<20	$\leq 0.05, \leq 0.01$		$2^+$
$10.346 \pm 6^{e,j})$	$35 \pm 5$	0.25, 0.47	$0.86 \pm 0.09$	$4^+; 0^+$
10.95				$0^-$
11.09 <sup>j)</sup>	<30	$\leq 0.06, \leq 0.03$	$0.31 \pm 0.03$ ( $J = 4^+$ )	$3^+ + 4^+$
11.59 ± 20	$700 \pm 100$	~0.4		$3^-; 0^-$
13.09	~230			$1^-$
$14.363 \pm 15^h)$	<120			$>5, \pi = \text{nat.}$
14.66 ± 20	$500 \pm 50$		$1.03 \pm 0.1$	$5^-; 0^-$
14.82	45 ± 10			$(6^+)$
$16.30 \pm 20^{e,k})$	$300 \pm 50$		$1.07 \pm 0.11$	$6^+; 0^+$
17.65 ± 50	$100 \pm 50$			
17.85 ± 50	~200			
(18.6) <sup>f)</sup>				$(5^-)$
19.30 ± 50	~200			
$20.8 \pm 100^e)$	$500 \pm 100$		$1.16 \pm 0.23$	$7^-; 0^-$
$21.6 \pm 100$	$\leq 100$		$0.67 \pm 0.14$	$6^+{}^s)$
$23.0 \pm 100$	~200			$(6^+)$
$23.8 \pm 100$	$1980 \pm 250$			$(6^+{}^s)$
$26.9 \pm 100$	$1700 \pm 250$			$(7^-{}^s)$
(29.3) <sup>s)</sup>				$(7^-{}^s)$

<sup>a)</sup>  $E_x$  quoted without errors are from table 16.11. The others are from (Be 78u:  $E(^6\text{Li}) = 42.1$  MeV) and (Be 80o:  $E(^6\text{Li}) = 90.2$  MeV). The states with  $E_x < 11.1$  MeV have also been studied by (Be 78t:  $E(^7\text{Li}) = 34$  MeV). Angular distributions are reported in both reactions for the first nine states. See also (Cu 78c).

<sup>b)</sup> See (Be 78t, Be 78u, Be 80o): line widths, not corrected for  $\alpha$ -penetrabilities.

<sup>c)</sup> Ratio of dimensionless reduced  $\alpha$ -width calculated at a channel radius of 5.4 fm, relative to that for <sup>16</sup>O\*(6.92). ( $N, L$ ) here are taken to be (2, 0) and (4, 1), respectively, for <sup>16</sup>O\*(0, 7.12). The first number listed is the value reported at  $E(^6\text{Li}) = 42$  MeV (Be 78u), the second at  $E(^6\text{Li}) = 90.2$  MeV (Be 80o).

<sup>d)</sup> (Cu 80b); d- $\alpha$  angular correlations. See also (Cu 77, Cu 78c, Ar 78h).

<sup>e)</sup> (Ba 71k).

<sup>f)</sup> (Ar 79e); d- $\alpha$  angular correlations.

<sup>g)</sup> (Ar 76h, Ar 76i); d- $\alpha$  angular correlations.

<sup>h)</sup> Observed at  $E(^7\text{Li}) = 38$  MeV (Co 76c) and M. E. Coborn, private communication.

<sup>i)</sup> On the basis of studies of the <sup>12</sup>C(<sup>6</sup>Li, d), <sup>12</sup>C(<sup>7</sup>Li, t), <sup>12</sup>C(<sup>10</sup>B, <sup>6</sup>Li) and <sup>19</sup>F(p,  $\alpha$ ) reactions, the energy of <sup>16</sup>O\*(9.6) is  $9619 \pm 15$  keV,  $\Gamma = 400 \pm 10$  keV (line width),  $\Gamma_R = 430 \pm 10$  keV as inferred from the best fit BW line shape. This value is corrected for penetrability (Ov 81 and F. Becchetti, private communication).

<sup>j)</sup> Angular distributions are reported at  $E(^6\text{Li}) = 35.5-35.6$  MeV to <sup>16</sup>O\*(10.35) and to the unresolved  $3^+$  and  $4^+$  states at 11.1 MeV. It appears that the  $4^+$  state is dominantly populated, and that two-step processes may be important in this reaction (Ke 80h, Cl 78f). See also (Cu 78c).

<sup>k)</sup> Angular distributions at  $E(^6\text{Li}) = 35.5$  MeV (Cl 78f) and 28 and 34 MeV (Cu 78c).

by ( $d, \alpha$ ) angular correlations has been unsuccessful: see (Ar 76i;  $E(^6\text{Li})=29.0$  and  $34.7$  MeV). The ratio  $R$  [ $=\theta_\alpha^2(7.1 \text{ MeV})/\theta_\alpha^2(9.6 \text{ MeV})$ ] is of astrophysical interest: an FRDW analysis by (Be 80o) leads to  $R = 0.7 \pm 0.2$  for  $s = 5.4$  fm. See also (Be 78u).


For analyzing power measurements see <sup>18</sup>F in (Aj 83); (Ma 78i;  $E(^6\text{Li})=20$  MeV; to <sup>16</sup>O states with  $E_x < 11.1$  MeV). See also (Be 79d, Fu 80c, Ma 80j), (Fo 77g, Be 78w, Fi 78a, Ma 79aa) and (St 76n, Ba 77mm, Ap 78, Cu 78g, Me 78d, Su 79g, Ze 79a, Ap 80; theor.).



This reaction has been studied extensively: see (Aj 77) for the earlier references. Observed states are displayed in table 16.14. Angular distributions have been reported recently at  $E(^7\text{Li})=34$  MeV (Be 78t; first nine states) and at  $68$  MeV (Br 79a; to <sup>16</sup>O\*(6.92, 10.35, 16.3)). The value of  $R$  [see reaction 9] derived by (Be 78t) is  $0.35 \pm 0.13$ . See also <sup>19</sup>F in (Aj 83), (Sa 76m, Cl 78f, Ro 79j, De 80d, De 81i) and (Bi 77e; theor.).



At  $E(^{10}\text{B})=18$  and  $45$  MeV angular distributions have been studied involving <sup>16</sup>O\*(0, 6.1, 7.1, 8.9, 9.9, 10.4) (Hi 70b). At  $E(^{10}\text{B})=68$  MeV angular distributions to <sup>16</sup>O\*(0, 6.1, 6.9, 10.4, 11.1, 14.7, 16.2, 20.9) are forward peaked and fairly structureless. <sup>16</sup>O\*(0, 6.9, 11.1) are weakly excited (Bi 81c). See also footnote (i) in table 16.14 and (Be 78j).



See (Be 78j). See also (Aj 77).



Angular distributions have been reported at  $E(^{12}\text{C})$  to  $63$  MeV: see (Aj 77). Angular correlations at  $E(^{12}\text{C})=78$  MeV confirm  $J^\pi = 4^+, 5^-, 6^+$  and  $7^-$  for <sup>16</sup>O\*(10.35, 14.59, 16.3, 20.9).  $\Gamma_{a_0}/\Gamma = 0.90 \pm 0.10$ ,  $0.75 \pm 0.15$  and  $0.90 \pm 0.10$ , respectively, for the first three of these states. In addition a state is reported at  $E_x = 22.5 \pm 0.5$  MeV which may be the  $8^+$  member of the  $K^\pi = 0^+$ , 4p-4h rotational band (Sa 79k, Sa 77j). For the decay of <sup>20</sup>Ne states see (Yo 79) and (Aj 83). For excitation function and fusion measurements see (Hi 77c, Wa 77e, Co 78e, Ja 78g, He 79a, Ko 80, Wa 80f, Wi 80b). See also (Ab 77e, To 78c; theor.).



Angular distributions are reported at  $E(^{14}\text{N})=53$  MeV involving <sup>16</sup>O\*(0, 6.05, 6.13, 6.92) and various states of <sup>10</sup>B (Ze 76) and at  $E(^{14}\text{N})=78.8$  MeV (Mo 77y, Mo 79z) involving <sup>16</sup>O<sub>gs..</sub>.

15. (a)  $^{12}\text{C}(^{17}\text{O}, ^{13}\text{C})^{16}\text{O}$   $Q_m = 0.8021$   
 (b)  $^{12}\text{C}(^{18}\text{O}, ^{14}\text{C})^{16}\text{O}$   $Q_m = 0.9341$

Angular distributions are reported at  $E(^{17}\text{O}) = 30.5$  and  $33.8 \text{ MeV}$  (Ch 78m). For reaction (b) see (Aj 77) and (Ta 78f; theor.).

16.  $^{12}\text{C}(^{19}\text{F}, ^{15}\text{N})^{16}\text{O}$   $Q_m = 3.1481$

Angular distributions have been measured at  $E(^{19}\text{F}) = 40, 60$  and  $68.8 \text{ MeV}$  involving different states in  $^{15}\text{N}$  and  $^{16}\text{O}^*(0, 6.1, 7.0, 10.4)$  (Sc 72c).

17.  $^{12}\text{C}(^{20}\text{Ne}, ^{16}\text{O})^{16}\text{O}$   $Q_m = 2.4310$

Angular distributions have been measured to  $E(^{20}\text{Ne}) = 147 \text{ MeV}$  see (Aj 77). Excitation functions are reported at  $E(^{20}\text{Ne}) = 45.3$  to  $93.3 \text{ MeV}$  (Do 78) and 150 to 250 MeV (Or 79:  $E_x = 10-15 \text{ MeV}$  and  $20-25 \text{ MeV}$ ). For an experiment studying  $^{16}\text{O} + \alpha$  coincidences see (Os 81g). See also (Pi 75f).

18.  $^{13}\text{C}(^3\text{He}, \gamma)^{16}\text{O}$   $Q_m = 22.7934$

The yield of capture  $\gamma$ -rays has been studied for  $E(^3\text{He})$  to  $16 \text{ MeV}$  [see (Aj 77)], as have angular distributions. Observed resonances are displayed in table 16.15. It is suggested that the structures at  $E_x \sim 26-29 \text{ MeV}$  are part of giant resonances built on the first few excited states of  $^{16}\text{O}$  (Ve 79).

19.  $^{13}\text{C}(^3\text{He}, \text{n})^{15}\text{O}$   $Q_m = 7.1295$   $E_b = 22.7934$

The excitation functions to  $E(^3\text{He}) = 11 \text{ MeV}$  are marked at low energies by complex structures and possibly by two resonances at  $E(^3\text{He}) = 1.55$  and  $2.0 \text{ MeV}$ : see table 16.15. See also (Aj 77) for polarization measurements.

20.  $^{13}\text{C}(^3\text{He}, \text{p})^{15}\text{N}$   $Q_m = 10.6658$   $E_b = 22.7934$

Excitation functions for  $E(^3\text{He}) = 3.6$  to  $6.6 \text{ MeV}$  have been measured for  $p_0$ ,  $p_{1+2}$ ,  $p_3$ : a resonance is reported at  $E(^3\text{He}) = 4.6 \text{ MeV}$  (Ch 78p). A resonance at  $6 \text{ MeV}$  has also been observed: see table 16.15. See also  $^{15}\text{N}$  in (Aj 81).

21.  $^{13}\text{C}(^3\text{He}, \text{d})^{14}\text{N}$   $Q_m = 2.0571$   $E_b = 22.7934$

See  $^{14}\text{N}$  in (Aj 81).

22. (a)  $^{13}\text{C}(^3\text{He}, \text{t})^{13}\text{N}$   $Q_m = -2.2392$   $E_b = 22.7934$   
 (b)  $^{13}\text{C}(^3\text{He}, ^3\text{He})^{13}\text{C}$

The excitation function for formation of  $^{13}\text{N}_{\text{g.s.}}$  has been studied for  $E(^3\text{He}) = 11$  to  $17 \text{ MeV}$ : see (Aj 77). Polarization measurements are reported by (Ba 81c) at  $E(^3\text{He}) = 33 \text{ MeV}$  ( $t_0, t_{2+3}$ ). For reaction (b) see table 16.15.

23.  $^{13}\text{C}(^3\text{He}, \alpha)^{12}\text{C}$   $Q_m = 15.6314$   $E_b = 22.7934$

TABLE 16.15  
Resonances in <sup>13</sup>C + <sup>3</sup>He

$E(^3\text{He})$ (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	<sup>16</sup> O*	$J^\pi; T$	Refs. <sup>a)</sup>
1.55	~80	$n_0, n_3$	24.05		
$1.55 \pm 100$	450	$\gamma_0$	24.1		
2.0	~250	$n_0$	24.4		
$2.6 \pm 100$		$\alpha\gamma_{15.1}$	24.9	( $T = 1$ )	
$2.87 \pm 50$	600	$\gamma_0$	25.12	$1^-$	(Ve 79)
~3.1		$\alpha_0, \alpha_2$	~25.3		
~3.5	~300	$\alpha_0$	~25.6	$(3^-)$	
~4	~300	$\alpha_0, \alpha_1,$ $\alpha_2$	~26	$(3^-)$	
$4.0 \pm 100$	<sup>b)</sup>	$\gamma_0, \gamma_{1+2},$ $\alpha\gamma_{15.1}$	26.0	$1^-; (1)$	(Ve 79)
$4.6 \pm 100^c)$	$720 \pm 160$	$\gamma_2, p_0$	26.5	$2^+, 4^+$	(Ch 77b, Ch 78p, Ve 79)
$5.2 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	27.0	( $T = 1$ )	
$5.6 \pm 100$	~600	$\gamma_0, \gamma_{1+2},$ $\alpha\gamma_{15.1}, {}^8\text{Be}$	27.3	$(1^-)$	(Ve 79)
~5.8	~2500	$\gamma_{3+4}$	27.5		
$6.0 \pm 100$	~500	$p_0, p_{1+2},$ ${}^3\text{He}, \alpha_1, \alpha_2$	27.7	$(3^-; 0)$	(Ve 79)
~6		$\gamma_0$	28		
$6.5 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	28.1	( $T = 1$ )	
$6.8 \pm 100$		$\alpha_0, \alpha_1, \alpha_2$	28.3	$(T = 0)$	
$7.1 \pm 200$		$\gamma_{1+2}$	28.6		
$7.5 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	28.9	$(T = 1)$	
$8.6 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	29.8	$(T = 1)$	
$9.4 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	30.4	$(T = 1)$	
$10.1 \pm 100$	<sup>b)</sup>	$\alpha\gamma_{15.1}$	31.0	$(T = 1)$	

<sup>a)</sup> For earlier references see tables 16.15 in (Aj 71) and 16.13 in (Aj 77).

<sup>b)</sup> Lab widths 0.5–1 MeV (Ta 69b).


<sup>c)</sup> Based on  $\Gamma_{\text{c.m.}} = 530 \pm 80$  keV [from <sup>15</sup>N(p,  $\gamma$ , see table 16.18],  $\Gamma_{p_0} = 150 \pm 45$  keV [ $J^\pi = 2^+$ ],  $110 \pm 35$  keV [ $4^+$ ],  $\Gamma_{p_0}/\Gamma = 0.29 \pm 0.10$  [ $2^+$ ],  $0.21 \pm 0.07$  [ $4^+$ ],  $\Gamma_{\gamma_2} = 740 \pm 240$  eV [ $2^+$ ],  $410 \pm 140$  eV [ $4^+$ ] (Ch 77b, Ch 78p).

Yields of  $\alpha_0, \alpha_1, \alpha_2$  and  $\gamma$ -rays from the decay of <sup>12</sup>C\*(12.71, 15.11) have been studied up to  $E(^3\text{He}) = 12$  MeV. Observed resonances are displayed in table 16.15. Those seen in the yield of  $\gamma_{15.1}$  are assumed to correspond to <sup>16</sup>O states which have primarily a  $T = 1$  character. Analyzing power measurements are reported at  $E(^3\text{He}) = 33$  MeV to <sup>12</sup>C\*(4.4) (Ka 81c). See (Aj 71, Aj 77) for the earlier references and <sup>12</sup>C in (Aj 80).

$$24. \quad ^{13}\text{C}(^3\text{He}, {}^8\text{Be}) {}^8\text{Be} \quad Q_m = 8.1728 \quad E_b = 22.7934$$

The excitation function for <sup>8</sup>Be(g.s.) has been studied for  $E(^3\text{He}) = 2$  to 6 MeV. It shows a strong resonance at  $E(^3\text{He}) = 5.6$  MeV corresponding to a state in <sup>16</sup>O

at  $E_x = 27.3$  MeV.  $J''$  appears to be  $2^+$  from angular distribution measurements (Ja 68h).



Angular distributions for the  $n_0$  group have been measured for  $E_\alpha = 12.8$  to 22.5 MeV: see (Aj 71). See also (Aj 77), (Ha 81d), (Sc 77i, Co 80q, Tr 80j; astrophys.) and (Ge 78i, Ba 80l; applications).



Angular distributions have been studied at  $E(^6\text{Li}) = 20$  and 28 MeV to  $^{16}\text{O}^*(0$  [at 28 MeV], 6.13, 7.0, 8.87, 9.85, 10.35, 11.09). At these energies the spectra are dominated by the triton groups to  $^{16}\text{O}^*(11.09, 14.30, 14.39, 14.82)$ . At  $E(^6\text{Li}) = 25$  MeV the excitation of  $^{16}\text{O}^*(14.52, 14.66)$  is also reported: see (Aj 77) and (Cu 80a). See also (Ma 77g) and  $^{19}\text{F}$  in (Aj 83).



Angular distributions have been measured at  $E(^{12}\text{C}) = 87$  MeV and at  $E(^{13}\text{C}) = 36$  MeV: see (Aj 77). At  $E(^{13}\text{C}) = 105$  MeV,  $^{16}\text{O}^*(6.05 + 6.13, 10.35, 16.3, 20.7)$  are strongly populated (Br 79a).



Angular distributions have been measured at  $E(^{17}\text{O}) = 29.8$  and 32.3 MeV (Ch 77m, Ch 78m).



At  $E(^3\text{He}) = 11$  to 16 MeV, neutron groups are observed to  $T = 2$  states at  $E_x = 22.717 \pm 0.008$  and  $24.522 \pm 0.011$  MeV ( $\Gamma < 30$  keV and  $< 50$  keV, respectively). These two states are presumably the first two  $T = 2$  states in  $^{16}\text{O}$ , the analog states to  $^{16}\text{C}^*(0, 1.75)$ .  $J''$  for  $^{16}\text{O}^*(24.52)$  is found to be  $2^+$  from angular distribution measurements (Ad 70a). See also (Aj 71).



See  $^{16}\text{C}$ .



The  $\gamma_0$  yield has been studied for  $E_d = 0.5$  to 5.5 MeV. Three resonances are reported: see table 16.16. See (Aj 77) for references.



For  $E_d = 0.66$  to 5.62 MeV, there is a great deal of resonance structure in the excitation curves with the anomalies appearing at different energies at different angles: the more prominent structures in the yield curves are displayed in

table 16.16. For polarization measurements see (Aj 77). See also <sup>15</sup>O in (Aj 81), (No 81) and (Yo 77k; applications).

33. <sup>14</sup>N(d, p)<sup>15</sup>N $Q_m = 8.6087$  $E_b = 20.7363$ 

TABLE 16.16  
Structure in <sup>14</sup>N+d

$E_d$ (MeV)	Resonant channel	$\Gamma_{c.m.}$ (keV)	$J^\pi; T$	$E_x$ (MeV)	Refs. <sup>a</sup> )
1.4	$n_0$	12 ± 3	0 <sup>+</sup> ; 2	22.0	(Ko 77h)
1.7 ± 0.1	$\gamma_0, \alpha_0 \rightarrow \alpha_3$			22.2	
1.85	$n_0, \alpha_0$			22.35	
2.0 ± 0.1	$\alpha_0, \alpha_3$			22.5	
2.272 ± 0.005 <sup>b</sup> )	$p_0, p_{1+2}, (p_3)$			22.722	
2.40 ± 0.05 <sup>c</sup> )	$p_4, p_5, \alpha_0, \alpha_2$			22.83	
2.5	$\gamma_0$ <sup>d</sup> )			22.9	
2.6	$\alpha_0$			23.0	
2.8	$(n_0), \alpha_1$			23.2	
3.24	$(n_0), d_0$			23.57	
4.2	$p_0, p_{1+2}, p_4, p_5,$	600	1 <sup>-</sup> ; 1	24.4	(Ko 77h)
4.58	$p_6, d_0, \alpha_3$			24.74	
4.9	$\gamma_0, (p_0), d_0, \gamma_{15.1}$			25.0	
5.95	$(p_0), d_0, \gamma_{15.1}$			25.9	
7.1	$n_0, p_0$			26.9	
7.4	$d_1, \gamma_{15.1}$			27.2	
7.7	$\gamma_{15.1}$			27.5	
(8.5)	$d_2$			(28.2)	
10.2	$d_1$			29.7	
	$(\gamma_{15.1})$				

<sup>a</sup>) For earlier references see table 16.14 in (Aj 77).<sup>b</sup>)  $(\Gamma_{d_0} \Gamma_i / \Gamma^2) \times 10^{-3}$  are greater than  $1.6 \pm 0.4$ ,  $0.27 \pm 0.13$ ,  $0.41 \pm 0.15$  and  $0.07 \pm 0.05$  for the  $\alpha_2$ ,  $p_0$ ,  $p_{1+2}$  and  $p_3$  groups.<sup>c</sup>) If this resonance is fitted with a single-level Breit-Wigner shape, penetrability effects could lower the resonance energy by as much as 50 keV, assuming  $l=1$  (We 72a).<sup>d</sup>) The angular distribution of  $\gamma_0$  is consistent with E1.

The yield of various proton groups for  $E_d < 5.0$  MeV shows some fluctuations and two resonances: see table 16.16 and (Aj 71, Aj 77) for the earlier references. Recent work includes that of (Va 76h; 0.309 to 0.638 MeV;  $p_0, p_{1+2}, p_3, p_4, p_5$ ), (Ni 80; 0.32 to 1.45 MeV;  $p_5$ ), (Us 79;  $E_d = 1.50$  to 3.00 MeV;  $p_5$ ) and (Ko 77h;  $E_d = 0.5$  to 5.0 MeV;  $p_0, p_{1+2}, p_3 \rightarrow p_8, p_{10}$ ). Polarization measurements are also reported by (Kr 80;  $E_d = 10$  MeV –  $p_0, p_{1+2}, p_3, p_7, p_8$ ; 9.5 to 10.75 MeV –  $p_4, p_5, p_6$ ). See also (Aj 77) and <sup>15</sup>N in (Aj 81).

34. <sup>14</sup>N(d, d)<sup>14</sup>N $E_b = 20.7363$

The yield of elastically scattered deuterons has been studied for  $E_d = 0.65$  to  $5.5$  MeV and for  $14.0$  to  $15.5$  MeV: see (Aj 71, Aj 77). There is indication of broad structure at  $E_d = 5.9$  MeV and of sharp structure at  $E_d = 7.7$  MeV in the total cross section of the  $d_1$  group to the  $T = 1$  (isospin-forbidden),  $J'' = 0^+$  state at  $E_x = 2.31$  MeV in  $^{14}\text{N}$ . The yield of deuterons ( $d_2$ ) to  $^{14}\text{N}^*(3.95)$  [ $J'' = 1^+$ ,  $T = 0$ ] shows gross structures at  $E_d = 7.4$  and  $10.2$  MeV (Du 70): see table 16.16. The yield of  $d_1$  has also been studied for  $E_d = 10.0$  to  $17.9$  MeV by (Ao 79). For polarization measurements, see (Aj 77) and (Kr 80:  $E_d = 10$  MeV;  $d_0$ ). See also  $^{14}\text{N}$  in (Aj 81) and (Iz 77, Iz 77b, Se 79; theor.).

35. (a) $^{14}\text{N}(d, t)^{13}\text{N}$	$Q_m = -4.2963$	$E_b = 20.7363$
(b) $^{14}\text{N}(d, {}^3\text{He})^{13}\text{C}$	$Q_m = -2.0571$	

For polarization measurements see (Aj 77) and (Ma 81). See also  $^{13}\text{C}$ ,  $^{13}\text{N}$  in (Aj 81).

36. $^{14}\text{N}(d, \alpha)^{12}\text{C}$	$Q_m = 13.5743$	$E_b = 20.7363$
---	-----------------	-----------------

There is a great deal of structure in the yields of various  $\alpha$ -particle groups for  $E_d = 0.5$  to  $12$  MeV: see table 16.16 in (Aj 71) for the earlier work, and (Ko 77h:  $0.5$  to  $5.0$  MeV;  $\alpha_0 \rightarrow \alpha_3$ ,  $\alpha_5 \rightarrow \alpha_7$ ). The latter group reports broad oscillations ( $\Gamma \sim 0.5$  MeV) in the  $\alpha_0$  and  $\alpha_1$  yields for  $E_d = 2.0$  to  $5.0$  MeV. In addition,  $^{16}\text{O}^*(23.54)$  is reflected in the  $\alpha_3$  yield: see table 16.16. The yield of  $15.11$  MeV  $\gamma$ -rays [from the decay of  $^{12}\text{C}^*(15.11)$ ,  $J'' = 1^+$ ,  $T = 1$ ] which is isospin-forbidden has been studied for  $E_d = 2.8$  to  $12$  MeV. Pronounced resonances are observed at  $E_d = 4.2$ ,  $4.58$  and  $5.95$  MeV and broader peaks occur at  $E_d = 7.1$  and, possibly, at  $8.5$  MeV. See also  $^{12}\text{C}$  in (Aj 80).

For polarization measurements, see (De 79m:  $E_d = 1.5$  to  $3.0$  MeV;  $\alpha_0$ ,  $\alpha_1$ ), (Kr 80h, Kr 81:  $E_d = 10$  MeV;  $\alpha_0 \rightarrow \alpha_5$ ) and (Aj 77). See also (Se 79; theor.).

37. $^{14}\text{N}(t, n)^{16}\text{O}$	$Q_m = 14.4790$
--	-----------------

See (Aj 77).

38. (a) $^{14}\text{N}({}^3\text{He}, p)^{16}\text{O}$	$Q_m = 15.2428$
(b) $^{14}\text{N}({}^3\text{He}, p\alpha)^{12}\text{C}$	$Q_m = 8.0808$

Observed proton groups are displayed in table 16.17. Angular distributions have been measured at  $E({}^3\text{He}) = 2.5$  to  $24.7$  MeV: see (Aj 77) for the earlier work and (Sl 79a:  $9.8$  MeV;  $p_0$ ), (Bi 78j, Fo 78r:  $15$  MeV; most states in table 16.17) and (Ma 77y:  $24.7$  MeV;  $p$  to  $^{16}\text{O}^*(0, 6.13, 16.21, 17.14)$ ). Branching ratios and  $\tau_m$  measurements are shown in tables 16.11 and 16.12.

At  $E({}^3\text{He}) = 8$  MeV a study of the protons in coincidence with  $4.4$  MeV  $\gamma$ -rays (reaction (b)) indicates that the reaction proceeds via  $^{16}\text{O}^*(12.51, 13.97, 14.39, 14.92, 15.82, 16.23, 17.82, 18.04)$  [ $\pm 40$  keV] (Ho 69h). In a kinematically

TABLE 16.17  
<sup>16</sup>O states from <sup>14</sup>N(<sup>3</sup>He, p)<sup>16</sup>O

$E_x^{a,b)}$ (MeV $\pm$ keV)	$\Gamma_{c.m.}^{a,b)}$ (keV)	$L^c)$	$J^\pi c)$
0		0 + 2 <sup>d)</sup>	
6.052 $\pm$ 5		(0) <sup>e)</sup>	
6.131 $\pm$ 4		1 + 3 <sup>d)</sup>	
6.916 $\pm$ 3		(0)	
7.115 $\pm$ 3		1 + 3 <sup>d)</sup>	
8.870 $\pm$ 3	<20	3 + 1	
9.614 $\pm$ 30	510 $\pm$ 60		
9.847 $\pm$ 3	<20	0(+2)	
10.356 $\pm$ 3	25 $\pm$ 5	<sup>e)</sup>	
10.957 $\pm$ 1	<12	1 <sup>d)</sup>	
11.080 $\pm$ 3	<12 <sup>f)</sup>	2 + 4 <sup>f)</sup>	
11.098 $\pm$ 2	<12 <sup>f)</sup>		
11.520 $\pm$ 4	64 $\pm$ 5	<sup>e)</sup>	
12.049 $\pm$ 2	<12	0	
12.438 $\pm$ 3	70 $\pm$ 10	1	
12.530 $\pm$ 2	<12	1 + 3 <sup>d)</sup>	
12.797 $\pm$ 4	40 $\pm$ 10	1	0 <sup>-</sup> ; $T = 1^{\text{g})}$
12.970 $\pm$ 1	<12	1 + 3	2 <sup>-</sup> ; $T = 1^{\text{g})}$
13.105 $\pm$ 15	160 $\pm$ 30	0 + 3 <sup>f)</sup>	
13.257 $\pm$ 2	20 $\pm$ 5	(1 + 3)	3 <sup>-</sup> ; $T = 1^{\text{g})}$
13.663 $\pm$ 4	63 $\pm$ 7	0	
13.869 $\pm$ 2	85 $\pm$ 20	(4) <sup>e)</sup>	
13.979 $\pm$ 2	14 $\pm$ 5	1(+3)	
14.302 $\pm$ 3	<20	<sup>e)</sup>	
14.399 $\pm$ 2	27 $\pm$ 5	(4)	
14.818 $\pm$ 3		2	
14.927 $\pm$ 2	60 $\pm$ 10	0(+2)	(0 $\rightarrow$ 4) <sup>+</sup> (0, 1, 2) <sup>+</sup> <sup>h)</sup>
15.103 $\pm$ 5			
15.196 $\pm$ 3		(0 + 2)	
15.409 $\pm$ 6		<sup>e)</sup>	
15.785 $\pm$ 5	40 $\pm$ 10	2(+4)	(2, 3, 4) <sup>+</sup> <sup>h)</sup>
16.114 $\pm$ 4 <sup>j)</sup>			
16.209 $\pm$ 2	40 $\pm$ 10	0 + 2	
16.350 $\pm$ 13			
16.440 $\pm$ 3	$\sim$ 30	0 + 2	
16.817 $\pm$ 2	70 $\pm$ 10		

<sup>a)</sup> (Br 64f):  $E(^3\text{He}) = 3.74$  and  $3.97$  MeV.<sup>b)</sup> (Bi 78j):  $E(^3\text{He}) = 15$  MeV. <sup>c)</sup> (Bi 78j, Fo 78l, Fo 78r).<sup>d)</sup> See also (We 71q).<sup>e)</sup> Mostly compound nucleus. <sup>f)</sup> Unresolved.<sup>g)</sup> (Fo 78r) have compared the cross-section ratios of these three  $T = 1$  states with their analogs in <sup>16</sup>N populated in the (t, p) reaction: only the  $2^-$  states have the expected cross-section ratio of 0.5 for (<sup>3</sup>He, p)/(t, p). The populations of the  $0^-$  and  $3^-$  states in <sup>16</sup>O are lower by a factor of two.<sup>h)</sup> (Fo 78l) suggest that these two states [<sup>16</sup>O\*(14.93, 15.79)] are  $1^+$  and  $3^+2p-2h$  states with  $T_p = T_h = 0$ .<sup>i)</sup> Very weak proton group. I am indebted to Prof. H. T. Richards for his comments.

complete experiment at  $E(^3\text{He}) = 13 \text{ MeV}$ ,  $^{16}\text{O}^*(14.94, 18.04)$  are strongly populated (Fr 80j; prelim.). See also (Ha 79h).

39. (a) $^{14}\text{N}(\alpha, d)^{16}\text{O}$	$Q_m = -3.1104$
(b) $^{14}\text{N}(\alpha, d\alpha)^{12}\text{C}$	$Q_m = -10.2724$

Angular distributions to states of  $^{16}\text{O}$  have been reported at many energies to  $E_\alpha = 48 \text{ MeV}$ : see (Aj 71, Aj 77). Among the states reported by (Lo 72b) [see table 16.17 in (Aj 77)] are  $^{16}\text{O}^*(11.094 \pm 3, 14.400 \pm 3, 14.815 \pm 2, 17.18 \pm 50) [\text{MeV} \pm \text{keV}]$ : the results of (Lo 72b) are consistent with  $J'' = 5^+, 6^+, 4^+$  for  $^{16}\text{O}^*(14.40, 14.82, 16.29)$  [2p-2h] and with  $6^+$  for  $^{16}\text{O}^*(16.30)$  [4p-4h]. (Cl 79d) report  $\Gamma_{\text{c.m.}} = 34 \pm 12, 27 \pm 5$  and  $70 \pm 8 \text{ keV}$ , respectively for  $^{16}\text{O}^*(14.31 \pm 10, 14.40 \pm 10, 14.81)$ . For reaction (b) see (Aj 77).

40. $^{14}\text{N}(^6\text{Li}, \alpha)^{16}\text{O}$	$Q_m = 19.2628$
---	-----------------

Angular distributions have been measured at  $E(^6\text{Li}) = 5.3$  to  $6.0 \text{ MeV}$  and  $E(^{14}\text{N}) = 27.6 \text{ MeV}$ : see (Aj 77).

41. $^{14}\text{N}(^{10}\text{B}, ^8\text{Be})^{16}\text{O}$	$Q_m = 14.7104$
--	-----------------

See (Wu 78c). See also (Hi 77).

42. (a) $^{14}\text{N}(^{11}\text{B}, ^9\text{Be})^{16}\text{O}$	$Q_m = 4.9204$
(b) $^{14}\text{N}(^{13}\text{C}, ^{11}\text{B})^{16}\text{O}$	$Q_m = 2.0576$

At  $E(^{14}\text{N}) = 50 \text{ MeV}$  angular distributions are reported to  $^{16}\text{O}^*(0, 6.1(\text{u}), 7.0(\text{u}), 8.87, 9.85, 10.3(\text{u}), 11.0 (\text{u}))$  [ $\text{u} = \text{unresolved}$ ] (Re 77b). For both reactions see (Pr 80c). See also (Aj 77).

43. $^{15}\text{N}(\text{p}, \gamma)^{16}\text{O}$	$Q_m = 12.1276$
--	-----------------

The yield of ground state radiation ( $\gamma_0$ ) has been measured for  $E_\text{p} = 0.15$  to  $27.4 \text{ MeV}$ : see table 16.18 in (Aj 77) and (Sn 79:  $E_\text{p} = 4$  to  $9 \text{ MeV}$ ), (Ch 77j: 6 to  $22 \text{ MeV}$ ; also  $\gamma_2$ ), (Ca 77l:  $E_\text{p} = 8.6$  to  $11 \text{ MeV}$ ) and (Oc 78a; 8.6 to  $18 \text{ MeV}$ ; also  $\gamma_{1+2+3+4}$  for  $12.7 \rightarrow 18 \text{ MeV}$ ). Table 16.18 displays the parameters of the observed resonances. The cross section shows a great deal of structure up to  $E_\text{p} = 17 \text{ MeV}$ . Above that energy the  $\gamma_0$  yield decreases monotonically. Besides the GDR which peaks at  $^{16}\text{O}^*(22.15)$  (Oc 78a), the giant M2 resonance, based on  $^{16}\text{O}^*(6.13)$ , is identified to be  $^{16}\text{O}^*(20.43)$  (Ch 77j), and there is evidence for the emergence of a giant structure (E2) with  $E_\text{x} = 24-29 \text{ MeV}$  in the  $\gamma_{1+2+3+4}$  yield (Oc 78a). (Sn 79) report that the total M1 transition strength, built on the 2p-2h ground state correlations,  $B(\text{M1}) \geq 0.24 \text{ n.m.}^2$  for  $^{16}\text{O}^*(16.22, 17.14, 18.8)$ .

A study of the M1 decays of  $^{16}\text{O}^*(16.22, 17.14)$  [both  $J'' = 1^+; 1$ ] to  $^{16}\text{O}^*(6.05)$  finds  $B(\text{M1}, 1^+ \rightarrow 0_2^+)/B(\text{M1}, 1^+ \rightarrow 0_1^+) \sim 0.7$  for both transitions. The M1 (E2)  $\gamma$ -decay of  $^{16}\text{O}^*(18.03, 18.98)$  [ $J'' = 3^-; 1$  and  $4^-; 1$ , respectively] to  $^{16}\text{O}^*(6.13)$  [ $J'' = 3^-$ ] are also reported by (Ik 80). See also (Sn 81).

TABLE 16.18  
Levels of  $^{16}\text{O}$  from  $^{15}\text{N}(\text{p}, \gamma)$ ,  $^{15}\text{N}(\text{p}, \text{p})$  and  $^{15}\text{N}(\text{p}, \alpha)$

No.	$E_p$ (keV)	$\Gamma_{\nu}^{(s)}$ (eV)	$\Gamma_{\nu}^{(s)}$ (eV)	$\Gamma_p^{(s)}$ (keV)	$\Gamma_p \Gamma_w / \Gamma$ (eV)	$\Gamma_{\alpha}^{(s)}$ (keV)	$\Gamma_{\alpha}^{(s)}$ (keV)	$\Gamma_{\text{lab}}$ (keV)	$J^{\pi}; T$	$E_x$ (MeV $\pm$ keV)	Refs. <sup>b</sup>	
1	338	$12 \pm 2$	$0.12 \pm 0.04$	1.2		95	0.025	96	$1^-; 0$	12.444		
2	$429 \pm 1$	$(21 \pm 6) \times 10^{-3}$	$2.1 \pm 2$	0.02	40	nr	0.90	0.9	$2^-; 0$	12.530		
3	$710 \pm 7$	$(78 \pm 16) \times 10^{-3}$		1.2		nr	$40 \pm 4$	$40 \pm 4$	$0^-; 1$	12.793	(Br 77s)	
4	$897.37 \pm 0.29$	$(78 \pm 16) \times 10^{-3}$	$32 \pm 5$	100	40	nr	$0.69 \pm 0.07$	$2.0 \pm 0.2$	$2^-; 1$	12.9685	(Br 77s)	
5	$1028 \pm 10$					40	$140 \pm 10$	$140 \pm 10$	$1^-; 1$	13.091	(Br 77s)	
6	$1050 \pm 150$					r			$2^+$	13.1		
7	$1210 \pm 3$			4.1		8.2 $\pm$ 1.1	$22.5 \pm 1$	$3^-; 1$	13.261			
8	$1640 \pm 3$	$=8.5$		10		59 $\pm$ 6	$68 \pm 3$	$1^+; 0$	13.664			
9	$1890 \pm 20$			0.5		r	$90 \pm 2$			13.90		
10	$1979 \pm 3$			r		nr	$23 \pm 2$	$2^-$		13.982		
11	$2977 \pm 10$			33		1.5	30	$69 \pm 10$	$2^+$	$14.917 \pm 5^c$	(Ja 77j, Fr 78b)	
12	$3170^{\text{j}}$					152	163	$327 \pm 100$	$0^+$	$15.10 \pm 50$	(Fr 78b)	
13	$3260^{\text{j}}$					nr	7	$62 \pm 10$	$2^-$	$15.188 \pm 30^c$	(Fr 78b)	
14	$3340^{\text{j}}$					12	182	$315 \pm 100$	$2^+; (0)$	$15.26 \pm 20^c$	(Fr 78b)	
15	$3520 \pm 40$					1	103	$167 \pm 20$	$3^-$	$15.43^c$	(Ja 77j, Fr 78b)	
16	$4380 \pm 20$	3.6	r	16	$2.65 \pm 0.22$			19 $\pm$ 3	$1^+; 1$	16.23	(Sn 79, Ik 80)	
17	$4620 \pm 20$					r				16.46	(Ja 77j)	
18	$5010 \pm 20$					nr	r	$75 \pm 10$	$3^+$	$16.82$	(Ja 77j)	
19	$5200$	r						$\sim 1500$	$1^-; 1$	17.0		
20	$5350 \pm 20$	6.5	r	26	$3.75 \pm 0.50$			$38 \pm 5$	$1^+; 1$	17.14	(Sn 79, Ik 80)	
21	$5490 \pm 20$	67	r					$\sim 110$	$1^-; 1$	17.27		
22	$6290 \pm 20$	nr	$5 \pm 2 [4 \pm 2\%]$		$\leq 15^f$			$\leq 40$	$3^-; 1$	18.02	(Ch 77j, Ik 80)	
23	$7310 \pm 20$		$\leq 2^{\circ\circ}$		$\leq 40^f$			$\leq 40$	$4^-; 1$	18.98	(Ch 77j, Ik 80)	
24	$7330 \pm 30$	38						$\sim 260$		18.99	(Sn 79)	
25	$7420$	r						$\sim 130$	$\geq 1.8 \pm 0.3$	$\sim 19.08$	$2^+; (1)$	

## ENERGY LEVELS OF LIGHT NUCLEI A = 16-17

26	7600 ± 30	nr	59	1.5 <sup>b</sup> )	(r)				100 350 80 ± 30 200 ± 40 160 320 ± 10	(2,3;1) 1-;1 3 2-;1 20.55 1-;1	19.25 19.47 19.89 20.43 <sup>a</sup> ) 20.55 20.945 ± 20	(Ch 77i) (Ch 77j) (Oc 78a, Ca 77i)
27	7840 ± 30	nr	59	17 ± 6 <sup>c</sup> )	25 ± 10 <sup>d</sup> )							
28	8290 ± 20	nr	120 ± 45 <sup>e</sup> )	86 <sup>f</sup> )								
29	8860 ± 30	nr	170			21 ± 1						
30	8990											
31	9410 <sup>g</sup> )											
32	10 000 <sup>h</sup> )											
33	10 180 <sup>j</sup> )											
34	10 700 <sup>i,m</sup> )	r		27 <sup>h</sup> )								
35	11 490 <sup>j</sup> )	120			488 ± 20							
36	12 740 <sup>j</sup> )	r		85 <sup>f</sup> )	69 ± 5							
37	13 490 ± 60			230 ± 90, or 130 ± 50 <sup>e</sup> )	130 ± 13							
38	13 870 <sup>j</sup> )	r		740 ± 240, or 410 ± 140 <sup>e</sup> )	651 ± 117							
39	15 250 ± 80			1070 ± 380, or 590 ± 10 <sup>e</sup> )	122 <sup>f</sup> )							
40	16 250 ± 100				206 <sup>f</sup> )							

<sup>a</sup>) nr = non-resonant; r = resonant.<sup>b</sup>) For earlier references see tables 16.21 in (Aj 71) and 16.19 in (Aj 77).<sup>c</sup>) The values for  $\Gamma_{\nu\nu}$ ,  $\Gamma$ ,  $J^*$  and  $E_x$  are from a multilevel R-matrix analysis of  $p_0$ ,  $\alpha_0$  and  $\alpha_1$  excitation curves (Fr 78b): in addition to these states, others at  $E_x = 14.0$  [0<sup>+1</sup>] (fixed), 14.6 [1<sup>-1</sup>] and 16.2 MeV [1<sup>-1</sup>],  $\Gamma = 0.44$ , 0.68 and 1.23 MeV were included in the analysis.<sup>d</sup>) Not observed in  $p_0$  channel.<sup>e</sup>)  $\Gamma_{\nu\nu}$  (eV) (Ch 77i).<sup>f</sup>)  $\Gamma_{\nu\nu}$  based on  $\Gamma_{c.m.}$  and values of  $\Gamma_{\nu\nu}/\Gamma$  assumed by (Ch 77i).<sup>g</sup>) This state is attributed to the giant M2 resonance based on  $^{16}\text{O}^*(6,13) [J^* = 3^-]$  (Ch 77j).<sup>h</sup>)  $\gamma_1 + \gamma_2$ .<sup>i</sup>) Resonant in  $P_2$  (Dr 71b).<sup>j</sup>) Nominal  $E_p$  calculated from  $E_x$ .<sup>k</sup>) Resonant in  $P_1$  (Dr 71b).<sup>l</sup>) Resonant in  $P_0$ ,  $P_1$ ,  $P_6$  (Dr 71b).<sup>m</sup>)  $\sigma = 12.9$  mb at peak of GDR (Oc 78a).<sup>n</sup>) Average of values obtained in this experiment and in  $^{12}\text{C}(\alpha, \gamma_2)$  (Ch 77i).<sup>o</sup>) Resonant in  $\gamma_2$ .<sup>p</sup>) Apparent resonance in yield of ( $p, \alpha\gamma_{1,2}$ ) (Oc 78a).

Below  $E_p = 0.4$  MeV capture to the ground state is dominant:  $S(0) = 64 \pm 6$  keV b, a value which makes the oxygen side cycle in CNO burning more important than previously thought. Study of the direct radiative capture process leads to a single-particle spectroscopic factor  $C^2 S = 1.8 \pm 0.4$  for <sup>16</sup>O<sub>g.s.</sub> (Ro 74s).

TABLE 16.19  
Resonances in <sup>15</sup>N(p, n)<sup>15</sup>O<sup>a)</sup>

$E_p$ (MeV ± keV)	$\Gamma_{c.m.}$ (keV)	$J^\pi; T^b)$	$E_x$ (MeV)
$4.37 \pm 15$	$19 \pm 6$	$1^{(+)}; 1$	16.22
$4.45 \pm 30$	$240 \pm 30$	$0^{(-)}$	16.30
$5.35 \pm 15$	$33 \pm 5$	$1^{(-)}; 1$	17.14
$5.52 \pm 15$	$90 \pm 10$	$1^-; 1$	17.30
$5.88 \pm 15$	$59 \pm 10$	$\geq 1; 1$	17.64
$6.12 \pm 15$	$101 \pm 10$	$\geq 1; 1$	17.86
$6.23 \pm 15$ <sup>c)</sup>	$\leq 50$	$T = 1$	17.96
$6.33 \pm 15$	$26 \pm 5$	$\geq 1; 1$	18.06
$6.43 \pm 30$	$\approx 300$		18.15
$6.76 \pm 25$	$\approx 160$		18.46
$7.03 \pm 30$	$260 \pm 30$		18.71
$7.59 \pm 25$	$90 \pm 10$	$2^-; 1$	19.24
$7.86 \pm 30$	$300 \pm 80$		19.49
$8.30 \pm 25$	$120 \pm 40$		19.90
$8.88 \pm 40$ <sup>d,e)</sup>	$200 \pm 50$	2	20.45
$9.08 \pm 40$ <sup>c)</sup>	$130 \pm 50$		20.63
$9.42 \pm 100$ <sup>c)</sup>	$235 \pm 45$		20.95
$10.73 \pm 100$ <sup>c)</sup>	$800 \pm 95$	1	22.18
$11.01 \pm 100$	$300 \pm 100$		22.44
$11.92 \pm 100$	$520 \pm 200$		23.29
$13.03 \pm 100$	$520 \pm 100$		24.33
$13.63 \pm 100$	$\sim 280$	2, 4	24.89
$15.12 \pm 100$	$610 \pm 140$	2, 4	26.29
$18.4 \pm 200$	$470 \pm 150$		29.4

<sup>a)</sup> First fourteen resonances are from (Ba 68qq); the higher energy resonances are from (Ch 78k; n<sub>0</sub>).

<sup>b)</sup> Assignments are from (p, n) and (p,  $\gamma$ ) results. The  $T$ -assignments are made on the basis of energy and width comparisons with states of <sup>16</sup>N.

<sup>c)</sup> Probably a doublet: see (Ba 68qq).

<sup>d)</sup> Values of  $(2J+1) \Gamma_{po} \Gamma_{no} / \Gamma^2$  are derived for this resonance and the ones below: see (Ch 78k).

<sup>e)</sup> See also (Ba 68qq).

For branching ratios and  $\Gamma_\gamma$  values for the low-energy resonances see table 16.12. See also (Wi 78a, Wi 79c), (Ha 75dd, Pa 75o, Ha 76t, Gl 79k, Ha 79k, Sn 79a, We 80d, Wi 81d), (Zi 75c, Bo 76cc, Ro 77aa, Ro 78b, Ba 80k; astrophysics) and (Ba 77f, Ho 77s, Ba 78aa, Ga 78b; theor.).

TABLE 16.20  
States in <sup>16</sup>O from <sup>15</sup>N(d, n), <sup>15</sup>N(<sup>3</sup>He, d), <sup>17</sup>O(d, t) and <sup>17</sup>O(<sup>3</sup>He,  $\alpha$ )

<sup>16</sup> O* (MeV)	$J^\pi; T$	$I^a)$	$I^b)$	$S_c$	$I^e)$	$C^2 S^e)$	$I^f)$	$S^f)$
0	$0^+; 0$	1	1	3.1	2	0.74	2	0.88
6.05	$0^+; 0$		1	<sup>d)</sup>			2	0.009
6.13	$3^-; 0$	2	2		1	0.46	1	0.37
6.92	$2^+; 0$	not direct	1+3	<sup>d)</sup>	obs.		(2+0)	0.022
7.12	$1^-; 0$	0	0+2		1	0.04	(3+1)	0.007
8.87	$2^-; 0$	2	2	0.72	1	0.33	1	0.26
9.63	$1^-; 0$		0	<sup>d)</sup>				
9.85	$2^+; 0$	1	not direct	<sup>d)</sup>			2	0.025
10.35	$4^+; 0$		3	<sup>d)</sup>			2	0.025
10.95	$0^-; 0$	0	0	0.76			(3+1)	0.008
11.08	$3^+; 0$	3	3	0.18			2	0.044 or 0.086
11.26	$0^+; 0$		broad					
12.44	$1^-; 0$	0	0	0.40				
12.53	$2^+; 0^j)$	2	2	0.72	1	0.07		
12.80	$0^-; 1$	0	0	0.44				
12.97	$2^-; 1^j)$	2	2	0.40	1	0.69	1	0.38
13.10	$1^-; 1$	(0)		0.58			1	0.10
13.13 <sup>a)</sup>	$3^-; 0$	(2)		0.32				
13.25	$3^-; 1$	2	2	0.46	1	0.70	1	0.34
15.22	$2^-; 0^e)$				1	0.12		
15.42	$3^-; 0^e)$				1	0.37		
17.14			obs.					
17.20	$2^+$		obs.					
17.788 $\pm$ 16 <sup>i)</sup>	$4^-; 0$					0.17		
18.033 $\pm$ 10 <sup>i)</sup>	$3^+; 1^h)$				(1)	0.12		
<sup>16</sup> O* (MeV $\pm$ keV)	$J^\pi; T$	$I^e)$	$C^2 S^e)$	$I^f)$	$S^f)$	$\Gamma^j)$ (keV)		
18.48	$T=1$	(1)	0.25					
18.975 $\pm$ 10 <sup>i)</sup>	$4^-; 1$	1	0.73					
19.206 $\pm$ 12 <sup>i)</sup>	$3^-; 1^h)$	1	0.50			68 $\pm$ 10		
19.802 $\pm$ 16 <sup>i)</sup>	$4^-; 0$	1	0.52			36 $\pm$ 5		
20.45	$(2, 4)^-; 1$	1	0.21					

<sup>a)</sup> <sup>15</sup>N(d, n);  $E_d = 4.8$  to 6 MeV; see (Aj 77) for references.

<sup>b)</sup> <sup>15</sup>N(<sup>3</sup>He, d);  $E(^3\text{He}) = 11, 16.0$  and 24.0 MeV; see (Aj 77).

<sup>c)</sup> "Best" values as discussed by (Bo 73o) [from (d, n) and (<sup>3</sup>He, d) data]. See table 16.22 in (Aj 77) for a more complete display.

<sup>d)</sup> Very small value of  $S$ ; see (Aj 77).

<sup>e)</sup> <sup>17</sup>O(d, t);  $E_d = 52$  MeV (Ma 78p). See also (Ma 81d;  $E_d = 52$  MeV).

<sup>f)</sup> <sup>17</sup>O(<sup>3</sup>He,  $\alpha$ );  $E(^3\text{He}) = 11$  MeV (Bo 71b).

<sup>g)</sup>  $\Gamma = 128$  keV.

<sup>h)</sup> I am indebted to Prof. H. T. Richards for an illuminating discussion of the evidence for the parameters of this state. See also (Ma 81d).

<sup>i)</sup> (Br 80j): <sup>17</sup>O(<sup>3</sup>He,  $\alpha$ ).  $\Gamma < 50$  keV for <sup>16</sup>O\* (17.79).

<sup>j)</sup> See text (Wa 77l).

Elastic scattering studies are reported for  $E_p = 0.6$  to 15 MeV: see table 16.20 in (Aj 71), 16.18 in (Aj 77), (Ja 77j; 2.50 to 5.14 MeV) and (Fr 78b: 2.88 to 3.64 MeV). See also (La 76j). Observed anomalies are displayed in table 16.18. For inelastic groups see (Dr 71b). See also (Fa 80) and (Do 80d, Ha 80p; theor.).

$$45. \ ^{15}\text{N}(\text{p}, \text{n})^{15}\text{O} \quad Q_m = -3.5363 \quad E_b = 12.1276$$

Excitation functions and cross sections have been measured for  $E_p = 3.8$  to 19.0 MeV: see (Aj 77) and (Ch 78k; 8.5 to 19.0 MeV;  $n_0$ ). For a listing of observed resonances see table 16.19. Angular distributions of  $n_0$  have been studied for  $E_p = 5.52$  to 9.26 MeV (Mu 81) and 9.05 to 15.62 MeV (By 81). Polarization measurements are reported for the  $n_0$  group at  $E_p = 7$  to 17 MeV: see (Mu 81b, Wa 81a). See also (By 77, By 80, By 81b), (Wa 76p, By 79a, Wa 80l, Co 81, Wa 81b) and (Ba 77, Ho 77s, Ab 79e, Do 80d, Ha 80p, Ph 81a; theor.).

$$46. \begin{array}{ll} (\text{a}) \ ^{15}\text{N}(\text{p}, \text{t})^{13}\text{N} & Q_m = -12.9050 \\ (\text{b}) \ ^{15}\text{N}(\text{p}, {^3\text{He}})^{13}\text{C} & Q_m = -10.6658 \end{array} \quad E_b = 12.1276$$

The yields of the first three triton and  $^3\text{He}$  groups have been measured for  $E_p = 24.0$  to 43.5 MeV (Pi 74b, Mi 75). Polarized protons with  $E_p = 43.8$  MeV have been used to study the transitions to  $^{13}\text{C}^*(0, 3.68, 7.55, 15.11)$  and  $^{13}\text{N}^*(0, 3.51, 7.38, 15.07)$  (Ma 74c). See also  $^{13}\text{C}$ ,  $^{13}\text{N}$  in (Aj 81).

$$47. \ ^{15}\text{N}(\text{p}, \alpha)^{12}\text{C} \quad Q_m = 4.9656 \quad E_b = 12.1276$$

Excitation functions for  $\alpha_0$  and  $\alpha_1$  particles [corresponding to  $^{12}\text{C}^*(0, 4.43)$ ] and of 4.43 MeV  $\gamma$ -rays have been measured for  $E_p = 93$  keV to 45 MeV: see tables 16.20 in (Aj 71), 16.18 in (Aj 77) and (Zy 79a: 93 to 418 keV;  $\alpha_0$ ), (Br 77s: 0.9 to 1.25 MeV;  $\alpha_0, \alpha_1$ ), (Ja 77j: 2.5 to 5.14 MeV;  $\alpha_0, \alpha_1$ ) and (Fr 78b: 2.88 to 3.64 MeV;  $\alpha_0, \alpha_1$ ). The yield of 15.11 MeV  $\gamma$ -rays has been measured for  $E_p = 12.5$  to 17.7 MeV (Oc 78a). Observed resonances are displayed in table 16.18.

A study of the  $\alpha_0$  cross section for  $E_p = 93$  to 418 keV leads to  $S(0) = 78 \pm 6 \text{ MeV} \cdot \text{b}$  (Zy 79a). For the  $\alpha_1 \gamma$ -channel  $S(0) = 0.1 \text{ keV} \cdot \text{b}$  (Ro 74s). See also (Aj 77), (Cl 79d), (Ro 77aa, Ba 80k; astrophysics), (Th 78d; applications) and (Ga 77g; theor.).

$$48. \ ^{15}\text{N}(\text{d}, \text{n})^{15}\text{O} \quad Q_m = 9.9029$$

Observed neutron groups,  $l$ -values and spectroscopic factors are displayed in table 16.20. Angular distributions are reported for  $E_d$  to 6 MeV: see (Aj 77).

$$49. \ ^{15}\text{N}(^3\text{He}, \text{d})^{16}\text{O} \quad Q_m = 6.6340$$

Angular distributions have been measured at  $E(^3\text{He}) = 11, 16.0$  and 24.9 MeV [see (Aj 77)]:  $l$ - and  $S$ -values are shown in table 16.20.

$$50. \ ^{15}\text{N}(\alpha, \text{t})^{16}\text{O} \quad Q_m = -7.6865$$

At  $E_\alpha = 65$  MeV, the population of many <sup>16</sup>O states with  $E_x < 22$  MeV is reported by (Ya 80a).

$$51. \begin{array}{ll} (a) {}^{15}\text{N}({}^{11}\text{B}, {}^{10}\text{Be}){}^{16}\text{O} & Q_m = 0.8988 \\ (b) {}^{15}\text{N}({}^{13}\text{C}, {}^{12}\text{B}){}^{16}\text{O} & Q_m = -5.406 \end{array}$$

Angular distributions are reported at  $E({}^{11}\text{B}) = 115$  MeV to <sup>16</sup>O\*(6.13) and at  $E({}^{13}\text{C}) = 105$  MeV to <sup>16</sup>O\*(6.13, 8.87) (Pr 80c, Ra 79c).

$$52. {}^{16}\text{N}(\beta^-){}^{16}\text{O} \quad Q_m = 10.418$$

TABLE 16.21  
Beta decay of the ground state of <sup>16</sup>N\*)

Final state		Branch (%)	$\log f_{1t}^b)$
<sup>16</sup> O* (MeV)	$J^\pi$		
0	$0^+$	$26 \pm 2^c)$	$9.10 \pm 0.04^b)$
6.05	$0^+$	$(1.2 \pm 0.4) \times 10^{-2}^d)$	$9.96 \pm 0.15^b)$
6.13	$3^-$	$68 \pm 2^c)$	$4.47^h)$
7.12	$1^-$	$4.9 \pm 0.4^c)$	$5.09^h)$
8.87	$2^-$	$1.0 \pm 0.2^c)$	$4.37^h)$
9.63	$1^-$	$(1.20 \pm 0.05) \times 10^{-3}^e)$	$6.21^h)$
9.85	$2^+$	$(6.5 \pm 2.0) \times 10^{-7}^f)$	$9.07 \pm 0.13^i)$

\*) See also reaction 1 in <sup>16</sup>N.

b)  $\tau_{1/2} = 7.13 \pm 0.02$  sec: see table 16.3 in (Aj 71).

c) (Wi 56, Al 58c, Al 59f).

d) (Wa 68n).

e) (Ka 61c).

f) (Ha 69s).

g) (To 71h):  $\log f_{1t}$ .

h) B. Zimmerman, private communication:  $\log f_{0t}$ .

i) E. K. Warburton, private communication:  $\log f_{1t}$ .

The ground state of <sup>16</sup>N decays to seven states of <sup>16</sup>O: reported branching ratios are listed in table 16.21. The ground state transition has the unique first-forbidden shape corresponding to  $\Delta J = 2$ , yes, fixing  $J^\pi$  of <sup>16</sup>N as  $2^-$ : see (Aj 59). For the  $\beta$ -decay of <sup>16</sup>N\*(0.12) see reaction 1 in <sup>16</sup>N.

The  $\alpha$ -decay of <sup>16</sup>O\*(8.87, 9.63, 9.85) has been observed: see (Aj 71). The parity-forbidden  $\alpha$ -decay from the  $2^-$  state <sup>16</sup>O\*(8.87) has been reported:  $\Gamma_\alpha = (1.03 \pm 0.28) \times 10^{-10}$  eV [ $E_\alpha = 1282 \pm 5$  keV] (Ne 74d).

Transition energies derived from  $\gamma$ -ray measurements are:  $E_x = 6130.43 \pm 0.05$  keV [ $E_\gamma = 6129.170 \pm 0.043$  keV (Sh 75h)] and  $7116.85 \pm 0.14$  keV [ $E_\gamma = 7115.15 \pm 0.14$  keV (Al 76d)].

53. (a)  $^{16}\text{O}(\gamma, \text{n})^{15}\text{O}$   $Q_m = -15.6639$   
 (b)  $^{16}\text{O}(\gamma, 2\text{n})^{14}\text{O}$   $Q_m = -28.8882$   
 (c)  $^{16}\text{O}(\gamma, \text{pn})^{14}\text{N}$   $Q_m = -22.9609$

The absorption cross section and the  $(\gamma, \text{n})$  cross section are marked by a number of resonances. On the basis of monoenergetic photon data, excited states of  $^{16}\text{O}$  are observed at  $E_x = 17.3$  [u], 19.3 [u] and 21.0 MeV [u = unresolved], followed by the giant resonance whose principal structures are at  $\approx 22.2$  and 24.1 MeV, with additional structures at 23 and 25 MeV: see (Be 76w, Aj 77). The cross section for  $(\gamma, \text{n}_0)$  decreases monotonically for  $E_\gamma = 25.5$  to 43.8 MeV. There is a significant E2 cross section in that region: it exhausts  $\sim 68\%$  of the isovector E2 energy-weighted sum rule (Ph 79d). The differential  $(\gamma, \text{n}_0)$  cross sections have been measured for  $E_\gamma = 60$  to 160 MeV (Go 80l).

The absorption cross section has been measured from  $E_{ba} = 10$  MeV to above the meson threshold (Ah 75b). The yield of reaction (b) has been studied from threshold to  $E_\gamma = 120$  MeV [see (Aj 77)] and for  $E_{ba} = 100$  to 800 MeV (Jo 77). The  $(\gamma, \text{Tn})$  cross section has recently been reported for  $E_\gamma = 15.9$  to 39.7 MeV (Ju 80). See also (Be 76w). The polarization of  $\text{n}_0$  has been studied for  $E_\gamma = 21.6$  to 25.7 MeV (Na 79e). See also (Aj 77). For the decay of  $^{16}\text{O}^*$  to  $^{15}\text{O}$  states, see  $^{15}\text{O}$  in (Aj 81). See also (Kn 75d), (Be 76w, Ki 76k, Da 77e, Ha 79k, Ca 80m, Sc 80d, Sc 81g) and (Ba 76kk, Ka 76q, Gr 77b, Ma 77oo, Co 78h, El 78a, Ka 78t, Sc 78e, Ma 79e, Ro 79d, Ro 79l, Bo 80o, Du 80d, Os 80c, Ta 80d, Ba 81a, Bo 81p, Ca 81d, Ko 81g, Ro 81j, We 81e; theor.).

54. (a)  $^{16}\text{O}(\gamma, \text{p})^{15}\text{N}$   $Q_m = -12.1276$   
 (b)  $^{16}\text{O}(\gamma, \pi^+)^{16}\text{N}$   $Q_m = -149.986$

The  $(\gamma, \text{p}_0)$  cross section derived from the inverse capture reaction (reaction 43) confirms the giant resonance structure indicated above, in reaction 53, as do also the direct  $(\gamma, \text{p}_0)$  measurements. The total  $(\gamma, \text{p})$  cross section is given by (De 68g). Recent measurements of the  $(\gamma, \text{p}_0)$  cross section are reported by (Fi 77b:  $E_{ba} = 40$  to 105 MeV), (Sc 80i:  $E_\gamma = 80$  MeV; forward angles) and (Ma 77:  $E_{ba} = 100$  to 280 MeV). See also (Aj 77) and (Kh 80, Ki 80e). The momentum distribution for  $p_{1/2}$  shell protons in  $^{16}\text{O}$  up to 930 MeV/c has been derived by (Fi 78). See also (Ma 79ff). Branching ratios for the decays of  $^{16}\text{O}$  states in the giant resonance region to various excited states in  $^{15}\text{N}$  are discussed in  $^{15}\text{N}$  (Aj 81).

The cross section for reaction (b) for  $E_{ba} = 175$  MeV is reported by (Bo 79ee). See also (Be 77aa, Ya 81). For other  $\pi$ -emitting photonuclear reactions see (Ep 79a, Bo 80q) and p. 23. See also (Ha 76t, We 80d, Ha 81l) and (Ba 76kk, Ka 76q, Ne 77d, Fi 78c, Ro 78j, Sc 78e, Ko 79p, Le 79h, Lo 79b, Wa 79g, Mo 80o, Bo 81h, Bo 81p, Ca 81d, Co 81m; theor.).

55. (a)  $^{16}\text{O}(\gamma, \text{d})^{14}\text{N}$   $Q_m = -20.7363$   
 (b)  $^{16}\text{O}(\gamma, \text{t})^{13}\text{N}$   $Q_m = -25.033$   
 (c)  $^{16}\text{O}(\gamma, \alpha)^{12}\text{C}$   $Q_m = -7.1620$

For the earlier work on these reactions see (Aj 71, Aj 77). For reaction (a) see also (Wi 79c). A study of the  $^{16}\text{O}(\gamma, \alpha_0)$  reaction at  $\theta = 45^\circ$  and  $90^\circ$  shows a  $2^+$  resonance at  $E_x = 18.2$  MeV with an E2 strength which is spread out over a wide energy interval. A strong resonance corresponding to an isospin-forbidden  $1^-$  state at  $E_x \approx 21.1$  MeV is also observed (Sk 75a). For spallation see (Di 78f). See also (Ch 81g).

### 56. $^{16}\text{O}(\gamma, \gamma)^{16}\text{O}$

(Ah 70a) report resonances at  $E_\gamma = 22.5 \pm 0.3$ ,  $25.2 \pm 0.3$ ,  $31.8 \pm 0.6$  and  $50 \pm 3$  MeV: the dipole sum up to 80 MeV exceeds the classical value  $60 \text{ NZ}/A \text{ MeV} \cdot \text{mb}$  by a factor 1.4. See also (Is 80e). The integrated cross section to 31 MeV for elastic scattering is  $0.62 \pm 0.07 \text{ MeV} \cdot \text{mb}$ ; the inelastic is  $\leq 5\%$  of the elastic cross section (Is 80b). For widths of excited states of  $^{16}\text{O}$  see table 16.22. The separation between the  $^{16}\text{O}^*(7.12, 6.92)$   $\gamma$ -lines is  $199.8 \pm 0.5$  keV (Sw 70). Based on  $7116.85 \pm 0.14$  keV (table 16.11),  $E_x$  for the lower state is  $6917.11 \pm 0.6$  keV.

- |   |                  |
|---|------------------|
| 57. (a) $^{16}\text{O}(\text{e}, \text{e})^{16}\text{O}$  |                  |
| (b) $^{16}\text{O}(\text{e}, \text{ep})^{15}\text{N}$     | $Q_m = -12.1276$ |
| (c) $^{16}\text{O}(\text{e}, \text{e}\pi^+)^{16}\text{N}$ | $Q_m = -149.986$ |
| (d) $^{16}\text{O}(\text{e}, \text{e}\pi^-)^{16}\text{F}$ | $Q_m = -154.996$ |

The  $^{16}\text{O}$  charge radius =  $2.710 \pm 0.015$  fm (Ki 78a). Form factors for transitions to the ground and to excited states of  $^{16}\text{O}$  have been reported in many studies: see (Aj 77) and (Mi 79d:  $q = 0.5$  to  $2.6 \text{ fm}^{-1}$ ; elastic). Table 16.22 lists the excited states observed from ( $\text{e}, \text{e}'$ ). The isospin-forbidden (E1) excitation of  $^{16}\text{O}^*(7.12)$  has been reported by (Mi 75i, Mi 75l): the isovector contribution interferes destructively with the isoscalar part and has a strength  $\approx 1\%$  of the  $T = 0$  amplitude (Mi 75i). The  $0^+$  states  $^{16}\text{O}^*(6.05, 12.05, 14.00)$  saturate  $\approx 19\%$  of an isoscalar monopole sum rule (Be 73gg). As for the E2 strength it is distributed over a wide energy region: see table 16.22. The effective mass of an intranuclear nucleon has been derived by (De 78s) from studies on this and on other light nuclei. See also (De 80gg).

Reaction (b) has been studied at  $E_e = 500$  MeV: two peaks at  $E = 11.5$  and 18 MeV carry most of the strength of the  $1p_{1/2}$  and  $1p_{3/2}$  hole states, respectively (Tu 77a). See also (Be 77bb, Ba 80t). The virtual photon shape and intensity-for ( $\text{e}, \text{e}'\pi$ ) has been deduced at  $E_e = 180.4$  MeV ( $E_\pi = 28$  MeV,  $\theta_\pi = 90^\circ$ ) (St 80d). The ratio  $\pi^-/\pi^+$  for  $E_e = 280$  MeV ( $E_\pi = 10$  MeV,  $\theta_\pi = 90^\circ$ ) is reported by (Je 79a). See also (Bu 76j, Sc 79b, Bu 80a, Bu 81a), (Tu 76a, Li 79h, Mo 79y, Ca 80h, Mo 80, Mo 81k) and (An 76c, Bh 76b, Fr 76e, Ga 76k, He 76k, Ra 76h, De 77f, He 77k, Wa 77o, Al 78f, Fu 78c, Ha 78n, Ha 78z, Ka 78r, Kr 78c, Ri 78, Bu 79j, Do 79o, In 79b, Kn 79, Ko 79ll, Li 79n, Wa 79g, Ro 79p, De 80z, Er 80b, Gr 80, Sa 80c, Sh 80n, Va 80b, Ag 81, Ba 81w, Bo 81m, Bu 81c, De 81p, Ka 81f, Or 81, Pe 81b; theor.).

TABLE 16.22  
Excited states observed in  $^{16}\text{O}(\text{e}, \text{e}')^{16}\text{O}^*$

$E_x$ (MeV $\pm$ keV)	$J''; T$	Mult.	$\Gamma$ (keV)	$\Gamma_\infty$ (eV)
6.05	$0^+$	E0		$3.55 \pm 0.21$ <sup>b)</sup>
6.13	$3^-$	E3		$(2.60 \pm 0.13) \times 10^{-5}$
6.92	$2^+$	E2		$0.130 \pm 0.009$
				$0.100 \pm 0.004$
7.12 <sup>c)</sup>	$1^-$	E1		$(4.6 \pm 2.3) \times 10^{-2}$
9.85	$2^+$	E2		$(8.8 \pm 1.7) \times 10^{-3}$
10.35	$4^+$	E4		$(5.6 \pm 2.0) \times 10^{-8}$
11.52	$2^+$	E2		$0.61 \pm 0.02$
12.05	$0^+$	E0		$4.03 \pm 0.09$ <sup>b)</sup>
12.53	$2^-$	M2		$0.021 \pm 0.006$
				$0.108 \pm 0.015$
12.97	$2^-$	M2		$0.071 \pm 0.002$
13.0	$2^+$	E2		0.89
$13.10 \pm 250$	$1^-; 1$	E1		$\leq 49 \pm 13$
$14.00 \pm 50$	$0^+$	E0	$170 \pm 50$	$3.3 \pm 0.7$ <sup>b)</sup>
$15.15 \pm 150$	$2^+$	E2	$500 \pm 200$	$1.0 \pm 0.5$
$16.21 \pm 30$	$1^+$	M1	$18 \pm 3$	$5.1 \pm 0.8$
$16.46 \pm 70$	$2^+$	E2	$35 \pm 5$	$0.5 \pm 0.2$
$16.80 \pm 100$	$(3^+)$		$\leq 100$	$(1.7 \pm 1.9) \times 10^{-3}$
17.14	$1^-; 1$	E1	$40 \pm 6$	$62 \pm 12$
$17.60 \pm 100$	$(2^-)$		$\leq 100$	$0.07 \pm 0.04$
<sup>d)</sup>				
18.50 $\pm$ 100	$2^+$	E2	$60 \pm 9$	
$19.00 \pm 100$	$1^-; 1$	E1	$300 \pm 100$	$41 \pm 20$
$19.04 \pm 50$	$2^-; 1$	M2	$400 \pm 50,$ $850 \pm 150$	$1.5 \pm 0.3$
$19.50 \pm 100$	$1^-; 1$	E1	$200 \pm 70$	$40 \pm 20$
$20.36 \pm 70$	$2^-$	M2	$500 \pm 100$	$2.9 \pm 1.0$
$20.95 \pm 50$	$1^-; 1$	E1	$270 \pm 70$	$180 \pm 50$
$21.34 \pm 250$	$(2^-)$	(M2)		
22.3				
23.0				
$23.7 \pm 250$	$(2^-; 1)$			
24.2				
$25.5 \pm 250$	$1^-; 1$	E1		
$26.7 \pm 250$	$1^+$	M1		
44.5	$(1^-; 1)$		$2000-3000$	5300
49	$(1^-; 1)$		$2000-3000$	19 000

<sup>a)</sup> See also table 16.26 in (Aj 71). For references see table 16.24 in (Aj 77).

<sup>b)</sup> Monopole matrix element in fm<sup>2</sup>.

<sup>c)</sup> See also text and (Mi 75I).

<sup>d)</sup> Measurements are also reported to  $^{16}\text{O}^*(17.79, 18.6, 18.98, 19.8)$  with  $J'' = 4^-$  (Hy 81; prelim.).

58. (a)  $^{16}\text{O}(\pi^+, \pi^+)^{16}\text{O}$   
 (b)  $^{16}\text{O}(\pi^-, \pi^-)^{16}\text{O}$

Angular distributions of elastically scattered pions have been studied at  $E_{\pi^+} = 20$  MeV (Ob 81a),  $E_{\pi^-} = 20, 30, 40$  and  $50$  MeV (Ka 81g),  $29.2$  MeV (Jo 79g),  $E_{\pi^+} = 30.0$  and  $49.7$  MeV (Pr 81),  $40$  and  $49.7$  MeV (Ma 78g, Bl 79e),  $46.8$  MeV (Dy 79a),  $79, 114, 163, 240$  and  $343$  MeV (Al 78a, In 78, Ja 78k ( $\pi^\pm$ ), Al 80g), at  $E_{\pi^-} = 114, 163$  and  $240$  MeV (In 78) and at  $E_{\pi^\pm} = 165$  and  $E_{\pi^\pm} = 315$  MeV (Na 80i). At  $E_{\pi^\pm} = 1$  GeV/c angular distributions of  $^{16}\text{O}^*(0, 6.1)$  are reported. See also (Bu 78e) and (Aj 77) for the earlier work. At  $E_{\pi^+} = E_{\pi^-} = 164$  MeV  $^{16}\text{O}^*(0, 6.1, 6.9 + 7.1, 11.5, 17.8, 19.0, 19.8)$  are relatively strongly populated. The  $\pi^+$  and  $\pi^-$  cross sections to  $^{16}\text{O}^*(17.8, 19.8)$  [ $J'' = 4^-; T = 0$ ] are substantially different while those to  $^{16}\text{O}^*(19.0)$  [ $4^-; 1$ ] are equal. Isospin mixing is suggested with off-diagonal charge-dependent mixing matrix elements of  $-147 \pm 25$  and  $-99 \pm 17$  keV (Ho 80c). [See also reaction 81]. See also (Mi 77e, Mi 80d, Ba 81k, Ba 81u, Ob 81), (De 80m, Cu 81b, Ta 81g; theor.) and p. 23.

59.  $^{16}\text{O}(n, n)^{16}\text{O}$

Angular distributions have been measured at energies to  $E_n = 24$  MeV: see tables 16.27 in (Aj 71) and 16.25 in (Aj 77) and (Be 75ee: 5.04, 6.25, 9.29 MeV), (No 78g: 8.85 MeV; 6.13 MeV  $\gamma$ -ray), (Gl 77: 9.25 to 15 MeV;  $n_0$ ) and (Gr 80j: 24 MeV;  $n_0, n_2, n_3$ ) [ $\beta_3 = 0.60, \beta_2 = 0.36$ ]. The energy of  $^{16}\text{O}^*(6.13)$  is  $E_x = 6129.1 \pm 1.2$  keV

TABLE 16.23

Recent \*)  $^{16}\text{O}(p, p)$ , (d, d), ( $^3\text{He}, ^3\text{He}$ ), ( $\alpha, \alpha$ ), ( $^6\text{Li}, ^6\text{Li}$ ), ( $^7\text{Li}, ^7\text{Li}$ ), ( $^9\text{Be}, ^9\text{Be}$ ), ( $^{10}\text{B}, ^{10}\text{B}$ ), ( $^{11}\text{B}, ^{11}\text{B}$ ), ( $^{12}\text{C}, ^{12}\text{C}$ ), ( $^{13}\text{C}, ^{13}\text{C}$ ), ( $^{14}\text{C}, ^{14}\text{C}$ ), ( $^{14}\text{N}, ^{14}\text{N}$ ), ( $^{16}\text{O}, ^{16}\text{O}$ ) measurements

$E_p$ (MeV)	Angular distribution of group	Refs. b)
5.82	$p_0$	(Sa 77gg)
6.9	$p_0$	(Pr 79a)
27.2	$p_1$	(Ce 77a)
30.1	$p_0$	(Vo 79a)
35.2	$p_0, p_3$	(Fa 80a)
42.5, 44.0, 49.3	$p_5$	(Pe 77b)
65 °	$p_0$	(Sa 79p, Sa 81c)
135	$p_0, p_2, p_4$	(Ke 80g)
135	$p_0, p_2, p_3, p_4, p_5$	(He 79g)
800 °	$p_0, p_2, p_3, p_4, p_5$	(Ad 79b)
$E_d$ (MeV)	Angular distribution of group	Refs. d)
12.00	$d_0$	(An 79o)
20.5 °	$d_0, d_2$	(Cl 80b)
52.0 °	$d_0$	(Ma 80e)
56.0 °	$d_0$	(Ha 80d)

TABLE 16.23—continued

$E(^3\text{He})$ (MeV)	Angular distribution of $^3\text{He}$ group to	Refs. <sup>e)</sup>
32.0 <sup>f)</sup> 41.0 44.04 130.0 132	g.s. g.s. g.s. see table 16.24 g.s.	(Lu 80) (Tr 80c) (Go 79d) (Bu 77a) (Ch 81b)
$E_\alpha$ (MeV)	Angular distribution of group	Refs.
6.9 → 10.2 13.05 → 14.12 14.6 → 20.4 20.1 → 23.2 23.7 65 75	$\alpha_0$ $\alpha_0$ $\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4, \alpha_5$ $\alpha_0$ $\alpha_0, \alpha_{1+2}, \alpha_{3+4}, \alpha_5, \alpha_{6+7}$ $\alpha_4^{*f)}$ $\alpha_4, \alpha$ to $^{16}\text{O}^*(12.44)$	(St 78a) (Ce 75a) (Bi 79) (En 77) (Fe 76e) (Ja 76n) see (Ha 80a)
$E(^6\text{Li})$ (MeV)	Angular distribution of group to	
32 50.6	g.s. g.s.	(An 80f) (Ch 76p)
$E(^7\text{Li})$ (MeV)	Angular distribution of group to	Refs.
68	g.s.	(Br 79b)
$E(^9\text{Be})$ (MeV) (Reaction 66)		
20, 26 27.4 $E(^{16}\text{O}) = 15 \rightarrow 25$ 25.7 → 29.5	g.s. g.s. g.s. g.s.	(Un 79) (Gr 78h) (Ba 70mm) (Fu 75c)
$E(^{10}\text{B})$ (MeV)		
33.7, 41.6, 49.5 65.8 100	g.s. g.s. $^{16}\text{O}^*(0, 6.1, 6.9)$	(Pa 80a) (Mo 77y, Mo 79z) (Na 75j, To 77)
$E(^{11}\text{B})$ (MeV) (Reaction 67) <sup>g)</sup>		
41.6, 49.5 115	g.s. g.s.	(Pa 80a) (Br 79b, Ra 79c)

TABLE 16.23—continued

$E(^{12}\text{C})$ or $E(^{16}\text{O})$ (MeV) (Reaction 68)	Angular distribution of group to	Refs. <sup>b</sup> )
$E(^{12}\text{C}) =$ 65 77 115	g.s. g.s. g.s.	(Bo 78e) (Mo 77y, Mo 79z) (Ra 79c)
$E(^{16}\text{O}) =$ 30.8, 31.9, 33.9 31.7, 46.0, 51.3, 52.7 46.0, 47.8, 49.7, 51.3, 52.7 33.2 → 35.2 52 55.3, 56.7, 65.8 $80 \rightarrow 122$ 140, 218, 315	g.s. g.s. $^{16}\text{O}^*(6.1, 7.0)$ g.s. $^{16}\text{O}^*(6.05, 6.13)$ $^{16}\text{O}^*(6.05, 6.13)$ g.s. g.s.	(Sc 78a) (Ma 78n) (Ma 78n) (Fr 80c) (Sh 77k) (Ka 78a) (Co 77r) (Br 81b)
$E(^{13}\text{C})$ (MeV) (Reaction 69)		
36 105	g.s. g.s.	(We 76j) (Br 79b, Ra 79c)
$E(^{16}\text{O})$ (MeV) (Reaction 69)	Angular distribution of group to	Refs. <sup>b</sup> )
14, 17, 20 20, 25, 30	g.s. g.s.	(Ba 70mm) (Sc 75q)
$E(^{14}\text{N})$ (MeV) (Reaction 70)		
76.2 79 155	g.s. g.s. $^{16}\text{O}^*(0, 6.1, 6.9)$	(Mo 77y, Mo 79z) (Mo 76) (Na 75j, To 77)
$E(^{16}\text{O})$ (MeV) (Reaction 71)		
35 → 80 95.2	g.s. g.s.	(Fe 78b) (Mo 77y, Mo 79z)

<sup>a</sup>) See also tables 16.27 in (Aj 71) and 16.25 in (Aj 77).<sup>b</sup>) See also (Da 76n, Se 77, Ma 79mm, Gl 79c).<sup>c</sup>) Polarized.<sup>d</sup>) See also (Ro 78e).<sup>e</sup>) See also (Ch 78r).<sup>f</sup>)  $^4\text{He}^*(20.1)$  [ $J'' = 0^+$ ].<sup>g</sup>) See also reaction 71 in (Aj 77).<sup>h</sup>) See also (Ma 78c, Do 79, Wi 80h).<sup>i</sup>) See also for population of  $^{12}\text{C}$  states.

TABLE 16.24

Excited states of <sup>16</sup>O from <sup>16</sup>O(p, p'), (d, d'), (<sup>3</sup>He, <sup>3</sup>He') and ( $\alpha$ ,  $\alpha'$ )

No.	$E_x$ <sup>b)</sup> (MeV $\pm$ keV)	$L$ <sup>b)</sup>	$E_x$ <sup>d)</sup> (MeV)	$E_x$ <sup>f)</sup> (MeV $\pm$ keV)	$E_x$ <sup>h)</sup> (MeV $\pm$ keV)	$L$ <sup>h,i</sup>	$\Gamma$ <sup>b)</sup> (keV)	$J^\pi; T$ <sup>b,i)</sup>
1			6.05					
2	6.13 <sup>a)</sup>		6.13	6.13 <sup>e,f)</sup>	6.13 <sup>i)</sup>	3		3 <sup>-</sup> ; 0
3	6.92 <sup>a)</sup>		6.92	6.92 <sup>f,g)</sup>	6.92 <sup>i)</sup>	2		2 <sup>+</sup> ; 0 <sup>f,g)</sup>
4	7.12 <sup>a)</sup>		7.12		7.12 <sup>i)</sup>	1		1 <sup>-</sup> ; 0
5	8.87 <sup>a)</sup>		8.87	8.87 $\pm$ 30 <sup>f,g)</sup>	8.87 <sup>a)</sup>	3 <sup>a)</sup>		
6	9.85 <sup>a)</sup>		9.85	9.84 $\pm$ 30	9.85 <sup>i)</sup>	2		2 <sup>+</sup> ; 0 <sup>f)</sup>
7	10.35 $\pm$ 20	4	10.34	10.35 $\pm$ 30	10.35 $\pm$ 30	4		4 <sup>+</sup> ; 0
8	10.95 $\pm$ 30	1	10.95					T=0
9	11.10 $\pm$ 20	4	11.1 <sup>e)</sup>	11.09 $\pm$ 30 <sup>e)</sup>	11.10 $\pm$ 30	4		4 <sup>+</sup> ; 0
10	11.52 $\pm$ 20	2	11.52	11.52 $\pm$ 30 <sup>f,g)</sup>	11.52 $\pm$ 30	2	74 $\pm$ 4 <sup>i)</sup>	2 <sup>+</sup> ; 0
11	12.05 $\pm$ 20		12.05	12.04 $\pm$ 30	12.05 $\pm$ 30	(0)		0 <sup>+</sup> ; 0
12			12.44		12.44 <sup>i)</sup>	1		1 <sup>-</sup> ; 0
13	12.53 $\pm$ 20	1	12.53		12.51 $\pm$ 30			
14	13.02 $\pm$ 20	2	13.1 <sup>e)</sup>	13.11 $\pm$ 30 <sup>a)</sup>	13.07 $\pm$ 20 <sup>e)</sup>	2		2 <sup>+</sup> ; 0 <sup>a)</sup>
15	13.26 $\pm$ 30	3						3 <sup>-</sup> ; 1
16			13.66					
17	13.95 $\pm$ 50	(0+4)		13.97 $\pm$ 30	13.95 $\pm$ 50 <sup>e)</sup>	4		4 <sup>+</sup> ; 0
18				14.94 $\pm$ 30	14.87 $\pm$ 100	6		6 <sup>+</sup>
19	15.26 $\pm$ 50	(3)		15.4				
20	15.50 $\pm$ 30	3			15.50 $\pm$ 50	3	200 $\pm$ 60	3 <sup>-</sup> ; 0
21	16.52 $\pm$ 50	2		16.46 $\pm$ 30	16.40 $\pm$ 100		<100	2 <sup>+</sup> <sup>f)</sup>
22	16.93 $\pm$ 50	(3)						
23	17.25 $\pm$ 50			17.19 $\pm$ 30 <sup>a)</sup>	17.25 $\pm$ 80	(2)	160 $\pm$ 60	2 <sup>+</sup> <sup>f,g)</sup>
24	17.79 $\pm$ 40 <sup>e)</sup>	(3)		17.8	17.83 $\pm$ 100		150 $\pm$ 60	4 <sup>-</sup> ; 0 <sup>c,j)</sup>
25	18.15 $\pm$ 50	(2)			18.0 $\pm$ 100 <sup>i)</sup>	2	300 $\pm$ 50	(2 <sup>+</sup> ); 0
26	18.40 $\pm$ 100	2		18.52 $\pm$ 30 <sup>a)</sup>	18.5 $\pm$ 100 <sup>i)</sup>	2	250 $\pm$ 50 <sup>i)</sup>	2 <sup>+</sup> ; 0 <sup>a)</sup>
27	18.60 $\pm$ 100				18.70 $\pm$ 100	(3)	280 $\pm$ 80 <sup>e)</sup>	
28	18.98 $\pm$ 40 <sup>e)</sup>	(3)		19.09 $\pm$ 30			<100	4 <sup>-</sup> ; 1 <sup>c,j)</sup>
29	19.35 $\pm$ 80	(1)						
30	19.56 $\pm$ 50				19.50 $\pm$ 100	(2, 3)	300 $\pm$ 50 <sup>i)</sup>	(2 <sup>+</sup> , 3 <sup>-</sup> ); 0
31	19.80 $\pm$ 40 <sup>e)</sup>	3					<100	4 <sup>-</sup> ; 0 <sup>j)</sup>
32				20.2 $\pm$ 200 <sup>c,a)</sup>	20.15 $\pm$ 100 <sup>i)</sup>	2	350 $\pm$ 50 <sup>i)</sup>	2 <sup>+</sup> ; 0 <sup>a)</sup>
33	20.56 $\pm$ 80	(1, 2)					370 $\pm$ 100	
34	21.05 $\pm$ 50	1			21.0 $\pm$ 100	2	320 $\pm$ 50	(2 <sup>+</sup> , 0)
35				21.6 $\pm$ 200 <sup>a)</sup>			1000 $\pm$ 300 <sup>a)</sup>	2 <sup>+</sup> <sup>a)</sup>
36	21.80 $\pm$ 80	1			21.85 $\pm$ 100 <sup>i)</sup>	2	400 $\pm$ 50 <sup>i)</sup>	(2 <sup>+</sup> , 0)
37	22.40 $\pm$ 80	(1, 2)			22.5 $\pm$ 100 <sup>i)</sup>		420 $\pm$ 100	1 <sup>-</sup> ; 1
38							400 $\pm$ 50	(2 <sup>+</sup> , 3 <sup>-</sup> ); 0
39	23.20 $\pm$ 80	1					600 $\pm$ 200	1 <sup>-</sup> ; 1
40				23.50 $\pm$ 150 <sup>a)</sup>	23.25 $\pm$ 100 <sup>i)</sup>	2	400 $\pm$ 50 <sup>i)</sup>	2 <sup>+</sup> ; 0 <sup>a)</sup>
41					23.85 $\pm$ 100 <sup>i)</sup>	(0)	400 $\pm$ 50	(2 <sup>+</sup> , 0 <sup>+</sup> ); 0
42	24.00 $\pm$ 100	(1, 2)					1200 $\pm$ 300	1 <sup>-</sup> ; 1
43					24.4 $\pm$ 100 <sup>i)</sup>		400 $\pm$ 50 <sup>i)</sup>	(2 <sup>+</sup> , 3 <sup>-</sup> ); 0
44				25.15 $\pm$ 300 <sup>a)</sup>			2800 $\pm$ 600 <sup>a)</sup>	2 <sup>+</sup> <sup>a)</sup>
45	25.50 $\pm$ 150	(1)					1300 $\pm$ 300	1 <sup>-</sup> ; 1

[ $E_\gamma = 6127.8 \pm 1.2$  keV] (Be 66aa). See also (Be 78d), (Aj 77), (No 77d, Ph 77b, Ta 78u, Th 78e; theor.) and <sup>17</sup>O.

60. (a)	<sup>16</sup> O(p, p) <sup>16</sup> O	
(b)	<sup>16</sup> O(p, 2p) <sup>15</sup> O	$Q_m = -12.1276$
(c)	<sup>16</sup> O(p, pd) <sup>14</sup> N	$Q_m = -20.7363$
(d)	<sup>16</sup> O(p, pt) <sup>13</sup> N	$Q_m = -25.033$
(e)	<sup>16</sup> O(p, p <sup>3</sup> He) <sup>13</sup> C	$Q_m = -22.7934$
(f)	<sup>16</sup> O(p, p $\alpha$ ) <sup>12</sup> C	$Q_m = -7.1620$

Angular distributions of elastically and inelastically scattered protons have been measured at many energies up to  $E_p = 1000$  MeV: see tables 16.27 in (Aj 71), 16.25 in (Aj 77) and 16.23 here. Parameters of the observed groups are displayed in table 16.24. At  $E_p = 135$  MeV, <sup>16</sup>O\*(17.79, 18.98, 19.80) [ $\pm 0.04$  MeV] are strongly populated: the angular distributions of the proton groups lead to an assignment of  $4^-$  for <sup>16</sup>O\*(17.79) (He 79g). Angular distributions of the protons to <sup>16</sup>O\*(8.88) [ $J'' = 2^-$ ] at  $E_p = 42.5, 44.0$  and 49.3 MeV lead to strength distributions which suggest the existence of an octupole (E3) giant resonance at  $E_x \sim 40$  MeV (Pe 77b). At  $E_p = 40$  MeV the spin-flip cross section to <sup>16</sup>O\*(8.88) is interpreted as showing dominance of multistep processes in the reaction (Mo 78m, Co 81g). See also (Ki 77m; 3 GeV; intensity of 6.13 MeV  $\gamma$ -ray).

For reaction (b) see (Ki 80a, Ki 76m;  $E_p = 200$  MeV). See also (Na 77s, Na 78k, Sa 80j), <sup>14</sup>N in (Aj 81) and <sup>17</sup>F. For reactions (c), (d) and (e) see (Gr 77a;  $E_p = 75$  MeV). For reaction (f) see (Ca 81;  $E_p = 101.5$  MeV; to <sup>12</sup>C<sub>g.s.</sub>) and <sup>12</sup>C in (Aj 80). For the cross section for production of pions at  $E_p = 585$  MeV see (Cr 79c). For antiproton scattering see (Au 81; theor.).

See also (No 76c, Ba 77ao, Kr 81a), (Ha 76t, Ba 77ab, Br 77z, Al 78i, De 79w, Li 79h, De 80p, Wh 80a, Mo 81, Ra 81b), (An 76d, Go 76n, Al 77d, Dy 77c, Ga 77h, Ko 77jj, Le 77c, Ma 77u, Ma 77qq, Mo 77w, Ph 77c, Va 77e, Vo 77b, Al 78k, Au 78b, Bi 78n, Br 78k, Br 78q, Ch 78y, Cu 78b, Go 78f, Ha 78f, Ha 78m, He 78g, Io 78a, Le 78c, Ma 78k, Ph 78, Pi 78a, Wr 78, Ab 79d, Bi 79b, He 79e, Ja 79d, Ki 79k, Ko 79e, Ko 79ff, Ko 79kk, Ku 79g, Ma 79p, Ma 79ii, Mi 79i, Ph 79c,

#### Footnotes to table 16.24

<sup>a</sup>) See also tables 16.28 and 16.29 in (Aj 71) and 16.26 in (Aj 77).

<sup>b</sup>) (p, p'): (Bu 75d, Bu 76e;  $E_p = 45$  MeV).

<sup>c</sup>) (p, p'): (He 79g;  $E_p = 135$  MeV).

<sup>d</sup>) (d, d'): (Du 74c;  $E_d = 81.6$  MeV). Energies are nominal ( $\pm 100$  to  $\pm 260$  keV); angular distributions reported to all but last state.

<sup>e</sup>) Unresolved states.

<sup>f</sup>) (<sup>3</sup>He, <sup>3</sup>He'): (Mo 74c;  $E(^3\text{He}) = 71$  MeV). Angular distributions are reported to states labelled by <sup>f</sup>).

<sup>g</sup>) (<sup>3</sup>He, <sup>3</sup>He'): (Bu 77a;  $E(^3\text{He}) = 130$  MeV); measured angular distributions.

<sup>h</sup>) ( $\alpha$ ,  $\alpha'$ ): (Bu 76e;  $E_\alpha = 60$  MeV).

<sup>i</sup>) ( $\alpha$ ,  $\alpha'$ ): (Ha 76n, Ha 76o;  $E_\alpha = 104$  MeV).

<sup>j</sup>) See also (Ba 81u; theor.).

## **<sup>16</sup>O**

60

F. AJZENBERG-SELOVE

Wa 79c, Wa 79g, Am 80e, Ay 80, De 80v, Ko 80cc, Ma 80ee, Mc 80, Wu 80a, Ba 81u, Mi 81a, Pe 81b, Ro 81m, Th 81b, Tu 81; theor.) and <sup>17</sup>F.

### 61. <sup>16</sup>O(d, d)<sup>16</sup>O

Angular distribution studies have been carried out for  $E_d$  up to 81.6 MeV: see tables 16.27 in (Aj 71), 16.25 in (Aj 77) and 16.23 here. Observed deuteron groups are displayed in table 16.24. See also <sup>18</sup>F in (Aj 83), (Te 77b) and (Co 76u, Le 76p, Dm 77, Le 78k, Ta 78u, Th 78e, Gr 79m, Ay 80, Ni 80d; theor.).

### 62. <sup>16</sup>O(t, t)<sup>16</sup>O

Angular distributions are reported for  $E_t$  to 20.01 MeV: see tables 16.27 in (Aj 71) and 16.25 in (Aj 77). See also (Le 78k, Kh 80a; theor.). See also <sup>19</sup>F in (Aj 78).

### 63. <sup>16</sup>O(<sup>3</sup>He, <sup>3</sup>He)<sup>16</sup>O

Angular distributions have been measured to  $E(^3\text{He}) = 132$  MeV: see tables 16.27 in (Aj 71), 16.25 in (Aj 77) and 16.23 here. Inelastic groups are shown in table 16.24. The total E2 strength exhausted to  $E_x = 30$  MeV is ~37% of the quadrupole EWSR (Bu 77a; see also for  $\beta_2 R$ ). See also (Se 77), (Au 75: S<sub>a</sub>), (Ha 76t, Ka 79q, Tr 80f) and (Le 77c, Le 78k, Ta 78u, Th 78e, Be 79dd, Le 79o; theor.).

### 64. (a) <sup>16</sup>O( $\alpha$ , $\alpha$ )<sup>16</sup>O

(b) <sup>16</sup>O( $\alpha$ , 2 $\alpha$ )<sup>12</sup>C

$$Q_m = -7.1620$$

Angular distributions of  $\alpha$ -particles have been measured up to  $E_\alpha = 146$  MeV: see tables 16.27 in (Aj 71), 16.25 in (Aj 77) and 16.23 here. Observed excited states are displayed in table 16.24.

The isoscalar ( $E2, T = 0$ ) giant resonance decays predominantly via the  $\alpha_1$  channel which contains ~40% of the E2 EWSR, rather than via the  $\alpha_0$  and  $p_0$  channels (Kn 78a;  $E_\alpha = 155$  MeV). See also (Ha 76o, Ha 80a, Ha 81g). Reaction (b) has also been studied by (Wa 80b;  $E_\alpha = 140$  MeV) and (Do 79e;  $E_\alpha = 700$  MeV): values of  $S_\alpha$  and  $N_{eff}$  are derived. See also (Wo 76c, Aj 77). For the ( $\alpha, \alpha p$ ) reaction see (Sa 80b).

See also <sup>20</sup>Ne in (Aj 83), (Vo 75i, Sh 78p, Sh 80), (Po 78c; applied), (Re 80; astrophysics), (Be 76bb, Ha 76t, Ma 77uu, Ch 78s, Be 79ee, Kn 79i, Ma 79kk, Sp 80d, We 80d, Ca 81b) and (Co 75p, Am 76b, Ba 76dd, Cu 76f, He 76j, Lo 76f, Mi 76i, Pa 76i, Ab 77, Ba 77l, Br 77c, Bu 77, Cl 77b, Dm 77, Ik 77, Ho 77k, Le 77c, Oh 77c, Sa 77v, Tr 77c, We 77l, Ch 78b, Le 78k, Ma 78ii, Ta 78u, Th 78e, Ze 78c, Co 79o, Gr 79m, Iz 79, Ja 79a, Ja 79g, La 79j, Le 79a, Le 79d, Ve 79c, Dm 80, Kh 80a, Le 80t, To 80f, Dy 81a; theor.).

### 65. (a) <sup>16</sup>O(<sup>6</sup>Li, <sup>6</sup>Li)<sup>16</sup>O

(b) <sup>16</sup>O(<sup>7</sup>Li, <sup>7</sup>Li)<sup>16</sup>O

For studies of the elastic scattering see tables 16.25 in (Aj 77) and 16.23 here. For studies of (d- $\alpha$ ) angular correlations see <sup>20</sup>Ne in (Aj 83). See also (Sa 77c, Kn 79k), (Og 76, Fi 78a) and (Me 78i, Pe 78a, Su 79g, Kh 80a, Me 81c, Dy 81a, Gu 81; theor.).

66. <sup>16</sup>O(<sup>9</sup>Be, <sup>9</sup>Be)<sup>16</sup>O

For angular distributions see table 16.23. For fusion cross sections see (Sw 77, Ch 78e). See also (Pa 78, Hu 79, Ma 79cc, Pa 79, Gu 81; theor.).

67. (a) <sup>16</sup>O(<sup>10</sup>B, <sup>10</sup>B)<sup>16</sup>O  
(b) <sup>16</sup>O(<sup>11</sup>B, <sup>11</sup>B)<sup>16</sup>O

Angular distributions are displayed in table 16.23. For fusion cross-section measurement [reaction (a)] see (Go 79i, Wi 80e, Th 81a). See also (Go 79n) and (Va 80c, Va 81b; theor.).

68. (a) <sup>16</sup>O(<sup>12</sup>C, <sup>12</sup>C)<sup>16</sup>O  
(b) <sup>16</sup>O(<sup>12</sup>C,  $\alpha$ <sup>12</sup>C)<sup>12</sup>C       $Q_m = -7.1620$ 

Angular distributions have been reported at many energies to  $E(^{16}\text{O}) = 315$  MeV: see reaction 64 (Aj 71) and tables 16.25 (Aj 77) and 16.23 here. Most of the studies of this reaction have involved yield and cross-section measurements, as they apply to compound structures in <sup>28</sup>Si, as well as fusion cross sections: see (Aj 77) and (Br 77w, Co 77r, Na 77f, Ta 77, Ch 78l, Fe 78b, Ja 78a, Ka 78a, Ma 78j, Ma 78n, Sc 78a, Ch 79c, Fu 79g, Ja 79w, Ko 79b, Ko 79y, Be 80c, Fr 80a, Fr 80c, Ja 80b, Fu 81b). See also (Fr 76d, Ci 78, Ma 78c, Sh 78a, Fr 79b, Lu 79a, Uz 79, Fr 80g, Wi 80h, Sc 81b, Wi 81f).

At  $E(^{16}\text{O}) = 77$  MeV reaction (b) mainly proceeds via <sup>16</sup>O\*(10.3 ± 0.4, 14.8 ± 0.4) [which are associated with the  $J'' = 4^+, 6^+$  states <sup>16</sup>O\*(10.35, 14.82)]. (Fu 79g). At  $E(^{16}\text{O}) = 140$  MeV the breakup is interpreted as proceeding via <sup>16</sup>O\*(11.6, 13.1, 15.8) (Ra 80e). See also (Sc 79d). For the excitation of the GQR in <sup>12</sup>C see (Do 79:  $E(^{16}\text{O}) = 315$  MeV).

See also (La 78, Ga 79c, Sa 79n, Ta 80a), (Ga 77n, Ga 78h, Ho 78n, Le 78g, Ma 78hh, Ta 78m, Ts 78a, Ga 79g, Er 80c, Ga 80b), (Ro 78b; astrophysics) and (Ba 76dd, Ch 76s, Ba 77r, Ba 77w, Ch 77g, Cl 77b, Fr 77b, Ja 77c, Pa 77j, Ro 77w, Ab 78a, Av 78, Ba 78y, Bh 78c, Bi 78, Fr 78n, Go 78k, Ma 78h, Ma 78l, Ma 78ii, Ta 78c, Ta 78e, Va 78g, Go 79t, Kr 79b, Na 79, St 79d, Ta 79c, Ta 79e, Ta 79f, Te 79, Ve 79e, Ab 80d, De 80aa, Dr 80d, Fu 80g, Ga 80i, Hu 80, Ko 80u, La 80g, La 80l, Ta 80e, Ta 80k, Va 80c, Vo 80e, Ga 81c, Gu 81, Ha 81k, Hu 81c, Sc 81d, Ta 81a, To 81b, Ub 81, Wi 81b; theor.).

69. (a) <sup>16</sup>O(<sup>13</sup>C, <sup>13</sup>C)<sup>16</sup>O  
(b) <sup>16</sup>O(<sup>14</sup>C, <sup>14</sup>C)<sup>16</sup>O

# <sup>16</sup>O

62

F. AJZENBERG-SELOVE

For elastic scattering studies see table 16.23. For reaction (a) see also (Ch 79c; fusion), (Ra 80e;  $\alpha$ -breakup) and (Du 76b, Ga 79c). For reaction (b) see (Be 78ff, Ko 81a). For theoretical papers see (Pa 78, Sc 78m, He 79d, Pa 79, Ha 81k).

70. (a)  $^{16}\text{O}(^{14}\text{N}, ^{14}\text{N})^{16}\text{O}$   
 (b)  $^{16}\text{O}(^{15}\text{N}, ^{15}\text{N})^{16}\text{O}$

For elastic scattering studies see table 16.23 and (Aj 77). For total fusion cross-section measurements (reaction a) see (Sw 77b, Vo 77e, Vo 81). See also (Av 78, Va 78g; theor.).

71.  $^{16}\text{O}(^{16}\text{O}, ^{16}\text{O})^{16}\text{O}$

The angular distributions for elastic scattering have been measured with  $E(^{16}\text{O})$  up to 140.4 MeV: see reaction 66 in (Aj 71), table 16.25 in (Aj 77) and 16.23 here. The angular distributions corresponding to the excitation of the first four excited states of <sup>16</sup>O have been studied at 51.5 and 140.4 MeV: see (Aj 77). For yield and total fusion cross sections see (Ko 77j, Fe 78b, Ts 78a, Ch 79k, Ko 79m, Ko 79y, Fr 80a, Hu 80g, La 81c). See also (Ch 80c, We 80a, Vo 81b).

See also (Co 77p, Ha 78aa, Ga 79c, La 79p, Sa 79m, Sa 79n, Co 80t, De 81r), (Ho 78n), (Ro 78b; astrophysics) and (Al 76k, Ar 76g, Ba 76dd, Ch 76s, Cu 76f, Lo 76f, Mi 76i, Mo 76m, Ba 77b, Br 77c, Ca 77d, Ch 77g, Cl 77b, Fl 77, Fr 77b, Fr 77e, Fr 77f, Fu 77l, Ga 77e, Ha 77n, Ho 77r, Ik 77b, Ko 77d, La 77j, Os 77a, Pl 77, Sa 77a, Sa 77g, Sa 77h, Sa 77cc, To 77d, We 77l, Zi 77c, An 78, An 78a, Ar 78g, Av 78, Ba 78j, Be 78b, Bo 78d, Br 78j, Ch 78o, Cu 78e, Cu 78f, Da 78p, Fl 78c, Fr 78i, Gu 78a, Ko 78h, Kr 78d, Le 78q, Pa 78a, Sc 78c, Sc 78l, Sc 78m, Si 78b, Su 78c, Va 78a, Va 78g, Vo 78b, Vo 78f, An 79g, Be 79t, Ch 79k, Dh 79, Fr 79k, Ja 79, Mu 79d, Ph 79, Si 79c, Si 79f, Ta 79e, Ve 79e, Ab 80d, An 80n, Ba 80jj, Be 80g, Fa 80d, Fl 80, Ga 80b, Ko 80c, Ko 80h, Ko 80k, Ko 80u, La 80k, Oh 80, Pr 80b, Si 80d, Si 80i, Ta 80c, Ta 80e, Ta 80n, To 80d, Va 80c, Vi 80a, Wo 80a, Ab 81a, Ba 81, De 81g, Dy 81a, Ha 81k, He 81, Ma 81g, Sa 81f, Sa 81i, Si 81e, Ta 81j, Ub 81, Va 81b; theor.).

72. (a)  $^{16}\text{O}(^{17}\text{O}, ^{17}\text{O})^{16}\text{O}$   
 (b)  $^{16}\text{O}(^{18}\text{O}, ^{18}\text{O})^{16}\text{O}$

Angular distributions of elastically scattered ions have been studied at  $E(^{16}\text{O}) = 24, 28$  and  $32$  MeV and  $E(^{17}\text{O}) = 53.0$  to  $66$  MeV [reaction a] and at  $E(^{16}\text{O}) = 24$  to  $54.8$  MeV and  $E(^{18}\text{O}) = 53.1$  to  $89.3$  MeV [reaction b]: see (Aj 77) and (Ca 81c;  $E(^{18}\text{O} = 35 \text{ MeV})$ ). Yields and fusion cross sections are reported by (Wi 80e) (reaction a) and (Fr 78f) (reaction b). See also (Ca 77j, Ra 79c), (Fi 74f, Ho 78n) and (Im 77b, Le 78q, Cr 79b, Le 79a; theor.).

73.  $^{16}\text{O}(^{19}\text{F}, ^{19}\text{F})^{16}\text{O}$

Elastic scattering angular distributions have been studied at  $E(^{16}\text{O}) = 21.4$  and 25.8 MeV and at  $E(^{19}\text{F}) = 33$  and 36 MeV: see (Aj 77). See also (Sc 77e, Oh 80; theor.).

74.  $^{16}\text{O}(^{20}\text{Ne}, ^{20}\text{Ne})^{16}\text{O}$

Elastic scattering angular distributions have been measured at  $E(^{16}\text{O}) = 40.7$ , 51.1 and 59.4 MeV (see Aj 77), 94.8 MeV (Mo 77y, Mo 79z; also excited states of  $^{20}\text{Ne}$ ) and at  $E(^{20}\text{Ne}) = 50$  MeV (St 76i; also excited states of  $^{20}\text{Ne}$ ). See also (Ga 80c). For yield and fusion cross-section measurements see (Sh 78i, Ga 78e, Ga 79a, Ga 79g, Ko 79t, Re 79d, Di 80a, Ga 80c, Ga 81c, Ga 81e; all abstracts). See also (Va 79k) and (Ja 79f, Le 79a, Ga 80b, Oh 80, An 81c; theor.).

75. (a)  $^{16}\text{O}(^{24}\text{Mg}, ^{24}\text{Mg})^{16}\text{O}$   
 (b)  $^{16}\text{O}(^{26}\text{Mg}, ^{26}\text{Mg})^{16}\text{O}$

Elastic angular distributions are reported at  $E(^{16}\text{O}) = 35$  to 50 MeV (Fo 79f), 46.3 and 60.3 MeV (Pa 80c; also  $2^+$  in  $^{24}\text{Mg}$ ) and 67 MeV (Ec 81; also various  $^{24}\text{Mg}$  states) for reaction (a), and at 35 to 50 MeV (Fo 79f) and 27.4 to 48.4 MeV (Ro 80g; also  $2^+$  in  $^{26}\text{Mg}$ ) for reaction (b). Yield and fusion measurements have been carried out by (Ho 78k, Ta 78b, Fo 79f, Le 79b, Pe 79c, Ko 80b, Pa 80c, Ro 80g, Dr 81b). See also (Ra 80f), (Ta 78m) and (Ho 78h, Ga 80i, Lo 80, Pa 80g, Sa 80h, De 81k; theor.).

76.  $^{16}\text{O}(^{27}\text{Al}, ^{27}\text{Al})^{16}\text{O}$

An elastic angular distribution has been measured at  $E(^{16}\text{O}) = 46.5$  MeV (Pe 78d). For yield and fusion cross-section measurements see (Ei 77, Ra 79i, Sh 80e, Le 81c). See also (Ke 78b, Va 79h, Mi 80b, Sa 80m, Ta 80a), (Sc 77o, Ho 78n, Ma 78hh, Hu 79d) and (Af 77, Bi 78, Va 78g, Ka 79m, Lo 80, Sa 80h, Ac 81, De 81k, Sa 81a, Va 81f; theor.).

77. (a)  $^{16}\text{O}(^{28}\text{Si}, ^{28}\text{Si})^{16}\text{O}$   
 (b)  $^{16}\text{O}(^{29}\text{Si}, ^{29}\text{Si})^{16}\text{O}$   
 (c)  $^{16}\text{O}(^{30}\text{Si}, ^{30}\text{Si})^{16}\text{O}$   
 (d)  $^{16}\text{O}(^{32}\text{S}, ^{32}\text{S})^{16}\text{O}$

Angular distributions for reaction (a) are reported at  $E(^{16}\text{O}) = 32$  MeV (Ca 78f; also  $^{28}\text{Si}^* 2^+$ ), 33 to 81 MeV [(Ga 73n) and see (Cr 76d)], 33, 36 and 38 MeV (Ec 77), 35.7 and 38.2 MeV (Re 79c), 41.23 to 63 MeV (Br 77e, Ge 78f, Sh 78f, Sh 80c, Sh 80l), 56 MeV (Me 79; also  $^{28}\text{Si}^*$ ), 60 MeV (Du 78a; also  $^{28}\text{Si}^*$ ), 141.5 MeV (Sa 78i; also  $^{28}\text{Si}^*$ ) and 215.2 MeV (Cr 76d). For reaction (c) see (Re 79c). For yield and fusion cross-section measurements see (Ba 78, Ba 79ee, Fo 79f, Jo 79h, Ku 79j, Ra 79i, Be 80b). See also (Br 79e).

See also (Ze 77d, St 78l, De 79t, He 79h, Zi 80a) and (Ud 77, De 78g, Du 78c, Fr 78o, Le 78s, Sh 78o, Ta 78j, Te 78a, Va 78g, An 79c, Fr 79a, Ka 79m, Ko 79h,

Kr 79c, La 79, Li 79i, Sa 79l, Ba 80oo, Cr 80, Dr 80d, Fa 80d, Fr 80e, Ga 80b, Go 80e, La 80k, Ma 80y, To 80f, Fr 81g, Me 81, Ra 81f, Vi 81; theor.).

78. (a)  $^{16}\text{O}({}^{40}\text{Ca}, {}^{40}\text{Ca})^{16}\text{O}$   
 (b)  $^{16}\text{O}({}^{44}\text{Ca}, {}^{44}\text{Ca})^{16}\text{O}$   
 (c)  $^{16}\text{O}({}^{48}\text{Ca}, {}^{48}\text{Ca})^{16}\text{O}$

Elastic angular distributions are reported on  ${}^{40}\text{Ca}$  at  $E({}^{16}\text{O}) = 50$  MeV (Ku 79b), 51.5 and 54 MeV (Al 81d; also  ${}^{40}\text{Ca}^*$ ), 56.8 to 62.8 MeV (Ge 78c), 60 MeV (Re 78b; also  ${}^{40}\text{Ca}^*$ ) and 55.6, 74.4, 103.6, 139.6 and 214.1 MeV (Vi 79c), while the elastic scattering on  ${}^{48}\text{Ca}$  has been studied at 60 MeV (Ko 78a; also  ${}^{40}\text{Ca}^*$ ). Yield and fusion cross-section measurements are reported by (Ge 78c, Ku 79b, Vi 79c, Ku 80g). See also (Ol 78a), (Ze 77d, He 79h) and (Af 77, We 77l, Ko 78j, Pa 78a, Pe 78c, Ba 79gg, Ba 79kk, Be 79t, Bo 79z, Le 79a, Pe 79d, Wo 79b, Be 80d, Iz 80, La 80k, Le 80g, Ma 80y, Rh 80, Wo 80f, Va 81f; theor.).

79.  ${}^{17}\text{O}(\gamma, \text{n})^{16}\text{O}$   $Q_m = -4.1443$

See  ${}^{17}\text{O}$ .

80.  ${}^{17}\text{O}(\text{p}, \text{d})^{16}\text{O}$   $Q_m = -1.9197$

Angular distributions for the ground state deuteron group have been studied at  $E_p = 8.62$  to 11.44 MeV (Cr 75e). At  $E_p = 31$  MeV, angular distributions are reported for the deuterons corresponding to  ${}^{16}\text{O}^*(0, 6.05 + 6.13, 7.12, 8.87, 10.35, 12.97, 13.26)$ . States at  $E_x = 15.22$  and 15.42 MeV were also observed. Spectroscopic factors were obtained from a DWBA analysis (Me 70).

81.  ${}^{17}\text{O}(\text{d}, \text{t})^{16}\text{O}$   $Q_m = 2.1130$

Information obtained from this reaction at  $E_d = 52$  MeV is displayed in table 16.20. Comparison of the (d, t) and (d,  ${}^3\text{He}$ ) reactions leads to assignments of analog states in  ${}^{16}\text{N}$  and in  ${}^{16}\text{O}$  [see reaction 25 in  ${}^{16}\text{N}$  and table 16.10] (Wa 77l, Ma 78p). See also (Ma 81d). A study of this reaction, the (d,  ${}^3\text{He}$ ) reaction, and reaction 82 below, suggests that there is more than 17% isospin mixing of the  $2^-$  states  ${}^{16}\text{O}^*(12.97, 12.53)$ : the corresponding mixing matrix element is  $\geq 155 \pm 30$  keV (Wa 77l). An isospin mixing matrix element of  $110 \pm 10$  keV for the  $4^-$  states  ${}^{16}\text{O}^*(17.79, 18.98, 19.80)$  is compatible with the results from this reaction and with pion scattering (Cu 81b). [See also reaction 58.] See also (Br 80j) below and  ${}^{19}\text{F}$  in (Aj 83).

82.  ${}^{17}\text{O}({}^3\text{He}, \alpha)^{16}\text{O}$   $Q_m = 16.4335$

Angular distributions are reported at  $E({}^3\text{He}) = 11$  MeV (Bo 71b). Angular correlations have been studied by (Br 80j) [see for spectroscopic factors relating to a number of states of  ${}^{16}\text{O}$  with  $E_x > 12.9$  MeV with the  ${}^{17}\text{O}_{g.s.}$  as well as with states

of  $^{15}\text{N}$ . Table 16.20 displays some of the information derived from this reaction. For polarization measurements see (Ka 81d, Ro 81c) and  $^{20}\text{Ne}$  in (Aj 83).

$$83. \ ^{18}\text{O}(\gamma, 2\text{n})^{16}\text{O} \quad Q_m = -12.1889$$

See (Ba 76ii) and  $^{18}\text{O}$  in (Aj 83).

$$84. \ ^{18}\text{O}(\text{p}, \text{t})^{16}\text{O} \quad Q_m = -3.7069$$

Angular distributions of tritons have been measured for  $E_p$  to 43.7 MeV: see (Aj 71, Aj 77) for the earlier references and (An 80j;  $E_p = 23$  MeV;  $t_0$ ). The population of  $^{16}\text{O}^*(22.7, 24.5)$  is consistent with  $L = 0$  and 2, respectively, and with assignments of  $T = 2, J'' = 0^+$  and  $2^+$  (Ce 64b). The decay of  $^{16}\text{O}^*(22.7), J''; T = 0^+; 2$ , is via  $\alpha_0$ ,  $\alpha_1$  and  $\alpha_2$  [ $^{12}\text{C}^*(0, 4.4, 7.7)$ ] with  $1.6 \pm 0.7$ ,  $1.9 \pm 0.7$  and  $14 \pm 2\%$  branches and  $\Gamma_i$  (eV) =  $190 \pm 100$ ,  $230 \pm 110$  and  $1680 \pm 550$  eV, respectively; via  $p_0, p_{1+2}, p_3$  with  $7 \pm 2$ ,  $11 \pm 2$  and  $5 \pm 2\%$  branches and  $\Gamma_i$  (eV) =  $840 \pm 343$ ,  $1320 \pm 454$  and  $600 \pm 300$  eV; and via  $n_{1+2}$  with a  $23 \pm 15\%$  branch [ $\Gamma_n = 2760 \pm 1970$  eV] (the  $n_0$  branch is  $< 15\%$ ) (Fr 79c) [ $\Gamma_i$  are based on a total width of  $12 \pm 3.5$  keV]. See also (Aj 77),  $^{18}\text{F}$  in (Aj 83) and (St 77n, Ku 78d, Am 79b, Be 79z, Fe 80a, We 80, We 80e, Be 81e; theor.).

$$85. \ ^{18}\text{O}(\alpha, ^6\text{He})^{16}\text{O} \quad Q_m = -11.215$$

Angular distributions have been measured at  $E_\alpha = 58$  MeV to  $^{16}\text{O}^*(0, 6.1, 6.92, 7.12)$ . Groups at  $E_x = 10.4$ ,  $13.3 \pm 0.1$  and  $16.3 \pm 0.1$  MeV were also observed (Va 75e).

$$86. \ ^{18}\text{O}(^{17}\text{O}, ^{19}\text{O})^{16}\text{O} \quad Q_m = -0.187$$

Angular distributions are reported at  $E(^{17}\text{O}) = 36$  MeV (Ka 77bb; to  $^{16}\text{O}_{\text{g.s.}}$  and  $^{19}\text{O}$  states). See also (Du 76b).

$$87. \ ^{18}\text{O}(^{18}\text{O}, ^{20}\text{O})^{16}\text{O} \quad Q_m = -0.628$$

Angular distributions involving  $^{16}\text{O}_{\text{g.s.}}$  and  $^{20}\text{O}$  states are reported at  $E(^{18}\text{O}) = 24$  to 36 MeV (Ka 77v) and at 52 MeV (Ku 79).

$$88. \ ^{19}\text{F}(\text{p}, \alpha)^{16}\text{O} \quad Q_m = 8.1137$$

Angular distributions have been reported at many energies to  $E_p = 44.5$  MeV [see (Aj 77)] and at  $E_p = 0.83$  to 0.87 MeV (Ca 741:  $\alpha_0$ ), 1.5 to 3.8 MeV (Ca 76s:  $\alpha_0$ ), 2.06 to 2.68 MeV (Cu 80d:  $\alpha_0, \alpha_1$ ) and 13.5 to 18.0 MeV (St 77:  $\alpha_0$ ). Observed states of  $^{16}\text{O}$  are displayed in table 16.31 in (Aj 71).

The E0 transition ( $6.05 \rightarrow 0; 0^+ \rightarrow 0^+$ ) has been investigated in some detail: the internal conversion to pair production ratio is  $(4.00 \pm 0.46) \times 10^{-5}$  (Le 63b). The ratio of double  $\gamma$ -emission to pair production  $\Gamma_{E1E1}/\Gamma_{E0(\pi)} = (2.5 \pm 1.1) \times 10^{-4}$  (Wa 75h).  $\tau_m$  for  $^{16}\text{O}^*(6.05) = 96 \pm 7$  psec (Bi 73q).  $^{16}\text{O}^*(6.13)$  has also been studied extensively:  $|g| = 0.55 \pm 0.04$ ,  $\tau_m = 26.6 \pm 0.7$  psec (Br 73m). A search for double

positron-electron pair creation by the 6.13 MeV  $\gamma$ -ray was unsuccessful: the ratio of the cross section for production of such a double pair to the cross section for formation of a single pair is  $-(2 \pm 5) \times 10^{-5}$  (Wi 72b). The previously reported ground state decay of  $^{16}\text{O}^*(10.95)$  [ $0^- \rightarrow 0^+$ ] is not observed; neither is a pair peak due to a  $(10.95 \rightarrow 7.12)$  decay (Al 78d). For  $\gamma$ -ray branching ratios and lifetimes see tables 16.10 and 16.11.


See also  $^{20}\text{Ne}$  in (Aj 83), (Yo 66m, Go 76q, He 76m), (Ak 78, An 78h, Ch 78t, Go 78c, Vo 78c, Di 80, Ke 80; applications) and (Le 79g; theor.).



Angular distributions have been measured to  $E(^3\text{He}) = 40.7$  MeV: see (Aj 77).



See (Aj 77).



For reaction (a) see  $^{20}\text{Ne}$  in (Aj 78, Aj 83). For reaction (b) see (Ca 81). For reaction (c) see (Aj 77) and (Ch 77a; theor.).



Angular distributions have been studied at  $E_d$  to 40 MeV [see (Aj 77)] and at 80 MeV (Oe 79a;  $^{16}\text{O}_{\text{g.s.}}$ ; relative  $S_\alpha$ ). See also (Be 78w) and (Bi 77e, Ka 78i, Sh 78k; theor.).



At  $E(^3\text{He}) = 30$  MeV angular distributions involving  $^{16}\text{O}^*(0, 6.1)$  and the first two states of  $^7\text{Be}$  are reported by (De 70e).



See (Sa 78c). See also (Ca 79j).



See (Sa 79i).



Angular distributions are reported at  $E_\alpha = 22.8$  to 25.4 MeV (So 78b;  $^{16}\text{O}_{\text{g.s.}}$ ) and at 90.3 MeV (Be 80a, Be 80f; to  $^{16}\text{O}^*(0, 6.1, 7.0, 8.8, 9.8, 10.3)$ ).



See (No 78a). See also (Le 80r; theor.).

98.  $^{28}\text{Si}(\text{e}, \text{e}' ^{12}\text{C})^{16}\text{O}$  $Q_m = -16.7542$ 

See (Sa 78c). See also (Sc 80e; theor.).

 $^{16}\text{F}$ 


(Figs. 4 and 5)

GENERAL<sup>†</sup>

See (La 77a, Si 77d, Wo 78f, Ha 80p, Os 81a).

1. (a)  $^{14}\text{N}(^3\text{He}, \text{n})^{16}\text{F}$   $Q_m = -0.969$   
 (b)  $^{14}\text{N}(^3\text{He}, \text{np})^{15}\text{O}$   $Q_m = -0.421$

Observed neutron groups and  $L$ -values derived from angular distribution measurements are displayed in table 16.26 (Bo 73hh). For the results from reaction (b) see table 16.26 (Ot 76a). See also (Aj 77).

Fig. 4. Energy levels of  $^{16}\text{F}$ . For notation see fig. 2.<sup>†</sup> See also (Aj 77).

# <sup>16</sup>F MASTER TABLE

68

F. AJZENBERG-SELOVE

TABLE 16.25  
Energy levels of <sup>16</sup>F

$E_x$ (MeV ± keV)	$J^\pi; T$	$\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	(0) <sup>-</sup> ; 1	40 ± 20	p	1 → 5
0.194 ± 9	(1) <sup>-</sup>	<40	p	1 → 5
0.424 ± 5	(2) <sup>-</sup>	40 ± 30	p	1 → 5
0.720 ± 6	(3) <sup>-</sup>	<15	p	1 → 5
3.762 ± 7	1 <sup>+</sup>	<40	p	1, 3, 4, 5
3.869 ± 7	(2) <sup>+</sup>	<20	p	1, 4, 5
4.372 ± 7		50 ± 20	p	1, 4, 5
4.654 ± 7	1 <sup>+</sup>	60 ± 20	p	1, 3, 4, 5
4.71 ± 20				5
4.973 ± 10	$\pi = +$			1, 5
5.264 ± 20				1
5.390 ± 14	$\pi = +$			1, 5
5.448 ± 20				1
5.529 ± 14	$\pi = +$			1, 4, 5
5.840 ± 40				1
6.05 ± 20				5
6.230 ± 50				1, 3
6.371 ± 20				1
6.678 ± 10				1, 4, 5
6.93 ± 20				5
7.110 ± 20				1
7.730 ± 40				1, 3



At  $E({}^{10}\text{B}) = 100$  MeV unresolved structures to  ${}^{16}\text{F}^*(0.4, 0.7, 5.5, 6.3, 6.8, 7.8)$  are reported (Ha 78a).



Observed neutron groups are listed in table 16.26 (Mo 71l). See also (An 80g, An 81, Ma 81k) and (Aj 77).



Observed triton groups are displayed in table 16.26. Comparisons of relative populations of the first four states in this reaction and in the analog reaction ( ${}^{16}\text{O}(\text{t}, {}^3\text{He})^{16}\text{N}$ ) to known states in  ${}^{16}\text{N}$  [see reaction 23 in  ${}^{16}\text{N}$  (Fl 74a)] suggest that  ${}^{16}\text{F}^*(0, 0.20)$  are  $0^-$  and  $1^-$  and that  ${}^{16}\text{F}^*(0.42, 0.72)$  are  $2^-$  and  $3^-$  (the relative ordering of the  $J^\pi$  is ambiguous). (Pe 65) suggest, on the basis of angular distribu-

TABLE 16.26  
 $^{16}\text{F}$  levels from  $^{14}\text{N}({}^3\text{He}, \text{n})$ ,  $^{16}\text{O}(\text{p}, \text{n})$ ,  $^{16}\text{O}({}^3\text{He}, \text{t})$  and  $^{19}\text{F}({}^3\text{He}, {}^6\text{He})$

$^{16}\text{F}^*{}^b)$ (MeV $\pm$ keV)	$L$ <sup>b)</sup>	$^{16}\text{F}^*{}^c)$ (MeV $\pm$ keV)	$J^\pi{}^d)$	$^{16}\text{F}^*{}^e)$ (MeV $\pm$ keV)	$^{16}\text{F}^*{}^f)$	$\Gamma_{\text{c.m.}}{}^g)$ (keV)	$J^\pi{}^h)$
0	1	0	(1 $^-$ )	0	0	$40 \pm 20$	(0) $^-$
$0.192 \pm 15$	1	$0.190 \pm 20$	(0 $^-$ )	$0.197 \pm 12$	$0.19 \pm 20$	40	(1) $^-$
$0.425 \pm 15$	3	$0.425 \pm 10$	( $\geq 2$ )	$0.424 \pm 5$	$0.425 \pm 20$	$40 \pm 30$	(2) $^-$
$0.722 \pm 10$	(3)	$0.725 \pm 10$	( $\geq 2$ )	$0.720 \pm 6$	$0.72 \pm 20$	<15	(3) $^-$
$3.751 \pm 10$	0	$3.775 \pm 10$	(1)	j)	$3.75 \pm 20$	<40	1 $^+$
$3.861 \pm 10$	2	$3.880 \pm 10$			$3.86 \pm 20$	<20	(2) $^+$
$4.370 \pm 10$		$4.375 \pm 10$	( $\geq 2$ )		$4.37 \pm 20$	$50 \pm 20$	
$4.646 \pm 10$	0	$4.661 \pm 10$		j)	$4.66 \pm 20$	$60 \pm 20$	1 $^+$
					$4.71 \pm 20$ <sup>i)</sup>		
					$4.97 \pm 20$ <sup>i)</sup>		$\pi = +$
$4.973 \pm 10$	2						
$5.264 \pm 20$							
$5.390 \pm 20$	2				$5.39 \pm 20$ <sup>i)</sup>		$\pi = +$
$5.448 \pm 20$							
$5.528 \pm 20$	2				$5.53 \pm 20$		$\pi = +$
$5.840 \pm 40$					$6.05 \pm 20$ <sup>i)</sup>		
$6.230 \pm 50$				j)			
$6.371 \pm 20$							
$6.678 \pm 10$					$6.68 \pm 20$		
$7.110 \pm 20$				j)	$6.93 \pm 20$ <sup>i)</sup>		
$7.730 \pm 40$							

<sup>a)</sup> See also table 16.33 in (Aj 71).

<sup>b)</sup>  $^{14}\text{N}({}^3\text{He}, \text{n})^{16}\text{F}$  (Bo 73hh;  $E({}^3\text{He}) = 13$  MeV).

<sup>c)</sup>  $^{14}\text{N}({}^3\text{He}, \text{np})^{15}\text{O}$  (Ot 76a;  $E({}^3\text{He}) = 6.5 \rightarrow 7.8$  MeV).

<sup>d)</sup> From angular correlation studies (Ot 76a).

<sup>e)</sup>  $^{16}\text{O}(\text{p}, \text{n})^{16}\text{F}$  (Mo 71);  $E = 23.9$  MeV.

<sup>f)</sup>  $^{16}\text{O}({}^3\text{He}, \text{t})$  and  $^{19}\text{F}({}^3\text{He}, {}^6\text{He})^{16}\text{F}$  (Na 77c;  $E({}^3\text{He}) = 35$  and 70 MeV, respectively).

<sup>g)</sup>  $^{14}\text{N}({}^3\text{He}, \text{n})^{16}\text{F}$  (Za 65, Ot 76a).

<sup>h)</sup> See (Pe 65, Bo 73hh, Fl 74a, Ot 76a, Na 77c).

<sup>i)</sup> Observed only in  $^{19}\text{F}({}^3\text{He}, {}^6\text{He})$ .


<sup>j)</sup> Strongly excited states at  $E_x = 7.6, 9.4$  and  $11.5$  MeV (with  $\Delta l = 1$  distributions) and weakly excited states at  $E_x = 3.76, 4.65$  and  $6.23$  MeV ( $l = 0$ ) are reported by (An 81: abstract).

tions, that the ordering is  $J^\pi = 0^-, 1^-, 2^-, 3^-$  for the first four states of  $^{16}\text{N}$ . At  $E({}^3\text{He}) = 35$  MeV,  $^{16}\text{F}^*(0.43, 0.72)$  are most strongly populated (Na 77c).

##### 5. $^{19}\text{F}({}^3\text{He}, {}^6\text{He})^{16}\text{F}$

$$Q_m = -14.842$$

Observed states of  $^{16}\text{F}$  are displayed in table 16.26. At  $E({}^3\text{He}) = 70$  MeV the transition strengths favor the sequence  $0^-, 1^-, 2^-, 3^-$  for the first four states of  $^{16}\text{F}$  (Na 77c).



**Fig. 5.** Isobar diagram,  $A = 16$ . The diagrams for individual isobars have been shifted vertically to eliminate the neutron-proton mass difference and the Coulomb energy, taken as  $E_C = 0.60Z(Z-1)/A^{1/3}$ . Energies in square brackets represent the (approximate) nuclear energy,  $E_N = M(Z, A) - AM(H) - NM(n) = E_C$ , minus the corresponding quantity for O; here  $M$  represents the atomic mass excess in MeV. Levels which are presumed to be isospin multiplets are connected by dashed lines.

**<sup>16</sup>Ne MASTER TABLE**ENERGY LEVELS OF LIGHT NUCLEI  $A = 16-17$ 

71

**<sup>16</sup>Ne**

(Fig. 5)

GENERAL<sup>†</sup>*Theoretical work:* (Gu 78d, Sp 78, Li 81h)*Reviews:* (Ce 77c, Al 79k, Tr 80d)

**Mass of <sup>16</sup>Ne:** The  $Q$ -values of the  $^{20}\text{Ne}(\alpha, ^8\text{He})$  and  $^{16}\text{O}(\pi^+, \pi^-)$  reactions lead to an atomic mass excess of  $24.02 \pm 0.04$  MeV for <sup>16</sup>Ne. <sup>16</sup>Ne is then unbound with respect to decay into  $^{14}\text{O} + 2\text{p}$  by 1.43 MeV and is bound with respect to decay into  $^{15}\text{F} + \text{p}$  by 0.04 MeV.

TABLE 16.27  
Energy levels of <sup>16</sup>Ne

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$0^+; 2$	$50^{+50}_{-45}$	$\text{p}$	1, 2
$1.69 \pm 0.07$	$(2^+); 2$			2



$$Q_0 = -28.785 \pm 0.045 \text{ MeV} \text{ (Bu 80b).}$$

At  $E_{\pi^+} = 180$  MeV <sup>16</sup>Ne<sub>g.s.</sub> is populated. The cubic factor,  $d$ , in the IMME is calculated to be  $2.5 \pm 3.7$  keV based on  $Q_0$  (Bu 80b). See also (Ho 77f, Bu 78e, Mi 80d, Gr 81a, Gr 81c).



$$Q_0 = -60.15 \pm 0.08 \text{ MeV}$$

At  $E_\alpha \approx 117.5$  MeV, <sup>16</sup>Ne\*(0,  $1.69 \pm 0.07$ ) are populated, the former with a differential cross section of  $5 \pm 3$  nb/sr at  $8^\circ$  (lab). The  $\Gamma_{\text{c.m.}}$  for the ground state group is  $200 \pm 100$  keV; applying penetrability corrections leads to a total decay width of 5–100 keV. The di-proton branching ratio is 10–90%, with the most probable value being 20%. The cubic term,  $d$ , in the IMME is  $8 \pm 5$  keV,  $15 \pm 6$  keV based, respectively, on the masses of <sup>16</sup>Ne\*(0, 1.69). The first  $T = 2$  states in <sup>16</sup>F[ $0^+, 2^+$ ] are predicted to lie at  $E_x = 10.08 \pm 0.02$  and  $11.87 \pm 0.03$  MeV (Ke 78c).

<sup>†</sup> See also (Aj 77).

$^{17}\text{Be}$ 

(Not illustrated)

This nucleus has not been observed. Its atomic mass excess is calculated to be 70.67 MeV. It is then unstable with respect to breakup into  $^{16}\text{Be} + \text{n}$  and  $^{15}\text{Be} + 2\text{n}$  by 3.37 and 3.34 MeV, respectively (Th 74).

 $^{17}\text{B}$ 

(Not illustrated)

$^{17}\text{B}$  has been observed in the 4.8 GeV proton bombardment of uranium: it is particle stable and its ground state  $J^\pi$  is probably  $\frac{3}{2}^-$  (Bo 74a). Its atomic mass excess is calculated to be 44.37 MeV (transverse form of the mass equation): it is then stable with respect to decay into  $^{15}\text{B} + 2\text{n}$  by 0.73 MeV (Th 74, Je 75). The  $E_{\beta^-}$  (max) for the decay to  $^{17}\text{C}$  would then be 23.3 MeV. See also (Wa 77) and (Se 81c; theor.).

 $^{17}\text{C}$ 

(Fig. 9)

The  $Q$ -value of the  $^{48}\text{Ca}(^{18}\text{O}, ^{17}\text{C})^{49}\text{Ti}$  reaction, studied at  $E(^{18}\text{O}) = 102$  MeV,  $\theta \sim 8^\circ$ , leads to an atomic mass excess of  $21.023 \pm 35$  keV. A group is also observed corresponding to an excited state of  $^{17}\text{C}$  with  $E_x = 292 \pm 20$  keV. Three closely spaced states with  $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+$  are predicted, based on systematics. The cross sections for formation of  $^{17}\text{C}^*(0, 0.29)$  are 3 and 9  $\mu\text{b}/\text{sr}$ , respectively, at  $\theta_{\text{c.m.}} = 10.5^\circ$  (No 77c). On the basis of the atomic mass excess given above,  $E_{\beta^-}$  (max) to  $^{17}\text{N}_{\text{g.s.}} = 13.15$  MeV.  $^{17}\text{C}$  is stable with respect to  $^{16}\text{C} + \text{n}$  by 0.74 MeV. The  $Q$ -value of the  $^{207}\text{Pb}(^{18}\text{O}, ^{17}\text{C})^{208}\text{Po}$  reaction is  $-26.87 \pm 0.22$  MeV, leading to a mass excess of  $21.10 \pm 0.22$  MeV for the ground state (doublet) of  $^{17}\text{C}$  (Ba 79aa). See also (Aj 77) and (Ar 77b, Al 80h).

 $^{17}\text{N}$ 

(Figs. 6 and 9)


GENERAL<sup>†</sup>

*Theoretical papers and reviews:* (Kr 78f, Al 79n, Be 79a, Bo 79gg, Mi 80h, Os 81a)

*Experimental papers:* (Ar 77b, Ba 77al, Ge 78b, Ko 78a, Ol 80, Gr 81b)

1. (a)  $^{17}\text{N}(\beta^-)^{17}\text{O}^* \rightarrow ^{16}\text{O} + \text{n}$   $Q_m = 4.536$   
(b)  $^{17}\text{N}(\beta^-)^{17}\text{O}$   $Q_m = 8.680$

<sup>†</sup> See also (Aj 77).

Fig. 6. Energy levels of <sup>17</sup>N. For notation see fig. 2.

The half-life of <sup>17</sup>N is  $4.173 \pm 0.004$  sec (Oh 76e, Al 72m). See also (Aj 71). The decay is principally [see table 17.2] to the neutron unbound states <sup>17</sup>O\*(4.55, 5.38, 5.94) [ $J^\pi = \frac{3}{2}^-, \frac{3}{2}^+, \frac{1}{2}^-$ ]. The nature of the decay is in agreement with  $J^\pi = \frac{1}{2}^-$  for <sup>17</sup>N<sub>g.s.</sub> (Po 73d, Al 76). For a comparison of the <sup>17</sup>N and <sup>17</sup>Ne decays see table 17.3 (Al 76, Oh 76e).

$$2. {}^{10}\text{Be}({}^9\text{Be}, \text{p}){}^{17}\text{N} \quad Q_m = 7.537$$

Angular distributions are reported at  $E({}^9\text{Be}) = 5.0$  MeV to <sup>17</sup>N\*(0, 1.37, 2.53) (Yo 77d).

<sup>†</sup> See (Aj 77) for several other reactions leading to <sup>17</sup>N.

**<sup>17</sup>N MASTER TABLE**

74

F. AJZENBERG-SELOVE

 TABLE 17.1  
 Energy levels of <sup>17</sup>N<sup>a)</sup>

$E_x$ in <sup>17</sup> N (MeV $\pm$ keV)	$J^\pi; T$	$\tau$	Decay	Reactions
0	$\frac{1}{2}^-; \frac{3}{2}$	$\tau_{1/2} = 4.173 \pm 0.004$ sec	$\beta^-$ b)	1 $\rightarrow$ 10
$1.3739 \pm 0.3$	$\frac{3}{2}^-$	$\tau_m = 93 \pm 35$ fsec	$\gamma$	2, 3, 4, 6, 9, 10
$1.8496 \pm 0.3$	$\frac{1}{2}^+$	$41^{+20}_{-9}$ psec	$\gamma$	3, 4, 9, 10
$1.9068 \pm 0.3$	$\frac{3}{2}^-$	$11 \pm 2$ psec	$\gamma$	3, 4, 6, 10
$2.5260 \pm 0.5$	$\frac{3}{2}^+$	$33 \pm 3$ psec	$\gamma$	2, 3, 4, 9, 10
$3.1289 \pm 0.5$	$\frac{7}{2}^-$	$275 \pm 80$ fsec	$\gamma$	3 $\rightarrow$ 6, 10
$3.2042 \pm 0.9$	$\frac{3}{2}^-$	$< 30$ fsec	$\gamma$	3, 4, 9, 10
$3.6287 \pm 0.7$	$(\frac{7}{2}, \frac{9}{2})^-$	$12 \pm 2$ psec	$\gamma$	3 $\rightarrow$ 6
$3.663 \pm 4$	$\frac{1}{2}^-$	$< 350$ fsec	$\gamma$	3, 4
$3.9060 \pm 2.0$	$(\frac{3}{2}, \frac{5}{2})^-$	$52 \pm 22$ fsec	$\gamma$	3, 4
$4.0064 \pm 2.0$	$\frac{3}{2}^-$	$< 15$ fsec	$\gamma$	3, 4, 9
$4.209 \pm 3$	$\frac{3}{2}^+$	$< 70$ fsec	$\gamma$	3, 4
$4.415 \pm 3$	$(\frac{3}{2}, \frac{5}{2})^-$	$(< 60$ fsec)	$\gamma$	3, 4
$5.170 \pm 2$	$(\frac{3}{2})^+$	$< 60$ fsec	$\gamma$	3, 4, 9
$5.195 \pm 3$	$(\frac{1}{2}, \frac{3}{2})^+$	$< 95$ fsec	$\gamma$	3, 4
$5.515 \pm 3$	$\frac{5}{2}^-$	$< 100$ fsec	$\gamma$	3, 4, 9
$5.772 \pm 3$	$\leq \frac{7}{2}$	$< 120$ fsec	$\gamma$	3, 4
( $6.08 \pm 30$ )				3
$E_x$ in <sup>17</sup> N (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma$	Decay	Reactions
$6.233 \pm 8$				3, 4
$6.449 \pm 3$				3, 4
$6.615 \pm 19$				3, 4
$6.938 \pm 15$				4
$6.981 \pm 20$	$(\frac{3}{2})^-$			3, 4, 9
$7.013 \pm 22$				3, 4, 9
$7.17 \pm 40$				3
$7.37 \pm 40$				3
$7.63 \pm 40$				3
$7.73 \pm 40$				3
$8.00 \pm 25$				3
$8.14 \pm 40$		broad		3
$8.55 \pm 40$		broad		3
$8.93 \pm 40$		broad		3
$9.26 \pm 40$		broad		3
$9.74 \pm 40$		broad		3
10.14	$(\frac{1}{2}, \frac{3}{2})^-$			9

<sup>a)</sup> See also table 17.5.<sup>b)</sup> See also tables 17.2 and 17.3.

TABLE 17.2  
Beta decay of <sup>17</sup>N

Decay to <sup>17</sup> O* (keV)	$J^\pi$	Branch (%)	$\log ft$
0	$\frac{5}{2}^+$	$1.6 \pm 0.5$ <sup>c)</sup>	$7.29 \pm 0.11$ <sup>c,f)</sup>
871	$\frac{1}{2}^+$	$3.0 \pm 0.5$ <sup>c)</sup>	$6.80 \pm 0.07$ <sup>c)</sup>
$3055.2 \pm 0.3$ <sup>a)</sup>	$\frac{1}{2}^-$	$0.34 \pm 0.06$ <sup>c)</sup>	$7.08 \pm 0.08$ <sup>c)</sup>
3841	$\frac{5}{2}^-$	$< 7 \times 10^{-3}$ <sup>c)</sup>	$> 8.5$ <sup>c)</sup>
$4551.2 \pm 1.3$ <sup>b)</sup>	$\frac{3}{2}^-$	$38.0 \pm 1.3$ <sup>d)</sup>	$4.41 \pm 0.02$ <sup>a)</sup>
$5083 \pm 21$ <sup>b)</sup>	$\frac{3}{2}^+$	$0.6 \pm 0.4$ <sup>c)</sup>	$5.9 \pm 0.5$ <sup>c)</sup>
$5389.0 \pm 1.2$ <sup>b,h)</sup>	$\frac{3}{2}^-$	$50.1 \pm 1.3$ <sup>d)</sup>	$3.86 \pm 0.02$ <sup>a)</sup>
5738	$(\frac{1}{2}^+)$	$< 0.23$ <sup>c)</sup>	$> 6.0$ <sup>c)</sup>
5868	$\frac{3}{2}^+$	$< 0.15$ <sup>c)</sup>	$> 6.0$ <sup>c)</sup>
$5951.8 \pm 1.9$ <sup>b,h)</sup>	$\frac{1}{2}^-$	$6.9 \pm 0.5$ <sup>d)</sup>	$4.35 \pm 0.03$ <sup>a)</sup>
6356	$\frac{1}{2}^+$	$< 0.08$ <sup>c)</sup>	$> 6.0$ <sup>c)</sup>

<sup>a)</sup> (Al 76): direct ground state decay  $< 1.5\%$ .

<sup>b)</sup> (Oh 76e): from neutron groups. [The  $E_n$  have been recalculated here on the basis of  $4144.3 \pm 0.8$  keV for  $E_b$  for a neutron in <sup>17</sup>O.]  $\Gamma_n$  for <sup>17</sup>O\*(4.55, 5.08, 5.38, 5.94) are, respectively,  $54.8 \pm 0.4$ ,  $113 \pm 55$ ,  $63.2 \pm 1.1$  and  $60.5 \pm 3.2$  keV (Oh 76e). See also table 17.12.

<sup>c)</sup> (Al 76). See also (Aj 77).

<sup>d)</sup> Calculated from the mean of the values from (Po 73d, Al 76, Oh 76e), renormalized here, together with the new branch to <sup>17</sup>O\*(5.08), to lead to a total neutron emission probability of  $95 \pm 1\%$  [100% less the branches to <sup>17</sup>O\*(0, 0.87, 3.06)].

<sup>e)</sup> (Oh 76e).

<sup>f)</sup>  $\log f_{1t} = 9.56 \pm 0.13$  (To 71h).

<sup>g)</sup> Calculated using the tables of (Go 71t).

<sup>h)</sup> See, however, tables 17.12 and 17.7.

TABLE 17.3  
Comparison of <sup>17</sup>N and <sup>17</sup>Ne  $\beta$ -decay<sup>a)</sup>

Final state in		$J^\pi$	$\Gamma_n$ <sup>b,c)</sup> (keV)	$\Gamma_p$ <sup>b)</sup> (keV)	$(ft)^{- d,e})$	$(ft)^{+ d})$	$\delta$ <sup>f)</sup>
<sup>17</sup> O	<sup>17</sup> F						
3.06	3.10	$\frac{1}{2}^-$	0	19	$(1.2 \pm 0.2) \times 10^7$	$(2.78 \pm 0.40) \times 10^6$	$-0.77 \pm 0.08$
4.55	4.70	$\frac{5}{2}^+$	55	230	$(2.57 \pm 0.13) \times 10^4$	$(3.92 \pm 0.18) \times 10^4$	$0.53 \pm 0.11$
5.38	5.52	$\frac{3}{2}^-$	63	69	$(7.2 \pm 0.3) \times 10^3$	$(7.22 \pm 0.15) \times 10^3$	$0.00 \pm 0.04$
5.94	6.04	$\frac{1}{2}^-$	61	28	$(2.24 \pm 0.16) \times 10^4$	$(2.61 \pm 0.07) \times 10^4$	$0.17 \pm 0.09$

<sup>a)</sup> (Al 76, Oh 76e). I am indebted to Dr. D. E. Alburger for his comments.

<sup>b)</sup>  $\Gamma_n$  and  $\Gamma_p$  are the neutron and proton widths of the <sup>17</sup>O and <sup>17</sup>F states, respectively.

<sup>c)</sup>  $\Gamma_n$  for <sup>17</sup>O\*(4.55, 5.08, 5.38, 5.94) are reported to be, respectively  $54.8 \pm 0.4$ ,  $113 \pm 55$ ,  $63.2 \pm 1.1$  and  $60.5 \pm 3.2$  keV (Oh 76e).

<sup>d)</sup>  $(ft)^-$  and  $(ft)^+$  are for the <sup>17</sup>N and <sup>17</sup>Ne decays, respectively.

<sup>e)</sup> See table 17.2.

<sup>f)</sup>  $\delta = [(ft)^+ / (ft)^-] - 1$ .

TABLE 17.4  
Excited states of <sup>17</sup>N from <sup>11</sup>B(<sup>7</sup>Li, p), <sup>18</sup>O(d, <sup>3</sup>He) and <sup>18</sup>O(t,  $\alpha$ ) <sup>a)</sup>

$E_x$ (keV)		$I^{\pi}$	$J^{\pi}$	$C^2 S^{\dagger}$
A	B			
1373.7 ± 0.5	0	1	$\frac{1}{2}^-$	2.02
1850.0 ± 0.5	$1374.1 \pm 0.4$ <sup>e)</sup>	1	$\frac{3}{2}^-$	0.38
1906.8 ± 0.4	$1849.5 \pm 0.3$ <sup>e)</sup>	0 <sup>h)</sup>	$\frac{1}{2}^+$	$0.41 \pm 0.14$
2526.3 ± 1.0	$1906.9 \pm 0.5$ <sup>e)</sup>	2	$\frac{5}{2}^-$	$0.53 \pm 0.17$
3128.7 ± 0.6	$2525.9 \pm 0.6$ <sup>e)</sup>		$\frac{5}{2}^+$	
3203 ± 2	$3129.2 \pm 0.6$ <sup>e)</sup>	1	$\frac{7}{2}^-$	0.05
3628.7 ± 0.7	$3204.4 \pm 0.9$ <sup>e)</sup>		$>\frac{3}{2}^+$	
3663 ± 4			$(\frac{1}{2}, \frac{3}{2})^-$	
3906.0 ± 2.0			$\leq \frac{7}{2}$	
4006.4 ± 2.0	4000 <sup>f)</sup>	(1)	$\frac{3}{2}^-$	0.04
4208 ± 3			$\leq \frac{5}{2}$	
4415 ± 3			$\leq \frac{7}{2}$	
5170 ± 2	5170 <sup>f)</sup>	(2)	$\frac{3}{2} \leq J \leq \frac{9}{2}^-$	0.08
5195 ± 3			$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$	
5514 ± 3	= 5523 <sup>f)</sup>	1	$\frac{3}{2}^-$	1.83
5770 ± 3			$\leq \frac{7}{2}$	
6080 ± 30 <sup>b,c)</sup>				
6240 ± 25 <sup>b,c)</sup>				
6430 ± 30 <sup>b,c)</sup>				
6610 ± 25 <sup>b,c)</sup>				
6990 ± 20 <sup>b,c)</sup>	6990 <sup>f,h)</sup>	1	$(\frac{3}{2})^-$	0.32
7170 ± 40 <sup>g)</sup>				
7370 ± 40 <sup>g)</sup>				
7630 ± 40 <sup>g)</sup>	7510 <sup>f)</sup>	(1)	$(\frac{1}{2}, \frac{3}{2})^-$	0.09
7730 ± 40 <sup>g)</sup>				
8000 ± 25 <sup>b,c)</sup>				
8140 ± 40 <sup>g)</sup>				
8550 ± 40 <sup>c,d)</sup>				
8930 ± 40 <sup>g)</sup>				
9260 ± 40 <sup>g)</sup>				
9740 ± 40 <sup>g)</sup>	10 140 <sup>f)</sup>	(1)	$(\frac{1}{2}, \frac{3}{2})^-$	0.5

A <sup>11</sup>B(<sup>7</sup>Li, p)<sup>17</sup>N: (Ro 74j), except for states labelled <sup>b)</sup> and <sup>g)</sup>.

B <sup>18</sup>O(t,  $\alpha$ )<sup>17</sup>N (Gu 76d) and <sup>18</sup>O(d, <sup>3</sup>He)<sup>17</sup>N (Ma 77k).

<sup>a)</sup> See also table 17.4 in (Aj 77).

<sup>b)</sup> (Ha 65c).

<sup>c)</sup> (Mc 66a).

<sup>d)</sup> This state and the ones below are broad.

<sup>g)</sup> (Gu 76d).

<sup>f)</sup> <sup>18</sup>O(d, <sup>3</sup>He)<sup>17</sup>N (Ma 77k);  $E_d = 52$  MeV; DWBA analysis.

<sup>h)</sup> (Ro 74j, Gu 76d, Ma 77k).

<sup>i)</sup> Unresolved.

<sup>j)</sup> Probably  $(\frac{7}{2}, \frac{5}{2})^-$  (Ro 74j).

<sup>k)</sup> Probably  $(\frac{7}{2}, \frac{5}{2})^+$  (Ro 74j); see, however, (Ma 77k).

<sup>l)</sup> <sup>18</sup>O(d, <sup>3</sup>He)<sup>17</sup>N (Ma 81);  $E_d = 52$  MeV.

TABLE 17.5  
Radiative transitions and lifetimes of <sup>17</sup>N states <sup>a)</sup>

$E_i$ (MeV)	$E_f$ (MeV)	Mpl.	Branch (%)	$\Gamma_r/\Gamma_w$ <sup>b)</sup> (W.u.)	$\tau_m$
1.37	0	M1	100	$0.13 \pm 0.05$	$93 \pm 35$ fsec
1.85	0	E1	$86.5 \pm 2.5$	$41^{+20}_{-9}$ psec	
	1.37	E1	$13.5 \pm 2.5$	$(3.2 \pm 1.5) \times 10^{-5}$	
1.91	0	E2	$77.0 \pm 2.5$	$0.9 \pm 0.2$	$11 \pm 2$ psec
	1.37	M1	$23.0 \pm 2.5$	$(5 \pm 1) \times 10^{-3}$ <sup>c)</sup>	
2.53	0	M2	$11 \pm 1$	$0.22 \pm 0.04$	$33 \pm 3$ psec
	1.37	E1	$34 \pm 3$	$(1.0 \pm 0.2) \times 10^{-5}$	
	1.85	E2	$12.0 \pm 1.5$	$8.1 \pm 1.6$	
	1.91	E1	$41.0 \pm 2.5$		
3.13 <sup>d)</sup>	1.91	M1	100	$0.06 \pm 0.02$	$275 \pm 80$ fsec
3.20 <sup>e)</sup>	0	M1	$88 \pm 4$	$>0.025$ <sup>f)</sup>	$<30$ fsec
	1.91	M1	$12 \pm 4$	$>0.05$	
3.63 <sup>g)</sup>	1.91	E2	$47 \pm 10$	$0.8 \pm 0.2$	$12 \pm 2$ psec
	3.13	M1	$53 \pm 10$	$0.010 \pm 0.03$	
3.66	1.85	E1	100	$>7 \times 10^{-4}$	$<350$ fsec
3.91	1.91	M1	100	$(8^{+5}_{-3}) \times 10^{-2}$ <sup>h)</sup>	$52 \pm 22$ fsec
4.01	1.85		$\leq 15 \pm 5$ <sup>i)</sup>		
	2.53	(M1)	$85 \pm 5$	0.55	$<15$ fsec
4.21	1.37		100		$<70$ fsec
4.42	1.91		100		$(<60$ fsec)
5.17	2.53	E2	$37 \pm 7$	$>15$	$<60$ fsec
	3.13		$63 \pm 7$		
5.20	1.85		$\sim 42$		$<95$ fsec
	1.91		$\sim 58$		
5.52	0		$\sim 50$		$<100$ fsec
	1.37		$\sim 50$		
5.77	1.37		$\sim 25$		$<120$ fsec
	1.91		$\sim 25$		
	4.01		$\sim 50$ <sup>j)</sup>		

<sup>a)</sup> See table 17.5 in (Aj 77) for references and additional detail.<sup>b)</sup> Assuming pure multipole transitions and  $J^\pi$  from table 17.1: see also table 2.<sup>c)</sup>  $\Gamma_r/\Gamma_w = 0.4^{+0.4}_{-1.3}$  (E2) (Gu 76d).<sup>d)</sup> Branches to <sup>17</sup>N\*(0, 1.37, 1.85, 2.53) are, respectively,  $<2$ ,  $<5$ ,  $<2$  and  $<3$  (Gu 76d).<sup>e)</sup> Branches to <sup>17</sup>N\*(1.37, 1.85, 2.53) are, respectively,  $<5$ ,  $<6$  and  $<3$  (Gu 76d).<sup>f)</sup>  $\delta = -0.06 \pm 0.08$  or  $2.1 \pm 0.4$  (Gu 76d). All other  $\delta$  reported by (Gu 76d) are consistent with 0.<sup>g)</sup> Branches to <sup>17</sup>N\*(0, 1.37, 1.85, 2.53, 3.20) are, respectively,  $<10$ ,  $<10$ ,  $<7$ ,  $<3$ ,  $<2$  (Ro 74k).<sup>h)</sup> This number appears to be in error: see table 2.<sup>i)</sup> This branch is uncertain.

**<sup>17</sup>N**

78

F. AJZENBERG-SELOVE

3.  $^{11}\text{B}({}^7\text{Li}, \text{p})^{17}\text{N}$   $Q_m = 8.417$

Observed proton groups and  $\gamma$ -rays are displayed in table 17.4. Table 17.5 shows branching ratio and lifetime measurements (Ro 74j, Ro 74k).

4.  $^{15}\text{N}(\text{t}, \text{p})^{17}\text{N}$   $Q_m = -0.108$

Observed proton groups are displayed in table 17.6. See also (Aj 71).

5.  $^{15}\text{N}(\alpha, 2\text{p})^{17}\text{N}$   $Q_m = -19.922$

At  $E_\alpha = 65 \text{ MeV}$   ${}^{17}\text{N}^*(3.13, 3.63)$  are the only states of  ${}^{17}\text{N}$  which are strongly populated (Ja 78i).

TABLE 17.6  
States of  ${}^{17}\text{N}$  from  ${}^{15}\text{N}(\text{t}, \text{p})^*$

$E_x$ (keV)	$L$	$J^\pi$
0 <sup>b)</sup>	0	$\frac{1}{2}^-$
$1372 \pm 6$ <sup>b)</sup>	2	$(\frac{3}{2}, \frac{5}{2})^+$
$1851 \pm 4$	1	$(\frac{1}{2}, \frac{3}{2})^+$
$1909 \pm 3$ <sup>b)</sup>	2	$(\frac{3}{2}, \frac{5}{2})^-$
$2524 \pm 4$	3	$(\frac{5}{2}, \frac{7}{2})^+$
$3127 \pm 6$ <sup>b)</sup>	4	$(\frac{7}{2}, \frac{9}{2})^-$
$3201 \pm 5$ <sup>b)</sup>	2	$(\frac{3}{2}, \frac{5}{2})^-$
$3625 \pm 6$ <sup>b)</sup>	4	$(\frac{7}{2}, \frac{9}{2})^-$
$3664 \pm 6$ <sup>b)</sup>	0	$\frac{1}{2}^-$
$3906 \pm 5$ <sup>b)</sup>	2	$(\frac{1}{2}, \frac{3}{2})^-$
$4011 \pm 6$	(1)	
$4213 \pm 6$	3	$\frac{5}{2}^+ \text{?})$
$4420 \pm 7$ <sup>b)</sup>	2	$(\frac{1}{2}, \frac{3}{2})^-$
$5179 \pm 4$ <sup>c)</sup>	5	$(\frac{9}{2}^+)$
	1	$((\frac{1}{2}, \frac{3}{2})^+)$
$5517 \pm 6$	(2)	
$5780 \pm 6$	(1)	
$6233 \pm 8$ <sup>d)</sup>	(2)	
$6449 \pm 3$	(4, 5)	
$6627 \pm 30$	weak	
$6938 \pm 15$		
$6981 \pm 20$	(3, 4)	
$7013 \pm 22$		

<sup>a)</sup> (Fo 79j):  $E_t = 15.0 \text{ MeV}$ ; DWBA analysis. See also (Mo 78g).

<sup>b)</sup> Predominantly 2p-1h states.

<sup>c)</sup> Unresolved states.

<sup>d)</sup>  ${}^{17}\text{N}^*(6.08)$  is not observed.

<sup>e)</sup> The  $\frac{7}{2}^+$  possibility can be eliminated because the  $4.21 \rightarrow 1.37 \text{ MeV}$  transition would then have too large an M2 strength ( $> 500 \text{ W.u.}$ ) [P. M. Endt, private communication].

6.  $^{15}\text{N}({}^{13}\text{C}, {}^{11}\text{C})^{17}\text{N}$   $Q_m = -15.293$

At  $E({}^{13}\text{C}) = 105 \text{ MeV}$   $^{17}\text{N}^*(3.13, 3.63)$  are strongly populated (Ra 79c).

7.  $^{17}\text{O}(\text{n}, \text{p})^{17}\text{N}$   $Q_m = -7.898$

See  ${}^{18}\text{O}$  in (Aj 78).

8.  ${}^{18}\text{O}(\gamma, \text{p})^{17}\text{N}$   $Q_m = -15.942$

See (Ba 76ii) and  ${}^{18}\text{O}$  in (Aj 83).

9.  ${}^{18}\text{O}(\text{d}, {}^3\text{He})^{17}\text{N}$   $Q_m = -10.449$

Observed groups of  ${}^3\text{He}$  ions are displayed in table 17.4: the  $\frac{1}{2}^-$  ground state contains the full  $1p_{1/2}$  strength. The strength of all other observed  $l=1$  transitions is assumed to originate from pickup from the  $1p_{3/2}$  shell (Ma 77k). Analyzing powers have been obtained at  $E_d = 52 \text{ MeV}$  to  $^{17}\text{N}^*(0, 1.37, 1.85, 2.53, 5.51, 6.99)$  (Ma 81).

10.  ${}^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$   $Q_m = 3.872$

At  $E_t = 3.5 \text{ MeV}$  (Gu 76d) have studied  $\alpha - \gamma$  angular correlations and  $\gamma$ -ray branching ratios for the first six excited states of  ${}^{17}\text{N}$ : see tables 17.4 and 17.5. See also (Aj 71).

## $^{17}\text{O}$

(Figs. 7 and 9)

### GENERAL<sup>†</sup>

*Shell model*: (Ba 76gg, Du 77b, Ha 77w, Po 77c, Ch 78z, Kr 78, Ka 79a, Br 80i, Va 80a).

*Collective and cluster models*: (Ch 78z, Ta 78u, Th 78e, Fu 80h).


*Special states*: (He 77h, Sh 77m, Al 78r, En 78c, Ta 78u, Ma 79bb, Za 79b, Ch 80o, Fu 80h, Hi 80c, Va 80a, Ta 81i).

*Electromagnetic transitions*: (Mc 76m, Al 77i, An 77a, Br 77, Ha 77w, Ho 77, Za 78, Su 79a, Ch 80o).

*Complex reactions involving  ${}^{17}\text{O}$* : (Du 76b, Le 76t, Ar 77b, Pe 77<sup>††</sup>, Re 77, Ko 78a, Se 78d, Sa 79l, La 80j, Ol 80, Sh 81e).

<sup>†</sup> See also (Aj 77).

<sup>††</sup> See also for spectroscopic factors.

Fig. 7. Energy levels of <sup>17</sup>O. For notation see fig. 2.

*Astrophysical questions:* (Bo 76cc, No 76l, Au 77h, Ca 77q, Co 77h, Co 77z, De 77p, Di 77e, En 77a, No 77, Pr 77h, Ra 77, Tr 77i, Wa 77t, Bu 78k, Cl 78g, Le 78t, Po 78b, St 78i, Tr 78i, Wo 78f, Ch 79i, Ja 79v, La 79f, Le 79m, Pe 79e, Ro 79m, Sw 79, Au 80e, Cl 80c, Cl 80d, Wa 80n, Gu 81b, Wi 81c, Wi 81i).

*Applications:* (Am 78c).

*Pion and other meson capture and reactions*<sup>†</sup>: (Ho 77p, Si 77, Ga 78c, Jo 79g, Do 80e, Os 81a).

*Other topics:* (Ba 76gg, Bi 76e, Of 76, Sa 76i, Ba 77v, Gr 77h, Sh 77m, Su 77e, En 78c, Ga 78c, Kr 78, Mc 78d, Sh 78q, Be 79a, Br 79j, Co 79c, He 79, Ka 79a, Ma 79bb, Pi 79g, Va 79b, Zo 79, Hi 80c, Zo 80, Ta 81i).

*Ground state of <sup>17</sup>O:* (Du 76b, Fu 76h, Jo 76c, Mc 76m, Sa 76i, An 77f, Br 77, Du 77b, Ha 77w, Ko 77w, No 77d, Po 77c, Sh 77i, An 78b, Ar 78a, Ar 78b, Ch 78z, Fo 78o, He 78k, Sl 78, Za 78b, Br 79i, Br 79j, Sa 79l, Su 79a, Va 79b, Bo 80b, Br 80i, Ch 80o, Hi 80c, Av 81b, Si 81b, Ta 81i, Va 81c).

$$\mu = -1.89379 \pm 0.00009 \text{ n.m. (see Le 78n).}$$

$$Q = -25.78 \text{ mb (Sc 69u).}$$

$$1. \ ^7\text{Li}(\text{<sup>14</sup>N}, \alpha)^{17}\text{O} \quad Q_m = 16.157$$

The angular distribution of the  $\alpha_{0+1}$  group has been measured at  $E(^{14}\text{N}) = 27.6 \text{ MeV}$  (Wa 64g).

$$2. \ ^9\text{Be}(\text{<sup>16</sup>O}, \text{<sup>8</sup>Be})^{17}\text{O} \quad Q_m = 2.479$$

Angular distributions have been studied at  $E(^{16}\text{O}) = 11, 15$  and  $18 \text{ MeV}$ , involving  $^{17}\text{O}^*(0, 0.87)$ : see (Aj 77). The cross section for the population of  $^{17}\text{O}^*(0.87)$  has been measured for  $E(^{16}\text{O}) = 5.5$  to  $22 \text{ MeV}$ : see (Sw 77, Ba 70v). See also (Ch 79f). For astrophysical questions see (Sw 77, Cu 80). At  $E(^9\text{Be}) = 50 \text{ MeV}$  the population of  $^{17}\text{O}^*(5.09, 5.7, 7.6)$  is also reported (St 77p).

3. (a) $^{10}\text{B}(^7\text{Li}, \text{p})^{16}\text{N}$	$Q_m = 13.989$	$E_b = 27.770$
(b) $^{10}\text{B}(^7\text{Li}, \text{d})^{15}\text{N}$	$Q_m = 13.723$	
(c) $^{10}\text{B}(^7\text{Li}, \text{t})^{14}\text{N}$	$Q_m = 9.147$	
(d) $^{10}\text{B}(^7\text{Li}, \alpha)^{13}\text{C}$	$Q_m = 21.4099$	

Cross sections to various of the final states have been measured at  $E(^7\text{Li}) = 5.20 \text{ MeV}$  (Mc 66a).

<sup>†</sup> See also reactions 38 and 68.

# <sup>17</sup>O MASTER TABLE

TABLE 17.7  
Energy levels of <sup>17</sup>O

$E_x$ in <sup>17</sup> O (MeV ± keV)	$J^\pi; T$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{1}{2}$		stable	1, 2, 5 → 9, 12, 13, 15, 16, 18 → 24, 29 → 32, 38 → 73
0.87081 ± 0.12	$\frac{1}{2}^+$	$\tau_m = 258.6 \pm 2.6$ psec	$\gamma$	1, 2, 5 → 7, 12, 13, 15, 16, 18 → 22, 29 → 32, 38, 39, 42 → 44, 46 → 48, 50, 52, 54, 56, 57, 61, 62, 63, 68 → 73
3.0552 ± 0.3	$\frac{1}{2}^-$	$\tau_m = 120^{+80}_{-60}$ fsec	$\gamma$	5 → 7, 12, 13, 18, 21, 22, 29, 31, 32, 38, 39, 42, 48, 50, 52, 61, 62, 70, 71
3.841 ± 3	$\frac{5}{2}^-$	$\tau_m \leq 25$ fsec	$\gamma$	5 → 7, 12 → 14, 18, 21, 22, 29, 30, 39, 42, 50, 61, 62, 70, 71
4.552 ± 2	$\frac{3}{2}^-$	$\Gamma = 40 \pm 5$	$\gamma, n$	5, 7, 12, 13, 21, 22, 29, 30, 33, 39, 42, 48, 49, 50, 61, 62, 71
5.085 ± 2	$\frac{3}{2}^+$	96 ± 5	$\gamma, n$	2, 6, 7, 12, 13, 21, 22, 29, 33, 39, 48, 49, 50, 61, 62
5.218	$(\frac{9}{2}^-)$	<0.1	$\gamma, n$	6, 7, 12 → 14, 21, 22, 29, 30, 33, 39, 50, 61, 71
5.378 ± 2	$\frac{3}{2}^-$	28 ± 7	$\gamma, n$	7, 21, 22, 29, 33, 39, 42, 48, 49, 50, 61, 62, 71
5.697 ± 2	$\frac{7}{2}^-$	3.4 ± 0.3	$\gamma, n$	2, 6, 12, 13, 21, 22, 29, 30, 33, 39, 49, 50, 62
5.733 ± 2		<1	n	2, 5, 6, 12, 13, 21, 22, 33, 39, 71
5.868 ± 2	$\frac{3}{2}^+$	6.6 ± 0.7	n	6, 7, 12, 13, 21, 22, 29, 33, 39, 71
5.939 ± 4	$\frac{1}{2}^-$	32 ± 3	$\gamma, n$	5, 6, 12, 13, 21, 22, 29, 33, 39, 48, 50, 62, 71
6.356 ± 8	$\frac{1}{2}^+$	124 ± 12	$\gamma, n$	5, 7, 21, 29, 33, 50
6.862 ± 2	$(\frac{1}{2}^-)$	<1	$\gamma, n$	5, 6, 7, 12, 13, 21, 22, 29, 33, 39, 50, 62, 71
6.972 ± 2	$(\frac{5}{2}^+)$	<1	$\gamma, n$	6, 7, 12, 13, 21, 22, 29, 33, 50, 71
7.1657 ± 0.8	$\frac{5}{2}^-$	1.38 ± 0.05	n, $\alpha$	5, 6, 7, 10, 12, 13, 21, 29, 33, 37
7.202 ± 10	$\frac{3}{2}^+$	280 ± 30	n, $\alpha$	12, 13, 21, 33, 37
7.3792 ± 1.0	$\frac{5}{2}^+$	0.64 ± 0.23	( $\gamma$ ), n, $\alpha$	5, 6, 7, 10, 12, 13, 29, 30, 33, 37, 50, 62, 71
7.3822 ± 1.0	$\frac{5}{2}^-$	0.96 ± 0.20	$\gamma, n, \alpha$	5, 7, 10, 12, 13, 21, 30, 33, 37, 49, 50, 62, 71
7.559 ± 20	$\frac{3}{2}^-$	500 ± 50	n, $\alpha$	33, 37, 39
7.576 ± 2	$\frac{7}{2}^-$	<0.1	$\gamma, n, \alpha$	5, 6, 10, 12, 13, 21, 29, 33, 50
7.6882 ± 0.9	$\frac{7}{2}^-$	14.4 ± 0.3	$\gamma, n, \alpha$	5, 6, 10, 12, 13, 29, 33, 37, 49
7.757 ± 9	$\frac{11}{2}^-$		$\gamma$	29, 30, 50

TABLE 17.7—continued

$E_x$ in <sup>17</sup> O (MeV ± keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
7.956 ± 6	$\frac{1}{2}^+$	90 ± 9	n, $\alpha$	10, 29, 33, 37
7.99 ± 50	$\frac{1}{2}^-$	270 ± 30	n, $\alpha$	33, 37
8.070 ± 10	$\frac{3}{2}^+$	85 ± 9	n, $\alpha$	10, 29, 33, 37
8.200 ± 7	$\frac{3}{2}^-$	60	$\gamma, n, \alpha$	10, 29, 33, 37, 49, 62
8.3424 ± 0.9	$\frac{1}{2}^+$	11.4 ± 0.5	n, $\alpha$	10, 29, 33, 37, 50
8.4023 ± 0.8	$\frac{5}{2}^+$	6.17 ± 0.13	n, $\alpha$	6, 10, 12, 13, 29, 33, 37, 50
8.4660 ± 0.8	$\frac{7}{2}^+$	2.13 ± 0.11	n, $\alpha$	5, 6, 10, 12, 13, 29, 33, 37, 50, 62
8.5007 ± 0.8	$\frac{5}{2}^-$	6.89 ± 0.22	$\gamma, n, \alpha$	6, 10, 12, 13, 29, 33, 37, 49, 50
8.6870 ± 1.0	$\frac{3}{2}^-$	55.3 ± 0.6	$\gamma, n, \alpha$	10, 29, 33, 37, 49, 62
8.897 ± 8	$\frac{3}{2}^+$	101 ± 3	n, $\alpha$	6, 10, 12, 13, 29, 30, 33, 37
8.9672 ± 1.7	$\frac{7}{2}^-$	26 ± 2	$\gamma, n, \alpha$	6, 10, 12, 13, 29, 33, 37, 49
9.147 ± 4	$\frac{1}{2}^-$	4 ± 3	n, $\alpha$	6, 10, 12, 13, 62
9.15 ± 20	$\frac{9}{2}^-$			29, 30
9.18	$\frac{7}{2}^-$	3	$\alpha$	10, 12, 13
9.1939 ± 0.8	$\frac{5}{2}^+$	3.53 ± 0.13	n, $\alpha$	10, 12, 13, 33
9.42	$\frac{3}{2}^-$	120	n	33
9.492 ± 4	$\frac{5}{2}^-$	15 ± 1	n, $\alpha$	5, 10, 13, 29, 33, 62
9.7119 ± 0.9	$\frac{7}{2}^+$	23.1 ± 0.3	n, $\alpha$	10, 13, 29, 33
9.7833 ± 0.9	$\frac{3}{2}^+$	11.7 ± 0.3	n, $\alpha$	10, 13, 33
9.8589 ± 0.9	( $\frac{5}{2}^-$ )	4.01 ± 0.23	n, $\alpha$	10, 13, 29, 33
9.8765 ± 1.3	( $\frac{1}{2}^-$ )	16.7 ± 1.7	n, $\alpha$	10, 13, 29, 33
9.976 ± 20	$\frac{5}{2}^+$	~80	n, $\alpha$	10
10.045 ± 20		~100	n, $\alpha$	10
10.1678 ± 1.0	$\frac{7}{2}^-$	49.1 ± 0.8	n, $\alpha$	10, 33
10.336 ± 15	$\frac{5}{2}^+, \frac{7}{2}^-$	150	n, $\alpha$	10, 29
10.423 ± 3		14 ± 3	n, $\alpha$	10
10.49	$\frac{5}{2}^+, \frac{7}{2}^-$	75 ± 30	n, $\alpha$	10
10.5591 ± 1.0	( $\frac{7}{2}^-$ )	42.5 ± 1.1	n, $\alpha$	10, 14, 29, 33, 34
10.777 ± 3	$\frac{1}{2}^+, \frac{7}{2}^-$	74 ± 3	n, $\alpha$	10, 13, 29, 34
10.9129 ± 2.8	( $\frac{5}{2}^+$ )	41.7 ± 1.4	n, $\alpha$	10, 29, 33, 34
11.036 ± 3	$T = \frac{1}{2}$	31 ± 3	n, $\alpha$	10, 29
11.0787 ± 0.9 *	$\frac{1}{2}^-, \frac{3}{2}^+$	2.4 ± 0.3	n, $\alpha$	10, 29, 33, 62, 63
11.238		80 ± 3	n, $\alpha$	5, 10
11.51	$\geq \frac{3}{2}$	190	n	33, 34
11.622		65 ± 2	n, $\alpha$	10
11.750 ± 10		40 ± 25	$\gamma, n, \alpha$	10, 50
11.815 ± 15		12 ± 3	n, $\alpha$	10
12.005 ± 15	$\geq \frac{3}{2}$	270	n, $\alpha$	10, 33, 34, 50
12.11 ± 20		150 ± 50	n, $\alpha$	10, 14, 34
12.274 ± 15		100 ± 30	n, $\alpha$	10
12.38 ± 20			n, $\alpha$	10, 33
12.420 ± 15			n, $\alpha$	10
12.4660 ± 1.0	$\frac{1}{2}^-, \frac{3}{2}^+$	6.9 ± 1.1	n, $\alpha$	10, 33, 34, 62, 63
12.595 ± 15		75 ± 30	n, $\alpha$	10

TABLE 17.7—continued

$E_x$ in <sup>17</sup> O (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma_{c.m.}$ (keV)	Decay	Reactions
12.669 $\pm$ 15		$\approx 5$	n, $\alpha$	10, 33, 34
12.81 $\pm$ 25			n, $\alpha$	10
12.93 $\pm$ 20		$\geq 150$	n, $\alpha$	10
12.944 $\pm$ 5	$\frac{1}{2}^+; \frac{3}{2}^+$	$6 \pm 2$	n, $\alpha$	10, 33, 34, 62, 63
12.9982 $\pm$ 1.0	$\frac{5}{2}^-; \frac{3}{2}^-$	$2.5 \pm 1.0$	n, $\alpha$	10, 33, 63
13.076 $\pm$ 15		$16 \pm 4$	n, $\alpha$	10
13.484 $\pm$ 15		$\approx 120$	n, $\alpha$	10
13.58 $\pm$ 20	( $\frac{11}{2}^-$ , $\frac{13}{2}^-$ )			12, 13
13.609 $\pm$ 15		$250 \pm 100$	n, $\alpha$	10
13.6353 $\pm$ 2.5 (13.67)	( $\frac{5}{2}^+$ , $\frac{3}{2}^-$ )	$9 \pm 5$	n, $\alpha$	33, 62, 63
14.15 $\pm$ 100	( $\frac{9}{2}^+$ , $\frac{11}{2}^+$ )	400	n	33
14.2303 $\pm$ 1.7	( $\frac{7}{2}^-$ , $\frac{5}{2}^-$ )	$\approx 100$		12
14.286 $\pm$ 3	$T = \frac{5}{2}$	$20.5 \pm 1.6$	n, $\alpha$	33, 63
14.451 $\pm$ 3		$7.5 \pm 4$	n, $\alpha$	33, 63
14.76 $\pm$ 100	( $\frac{\geq 3}{2}^-$ )	$40 \pm 6$	n, $\alpha$	33
14.791 $\pm$ 3	( $\frac{1}{2}^-; \frac{3}{2}^-$ )	340	$\gamma, n$	33, 50
15.00		$36 \pm 13$	n, $\alpha$	33
15.1 $\pm$ 100	( $\frac{9}{2}^+, \frac{11}{2}^+$ )	180	n, d, $\alpha$	28, 33
15.199 $\pm$ 3	( $\frac{3}{2}^+, \frac{5}{2}^+$ )	$\approx 500$		12
15.368 $\pm$ 3 (15.6)	( $\frac{3}{2}^+; \frac{5}{2}^+$ )	$52 \pm 14$	$\gamma, n, d, \alpha$	28, 33, 50
15.95 $\pm$ 150	( $\frac{9}{2}^+, \frac{11}{2}^+$ )	$40 \pm 6$	n, d, $\alpha$	27, 33
16.243 $\pm$ 4	( $\frac{5}{2}^+; \frac{3}{2}^-$ )	$\approx 300$	p, d, $\alpha$	26, 27, 28
16.58 $\pm$ 10	( $\frac{1}{2}^-, \frac{3}{2}^-$ )	$\approx 700$		12
16.6 $\pm$ 150	( $\frac{11}{2}^-, \frac{13}{2}^-$ )	$21 \pm 10$	n, p, d, $\alpha$	26, 33
17.1 $\pm$ 150	( $\frac{11}{2}^-, \frac{13}{2}^-$ )			62
17.436 $\pm$ 11	( $T = \frac{3}{2}$ )	$66 \pm 20$	n, $\alpha$	12
18.110 $\pm$ 4	$\frac{3}{2}^-, \frac{5}{2}^-$	$46 \pm 12$	n, $\alpha$	33, 62
19.6 $\pm$ 150	( $\frac{13}{2}^+, \frac{15}{2}^+$ )	$\approx 250$		12
20.2 $\pm$ 150	( $\frac{13}{2}^+, \frac{15}{2}^+$ )	$\approx 250$		12
21.2	( $\frac{13}{2}^+, \frac{15}{2}^+$ )			12
21.7 $\pm$ 100	$\frac{5}{2}^+$	$\approx 750$	$\gamma, {}^3\text{He}, \alpha$	16, 17
22.1 $\pm$ 100	$\frac{7}{2}^-$	$\approx 750$	$\gamma, n, {}^3\text{He}, \alpha$	12, 16, 17
22.5 $\pm$ 200	$\frac{3}{2}^{(-)}$	$\approx 1000$	$\gamma, {}^3\text{He}$	16
23	$\frac{1}{2}^+$	$\approx 6000$	$\gamma, n$	49, 50
23.0	$\frac{1}{2}^+$	$\approx 400$	$\gamma, {}^3\text{He}$	16, 17
23.5			$\gamma, {}^3\text{He}$	16
24.4			$\gamma, {}^3\text{He}$	16

\*) See also table 17.11, and see table 17.6 in (Aj 77).

4. (a)  ${}^{11}\text{B}({}^6\text{Li}, p){}^{16}\text{N}$        $Q_m = 9.785$        $E_b = 23.565$   
 (b)  ${}^{11}\text{B}({}^6\text{Li}, d){}^{15}\text{N}$        $Q_m = 9.5178$   
 (c)  ${}^{11}\text{B}({}^6\text{Li}, t){}^{14}\text{N}$        $Q_m = 4.9418$   
 (d)  ${}^{11}\text{B}({}^6\text{Li}, \alpha){}^{13}\text{C}$        $Q_m = 17.205$

Cross sections to various of the final states have been measured at  $E(^6\text{Li}) = 4.72 \text{ MeV}$  (Mc 66a).

5.  $^{12}\text{C}(^6\text{Li}, \text{p})^{17}\text{O}$   $Q_m = 7.608$

Angular distributions have been studied for  $E(^6\text{Li}) = 3$  to  $20 \text{ MeV}$  (see Aj 71) and at  $28 \text{ MeV}$  (Wa 80i:  $p_0, p_1, p_2$  and  $p$  to  $^{17}\text{O}^*(8.47, 11.8, 12.4)$ ). See also  $^{18}\text{F}$  in (Aj 78, Aj 83).

6.  $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$   $Q_m = 2.582$

Angular distributions have been measured at  $E(^7\text{Li}) = 3.24 \rightarrow 3.64 \text{ MeV}$  and at  $21.1 \text{ MeV}$ : see table 17.7 in (Aj 77). See also  $^{19}\text{F}$  in (Aj 78, Aj 83).

7.  $^{12}\text{C}(^9\text{Be}, \alpha)^{17}\text{O}$   $Q_m = 9.733$

Angular distributions of the  $\alpha_0$  and  $\alpha_1$  groups are reported at  $E(^9\text{Be}) = 20 \text{ MeV}$  (Ja 79t) and  $16.1, 17.0$  and  $17.9 \text{ MeV}$  (De 81h; also  $\alpha_2$ ). The yield of  $\alpha$ -particles has been studied for  $E(^9\text{Be}) = 9.0$  to  $20 \text{ MeV}$  (De 81h;  $\alpha$  to  $^{17}\text{O}^* < 7.5 \text{ MeV}$ ),  $10.5$  to  $26.3 \text{ MeV}$  (Hu 80e;  $\alpha_0$ ) and  $17.5$  to  $26.3 \text{ MeV}$  (Ma 78v;  $\alpha_0 \rightarrow \alpha_3$ ). See also (Aj 77), (Bo 79bb) and (Br 80q; theor.).

8.  $^{12}\text{C}(^{18}\text{O}, ^{13}\text{C})^{17}\text{O}$   $Q_m = -3.098$

See (Ch 78m).

9.  $^{12}\text{C}(^{20}\text{Ne}, ^{15}\text{O})^{17}\text{O}$   $Q_m = -9.089$

See (Or 79).

10. (a) $^{13}\text{C}(\alpha, \text{n})^{16}\text{O}$	$Q_m = 2.2156$	$E_b = 6.3599$
(b) $^{13}\text{C}(\alpha, \alpha)^{13}\text{C}$		

The yield of neutrons increases monotonically for  $E_\alpha = 0.475$  to  $1 \text{ MeV}$ :  $S(E) = [(5.48 \pm 1.77) + (12.05 \pm 3.91)E] \times 10^5 \text{ MeV} \cdot \text{b}$  (Da 68;  $E_\alpha = 0.48$  to  $0.70 \text{ MeV}$ ),  $[(4.87 \pm 1.28) + (10.86 \pm 2.46)E] \times 10^5 \text{ MeV} \cdot \text{b}$  (Ra 76g, Ra 77;  $E_\alpha = 0.60$  to  $0.90 \text{ MeV}$ ). Astrophysical considerations are discussed by (Ra 76g, Co 77h, Ra 77) and see (Aj 77). Yield curves for reaction (a) have been measured for  $E_\alpha = 1.0$  to  $22.5 \text{ MeV}$ : see (Aj 71, Aj 77) and (Be 75bb;  $\alpha\gamma_{6.13}$ ; thresh. to  $7.2 \text{ MeV}$ ). Elastic scattering studies [(reaction b)] have been studied at  $E_\alpha = 2.0$  to  $26.6 \text{ MeV}$ : see (Aj 71, Aj 77). Observed resonances are displayed in table 17.8..

11. (a) $^{13}\text{C}(\alpha, \text{p})^{16}\text{N}$	$Q_m = -7.421$	$E_b = 6.3599$
(b) $^{13}\text{C}(\alpha, \text{d})^{15}\text{N}$	$Q_m = -7.6874$	
(c) $^{13}\text{C}(\alpha, \text{t})^{14}\text{N}$	$Q_m = -12.2634$	

For reaction (b) see (Aj 77). For all three reactions see  $^{16}\text{N}$ , and  $^{15}\text{N}$  and  $^{14}\text{N}$  in (Aj 81).

12.  $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$   $Q_m = 4.886$

Angular distributions of deuteron groups to many states of <sup>17</sup>O have been measured earlier at  $E(^6\text{Li}) = 18$  and 25.6 MeV: see table 17.7 in (Aj 77). At  $E(^6\text{Li}) = 35.5$  MeV angular distributions are reported to <sup>17</sup>O\*(13.58±0.02) which is strongly populated. Comparisons with <sup>12</sup>C(^6Li, d)<sup>16</sup>O\*(16.29) and with the results of reaction 13 here suggest that the peak corresponding to <sup>17</sup>O\*(13.58) contains a state or states of spin  $\frac{11}{2}^-$ ,  $\frac{13}{2}^-$ , or both, based on <sup>16</sup>O\*(16.29) (Cl 78f). (d,  $\alpha$ ) angular correlations [ $E(^6\text{Li}) = 26, 29$  and 34 MeV] indicate the involvement of <sup>17</sup>O states at  $13.6 \pm 0.1$  [ $J = 6$ ],  $14.15 \pm 0.1$  [5],  $15.1 \pm 0.1$  [5],  $15.95 \pm 0.15$  [5],  $16.6 \pm 0.15$  [6],  $17.1 \pm 0.15$  [6],  $19.6 \pm 0.15$  [7],  $20.2 \pm 0.15$  [7], 21.2 [7] and 22.1 MeV.  $\Gamma \approx 0.1, 0.5, 0.7, 0.25$  and  $0.25$  MeV for <sup>17</sup>O\*(14.2, 15.1, 16.0, 19.6, 20.2) (Ar 78j). See also (St 76n; theor.).

TABLE 17.8  
Resonances in <sup>13</sup>C( $\alpha, n$ ) and <sup>13</sup>C( $\alpha, \alpha$ )<sup>a)</sup>

$E_{\text{res}}$ (MeV ± keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\Gamma_{\alpha}/\Gamma$	$J^\pi$	$E_x$ (MeV)
$1.0563 \pm 1.5^b)$	$1.5 \pm 0.2$		$\frac{5}{2}^-$	7.1676
$1.3367 \pm 1.5$	$0.6^{+0.2}_{-0.1}$			7.3820
$1.3406 \pm 1.5$	$0.8^{+0.3}_{-0.2}$			7.3849
$1.590 \pm 2$	$\leq 1$		$\frac{7}{2}^-$	7.576
$1.745 \pm 6$	$\leq 15$		$\frac{5}{2}^+$	7.694
$2.083 \pm 8$	75	0.03	$\frac{1}{2}^-$	7.953
$2.250 \pm 8$	110	0.05	$\frac{3}{2}^+$	8.080
$2.407 \pm 8$	70	0.11	$\frac{3}{2}^-$	8.200
$2.604 \pm 4$	$9 \pm 3$	0.44	$\frac{1}{2}^+$	8.351
$2.680 \pm 3$	$4 \pm 3$	0.08	$\frac{5}{2}^+$	8.409
$2.763 \pm 3$	$7 \pm 3$	0.97	$\frac{7}{2}^+$	8.472
$2.808 \pm 3$	$5 \pm 3$	0.26	$\frac{5}{2}^-$	8.507
$3.059 \pm 5$	$50 \pm 3$	0.06	$\frac{3}{2}^-$	8.699
(3.1)	broad		$\frac{1}{2}^-$	(8.7)
$3.318 \pm 8$	$101 \pm 3$	0.50	$\frac{3}{2}^+$	8.897
$3.415 \pm 4$	$21 \pm 3$	0.04	$\frac{7}{2}^-$	8.971
$3.645 \pm 4$	$4 \pm 3$	0.45	$\frac{1}{2}^-$	9.147
(3.69)	3	1.00	$\frac{7}{2}^-$	(9.18)
$3.714 \pm 4$	$5.5 \pm 1$	0.20	$\frac{5}{2}^+$	9.200
$4.096 \pm 4$	$15 \pm 1$	0.85	$\frac{5}{2}^-$	9.492
(4.3)			$\frac{3}{2}^+$	(9.6)
$4.394 \pm 5$	$16 \pm 1$	0.70	$\frac{7}{2}^+$	9.719
$4.465 \pm 15$	$\approx 25$	0.90	$\frac{5}{2}^+$	9.774
$4.583 \pm 5$	14			9.864
$4.600 \pm 15$	$\approx 10$			9.877
$4.730 \pm 20$	$\approx 80$	0.78	$\frac{5}{2}^+$	9.976
$4.820 \pm 20$	$\approx 100$			10.045
(4.94)	138	0.85	$\frac{5}{2}^+$	(10.14)
$4.993 \pm 5$	45	0.15	$\frac{7}{2}^-$	10.177
(5.08)	122	0.60	$\frac{7}{2}^+$	(10.2)
$5.200 \pm 15$	150		$\frac{5}{2}^+, \frac{7}{2}^-$	10.336
$5.315 \pm 3^c)$	$14 \pm 3$		$\frac{5}{2}^+, \frac{7}{2}^-$	10.423
5.40	$75 \pm 30$		$\frac{5}{2}^+, \frac{7}{2}^-$	10.49

TABLE 17.8—continued

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\Gamma_a/\Gamma$	$J^\pi$	$E_x$ (MeV)
$5.492 \pm 3^{\circ}$ (5.68)	$51 \pm 2$ $\leq 25$	1.00	$\frac{7}{2}^-, \frac{9}{2}^+$ $(\frac{7}{2}^+)$	10.559 (10.70)
$5.778 \pm 3^{\circ}$	$74 \pm 3$		$\frac{1}{2}^+, \frac{7}{2}^-$	10.777
$5.945 \pm 3^{\circ}$	$46 \pm 2$		$\frac{5}{2}^-$	10.905
$6.117 \pm 3^{\circ}$	$31 \pm 3$			11.036
6.167	$5.0 \pm 1.1$		$\frac{1}{2}^-; T = \frac{3}{2}$	$11.075 \pm 0.005$
$6.380 \pm 3^{\circ}$	$80 \pm 3$			11.238
$6.883 \pm 3^{\circ}$	$65 \pm 2$			11.622
$7.051 \pm 10$	$40 \pm 25$			11.750
$7.136 \pm 15$	$12 \pm 3$			11.815
$7.384 \pm 15$				12.005
$7.52 \pm 20$	$150 \pm 50$			12.11
$7.736 \pm 15$	$100 \pm 30$			12.274
$7.88 \pm 20$				12.38
$7.927 \pm 15$				12.420
7.975	$8 \pm 2$		$\frac{3}{2}^-; T = \frac{3}{2}$	$12.457 \pm 0.005$
$8.156 \pm 15$	$75 \pm 30$			12.595
$8.253 \pm 15$	$\approx 5$			12.669
$8.44 \pm 25$				12.81
$8.59 \pm 20$	$\geq 150$			12.93
8.611	$6 \pm 2$		$\frac{1}{2}^+; T = \frac{1}{2}$	$12.943 \pm 0.006$
8.675	$\leq 3$		$\frac{5}{2}^-; T = \frac{3}{2}$	$12.992 \pm 0.006$
$8.72 \pm 20$				13.03
$8.785 \pm 15$	$16 \pm 4$			13.076
$9.319 \pm 15$	$\approx 120$			13.484
9.483 $\pm 15$	$250 \pm 100$			13.609

<sup>a</sup>) See references listed in tables 17.8 of (Aj 77) and 17.6 of (Aj 71). See also table 17.12 here.

<sup>b</sup>) See also (Ra 76g).

<sup>c</sup>) (Be 75bb).

### 13. $^{13}\text{C}(^7\text{Li}, t)^{17}\text{O}$

$$Q_m = 3.893$$

Angular distributions of triton groups to many states of  $^{17}\text{O}$  have been measured earlier at  $E(^7\text{Li}) = 17, 20.5$  and  $30.1$  MeV: see table 17.7 in (Aj 77). At  $E(^7\text{Li}) = 35.7$  MeV angular distributions are reported to  $^{17}\text{O}^*(3.06)$  and to  $^{17}\text{O}^*(13.58)$  which is preferentially populated (see discussion in reaction 12). Narrow states at  $E_x = 14.86, 18.17$  and  $19.24$  MeV are also strongly excited (Cl 78f). See also (Ma 78b).

### 14. $^{13}\text{C}(^{13}\text{C}, ^9\text{Be})^{17}\text{O}$

$$Q_m = -4.288$$

At  $E(^{13}\text{C}) = 105$  MeV states of  $^{17}\text{O}$  with  $E_x = 3.9, 5.2, 5.8 \pm 0.1, 7.2, 7.6, 8.4 \pm 0.06, 8.9, 9.8 \pm 0.07, 10.55 \pm 0.06, 12.1 \pm 0.06, 13.3, 14.6$  and  $18.9 \pm 0.14$  MeV are reported (Br 79a).

**<sup>17</sup>O**

88

F. AJZENBERG-SELOVE

15. (a)  $^{13}\text{C}(^{16}\text{O}, ^{12}\text{C})^{17}\text{O}$   $Q_m = -0.8021$   
 (b)  $^{13}\text{C}(^{18}\text{O}, ^{14}\text{C})^{17}\text{O}$   $Q_m = 0.132$

Angular distributions for reaction (a) have been studied at  $E(^{16}\text{O}) = 12$  to 25 MeV [ $^{17}\text{O}^*(0, 0.87)$ : see (Aj 77)] and at 13 and 14 MeV (Du 76b; see also for  $r_n$ ). For reaction (b) see (Ch 78m). See also (Go 80j, Pa 80; theor.).

16.  $^{14}\text{C}(^3\text{He}, \gamma)^{17}\text{O}$   $Q_m = 18.7611$

The capture cross sections at 90° for  $\gamma_0$  and for  $\gamma_1$  have been studied for  $E(^3\text{He}) = 3.2$  to 7.5 MeV and angular distributions of the  $\gamma$ -rays have been studied at the six observed resonances: see table 17.9 (Ch 76).

17. (a)  $^{14}\text{C}(^3\text{He}, n)^{16}\text{O}$   $Q_m = 14.6168$   $E_b = 18.7611$   
 (b)  $^{14}\text{C}(^3\text{He}, p)^{16}\text{N}$   $Q_m = 4.981$   
 (c)  $^{14}\text{C}(^3\text{He}, d)^{15}\text{N}$   $Q_m = 4.7139$   
 (d)  $^{14}\text{C}(^3\text{He}, ^3\text{He})^{14}\text{C}$   
 (e)  $^{14}\text{C}(^3\text{He}, \alpha)^{13}\text{C}$   $Q_m = 12.4013$

See (Aj 77) and table 17.9. See also  $^{16}\text{N}$ ,  $^{16}\text{O}$ , and  $^{15}\text{N}$ ,  $^{14}\text{C}$  and  $^{13}\text{C}$  in (Aj 81).

18.  $^{14}\text{C}(\alpha, n)^{17}\text{O}$   $Q_m = -1.8167$

The upper limits to the decays  $3.06 \rightarrow 0$  and  $3.84 \rightarrow 0.87$  are, respectively, 2 and 5%. A study of  $n-\gamma$  correlations leads to  $J'' = \frac{1}{2}^-$  and  $(\frac{5}{2}^-)$  for  $^{17}\text{O}^*(3.06, 3.84)$  (Al 64g).

TABLE 17.9  
 States of  $^{17}\text{O}$  from  $^{14}\text{C} + ^3\text{He}^*$ )

$E_{res}$ (MeV)	Resonant for	$E_{c.m.}$ (MeV)	$E_x$ (MeV)	$J''$
$3.6 \pm 0.1$	$\gamma_0, (\gamma_1), \alpha_0, \alpha_1$	0.75	21.7	$\frac{5}{2}^+$
$4.1 \pm 0.1$	$\gamma_0, n_0, n_{3+4}, \alpha_0, \alpha_1$	0.75	22.1	$\frac{5}{2}^-$
$4.6 \pm 0.2$	$\gamma_1$	$\approx 1$	22.5	$\frac{3}{2}(-)$
$5.1 \pm 0.1$	$\gamma_0, ^3\text{He}$	$\approx 0.4$	23.0	$\frac{5}{2}^+$
$5.7 \pm 0.1$	$\gamma_1$		23.5	$\frac{1}{2}^+$
$6.9 \pm 0.1$	$\gamma_1$		24.4	

\*) For references see table 17.9 in (Aj 77).

19.  $^{14}\text{C}(^{16}\text{O}, ^{13}\text{C})^{17}\text{O}$   $Q_m = -4.0322$

Angular distributions have been measured at  $E(^{16}\text{O}) = 20$ , 25 and 30 MeV ( $^{17}\text{O}^*(0, 0.87)$ ): see (Aj 77). See also (We 77k; theor.).

20. (a)  $^{14}\text{N}(t, \gamma)^{17}\text{O}$   $Q_m = 18.6233$   
 (b)  $^{14}\text{N}(t, p)^{16}\text{N}$   $Q_m = 4.843$   $E_b = 18.6233$

- (c)  $^{14}\text{N}(\text{t}, \text{d})^{15}\text{N}$   $Q_m = 4.5760$   
 (d)  $^{14}\text{N}(\text{t}, \text{t})^{14}\text{N}$   
 (e)  $^{14}\text{N}(\text{t}, \alpha)^{13}\text{C}$   $Q_m = 12.2634$

The excitation functions for  $\gamma_0$  and  $\gamma_1$  have been measured for  $E_t = 0.8$  to 3.3 MeV; broad resonances are observed at 2.2 and 2.8 MeV in the  $\gamma_0$  cross section, and at 2.4 and 2.8 MeV in the  $\gamma_1$  cross section. Both also exhibit a structure at 1.5 MeV. The data are consistent with states at  $E_x = 19.76 \pm 0.06$  [ $J = \frac{3}{2}$ ], 20.39  $\pm 0.05$  [ $\frac{5}{2}, \frac{7}{2}$ ], 20.58  $\pm 0.05$  [ $\frac{1}{2}$ ] and 21.05  $\pm 0.05$  MeV [ $\frac{3}{2}$ ] with  $\Gamma = 0.55 \pm 0.05$ , 0.66  $\pm 0.07$ , 0.57  $\pm 0.08$  and 0.47  $\pm 0.06$  MeV, and possibly with a state at  $\sim 19.3$  MeV.  $\Gamma_{\gamma_0} > 1.0$ , 4.3 and 5.8 eV for  $^{17}\text{O}^*(19.8, 20.4, 21.1)$  and  $\Gamma_{\gamma_1} > 2.3$ , 5.1 and 6.5 eV for  $^{17}\text{O}^*(19.8, 20.6, 21.1)$  (Li 80b). For reactions (b)  $\rightarrow$  (e) see (Aj 77). See also  $^{16}\text{N}$ , and  $^{15}\text{N}$ ,  $^{14}\text{N}$  and  $^{13}\text{C}$  in (Aj 81).

21. (a)  $^{14}\text{N}(\alpha, \text{p})^{17}\text{O}$   $Q_m = -1.1908$   
 (b)  $^{14}\text{N}(\alpha, \alpha\text{p})^{13}\text{C}$   $Q_m = -7.55063$

Angular distributions have been measured for  $^{17}\text{O}$  states with  $E_x < 7.6$  MeV in the range  $E\alpha = 8.1 \rightarrow 33.3$  MeV: see a listing of the references in (Aj 71). The sequential decay [reaction (b)] appears to take place via  $^{17}\text{O}$  states with  $8.46 \leq E_x \leq 13.57$  MeV. Those involved are believed to have  $J \geq \frac{5}{2}$ ,  $\Gamma_\alpha/\Gamma \geq 0.6$  (Ba 69n).

22.  $^{14}\text{N}(^{6}\text{Li}, ^3\text{He})^{17}\text{O}$   $Q_m = 2.829$

At  $E(^6\text{Li}) = 18$  MeV, the  $^3\text{He}$  groups in this reaction and the triton groups in the mirror reaction (see  $^{17}\text{F}$ , reaction 8) have been compared:  $^{17}\text{O}^*(3.84, 4.55, 5.22, 5.70 + 5.73)$  are strongly excited.  $^{17}\text{O}^*(0, 0.87, 3.06, 5.09, 5.38, 5.87 + 5.94, 6.86, 6.97)$  are also populated (Bi 73a). See also (Ma 77g, Ma 78b).

23.  $^{14}\text{N}(^{10}\text{B}, ^7\text{Be})^{17}\text{O}$   $Q_m = -0.045$

See (Mo 77y).

24.  $^{15}\text{N}(\text{d}, \gamma)^{17}\text{O}$   $Q_m = 14.0473$

See (Aj 77) [paper quoted there has not been published].

25.  $^{15}\text{N}(\text{d}, \text{n})^{16}\text{O}$   $Q_m = 9.9029$   $E_b = 14.0473$

Excitation functions have been measured for a number of neutron groups for  $E_d = 0.5$  to 5.9 MeV: a great deal of unresolved structure is apparent. Polarization measurements are reported for  $E_d = 1.6$  to 5.5 MeV and at 10.0 and 11.8 MeV. See (Aj 77) for a listing of the work. See also  $^{16}\text{O}$ .

26.  $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$   $Q_m = 0.267$   $E_b = 14.0473$

Excitation functions have been obtained for  $E_d = 0.3$  to 6.3 MeV: [see (Aj 77)] and at  $E_d = 1.4$  to 2.7 MeV (Ca 77b;  $p_0 \rightarrow p_3$ ). Structures are reported at  $E_d = 1.8$  [ $p_0, p_1, p_3$ ] and 2.4 MeV [ $p_2$ ] [ $^{17}\text{O}^*(15.6, 16.2)$ ] (Ca 77b).

**<sup>17</sup>O**

90

F. AJZENBERG-SELOVE



$E_b = 14.0473$


Excitation functions for  $d_0$  have been measured for  $E_d = 4.25$  to  $6.25$  MeV [see (Aj 77)] and  $1.4$  to  $2.7$  MeV (Ca 77b). The latter report structures at  $\sim 1.4$  and  $1.8$  MeV.



$Q_m = 7.6874$


$E_b = 14.0473$

Yield curves have been measured for  $E_d = 0.8$  to  $2.5$  MeV [see (Aj 77)] and  $1.2$  to  $2.7$  MeV (Ca 76r:  $\alpha_0 \rightarrow \alpha_3$ ): structures have been reported at  $E_d = 1.06$  and  $1.25$  MeV, and at  $\sim 1.8$  MeV. The latter has  $\Gamma \sim 300$  keV (Ca 76r).



$Q_m = 8.5537$

Observed proton groups are displayed in table 17.10. See also (Aj 77). For the parameters of the first  $T = \frac{3}{2}$  state see table 17.11.



$Q_m = -9.7995$

TABLE 17.10  
Levels of  $^{17}\text{O}$  from  $^{15}\text{N}(^3\text{He}, \text{p})^{17}\text{O}$ <sup>a)</sup>

$E_x$ <sup>b)</sup> (MeV)	$L$	$E_x$ <sup>b)</sup> (MeV)	$L$
0	(1+3)	8.192	0
0.874	1	8.322	
3.053	0	8.390	
3.845	2	8.492	(2)
4.549	0	8.682	
5.081	(1)	8.900	
5.215	(4)	8.955	
5.381	0	9.16	
5.698	2	9.495	
5.873	(1)	9.712	
5.938	0	9.856	
6.37		(10.24)	
6.861	(0)	10.33	
6.973	(1+3)	10.57	
7.162	2	10.782	
7.382	2	10.913	
7.561		$11.032 \pm 0.004$ <sup>c)</sup>	
7.687		$11.075 \pm 0.004$ <sup>d)</sup>	
7.761	4		
7.938			
8.054	(1)		

<sup>a)</sup> (Le 72c).  $E(^3\text{He}) = 18$  MeV.<sup>b)</sup>  $\pm 10$  keV, except where shown otherwise.<sup>c)</sup> See also (Mc 70b):  $T = \frac{1}{2}$ .<sup>d)</sup>  $J'' = \frac{1}{2}^-$ ;  $T = \frac{3}{2}$ : see table 17.11 (Le 72c, Ad 73b).

TABLE 17.11

Decay properties of the lowest  $T = \frac{3}{2}$  states in  $A = 17$ <sup>a)</sup>

	<sup>17</sup> O*(11.0787 ± 0.0008) <sup>b)</sup>	<sup>17</sup> F*(11.1928 ± 0.0021) <sup>c)</sup>
$J^\pi$ $\Gamma_{c.m.}$ (keV)	$\frac{1}{2}^-$ $2.4 \pm 0.3$ <sup>b)</sup>	$\frac{1}{2}^-$ $0.20 \pm 0.04$
Branching ratio (%) to <sup>16</sup> O*(MeV) $J^\pi$		
0      0 <sup>+</sup>	81 ± 6 <sup>c)</sup>	9.3 ± 1.3
6.05    0 <sup>+</sup>		<3
6.13    3 <sup>-</sup>	5 ± 2	22 ± 2
6.92    2 <sup>+</sup>		24 ± 6
7.12    1 <sup>-</sup>		44 ± 4
<sup>13</sup> C + $\alpha_0$ or <sup>13</sup> N + $\alpha_0$	6	<7
Partial widths [ $\Gamma_p$ or $\Gamma_n$ ] to		
<sup>16</sup> O(0)	$1.88 \pm 0.12$ keV	$19 \pm 3$ eV
<sup>16</sup> O*(6.05)		<8 eV
<sup>16</sup> O*(6.13)	$0.12 \pm 0.05$ keV	$45 \pm 14$ eV <sup>d)</sup>
<sup>16</sup> O*(6.92)		$49 \pm 19$ eV <sup>d)</sup>
<sup>16</sup> O*(7.12)		$90 \pm 27$ eV <sup>d)</sup>
$\Gamma_{\alpha_0}$	0.14 keV	<19 eV <sup>d)</sup>
$\Gamma_{\gamma^1}$		$6.0 \pm 2.5$ eV
$\theta^2(\text{g.s.})/\theta^2(6.13)$	$0.31 \pm 0.14$	$0.065 \pm 0.019$

<sup>a)</sup> See also table 2 in (Ad 73b) and reaction 63. See also (Mc 78d).<sup>b)</sup> (Hi 81a):  $\Gamma_{n_0} = 1.88 \pm 0.12$  keV. See also for IMME parameters for six  $T = \frac{3}{2}$  states.<sup>c)</sup> (Ha 71b, Ad 73b, Sk 74, Ha 75g, Hi 76d).<sup>d)</sup> Note that the total width is  $200 \pm 40$  eV.<sup>e)</sup> Weighted mean of  $91 \pm 15$  (Ad 73b) and  $79 \pm 7\%$  (Hi 81a, and F. Hintenberger, private communication).

At  $E_\alpha = 45.4$  MeV, the deuteron spectrum is dominated by the groups corresponding to states with  $E_x = 7.742 \pm 0.020$  and  $9.137 \pm 0.030$  MeV. These states are assigned  $J^\pi = (\frac{1}{2}^-)$  and  $(\frac{3}{2}^-)$  and arise from a dominant  $(d_{5/2})^2 p_{1/2}^{-1}$  configuration. Angular distributions were measured as well for the deuterons corresponding to <sup>17</sup>O<sub>g.s.</sub> and to states with  $E_x = 0.87 \pm 0.05$ , 5.22, 5.70, 7.38, 8.46, 8.89 and 9.81 [ $\pm 0.03$ ] MeV. In addition the population of <sup>17</sup>O\*(3.84, 4.55,  $8.15 \pm 0.03$ ) is also reported (Lu 69a).

31. <sup>15</sup>N(<sup>11</sup>B, <sup>9</sup>Be)<sup>17</sup>O

$$Q_m = -1.769$$

At  $E(^{11}\text{B}) = 115$  MeV angular distributions have been studied to <sup>17</sup>O\*(5.23, 7.75, 9.18) [ $J^\pi = \frac{9}{2}^-, \frac{11}{2}^-, \frac{9}{2}^-$ ] (Pr 80c, Ra 79c). See also (Aj 77).

32. <sup>16</sup>O(n,  $\gamma$ )<sup>17</sup>O

$$Q_m = 4.1443$$

$$\sigma_{\text{capt.}} = 202 \pm 28 \mu\text{b}$$
 (Mc 77e).

At thermal energies the branchings via <sup>17</sup>O\*(0.87, 3.05) are  $(18 \pm 3)$  and  $(82 \pm 3)\%$ ;  $E_\gamma = 870.89 \pm 0.22$  keV. The cross section for two-photon emission  $\sigma_{2\gamma} < 3 \pm 19$   $\mu\text{b}$  for  $1200 < E_\gamma < 2943$  keV. The two-photon branching ratio is  $(1.6 \pm 10) \times 10^{-2}$  (Mc 77e). See also (Le 76r; theor.).

33. <sup>16</sup>O(n, n)<sup>16</sup>O

$$E_b = 4.1443$$

The scattering amplitude (bound)  $a = 5.805 \pm 0.005$  fm,  $\sigma_{\text{free}} = 3.761 \pm 0.007$  b (Ko 79w). High resolution cross-section measurements and analyses of the elastic scattering and of the (n,  $\alpha$ ) and <sup>13</sup>C( $\alpha$ , n) data have led to a much better understanding of the <sup>17</sup>O structure below  $E_x = 18.2$  MeV: see table 17.12 (Fo 73g, Jo 73, Ci 80, Hi 81a). (Jo 73) has performed a multilevel two-channel  $R$ -matrix analysis. Five states contain nearly 100% of the  $1d_{3/2}$  strength and have their eigenenergy at  $E_x \approx 5.7$  MeV [the dominant state is <sup>17</sup>O\*(5.08)]. Spectroscopic factors are deduced for 26 states in <sup>17</sup>O for  $4.5 < E_x < 9.5$  MeV [see table 17.12 in (Aj 77)]: the sum of these factors is 1% for  $J^\pi = \frac{1}{2}^+$ , 5% for  $\frac{1}{2}^-$ , 12% for  $\frac{3}{2}^-$ , 99% for  $\frac{3}{2}^+$ , 0.1% for  $\frac{5}{2}^+$ , 1% for  $\frac{5}{2}^-$  and 14% for  $\frac{7}{2}^-$  (Jo 73).  $T = \frac{3}{2}$  resonances are discussed by (Hi 81a): see tables 17.11 and 17.12.

Cross-section measurements are listed in table 17.10 of (Aj 71) and in (Aj 77). At  $E_n = 23.5$  keV,  $\sigma_t = 3.736 \pm 0.007$  b (Bl 75i; elemental O).  $\sigma_t$  has also been

TABLE 17.12  
Resonances \*) in <sup>16</sup>O(n, n) and <sup>16</sup>O(n,  $\alpha$ )

$E_n$ (keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\Gamma_{\lambda n}$ (keV)	$\Gamma_{\lambda\alpha}$ (keV)	$J^\pi$	$E_x$ (keV)
$433 \pm 2$ b)	45	45 <sup>b)</sup>		$\frac{3}{2}^-$	4552
$1000 \pm 2$	96	96 <sup>b)</sup>		$\frac{3}{2}^+$	5085
$1140$ c)	<0.1				5218
$1312 \pm 2$	42	41.5 <sup>b)</sup>		$\frac{3}{2}^-$	5378
$1651 \pm 2$	$3.4 \pm 0.3$	3.4		$\frac{7}{2}^-$	5697
$1689 \pm 2$	<1			$\frac{5}{2}^-$	5733
$1833 \pm 2$	$6.6 \pm 0.7$	6.6		$\frac{3}{2}^+$	5868
$1908 \pm 4$	$32 \pm 3$	31.5 <sup>b)</sup>		$\frac{1}{2}^-$	5939
$2351 \pm 8$	$124 \pm 12$	124		$\frac{1}{2}^+$	6356
$2889 \pm 2$	<1			$\frac{5}{2}^-$	6862
$3006 \pm 2$	<1			$\frac{5}{2}^+$	6972
$3211.70 \pm 0.17$	$1.38 \pm 0.05$	$1.38 \pm 0.05$ *)	0.0033	$\frac{5}{2}^-$	7165.2
$3250 \pm 10$	$280 \pm 30$	280	0.07	$\frac{3}{2}^+$	7202
$3438.38 \pm 0.19$	$0.64 \pm 0.23$	$0.64 \pm 0.23$ *)	0.01	$\frac{5}{2}^+$	7378.4
$3441.73 \pm 0.14$	$0.96 \pm 0.20$	$0.96 \pm 0.20$ *)	0.003	$\frac{3}{2}^-$	7381.5
$3630 \pm 20$	$500 \pm 50$	500	0.08	$\frac{3}{2}^-$	7559
$3647$ c)	<0.1				7576
$3767.76 \pm 0.22$	$14.4 \pm 0.3$	$13.0 \pm 0.6$ *)	0.01	$\frac{7}{2}^-$	7688.2
$4053 \pm 8$	$90 \pm 9$	84	6.7	$\frac{1}{2}^+$	7958
$4090 \pm 50$	$270 \pm 30$	250	16	$\frac{1}{2}^-$	7990
$4162 \pm 8$	$85 \pm 9$	71	15	$\frac{3}{2}^+$	8059
$4290 \pm 20$	$69 \pm 7$	68	0.8	$\frac{1}{2}^-$	(8180)
$4310 \pm 10$	52	48	4.0	$(\frac{3}{2}^-)$	8199

TABLE 17.12—continued

$E_n$ (keV)	$\Gamma_{c.m.}$ (keV)	$\Gamma_{\alpha n}$ (keV)	$\Gamma_{\alpha \alpha}$ (keV)	$J^\pi$	$E_x$ (keV)
4463.41 ± 0.26	11.4 ± 0.5	8.1 ± 0.3	2.2	$\frac{1}{2}^+$	8342.4
4527.12 ± 0.07	6.17 ± 0.13	4.75 ± 0.11	0.54	$\frac{3}{2}^+$	8402.3
4594.83 ± 0.09	2.13 ± 0.11	1.18 ± 0.04	(7.6)	$\frac{7}{2}^+$	8466.0
4631.78 ± 0.12	6.89 ± 0.22	2.86 ± 0.08	1.9	$\frac{5}{2}^-$	8500.7
4829.9 ± 0.4	55.3 ± 0.6	48.9 ± 1.1	1.8	$\frac{3}{2}^+$	8687.0
5050	78	68	9.5	$\frac{1}{2}^-$	8895
5127.0 ± 1.6	26.3 ± 1.9	23.5 ± 1.9		$\frac{3}{2}^+$	8966.4
5368.90 ± 0.09	3.53 ± 0.13	2.37 ± 0.08		$\frac{3}{2}^-$	9193.9
5610	120	120		$\frac{5}{2}^+$	9420
5640	140			$\frac{7}{2}^+$	9450
5919.67 ± 0.14	23.1 ± 0.3	18.0 ± 0.6		$\frac{7}{2}^+$	9711.9
5995.68 ± 0.15	11.7 ± 0.3	10.3 ± 0.3		$\frac{5}{2}^+$	9783.3
6076.08 ± 0.15	4.01 ± 0.23	3.37 ± 0.23		$(\frac{5}{2}^-)$	9858.9
6094.8 ± 1.0	16.7 ± 1.7	10.9 ± 1.2		$(\frac{3}{2}^-)$	9876.5
6404.6 ± 0.5	49.1 ± 0.8	22.3 ± 0.6		$(\frac{5}{2}^-)$	10 167.8
6820.7 ± 0.6	42.5 ± 1.1	17.2 ± 0.7 <sup>a</sup> )		$(\frac{5}{2}^-)$	10 559.1
7199.3 ± 1.3	41.7 ± 1.4	26.4 ± 0.9 <sup>a</sup> )		$(\frac{3}{2}^+)$	10 915.1
7373.31 ± 0.18	2.4 ± 0.3	1.88 ± 0.12 <sup>a</sup> )		$(\frac{1}{2}^-)$	11 078.7
7830	190			$\frac{5}{2}^3$	11 509
8320	270			$\frac{7}{2}^3$	11 970
8740	130				12 365
8848.8 ± 0.6	6.9 ± 1.1	1.27 ± 0.14 <sup>a</sup> )		$\frac{3}{2}^-$	12 466.0
9050	95				12 656
9353 ± 6	6 ± 2 <sup>b</sup> )	0.21 ± 0.14 <sup>a</sup> )		$\frac{1}{2}^5$	12 940 ± 6
9414.9 ± 0.6	2.5 ± 1.0	0.40 ± 0.06 <sup>a</sup> )		$\frac{3}{2}^-$	12 998.2
10 092.5 ± 2.4	9 ± 5	0.24 ± 0.09 <sup>a</sup> )		$(\frac{5}{2}^+)$	13 635.3
10 130	400				13 672
10 725.5 ± 1.5	20.5 ± 1.6	2.07 ± 0.16 <sup>a</sup> )		$(\frac{7}{2}^-)$	14 230.3
10 785 ± 3	7.5 ± 4	0.80 ± 0.16 <sup>b</sup> )		$\frac{5}{2}^5$	14 286
10 960 ± 3	40 ± 6	13 ± 6 <sup>b</sup> )			14 451
11 140	340			$(\geq \frac{3}{2})$	14 621
11 322 ± 3	36 ± 13	3.2 ± 1.0 <sup>b</sup> )		$(\frac{1}{2}^-)$	14 791
11 540	180				14 997
11 756 ± 3	52 ± 14	11 ± 3 <sup>b</sup> )		$(\frac{3}{2})$	15 199
11 936 ± 3	40 ± 6	7 ± 1 <sup>b</sup> )		$(\frac{5}{2}^+)$	15 368
12 867 ± 4	21 ± 10	2 ± 0.5 <sup>b</sup> )		$(\frac{3}{2}^+)$	16 243
14 136 ± 11	66 ± 20	8.0 ± 2.4 <sup>b</sup> )		$\frac{5}{2}^5$	17 436
14 853 ± 4	43 ± 12	1.0 ± 0.3 <sup>a</sup> )		$\frac{3}{2}^-$	18 110

<sup>a</sup>) (Fo 73g, Jo 73, Ci 80, Hi 81a). See also table 17.12 in (Aj 77).<sup>b</sup>)  $\Gamma_\gamma < 4.0$  eV,  $\Gamma_n = 60 \pm 15$  keV (Al 71c).<sup>c</sup>) Not observed in  $\sigma_i$ ; see (Fo 73g).<sup>d</sup>) Not  $\frac{1}{2}^+$  (Fo 73g).<sup>e</sup>)  $\Gamma_{\alpha n}$  (Ci 80, Hi 81a, and F. Hintenberger, private communication).<sup>f</sup>) (Mc 76i).<sup>g</sup>)  $T = \frac{3}{2}$ .<sup>h</sup>) ( $J \pm \frac{1}{2}$ )  $\Gamma_{\alpha n}$  (Hi 81a).<sup>i</sup>)  $J^\pi$  assignment by comparison with <sup>17</sup>N states presumed to be analogs; then  $T = \frac{3}{2}$  (Hi 81a).<sup>j</sup>) See also table 17.2.

measured with high resolution for  $E_n = 4.5$  to 8 MeV (Ci 80). At  $E_n = 40.3$  and 50.4 MeV,  $\sigma_{\text{non}}$  is reported for elemental O (Za 81). See also  $^{16}\text{O}$ , (Fi 75e, Su 78e, La 79o, Fr 80k, La 80b, Pa 80e), (Ci 76, Ki 76k), (Li 77i, Jo 79d; applications) and (Le 76p, Ba 77oo, Ho 77c, Th 77d, Fe 78c, Le 78f, Ba 79dd; theor.).

$$34. \ ^{16}\text{O}(\text{n}, \text{n}')^{16}\text{O}^* \quad E_b = 4.1443$$

A number of resonances have been observed in the cross sections for production of 6.13 and  $(6.92 + 7.12)$   $\gamma$ -rays: see table 17.13 in (Aj 77) and (No 78g). For cross-section measurements see table 17.10 in (Aj 71), (Aj 77) and (No 78g;  $E_n = 6.95$  to 10.25 MeV;  $\gamma_{6.13}$ ,  $\gamma_{6.92+7.12}$ ). See also (Ya 78b).

$$35. \ ^{16}\text{O}(\text{n}, 2\text{n})^{15}\text{O} \quad Q_m = -15.6639 \quad E_b = 4.1443$$

See (Aj 77).

$$\begin{array}{lll} 36. \ (a) \ ^{16}\text{O}(\text{n}, \text{p})^{16}\text{N} & Q_m = -9.636 & E_b = 4.1443 \\ (b) \ ^{16}\text{O}(\text{n}, \text{d})^{15}\text{N} & Q_m = -9.9029 & \\ (c) \ ^{16}\text{O}(\text{n}, \text{t})^{14}\text{N} & Q_m = -14.4790 & \end{array}$$

For cross sections of reaction (a) see (Aj 77). See also (Sl 76a). For reaction (b) see  $^{15}\text{N}$  in (Aj 81) and (Sl 76a). For triton emission see (Qa 78).

$$37. \ ^{16}\text{O}(\text{n}, \alpha)^{13}\text{C} \quad Q_m = -2.2156 \quad E_b = 4.1443$$

Table 17.12 displays the results from a multilevel two-channel  $R$ -matrix analysis of the data from this reaction and from the elastic scattering of neutrons (Jo 73). For cross-section measurements see table 17.10 in (Aj 71), (Aj 77) and (No 78g;  $E_n = 8.06$  to 10.25 MeV;  $\gamma_{3.09}$ ,  $\gamma_{3.68+3.85}$ ). See also (Su 78, Ya 78b, Ro 79b) and (Sl 76a).

$$38. \ ^{16}\text{O}(\text{p}, \pi^+)^{17}\text{O} \quad Q_m = -136.205$$

At  $E_p = 185$  MeV (Da 74i) and 800 MeV (Ho 79i) angular distributions of pions to  $^{17}\text{O}^*(0, 0.87, 3.05)$  are reported. The cross section for  $\pi^+$  production near threshold has been studied by (Ma 79gg, Ma 79hh). For polarization measurements see (Sj 81: to  $^{17}\text{O}^*(0, 0.87)$ ). See also (So 79a, Sj 80a), (Jo 79a, Au 81a) and (Br 77a; theor.).

$$39. \ ^{16}\text{O}(\text{d}, \text{p})^{17}\text{O} \quad Q_m = 1.9197$$

Observed proton groups are displayed in table 17.14 of (Aj 77). Angular distributions have been measured at many energies in the range of  $E_d = 0.3$  to 63.2 MeV: see table 17.13 in (Aj 71) and (Aj 77), and at  $E_d = 698$  MeV (Bo 81a:  $p_0$ ,  $p_1$ ). At  $E_d = 12$  MeV angular distributions are reported for the proton groups to  $^{17}\text{O}^*(0, 0.87, 5.09)$ :  $\Gamma_n$  for  $^{17}\text{O}^*(5.09) = 97 \pm 5$  keV (An 79o). Reported parameters for  $^{17}\text{O}^*(0.87)$  are  $\tau_m = 258.6 \pm 2.6$  psec [see table 17.7 in (Aj 71)] and

$E_\gamma = 870.81 \pm 0.22$  keV [see (Aj 77)],  $E_\gamma = 870.725 \pm 0.020$  keV (Wa 80k) [ $E_x = 870.749 \pm 0.020$  keV (E. K. Warburton, private comm.)].

See also  $^{18}\text{F}$  in (Aj 83), (Ro 78e, So 79c, Fl 80b), (Am 78c, Bo 78l, Hi 78c, Ni 78; applications), (Te 77b) and (Co 76u, Gr 77g, Ka 77k, Ko 77mm, Mu 77, Ka 78u, Gr 79m, Am 80a, Ay 80, Kr 80i, Sh 81e; theor.).



The angular distribution of the  $d_0$  group has been studied at  $E_t = 5.5$  MeV (Ba 61c).



See (Aj 77) and (Gr 79m; theor.).



The angular distribution involving  $^{17}\text{O}_{\text{g.s.}}$  has been studied at  $E(^7\text{Li}) = 36$  MeV. The population of  $^{17}\text{O}^*(0.87, 3.06, 3.84, 4.55, 5.38)$  is also reported (Sc 73m).



Angular distributions are reported at  $E(^{11}\text{B}) = 115$  MeV to  $^{17}\text{O}_{\text{g.s.}}$  (Ra 79c). See also (Aj 77).




Angular distributions have been measured at  $E(^{13}\text{C}) = 36$  MeV [(We 76j); to  $^{17}\text{O}^*(0.87)$ ] and 105 MeV [(Ra 79c); to  $^{17}\text{O}^*(0, 0.87)$ ]. See also (Aj 77) and (Du 77b, Si 80g; theor.). For reaction (b) see (Ze 77c; theor.).



See (Ko 81a).



At  $E(^{14}\text{N}) = 79$  MeV angular distributions involving  $^{17}\text{O}^*(0, 0.87)$  have been studied: an anomaly is observed in the phase behavior of the distribution to the excited state. From this and other studies it is concluded that a multistep process via inelastic scattering is unlikely to occur in the excitation of  $2s_{1/2}$  states (Mo 76). An angular distribution involving  $^{16}\text{O}^*(0 + 0.87)$  has also been reported at  $E(^{14}\text{N}) = 155$  MeV (Na 76h). See also (Iz 80a; theor.).



At  $E(^{18}\text{O}) = 42$  and 52 MeV, the angular distributions involving  $^{17}\text{O}^*(0, 0.87)$  have been studied (Re 75d).

48.  $^{17}\text{N}(\beta^-)^{17}\text{O}$   $Q_m = 8.680$

The decay is principally to  $^{17}\text{O}^*(4.55, 5.38, 5.94)$ : see table 17.2.

49. (a) $^{17}\text{O}(\gamma, n)^{16}\text{O}$	$Q_m = -4.1443$
(b) $^{17}\text{O}(\gamma, 2n)^{15}\text{O}$	$Q_m = -19.808$
(c) $^{17}\text{O}(\gamma, \alpha)^{13}\text{C}$	$Q_m = -6.3599$

Forty-five resonances are reported for  $E_x$  up to 26.9 MeV in the differential cross section for  $(\gamma, n_0)$ .  $T = \frac{1}{2}$  components of the giant resonance are observed at  $E_x = 19.8, 21.5$  and  $\sim 26$  MeV (Jo 79; brems.). Monoenergetic photons with  $E_\gamma = 8.5$  to 39.7 MeV have been used to measure the  $(\gamma, n)$  and the  $(\gamma, 2n)$  [above 10 MeV] cross sections. The giant dipole resonance, 6 MeV broad, is centered at 23 MeV; a pigny resonance is also observed at 13 MeV. Most of the GDR strength decays to  $T = 1$  states in  $^{16}\text{O}$ : this implies at  $T = \frac{3}{2}$  assignment for the main part of the GDR. A broad structure, of  $T = \frac{1}{2}$  nature, with  $28 < E_x < 36$  MeV is also reported (Ju 80). For radiative widths see table 17.13. See also (Aj 77).

TABLE 17.13  
Radiative widths from resonances in  $^{17}\text{O}(\gamma, n)$

$E_x$ <sup>a)</sup> (MeV)	$J^\pi$ <sup>a)</sup>	$\Gamma_{\gamma 0}$ <sup>b)</sup> (eV)	$\Gamma_{\gamma 0}$ <sup>c)</sup> (eV)
4.55	$\frac{3}{2}^-$	0.42	
5.09	$\frac{3}{2}^+$	1.0	
5.38	$\frac{3}{2}^-$	0.06	$0.7 \pm 0.4$
5.70	$\frac{7}{2}^-$	0.4	$1.1 \pm 0.4$
6.36	$\frac{1}{2}^+$	<0.07	
7.38	$\frac{5}{2}^-$		$0.8 \pm 0.4$
7.69	$\frac{7}{2}^-$		$1.5 \pm 0.5$
8.20	$\frac{9}{2}^-$		$1.4 \pm 0.5$
8.50	$\frac{9}{2}^-$		$6.6 \pm 1.8$
8.69	$\frac{9}{2}^-$		$1.2 \pm 0.6$
8.97	$\frac{7}{2}^-$		$4.1 \pm 0.8$

<sup>a)</sup> Values from table 17.7.

<sup>b)</sup> (Ho 78i).

<sup>c)</sup> (Jo 79; bremsstrahlung radiation): thirty-seven additional resonances are reported with  $E_x$  to 26.9 MeV.

50.  $^{17}\text{O}(e, e)^{17}\text{O}$

The  $^{17}\text{O}$  charge radius,  $\langle r^2 \rangle^{1/2} = 2.710 \pm 0.015$  fm, based on  $\langle r_{17}^2 \rangle^{1/2} / \langle r_{16}^2 \rangle^{1/2} = 1.0015 \pm 0.0025$  (Ki 78a). (Mi 79d) report  $\langle r_{17}^2 \rangle^{1/2} - \langle r_{16}^2 \rangle^{1/2} = -0.008 \pm 0.007$  fm. Considerable deviation from single-particle predictions is observed in the elastic scattering for an effective momentum transfer range of 0.55 to  $2.8 \text{ fm}^{-1}$  (Hy 79). Inelastic scattering is reported to a number of  $^{17}\text{O}$  states: see tables 17.14 and

TABLE 17.14

Transition probabilities and ground state radiative widths from  $^{17}\text{O}(\text{e}, \text{e})^*$ 

$E_x$ (MeV)	$J^\pi$ <sup>a)</sup>	Mtp	$\Gamma_{\gamma 0}(\text{C}\lambda)$ (eV)	$B(\text{C}\lambda \uparrow)$ ( $e^2 \cdot \text{fm}^{2\lambda}$ )	Mtp	$\Gamma_{\gamma 0}(\text{M}\lambda)$ (eV)	$B(\text{M}\lambda \uparrow)$ ( $e^2 \cdot \text{fm}^{2\lambda}$ )
0.87	$\frac{1}{2}^+$	C2	$(8.7 \pm 1.7) \times 10^{-8}$	$31 \pm 6$			
3.06	$\frac{1}{2}^-$	C3	$(7.1 \pm 0.3) \times 10^{-7}$	$153 \pm 6$	M2	$(4.6 \pm 1.8) \times 10^{-3}$	$(5 \pm 2) \times 10^{-2}$
3.84	$\frac{1}{2}^-$	C3	$(2.2 \pm 0.2) \times 10^{-6}$	$98 \pm 8$	M2	$(1.8 \pm 0.7) \times 10^{-2}$	$(5.4 \pm 2.1) \times 10^{-2}$
4.55	$\frac{1}{2}^+$	C3	$(1.0 \pm 0.3) \times 10^{-2}$	$2.5 \pm 0.7$			
5.09	$\frac{1}{2}^+$	C2	$(8.5 \pm 0.3) \times 10^{-6}$	$360 \pm 11$	M2	$< 1 \times 10^{-2}$	$< 4 \times 10^{-2}$
5.22	$(\frac{5}{2}^-)$	C3	$(3.3 \pm 0.9) \times 10^{-6}$	$45 \pm 12$	M2	$(4.5 \pm 2.2) \times 10^{-2}$	$(6 \pm 3) \times 10^{-2}$
5.38	$\frac{3}{2}^-$	C3	$(1.5 \pm 0.2) \times 10^{-5}$	$270 \pm 32$	M2	$0.15 \pm 0.10$	$0.3 \pm 0.2$
5.70	$\frac{7}{2}^-$	C3	$(5.0 \pm 2.9) \times 10^{-6}$	$17 \pm 10$			
5.94	$\frac{1}{2}^-$	C3	$(5.3 \pm 3.3) \times 10^{-2}$	$2.1 \pm 1.3$			
6.36	$\frac{1}{2}^+$	C2	$(1.2 \pm 0.3) \times 10^{-4}$	$147 \pm 34$			
6.86 <sup>c)</sup>	$(\frac{1}{2}^-)$	C3	$(2.5 \pm 1.3) \times 10^{-2}$	$1.9 \pm 1.0$			
6.97 <sup>c)</sup>	$(\frac{3}{2}^+)$	C2		$5.5 \pm 1.0$			
7.38 <sup>c)</sup>	$\frac{1}{2}^+$	CO, or		$3.6 \pm 1.0$			
			$(6.3 \pm 1.8) \times 10^{-2}$	$47 \pm 38$			
	$\frac{5}{2}^-$	C3	$(2.1 \pm 1.7) \times 10^{-5}$				
7.58	$\frac{7}{2}^-$	C1	$(7.8 \pm 2.0) \times 10^{-2}$	$26 \pm 7$			
		C3	$(4.3 \pm 1.0) \times 10^{-5}$	$109 \pm 26$			
7.76	$(\frac{11}{2}^-)$	C3	$(1.16 \pm 0.05) \times 10^{-4}$	$369 \pm 15$			
8.35 <sup>c)</sup>	$\frac{1}{2}^+$	CO, or		$7.6 \pm 1.4$			
8.40 <sup>c)</sup>	$\frac{5}{2}^+$	C2		$8.3 \pm 2.6$			
8.47 <sup>c)</sup>	$\frac{7}{2}^+$						
8.50 <sup>c)</sup>	$\frac{9}{2}^-$						

<sup>a)</sup> (Ki 78a). See also table 17.15 in (Aj 77).<sup>b)</sup> Used to evaluate the widths.<sup>c)</sup> These levels were unresolved and were analyzed as a group.

17.15. The data of (No 77b) show a broad dipole resonance centered at 22–23 MeV with strength extending down to 10–12 MeV. A smaller resonance, with a form factor consistent with C2, is found between  $E_x = 17.5$  and 19.6 MeV (No 77b). The transitions to  $^{17}\text{O}^*(0.87, 5.08)$  can be described as single-particle transitions (Ki 78a). See also (Hy 80) and (Ar 78b, Ar 79n, Za 79, Bo 80b, Li 81j; theor.).

### 51. $^{17}\text{O}(\text{n}, \text{n})^{17}\text{O}$

See (No 77d; theor.).

### 52. $^{17}\text{O}(\text{p}, \text{p})^{17}\text{O}$

Angular distributions of elastically scattered protons have been measured at a number of energies in the range  $E_p = 8.6$  to 65.8 MeV [see (Aj 77)] and at  $E_p = 35.2$  MeV (Fa 80a). See also  $^{18}\text{F}$  in (Aj 83), (No 78e, Se 78f, Ke 80c, Pu 81) and (Co 76u, To 77i, Am 78b, Ka 78u, Gr 79m, Am 80a; theor.).

TABLE 17.15  
Inelastic groups observed in  $^{17}\text{O}(\text{e}, \text{e}')^a)$

$E_x$ (MeV)	$\Gamma$ (keV)
$11.71 \pm 0.05$	narrow
$11.95 \pm 0.05$	$\sim 250$
$12.66 \pm 0.05$	$\sim 90$
$12.96 \pm 0.05$	$\sim 200$
$13.56 \pm 0.05$	$\sim 150$
$14.14 \pm 0.10$	$\sim 100$
$14.76 \pm 0.10$	$> 300$
$15.24 \pm 0.10$	$\sim 200$
$16.52 \pm 0.05$	$\sim 300$
$17.09 \pm 0.05$	narrow
17.5–19.6 <sup>b)</sup>	
20.5	
22.0 <sup>c)</sup>	
23.0 <sup>c)</sup>	

<sup>a)</sup> (No 77b). Other inelastic groups are displayed in table 17.14.

<sup>b)</sup> C2.

<sup>c)</sup> C1.

### 53. $^{17}\text{O}(\text{d}, \text{d})^{17}\text{O}$

The angular distribution of elastically scattered deuterons has been studied at  $E_{\text{d}} = 18$  MeV (Li 76).

### 54. (a) $^{17}\text{O}(^3\text{He}, ^3\text{He})^{17}\text{O}$ (b) $^{17}\text{O}(\alpha, \alpha)^{17}\text{O}$

Elastic angular distributions have been measured at  $E(^3\text{He}) = 11.0$  and 17.3 MeV [see (Aj 77)] and at  $E(^3\text{He}) = 33.3$  MeV (Le 81f; also polarization; to both  $^{17}\text{O}^*(0, 0.87)$ ). In a study of the ( $^3\text{He}, \text{p}$ ) reaction, (Fo 78o) find 3–4% core excitation [3p–2h configuration] in the ground state of  $^{17}\text{O}$ . For reaction (b) see (Kn 79i, Kn 77c). See also (Ma 77uu).

### 55. (a) $^{17}\text{O}(^9\text{Be}, ^9\text{Be})^{17}\text{O}$ (b) $^{17}\text{O}(^{10}\text{B}, ^{10}\text{B})^{17}\text{O}$

For reaction (a) see (Pa 79; theor.). For fusion cross sections in reaction (b) see (Wi 80e, Ch 81c).

### 56. (a) $^{17}\text{O}(^{12}\text{C}, ^{12}\text{C})^{17}\text{O}$ (b) $^{17}\text{O}(^{13}\text{C}, ^{13}\text{C})^{17}\text{O}$

Elastic angular distributions are reported at  $E(^{17}\text{O}) = 30.5$  and 33.8 MeV for reaction (a) and 29.8 and 32.3 for reaction (b) [also  $^{17}\text{O}^*(0.87)$ ] (Ch 78f, Ch 77m).

For fusion cross section measurements see (He 78j; reaction a) and (Wi 80e; reactions a and b). See also (Pa 78, Va 78g, Pa 79; theor.).

57. (a)  $^{17}\text{O}(^{16}\text{O}, ^{16}\text{O})^{17}\text{O}$   
 (b)  $^{17}\text{O}(^{18}\text{O}, ^{18}\text{O})^{17}\text{O}$

Angular distributions involving  $^{17}\text{O}^*(0, 0.87)$  in reaction (a) have been studied at  $E(^{16}\text{O}) = 22$  to 32 MeV and  $E(^{17}\text{O}) = 25.7$  to 32.0 MeV: see (Aj 77). The elastic scattering angular distribution in reaction (b) has been reported at  $E(^{17}\text{O}) = 36$  MeV (Ka 77bb). For fusion cross sections see (Wi 80e; reaction a). See also (Fi 74f, Ho 78n) and (Im 77b, Cr 79b, Le 79a, Sa 79l; theor.).

58. (a)  $^{17}\text{O}(^{24}\text{Mg}, ^{24}\text{Mg})^{17}\text{O}$   
 (b)  $^{17}\text{O}(^{27}\text{Al}, ^{27}\text{Al})^{17}\text{O}$   
 (c)  $^{17}\text{O}(^{40}\text{Ca}, ^{40}\text{Ca})^{17}\text{O}$

For reaction (a) see (Pa 80g; theor.). For reaction (b) see (Ei 77) for fusion cross sections. See also (Sa 80m) and (Va 78g; theor.). For reaction (c) see (La 80f; elastic  $\sigma(\theta)$ ;  $E(^{17}\text{O}) = 61.1$  MeV).

59.  $^{17}\text{F}(\beta^+)^{17}\text{O}$   $Q_m = 2.762$

See  $^{17}\text{F}$ .

60.  $^{18}\text{O}(\gamma, n)^{17}\text{O}$   $Q_m = -8.0446$

See (Ba 76ii) and  $^{18}\text{O}$  in (Aj 78, Aj 83).

61.  $^{18}\text{O}(p, d)^{17}\text{O}$   $Q_m = -5.820$

Angular distributions have been measured at a number of energies for  $E_p = 17.6$  to 43.6 MeV [see (Aj 77)], at  $E_p = 24.4$  MeV (Pi 73d:  $d_0 \rightarrow d_7$ ) and at  $E_p = 51.9$  MeV (Oh 77:  $d_0 \rightarrow d_2$ ). See also  $^{19}\text{F}$  in (Aj 83).

62.  $^{18}\text{O}(d, t)^{17}\text{O}$   $Q_m = -1.787$

Angular distributions of tritons have been measured at  $E_d = 17$  MeV [see (Aj 77)] and at  $E_d = 17$  and 52 MeV: see table 17.16. Comparisons of the analog reactions ( $d, t$ ) and ( $d, ^3\text{He}$ ) have been made at  $E_d = 52$  MeV (Ma 77k): see also reaction 9 in  $^{17}\text{N}$ .

63.  $^{18}\text{O}(^3\text{He}, \alpha)^{17}\text{O}$   $Q_m = 12.533$

The  $T = \frac{3}{2}$  states reported by (De 69c) are displayed in table 17.17 [the isospin identification is based on the enhanced excitation and the narrow widths of these states]. The branching ratios for the various decays of  $^{17}\text{O}^*(11.08)$  [the lowest  $T = \frac{3}{2}$  state in  $^{17}\text{O}$ ] and for the analog state in  $^{17}\text{F}$  are displayed in table 17.11: the decay width of the  $^{17}\text{O}$  state is approximately 200 times greater than that of the  $^{17}\text{F}$  state (Ad 73b).

TABLE 17.16  
States of <sup>17</sup>O from <sup>18</sup>O(d, t) <sup>a)</sup>

$E_x$ <sup>b)</sup> (MeV)	$J^\pi; T$ <sup>b)</sup>	$l$	$C^2 S$
0	$\frac{5}{2}^+; \frac{1}{2}$	2	1.53 <sup>d)</sup>
0.87	$\frac{1}{2}^+; \frac{1}{2}$	0	0.21 <sup>d)</sup>
3.06	$\frac{1}{2}^-; \frac{1}{2}$	1	1.08
3.84	$\frac{5}{2}^-; \frac{1}{2}$	>2	
4.55	$\frac{3}{2}^+; \frac{1}{2}$	1	0.12
5.09	$\frac{3}{2}^+; \frac{1}{2}$	2	0.10
5.38	$\frac{3}{2}^+; \frac{1}{2}$	1	0.53
5.70	$\frac{7}{2}^+; \frac{1}{2}$		
5.94	$\frac{1}{2}^-; \frac{1}{2}$	1	0.06
6.86		≠1	
7.38 <sup>c)</sup>	$\frac{5}{2}^+ + \frac{5}{2}^-$	≠2	
8.20	$\frac{5}{2}^+; \frac{1}{2}$	1	0.15
8.47	$\frac{7}{2}^+; \frac{1}{2}$		
8.69	$\frac{3}{2}^-; \frac{1}{2}$	1	0.10
9.15	$\frac{1}{2}^-; \frac{1}{2}$	1	0.10
9.49	$\frac{5}{2}^-; \frac{1}{2}$		
11.08	$\frac{1}{2}^-; \frac{3}{2}$	1	0.96
11.41 ± 0.01 <sup>a)</sup>	$T = \frac{1}{2}^+$	(1)	0.04
12.12 ± 0.01 <sup>a)</sup>	$T = \frac{1}{2}^+$	(1)	0.24
12.47	$\frac{3}{2}^+; \frac{3}{2}$	1	0.24
12.76 ± 0.01 <sup>a)</sup>	$T = \frac{1}{2}^+$	(1)	0.17
12.94	$\frac{1}{2}^+; T = \frac{3}{2}$	0	0.19 ± 0.05
13.64	$(\frac{5}{2}^+; \frac{3}{2})$	2	0.29 ± 0.12
16.58 ± 0.01 <sup>a)</sup>	$(\frac{1}{2}, \frac{3}{2})^-; \frac{3}{2}^+$	1	0.93
18.14 ± 0.01 <sup>a)</sup>	$(\frac{1}{2}, \frac{3}{2})^-; \frac{3}{2}^+$	1	0.17

<sup>a)</sup> (Ma 77k):  $E_d = 52$  MeV; DWBA analysis.<sup>b)</sup> From table 17.7, unless footnote is shown.<sup>c)</sup> Unresolved.<sup>d)</sup> (Fo 78d;  $E_d = 17$  MeV) report spectroscopic factors of  $1.48 \pm 0.27$  and  $0.29 \pm 0.05$  (DWBA), 1.30 and 0.31 (CCBA), respectively, for <sup>17</sup>O\*(0, 0.87).64. <sup>18</sup>O(<sup>9</sup>Be, <sup>10</sup>Be)<sup>17</sup>O

$$Q_m = -1.233$$

Angular distributions have been studied at  $E(^{18}\text{O}) = 16$  and 20 MeV: see (Aj 77).65. (a) <sup>18</sup>O(<sup>10</sup>B, <sup>11</sup>B)<sup>17</sup>O

$$Q_m = 3.411$$

(b) <sup>18</sup>O(<sup>11</sup>B, <sup>12</sup>B)<sup>17</sup>O

$$Q_m = -4.675$$

Angular distributions (reaction a) have been measured at  $E(^{18}\text{O}) = 20$  and 24 MeV: see (Aj 77). For S-factor measurements see (Sw 74a). Cross sections for reaction (b) are several orders of magnitude less than those for reaction (a) for  $E(^{18}\text{O})_{c.m.} = 3 \rightarrow 7.7$  MeV (Sw 74a).

TABLE 17.17  
 $T = \frac{3}{2}$  states of  $^{17}\text{O}$  from  $^{18}\text{O}(\text{He}^3, \alpha)^{17}\text{O}$ <sup>a,b</sup>

$E_n$ (MeV $\pm$ keV)	$l_n$	$J^\pi$	$C^2 S$ <sup>c</sup>
11.082 $\pm$ 6	1	( $\frac{1}{2}$ ) <sup>-</sup>	0.49
12.471 $\pm$ 5	1	( $\frac{3}{2}$ ) <sup>-</sup>	0.27
12.950 $\pm$ 8	0	$\frac{1}{2}^+$	0.096
12.994 $\pm$ 8			
13.640 $\pm$ 5	2	( $\frac{5}{2}$ ) <sup>+</sup>	0.39
14.219 $\pm$ 8			
14.282 $\pm$ 12			
15.101 $\pm$ 8			

<sup>a</sup>) See also table 17.11.<sup>b</sup>) (De 69c).<sup>c</sup>) Calculated assuming  $C^2 S = 4$  for  $^{15}\text{O}^*(6.18)$  in  $^{16}\text{O}(\text{He}^3, \alpha)^{15}\text{O}$ .

66. (a)  $^{18}\text{O}(\text{He}^3, \text{C}^{12})^{17}\text{O}$   $Q_m = -3.098$   
 (b)  $^{18}\text{O}(\text{He}^3, \text{C}^{13})^{17}\text{O}$   $Q_m = 0.132$   
 (c)  $^{18}\text{O}(\text{He}^3, \text{N}^{14})^{17}\text{O}$   $Q_m = 2.789$

For reactions (a) and (b) see (Aj 77). For reaction (c) see (Sw 74a).

67.  $^{18}\text{O}(\text{He}^3, \text{O}^{19})^{17}\text{O}$   $Q_m = -4.088$

See (Ka 77v). See also (Aj 77).

68.  $^{19}\text{F}(\pi^-, 2\text{n})^{17}\text{O}$   $Q_m = 122.747$

The 0.87 MeV  $\gamma$ -ray is observed (En 76).

69.  $^{19}\text{F}(\text{n}, \text{t})^{17}\text{O}$   $Q_m = -7.5560$

Angular distributions of the  $t_0$  and  $t_1$  groups are reported at  $E_n = 14.4$  MeV: see (Aj 77).


70.  $^{19}\text{F}(\text{p}, \text{He}^3)^{17}\text{O}$   $Q_m = -8.320$

Angular distributions have been reported at  $E_p = 30.5$  and  $42.4$  MeV: see (Aj 77). See also  $^{20}\text{Ne}$  in (Aj 78).

71.  $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$   $Q_m = 10.033$

Observed  $\alpha$ -groups are displayed in table 17.14 of (Aj 77). Angular distributions have been measured at many energies in the range  $E_d = 0.3$  to  $27.5$  MeV: see table 17.16 in (Aj 71) and (Aj 77). See also (An 79r).

72.  $^{19}\text{F}(\alpha, \text{Li}^6)^{17}\text{O}$   $Q_m = -12.340$

Fig. 8. Energy levels of  $^{17}\text{F}$ . For notation see fig. 2.

Angular distributions are reported at  $E_\alpha = 28$  and 42 MeV, involving  $^{17}\text{O}^*(0, 0.87)$ : see (Aj 77).



At  $E_n = 14.1$  MeV angular distributions of  $\alpha_0$  and  $\alpha_1$  are reported: see (Aj 77).



See (Si 80g; theor.).

### $^{17}\text{F}$

(Figs. 8 and 9)

#### GENERAL<sup>†</sup>

*Shell and cluster models*: (Du 77b, Ha 77w, Po 77c, Th 78e).

*Special states*: (He 77h, En 78c, Ch 80o, Ta 81i).

*Electromagnetic transitions*: (Mc 76m, Br 77, Ha 77w, Ho 77, Ch 80o).

*Complex reactions involving  $^{17}\text{F}$* : (Ar 77b).

*Astrophysical questions*: (Si 77d, Wo 78f).

*Reactions involving pions*: (Os 81a, Pu 81).

*Other topics*: (Sa 76i, Ba 77v, En 78c, Sh 78q, Sl 78, Be 79a, Ta 81i).

*Ground state of  $^{17}\text{F}$* : (Jo 76c, Mc 76m, Sa 76i, An 77b, Br 77, Du 77b, Ha 77w, Ko 77w, Po 77c, Sh 77i, Ar 78a, Sl 78, Ch 80o, Ta 81i).

$$\mu = 4.7223 \pm 0.0012 \text{ n.m. (Le 78n).}$$

$$Q = 0.10 \pm 0.02 \text{ b (Mi 74l).}$$



Two recent measurements of the half-life of  $^{17}\text{F}$  are  $64.31 \pm 0.09$  sec (Az 77) and  $64.80 \pm 0.12$  sec (Al 77g): the weighted mean of these is  $64.49 \pm 0.16$  sec;  $\log ft = 3.488 \pm 0.001$ . Earlier values are listed in (Aj 71, Aj 77). The upper limit for the  $\beta^+$  decay to  $^{17}\text{O}^*(0.87)$  is  $< 3.4 \times 10^{-4}$  per decay (Ga 69e) [ $\log ft > 8.6$ ]. See also (Ra 78f) and (Az 77a, Br 78d, Ba 79j, Os 79, To 79; theor.).



See (Aj 71).

<sup>†</sup> See also (Aj 77).




Fig. 9. Isobar diagram,  $A = 17$ . For notation see fig. 5.

**<sup>17</sup>F MASTER TABLE**

## ENERGY LEVELS OF LIGHT NUCLEI A = 16-17

105

 TABLE 17.18  
 Energy levels of <sup>17</sup>F

$E_x$ in <sup>17</sup> F (MeV $\pm$ keV)	$J^\pi; T$	$\tau$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0 $0.49533 \pm 0.10$	$\frac{5}{2}^+; \frac{1}{2}$ $\frac{1}{2}^+$	$\tau_{1/2} = 64.49 \pm 0.16$ sec $\tau_m = 412 \pm 9$ psec	$\beta^+$ $\gamma$	$1 \rightarrow 5, 7 \rightarrow 12, 18 \rightarrow 32$ 3, 5, 8, 10, 12, 18, 19, 22 $\rightarrow$ 28, 30, 32
$3.104 \pm 3$	$\frac{1}{2}^-$	$\Gamma = 19 \pm 1$	$\gamma, p$	5, 8, 10, 12, 13, 18, 19, 28, 30
$3.857 \pm 4$	$\frac{5}{2}^-$	$1.5 \pm 0.2$	$\gamma, p$	5, 8, 10, 12, 13, 18, 19, 30
$4.696 \pm 10$	$\frac{5}{2}^+$	225	$p$	8, 10, 13, 18, 28
$5.103 \pm 10$	$\frac{5}{2}^+$	1530	$p$	12
$5.22 \pm 10$	$(\frac{9}{2}^-)$			8, 10, 11
$5.521 \pm 10$	$\frac{3}{2}^-$	68	$p$	8, 10, 13, 28
$5.672 \pm 10$	$\frac{7}{2}^-$	40	$p$	8, 10, 13
$5.682 \pm 10$	$\frac{1}{2}^+$	$<0.6$	$p$	8, 10, 13
$5.817 \pm 10$	$\frac{3}{2}^+$	180	$p$	8, 13
$6.036 \pm 10$	$\frac{1}{2}^-$	30	$p$	8, 10, 13, 28
$6.556 \pm 10$	$\frac{1}{2}^+$	200	$p$	13
$6.7087 \pm 0.3$	$(\frac{5}{2}, \frac{3}{2})^+$	$<3$	$p$	8, 10, 13
$6.774 \pm 10$	$\frac{3}{2}^+$	4.5	$p$	13
$7.027 \pm 10$	$\frac{5}{2}^-$	3.8	$p$	10, 13
$7.356 \pm 10$	$\frac{3}{2}^+$	$10 \pm 2$	$p, \alpha$	10, 13, 17
$7.448 \pm 7$		$\leq 5$	$p$	13
$7.454 \pm 7$		$7 \pm 2$	$p, \alpha$	13, 17
$7.471 \pm 7$		$5 \pm 2$	$p$	13
$7.479 \pm 10$		795	$p$	13
$7.546 \pm 10$	$\frac{5}{2}^-$	30	$p$	13
$7.75 \pm 20$	$\frac{1}{2}^+$	$179 \pm 3$	$p, \alpha$	13, 17, 28
$7.95 \pm 15$		$10 \pm 3$	$p$	13
$8.01 \pm 20$		$50 \pm 20$	$p, \alpha$	13, 17
$8.075 \pm 10$	$\frac{5}{2}^+$	$100 \pm 20$	$p, \alpha$	10, 13, 17, 28
8.2	$\frac{3}{2}^-$	$700 \pm 250$	$p, \alpha$	13, 17
$8.383 \pm 5$	$\frac{5}{2}^-$	$11 \pm 5$	$p, \alpha$	13, 17
$8.416 \pm 10$	$\frac{7}{2}^+$	$45 \pm 10$	$p, \alpha$	13, 17, 28
$8.75 \pm 30$	$\frac{5}{2}^+$	$170 \pm 30$	$p, \alpha$	13, 17
8.76	$\frac{5}{2}^+$	$90 \pm 20$	$p$	13
$8.98 \pm 20$	$\frac{7}{2}^-$	$165 \pm 30$	$p, \alpha$	13, 17
$9.17 \pm 60$	$\frac{5}{2}^-$	$140 \pm 30$	$p, \alpha$	13, 17
9.92	$\frac{9}{2}^+$	$90 \pm 30$	$p, \alpha$	13, 17
$10.04 \pm 20$	$\frac{7}{2}$	$280 \pm 100$	$p$	13
$10.22 \pm 20$		$250 \pm 80$	$\alpha$	17
$10.40 \pm 20$	$(\frac{5}{2})^+$	$160 \pm 40$	$p$	13
$10.499 \pm 15$	$\frac{7}{2}^-$	$165 \pm 25$	$p, \alpha$	13, 17
$10.79 \pm 20$		$120 \pm 40$	$p, (\alpha)$	13, 17
$10.91 \pm 100$	$\frac{1}{2}^-$	$560 \pm 100$	$p$	13
$10.95 \pm 20$		$190 \pm 50$	$p, (\alpha)$	13, 17
$11.1928 \pm 2.3$	$\frac{1}{2}; \frac{3}{2}$	$0.20 \pm 0.04$	$\gamma, p, \alpha$	10, 12, 13, 17, 28
$11.43 \pm 20$		$240 \pm 50$	$p, \alpha$	13, 17
$11.58 \pm 40$		$160 \pm 30$	$p$	13
$12.00 \pm 20$		$120 \pm 40$	$p, \alpha$	13, 17
$12.25 \pm 20$	$\frac{3}{2}^-$	$300 \pm 30$	$p$	13

TABLE 17.18—continued

$E_x$ in <sup>17</sup> F (MeV ± keV)	$J^\pi; T$	$\Gamma_{c.m.}$ (keV)	Decay	Reactions
12.355 ± 10	$\frac{1}{2}^-$	190 ± 20	p	13
~12.50	$\frac{7}{2}^-$	~600	p	13
12.5501 ± 0.9	$\frac{3}{2}^-, \frac{5}{2}^+$	2.83 ± 0.12	$\gamma, p, \alpha$	10, 12, 13, 17
13.061 ± 4	$\frac{5}{2}^-, \frac{7}{2}^-$	2 ± 1	$\gamma, p, \alpha$	10, 12, 13, 17
13.080 ± 4	$(\frac{1}{2}^+); \frac{3}{2}^-$	2 ± 1	p, $\alpha$	13, 17
13.13 ± 100	$\frac{5}{2}^-$	520 ± 50	p	13
13.781 ± 4	$\frac{5}{2}^+, \frac{3}{2}^-$	12 ± 5	$q, \alpha$	13, 17
14.00 ± 50	$\frac{7}{2}^-$	260 ± 30	p	13
14.176 ± 6	$\frac{3}{2}^-, \frac{5}{2}^+$	30 ± 5	$\gamma, p$	12, 13
14.3037 ± 3.1	$\frac{7}{2}^-, \frac{3}{2}^-$	19.3 ± 1.6	$\gamma, p, \alpha$	12, 13, 17
14.38 ± 50	$\frac{5}{2}^-$	610 ± 50	p	13
14.71 ± 100	$\frac{1}{2}^-$	470 ± 100	p	13
14.809 ± 20	$\frac{1}{2}^+$	190 ± 25	p	13
15.6		~550	p	13
17.1	$\frac{5}{2}^-$	1500	p	13
20.1 ± 200		1070 ± 160	$\gamma, {}^3\text{He}$	5
20.4 ± 100		700 ± 100	$\gamma, {}^3\text{He}$	5
20.9	$\frac{9}{2}^+$	600	p	13
21.3 ± 100		900 ± 100	$\gamma, {}^3\text{He}$	5
21.8	$(\frac{9}{2}^+)$	400	p	13
22.7	$\frac{7}{2}^+$	600	p	13
23.8	$\frac{7}{2}^+$	600	p	13
25.4		1500	p	13
27.2		1500	p	13
28.9	$\frac{5}{2}^{++}$	2000	p	13

TABLE 17.19  
States of <sup>17</sup>F from <sup>14</sup>N(<sup>3</sup>He,  $\gamma$ )<sup>a)</sup>

$E({}^3\text{He})$ (MeV)	$\Gamma_{c.m.}$ (MeV)	Res. in	$(2J+1)\Gamma_{{}^3\text{He}}\Gamma_\gamma$ (keV <sup>2</sup> )	$E_x$ (MeV)
(4.5) <sup>b)</sup>	1.0 ± 0.3	$\gamma_1, \alpha^c)$	32 ± 20	(19.5 ± 0.2)
5.2	1.07 ± 0.16	$\gamma_2$	63 ± 26	20.1 ± 0.2
(5.4) <sup>b)</sup>	1.2 ± 0.4	$\gamma_0$	46 ± 32	(20.2 ± 0.3)
5.6	0.7 ± 0.1	$\gamma_1$	33 ± 11	20.4 ± 0.1
6.6	0.9 ± 0.1	$\gamma_1$	74 ± 18	21.3 ± 0.1
(7.8) <sup>b)</sup>	2.0 ± 0.5	$\gamma_0$	260 ± 140	(22.2 ± 0.3)

<sup>a)</sup> (Wa 81). See also (Ch 79h, Ch 80h). See, however, (Mo 73h).<sup>b)</sup> Uncertain: J. Lowe, private communication.<sup>c)</sup> See reaction 6.

3.  $^{12}\text{C}({}^{14}\text{N}, {}^9\text{Be})^{17}\text{F}$   $Q_m = -10.4362$

Angular distributions are reported to  ${}^{17}\text{F}^*(0, 0.50)$  at  $E({}^{14}\text{N}) = 53$  MeV (Ze 76) and 78.8 MeV (Mo 77y; unresolved).

4.  $^{12}\text{C}({}^{20}\text{Ne}, {}^{15}\text{N})^{17}\text{F}$   $Q_m = -9.0962$

See (Or 79).

5.  ${}^{14}\text{N}({}^3\text{He}, \gamma)^{17}\text{F}$   $Q_m = 15.8431$

Excitation functions for  $\gamma_0$ ,  $\gamma_1$ ,  $\gamma_2$  and  $\gamma_3$  have been studied for  $E({}^3\text{He}) = 3$  to 19 MeV: observed resonances are displayed in table 17.19 (Wa 81). [The earlier work by (Mo 73h), who studied the  $\gamma_{0+1}$  yield at 90°, indicated several sharp resonances].

6. (a) ${}^{14}\text{N}({}^3\text{He}, \text{n})^{16}\text{F}$	$Q_m = -0.969$	$E_b = 15.8431$
(b) ${}^{14}\text{N}({}^3\text{He}, \text{p})^{16}\text{O}$	$Q_m = 15.2428$	
(c) ${}^{14}\text{N}({}^3\text{He}, \text{d})^{15}\text{O}$	$Q_m = 1.8035$	
(d) ${}^{14}\text{N}({}^3\text{He}, {}^3\text{He})^{14}\text{N}$		
(e) ${}^{14}\text{N}({}^3\text{He}, \alpha)^{13}\text{N}$	$Q_m = 10.024$	

Excitation functions for protons have been measured for  $E({}^3\text{He}) = 2.5$  to 18 MeV [see (Aj 77)]: some large structures are observed. The polarization of the  $p_0$  group has been studied at  $E({}^3\text{He}) = 9.8$  MeV. The elastic scattering of  ${}^3\text{He}$  (reaction d) and the yield of  $\alpha$ -particles (reaction e), studied in the ranges  $E({}^3\text{He}) = 4$  to 7 MeV and 2.5 to 8.5 MeV, respectively, show some evidence of structures (Kn 70a, Mo 73h): see the discussion in (Aj 77). See also  ${}^{16}\text{O}$ ,  ${}^{16}\text{F}$ , and  ${}^{13}\text{N}$ ,  ${}^{14}\text{N}$  and  ${}^{15}\text{O}$  in (Aj 81).

7.  ${}^{14}\text{N}(\alpha, \text{n})^{17}\text{F}$   $Q_m = -4.7347$

See  ${}^{18}\text{F}$  in (Aj 78).

8.  ${}^{14}\text{N}({}^6\text{Li}, \text{t})^{17}\text{F}$   $Q_m = 0.049$

At  $E({}^6\text{Li}) = 18$  MeV  ${}^{17}\text{F}^*(3.86, 5.22 \pm 0.01, 5.67 + 5.68)$  are strongly excited.  $J'' = \frac{9}{2}^-$  is suggested for  ${}^{17}\text{F}^*(5.22)$  (Bi 73a). See also (Ma 77g).

9.  ${}^{14}\text{N}({}^{10}\text{B}, {}^7\text{Li})^{17}\text{F}$   $Q_m = -1.945$

See (Aj 77).

10.  ${}^{15}\text{N}({}^3\text{He}, \text{n})^{17}\text{F}$   $Q_m = 5.0097$

Angular distributions have been reported to most of the states of  ${}^{17}\text{F}$  with  $E_x < 8.1$  MeV at  $E({}^3\text{He}) = 3.8$  and 4.8 MeV: see (Aj 77). Neutron groups have also been reported to  ${}^{17}\text{F}$  states at  $E_x = 11.195 \pm 0.007$ ,  $12.540 \pm 0.010$  and  $13.059$  MeV, with  $\Gamma < 20$ ,  $< 25$  and  $< 25$  keV, respectively. Angular distributions at  $E({}^3\text{He}) =$

10.36 and 11.88 MeV lead to  $J'' = \frac{1}{2}^-$  for  $^{17}\text{F}^*(11.20)$  [ $L = 0$ ],  $\frac{3}{2}^-$  or  $\frac{5}{2}^-$  for  $^{17}\text{F}^*(12.54)$  and  $(\frac{3}{2}^-, \frac{5}{2}^-)$  for  $^{17}\text{F}^*(13.06)$ . These three states are probably the first three  $T = \frac{3}{2}$  states in  $^{17}\text{F}$  (Ad 69). The branching ratios for transitions to  $^{16}\text{O}^*(0, 6.05, 6.13)$  for  $^{17}\text{F}^*(11.20)$  and for the analog  $T = \frac{3}{2}$  state in  $^{17}\text{O}$  are displayed in table 17.11: the ratios of the reduced widths are quite different in the two mirror nuclei (Mc 70b, Ad 73b).

$$11. \ ^{15}\text{N}(^{11}\text{B}, ^9\text{Li})^{17}\text{F} \quad Q_m = -18.138$$

At  $E(^{11}\text{B}) = 115$  MeV  $^{17}\text{F}^*(0, 5.22)$  are strongly populated (Ra 79c).

$$12. \ ^{16}\text{O}(\text{p}, \gamma)^{17}\text{F} \quad Q_m = 0.6004$$

$$Q_0 = 600.35 \pm 0.28 \text{ keV (Ro 76t)}$$

At low energies the direct capture to  $^{17}\text{F}^*(0, 0.50)$  is observed: see (Aj 71, Aj 77). Extrapolation of cross-section data leads to  $S(0) \approx 8 \text{ keV} \cdot \text{b}$ : see (Aj 77). In addition to two  $T = \frac{1}{2}$  resonances, five resonances corresponding to  $T = \frac{3}{2}$  states are observed in the  $\gamma_1$  and  $\gamma_0 + \gamma_1$  yields: see table 17.20 for the reported parameters. The lowest  $T = \frac{3}{2}$  states of even parity at  $E_x = 13.27$  and 14.02 MeV [ $J'' = (\frac{1}{2}^+)$  and  $\frac{5}{2}^+$ ] (see table 17.21) are not observed here:  $\Gamma_\gamma \leq 7$  and  $\leq 11.8 \text{ eV}$ , respectively (Ha 75g). The (E1) values for the  $T = \frac{3}{2}$  states are in good agreement with shell model 2p-1h calculations using realistic Kuo-Brown interaction matrix elements (Ha 75g).

TABLE 17.20  
Resonances in  $^{16}\text{O}(\text{p}, \gamma)^{17}\text{F}^*$

$E_p$ (MeV $\pm$ keV)	Resonant <sup>b</sup> ) in	$\Gamma_\gamma$ (eV)	$\Gamma$ (keV)	$E_x$ (MeV)	$J''; T$	Refs.
2.66	$\gamma_1$	$(12 \pm 2) \times 10^{-3}$	$< 1.5$	3.11	$\frac{1}{2}^-; \frac{1}{2}$	A, (Ro 73aa) <sup>a</sup> )
3.47	$\gamma_0$	$0.11 \pm 0.02$	$\leq 1.6$	3.86	$\frac{5}{2}^-; \frac{1}{2}$	(Se 63e)
$11.275 \pm 6$	$\gamma_1$	$6.0 \pm 2.5^\circ$	$\leq 1.6$	11.204	$\frac{1}{2}^-; \frac{3}{2}$	(Ha 75g)
$12.707 \pm 1$	$\gamma_0 + \gamma_1$	$11.3 \pm 3.4^\circ$	$1.8 \pm 0.5$	12.550	$\frac{3}{2}^-; \frac{3}{2}$	(Ha 75g, Kh 79b)
$13.255 \pm 6$	$\gamma_0 + \gamma_1$	$2.8 \pm 1.8^\circ$	$5.0 \pm 1.5$	13.065	$\frac{5}{2}^-; \frac{3}{2}$	(Ha 75g)
$14.435 \pm 10$	$\gamma_0$	$72 \pm 37^\circ$	$41 \pm 10$	14.174	$\frac{3}{2}^-; \frac{3}{2}$	(Ha 75g, Kh 79b)
$14.583 \pm 6^\circ$	$\gamma_0 + \gamma_1$	$13.4 \pm 7.0^\circ$	$28 \pm 5$	14.313	$\frac{7}{2}^-; \frac{3}{2}$	(Ha 75g)

<sup>a</sup> See (Aj 71).

<sup>b</sup>) See also table 17.21.

<sup>b</sup>)  $\gamma_0$  and  $\gamma_1$  correspond to transitions to  $^{17}\text{F}^*(0, 0.50)$ , respectively.

<sup>c</sup>) These  $\Gamma_\gamma$  are based on  $J''$  and  $\Gamma_{\text{p}}/\Gamma$  determinations by (Sk 74) and R. G. Van Bree (unpublished) [quoted by (Ha 75g)]. The B(E1) values for these four states are  $4.7 \pm 2.0$ ,  $5.4 \pm 1.6$ ,  $1.2 \pm 0.8$  and  $4.4 \pm 2.3 [ \times 10^{-3} ] e^2 \cdot \text{fm}^2$ .

<sup>d</sup>) See the text of reaction 12 for discussion of the observed pigny and giant resonances (Ha 75h).

<sup>e</sup>) See also table 17.18 in (Aj 77).

<sup>f</sup>)  $\Gamma_{\gamma_1}/\Gamma_{\gamma_0} \leq 0.14$ .  $(2J+1)\Gamma_{\text{p}}\Gamma_\gamma/\Gamma = 11.4 \pm 2.6 \text{ eV}$ .

<sup>g</sup>)  $C^2 S = 0.90$  and 1.00, respectively, for  $^{17}\text{F}^*(0, 0.50)$ .

TABLE 17.21  
Resonances in  $^{16}\text{O}(\text{p}, \text{p})^{16}\text{O}$  and  $^{16}\text{O}(\text{p}, \alpha)^{13}\text{N}^a$ )

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$\Gamma_{\text{p}_0}/\Gamma$	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$
2.663 $\pm$ 7	19 $\pm$ 1	$\text{p}_0$		3.105	$\frac{1}{2}^-$
3.47	1.53 $\pm$ 0.2	$\text{p}_0$		3.86	$\frac{5}{2}^-$
4.354 $\pm$ 10	225	$\text{p}_0$		4.696	$\frac{3}{2}^-$
4.787 $\pm$ 10	1530	$\text{p}_0$		5.103	$\frac{3}{2}^+$
5.231 $\pm$ 10	68	$\text{p}_0$		5.521	$\frac{3}{2}^-$
5.392 $\pm$ 10	40	$\text{p}_0$		5.672	$\frac{1}{2}^-$
5.402 $\pm$ 10	<0.6	$\text{p}_0$		5.682	$\frac{1}{2}^+$
5.546 $\pm$ 10	180	$\text{p}_0$		5.817	$\frac{3}{2}^+$
5.779 $\pm$ 10	30	$\text{p}_0$		6.036	$\frac{1}{2}^-$
6.332 $\pm$ 10	200	$\text{p}_0$		6.556	$\frac{1}{2}^+$
6.4944 $\pm$ 0.1 <sup>b)</sup>	<3	$\text{p}_0$		6.7087	(j)
6.564 $\pm$ 10	4.5	$\text{p}_0$		6.774	$\frac{3}{2}^+$
6.833 $\pm$ 10	3.8	$\text{p}_0, \gamma_{6.13}$		7.027	$\frac{5}{2}^-$
7.183 $\pm$ 10	10 $\pm$ 2	$\text{p}_0, \text{p}_2, \alpha_0$		7.356	$\frac{3}{2}^+$
7.280 $\pm$ 7	$\leq$ 5	$\text{p}_0$		7.448	
7.287 $\pm$ 7	7 $\pm$ 2	$\text{p}_0, \text{p}_1, \text{p}_2, \alpha$		7.454	
7.305 $\pm$ 7	5 $\pm$ 2	$\text{p}_0, \text{p}_2$		7.471	
7.313 $\pm$ 10	795	$\text{p}_0$		7.479	$\frac{3}{2}^+$
7.385 $\pm$ 10	30	$\text{p}_0, \text{p}_2,$ $\gamma_{6.13}$		7.546	$\frac{7}{2}^-$
7.60 $\pm$ 20	179 $\pm$ 3	$\text{p}_0, \text{p}_1,$ $\alpha_0$		7.75	$\frac{1}{2}^+$
7.81 $\pm$ 15	10 $\pm$ 3	$\text{p}_2$		7.95	
7.88 $\pm$ 20	50 $\pm$ 20	$\text{p}_0,$ $\gamma_{6.13},$ $\gamma_{6.92},$ $\alpha_0$		8.01	
7.94 $\pm$ 15	100 $\pm$ 20	$\text{p}_0, \text{p}_1,$ $\alpha_0$		8.07	$\frac{5}{2}^+$
8.1	700 $\pm$ 250	( $\text{p}_0),$ $\text{p}_1, \alpha_0$		8.2	$\frac{3}{2}^-$
8.275 $\pm$ 4	11 $\pm$ 5	$\text{p}_0 \rightarrow \text{p}_3, \alpha_0$		8.383	$\frac{5}{2}^-$
8.310 $\pm$ 10	45 $\pm$ 10	$\text{p}_0 \rightarrow \text{p}_3,$ $\gamma_{6.13},$ $\gamma_{6.92},$ $\alpha_0$		8.416	$\frac{7}{2}^+$
8.66 $\pm$ 30	170 $\pm$ 30	$\text{p}_2, \text{p}_3,$ $\text{p}_4, \alpha_0$		8.75	$\frac{5}{2}^+$
8.68	90 $\pm$ 20	$\text{p}_0$	0.2	8.76	$\frac{3}{2}^+$
8.91 <sup>c)</sup>	165 $\pm$ 30	$\text{p}_0 \rightarrow \text{p}_4,$ $\gamma_{6.13},$ $\gamma_{6.92},$ $\alpha_0$	0.34 $\pm$ 0.05	8.98 $\pm$ 0.02	$\frac{7}{2}^-$
9.11 <sup>c)</sup>	140 $\pm$ 30	$\text{p}_0 \rightarrow \text{p}_4,$ $\gamma_{6.13},$ $\gamma_{6.92},$ $\alpha_0$	0.55 $\pm$ 0.05	9.17 $\pm$ 0.06 <sup>d)</sup>	$\frac{3}{2}^-$

TABLE 17.21—continued

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$\Gamma_{\text{po}}/\Gamma$	<sup>17</sup> F* (MeV)	$J^\pi; T$
9.91 <sup>c</sup> )	90 $\pm$ 30	p <sub>0</sub> , p <sub>2</sub> , $\alpha_0$	0.095 $\pm$ 0.005	9.92	$\frac{9}{2}^+$
10.04 $\pm$ 20	280 $\pm$ 100	p <sub>0</sub> , p <sub>1</sub>		10.04	$\frac{7}{2}^-$
10.23 $\pm$ 20	250 $\pm$ 80	$\alpha_0$		10.22	
10.42 $\pm$ 20	160 $\pm$ 40	p <sub>0</sub> , p <sub>1</sub> , p <sub>3</sub>		10.40	( $\frac{5}{2}^+$ )
10.525 $\pm$ 15	165 $\pm$ 25	p <sub>0</sub> , p <sub>2</sub> , $\alpha_0$	~0.28 $\pm$ 0.03	10.499	$\frac{7}{2}^-$
(10.75 $\pm$ 50)		p <sub>0</sub> , p <sub>1</sub> , $\alpha_0$		(10.71)	( $\frac{7}{2}^-$ )
10.83 $\pm$ 20	120 $\pm$ 40	p <sub>0</sub> , p <sub>2</sub> , (p <sub>3</sub> ), ( $\alpha_0$ )		10.79	
10.96 $\pm$ 100	560 $\pm$ 100	p <sub>0</sub>	0.25 $\pm$ 0.07	10.91	$\frac{1}{2}^+$
11.00 $\pm$ 20	190 $\pm$ 50	(p <sub>2</sub> ), p <sub>3</sub> , ( $\alpha_0$ )		10.95	
11.2636 $\pm$ 2.0 <sup>e</sup> )	0.20 $\pm$ 0.04	p <sub>0</sub> , p <sub>2</sub> , p <sub>4</sub> , $\alpha_0$	0.093 $\pm$ 0.013	11.1928 $\pm$ 2.1	$\frac{1}{2}^-$ ; $\frac{3}{2}^-$
11.52 $\pm$ 20	240 $\pm$ 50	p <sub>2</sub> , $\alpha_0$		11.43	
11.67 $\pm$ 40	160 $\pm$ 30	p <sub>0</sub> , p <sub>3</sub>		11.58	
12.12 $\pm$ 20	120 $\pm$ 40	p <sub>2</sub> , $\alpha_0$		12.00	
12.39 $\pm$ 20	300 $\pm$ 30	p <sub>0</sub> , p <sub>2</sub>	0.26 $\pm$ 0.03	12.25	$\frac{3}{2}^-$
12.500 $\pm$ 10	190 $\pm$ 20	p <sub>0</sub> , p <sub>1</sub> , p <sub>4</sub>	0.31 $\pm$ 0.03	12.355	$\frac{1}{2}^-$ $\frac{3}{2}^-$
=12.65	=600	p <sub>0</sub>	=0.09	=12.50	$\frac{7}{2}^-$
12.7077 $\pm$ 2.0 <sup>f</sup> )	2.83 $\pm$ 0.12	p <sub>0</sub> , p <sub>2</sub> , p <sub>4</sub> , p <sub>5</sub> , $\alpha_0$ , $\alpha_1$	0.332 $\pm$ 0.018	12.5504 $\pm$ 2.3	$\frac{5}{2}^-$ ; $\frac{3}{2}^-$
(13.06 $\pm$ 100)		p <sub>0</sub>		(12.88)	( $\frac{7}{2}^-$ )
(13.06 $\pm$ 50)		p <sub>0</sub>		(12.88)	( $\frac{1}{2}^+$ )
13.250 $\pm$ 4	2 $\pm$ 1	p <sub>0</sub> , p <sub>1+2</sub> , p <sub>3+4</sub> , p <sub>5</sub> , $\alpha_0$	0.15 $\pm$ 0.04	13.060	$\frac{5}{2}^-$ ; $\frac{3}{2}^-$
13.271 $\pm$ 4	2 $\pm$ 1	p <sub>0</sub> $\rightarrow$ p <sub>4</sub> , $\alpha_0$	0.04 $\pm$ 0.02	13.080	( $\frac{1}{2}^+$ ); $\frac{3}{2}^-$
13.32 $\pm$ 100	520 $\pm$ 50	p <sub>0</sub>	0.163 $\pm$ 0.016	13.13	$\frac{5}{2}^-$
14.017 $\pm$ 4	12 $\pm$ 5	p <sub>0</sub> , p <sub>1+2</sub> , p <sub>3+4</sub> , $\alpha_0$	0.02 $\pm$ 0.01	13.781	$\frac{5}{2}^+$ ; $\frac{3}{2}^-$
(14.20 $\pm$ 50)		p <sub>0</sub>		(13.95)	( $\frac{1}{2}^+$ )
14.25 $\pm$ 50	260 $\pm$ 30	p <sub>0</sub>	0.08 $\pm$ 0.01	14.00	$\frac{7}{2}^-$
14.438 $\pm$ 6	27 $\pm$ 5	p <sub>0</sub> ,	0.04 $\pm$ 0.02	14.177	$\frac{5}{2}^+$ ; $\frac{3}{2}^-$
14.5730 $\pm$ 3.0 <sup>g</sup> )	19.3 $\pm$ 1.6	p <sub>3+4</sub>	0.085 $\pm$ 0.008	14.3037 $\pm$ 3.1	$\frac{7}{2}^-$
		p <sub>0</sub> , p <sub>1+2</sub> , p <sub>3+4</sub> , p <sub>5</sub> , $\alpha_0$			$\frac{3}{2}^-$
14.65 $\pm$ 50	610 $\pm$ 50	p <sub>0</sub>	0.10 $\pm$ 0.01	14.38	$\frac{5}{2}^-$

TABLE 17.21—continued

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	$\Gamma_{\text{po}}/\Gamma$	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$
(14.94 $\pm$ 100)		$p_0$			$(\frac{3}{2}^-)$
15.00 $\pm$ 100	470 $\pm$ 100	$p_0$	0.25 $\pm$ 0.03	14.71	$\frac{1}{2}^-$
15.110 $\pm$ 20	190 $\pm$ 25	$p_0$	0.150 $\pm$ 0.015	14.809	$\frac{1}{2}^+$
(15.245 $\pm$ 100)		$p_0$		(14.94)	$(\frac{5}{2}^+)$
(15.30 $\pm$ 50)		$p_0$		(14.98)	$(\frac{3}{2}^+)$
(15.37 $\pm$ 100)		$p_0$		(15.05)	$(\frac{3}{2}^-)$
(15.545 $\pm$ 100)		$p_0$		(15.22)	$(\frac{7}{2}^-)$
15.9 <sup>h)</sup>	$\approx 550$	$p_0$		15.6	
		$p_0$			
17.6	1500	$p_{1+2}$		17.1	$\frac{5}{2}^-$
		$p_0$			
20.4	600	$p_{3+4}$		19.8	$\frac{3}{2}^+$
21.6	600	$p_0, (\alpha)$		20.9	$\frac{9}{2}^+$
22.6	400	$p_0, (\alpha)$		21.8	$(\frac{5}{2}^+)$
23.5	600	$p_0, p_5$		22.7	$\frac{7}{2}^+$
24.7	600	$p_0, (\alpha)$		23.8	$\frac{7}{2}^+$
26.4	1500	$p_0, (\alpha)$		25.4	$\frac{7}{2}^-$
28.3	1500	$p_0$		27.2	$\frac{5}{2}^-$
30.1	2000	$p_0$		28.9	$\frac{5}{2}^+$

<sup>a)</sup> See earlier references in tables 17.20 (Aj 71) and 17.19 (Aj 77). See also table 17.20 here.<sup>b)</sup> Preliminary value (Ba 80h). See also (St 81b).<sup>c)</sup> (Kr 81a): polarized protons; preliminary results.  $\Gamma_p = 51 \pm 8, 143 \pm 30$  and  $11.4 \pm 2$  keV for  $^{17}\text{F}^*(8.98, 9.17, 9.92)$ .<sup>d)</sup> (Kr 81a) report that there are indications of two very sharp resonances corresponding to  $^{17}\text{F}^*(9.10, 9.13)$  with  $J^\pi = \frac{1}{2}^-$  and  $\frac{3}{2}^-$ .<sup>e)</sup>  $\Gamma_{\text{po}} = 19 \pm 3$  eV (Hi 76d).<sup>f)</sup>  $\Gamma_{\text{po}} = 0.94 \pm 0.06$  keV,  $\Gamma_{\alpha_0} = 62 \pm 16$  eV,  $\Gamma_{\alpha_1} = 53 \pm 22$  eV (Hi 76d); J. Lowe, private communication.<sup>g)</sup>  $\Gamma_{\text{po}} = 1.65 \pm 0.12$  keV,  $\Gamma_{\alpha_0} = 2.6 \pm 0.7$  keV (Hi 76d).<sup>h)</sup> See also table 17.20 of (Aj 71), for possible other resonances.<sup>i)</sup> (Da 79f) have confirmed  $J^\pi = \frac{3}{2}^-$ .<sup>j)</sup>  $J^\pi = (\frac{3}{2}, \frac{5}{2})^+$  (G. Bauer and H. T. Richards, private communication).

The ( $\gamma_0 + \gamma_1$ ) yield at  $90^\circ$  has been studied for  $E_p = 15.75$  to  $31.66$  MeV: it shows the giant dipole resonance centered at  $E_x = 22$  MeV with a width of  $\approx 5$  MeV and a pigmy resonance centered at  $17.5$  MeV. The integrated strength of the, mainly  $T = \frac{1}{2}$ , giant resonance is  $10$  MeV · mb: the observed strength distribution is in good agreement with odd parity 2p-1h, 1p excitation calculations. The pigmy resonance is due to  $f_{7/2} \rightarrow d_{5/2}$ : the main  $f_{7/2}$  strength lies in two states at  $E_x = 16.9$  and  $18.0$  MeV (Ha 75h). See also (Bl 79d), (Ca 81b) (Zi 75c; astrophys.) and (Al 77i; theor.).

13. (a)  $^{16}\text{O}(p, p)^{16}\text{O}$   
(b)  $^{16}\text{O}(p, 2p)^{15}\text{N}$

$$E_b = 0.6004$$

$$Q_m = -12.1276$$

(c) $^{16}\text{O}(\text{p}, \text{pn})^{15}\text{O}$	$Q_m = -15.6639$
(d) $^{16}\text{O}(\text{p}, \text{p}\alpha)^{12}\text{C}$	$Q_m = -7.1620$

Yield curves for elastic protons, protons scattered to  $^{16}\text{O}^*(6.05, 6.13, 6.92, 7.12, 8.87)$  and for  $\gamma$ -rays from  $^{16}\text{O}^*(6.13, 6.92)$  have been studied at many energies up to  $E_p = 46$  MeV: see table 17.19 in (Aj 71) and (Aj 77) for the earlier work and (Kr 81a:  $E_{\bar{p}} = 8.5$  to 10.6 MeV;  $p_0$ ), (Da 79f:  $E_{\bar{p}} \approx 14.4$  MeV;  $p_0$ ). The observed resonances are displayed in table 17.21.

Polarization results have been reported at  $E_p = 3.47$  to 49.4 MeV [see (Aj 77)] and at  $E_p = 40$  MeV (Mo 78m;  $p_5$ ), 42.5, 44.0 and 49.3 MeV (Pe 77b;  $p_5$ ), 65 MeV (Sa 79p, Sa 81c;  $p_0$ ), 135 MeV (Ke 80g;  $p_2, p_4$ ), 800 MeV (Ad 79b, Gl 79c;  $p_0, p_{1+2}, p_3, p_4, p_5$ ) and 1 GeV (Al 78q, Al 78t, Al 80b). For total reaction cross sections see (De 79z, Dy 81). See also (Aj 77).

Polarization in quasi-elastic scattering (reaction b) has been studied at  $E_p = 200$  MeV by (Ki 80a, Ki 76m) and 635 MeV (Na 78k, Na 77s). For reactions (c) and (d) see (Aj 77). For reaction (d) see (Dy 81). For spallation measurements see (In 76d, Mo 77k) and (Vd 79). For antiproton scattering see (Au 81; theor.).

See also  $^{16}\text{O}$  and (Go 78p, Sa 78k, Wi 79, Wi 79b, Le 80u), (Ra 79h; astrophys.), (Pl 77a, De 79w, Vo 79a, Mc 80, Wh 80a, Ra 81b) and (Je 76b, Le 76p, Ko 77jj, Ph 77c, Ab 78b, Al 78k, Bi 78n, Br 78q, Gr 78g, Ph 78, Re 78e, Sc 78g, Wu 78b, Bi 79b, Ch 79p, Er 79, Ko 79e, Ma 79mm, Ph 79c, Ra 79k, Au 80d, Ay 80, De 80ff, Dm 80, Ma 80cc, Ma 80ee, Ra 80, De 81n; theor.).

$$14. \quad ^{16}\text{O}(\text{p}, \text{n})^{16}\text{F} \quad Q_m = -16.211 \quad E_b = 0.6004$$

For analyzing power measurements see (Ma 81e, Ma 81k;  $E_{\bar{p}} = 135$  MeV). See also  $^{16}\text{F}$ .

$$15. \quad ^{16}\text{O}(\text{p}, \text{d})^{15}\text{O} \quad Q_m = -13.4392 \quad E_b = 0.6004$$

The excitation function for  $d_0$  ( $\theta = 70^\circ$ ) has been measured for  $E_p = 21$  to 38.5 MeV. A strong resonance is observed at  $E_p = 24$  MeV: see table 17.21. Polarization measurements are reported at  $E_p = 65$  MeV (Ho 81;  $d_0$ ), 30.3 MeV [see (Aj 77)] and  $E_{\bar{p}} = 200$  MeV (Li 81i). See also  $^{15}\text{O}$  in (Aj 81) and (Ca 79e).

$$16. \quad (a) \quad ^{16}\text{O}(\text{p}, \text{t})^{14}\text{O} \quad Q_m = -20.4062 \quad E_b = 0.6004 \\ (b) \quad ^{16}\text{O}(\text{p}, {^3\text{He}})^{14}\text{N} \quad Q_m = -15.2428$$

The excitation function ( $\theta = 70^\circ$ ) for tritons has been measured for  $E_p = 32$  to 39.5 MeV: no structure is observed [see (Aj 77)]. Differential cross sections and analyzing powers have been studied at  $E_p = 43.8$  MeV for the transitions to  $^{14}\text{O}^*(0, 5.17, 6.29, 6.59, 7.78, 9.72)$  and  $^{14}\text{N}^*(0, 2.31, 3.95, 5.11, 7.03, 9.17)$ : attempts to fit the analyzing powers with zero-range DWBA were only successful for the first pair of analog states [ $^{14}\text{O}_{g.s.}, {^{14}\text{N}^*(2.31)}$ ] (Ma 74c). For other polarization measurements see (Aj 77). See also  $^{14}\text{N}$  and  $^{14}\text{O}$  in (Aj 81) and (Ha 77v).

$$17. \ ^{16}\text{O}(\text{p}, \alpha)^{13}\text{N} \quad Q_m = -5.2185 \quad E_b = 0.6004$$

Excitation functions of various  $\alpha$ -groups and activation functions have been measured from threshold to  $E_p = 44$  MeV: see table 17.19 in (Aj 71) and (Aj 77) for the earlier work and (Gr 77d; 6.6 to 10.4 MeV;  $\sigma_t$ ). Observed resonances are displayed in table 17.21. Some broad structures have been reported above  $E_p \approx 15$  MeV; particularly strong peaks appear at  $E_p \sim 22$  and 25.5 MeV: see (Aj 77).

This reaction is involved in explosive burning in stars: see (Aj 77) for the earlier work and (Gr 77d:  $S_0 = 4 \pm 3$  b at  $E_p(\text{c.m.}) = 6.5$  MeV). See also (Cl 77c).

$$18. \ ^{16}\text{O}(\text{d}, \text{n})^{17}\text{F} \quad Q_m = -1.6243$$

Slow neutron thresholds have been observed corresponding to the ground and first excited states of  $^{17}\text{F}$ : see (Aj 71). The  $E_x$  of  $^{17}\text{F}^*(0.50)$  is  $495.33 \pm 0.10$  keV (Al 66g; from  $\gamma$ -measurement);  $\tau_m = 407 \pm 9$  psec (Be 64f). Neutron groups have been observed corresponding to  $^{17}\text{F}^*(0, 0.50, 3.10, 3.86, 4.70)$ . Angular measurements have been obtained for the  $n_0$  and  $n_1$  groups ( $l_p = 2$  and 0, respectively:  $J'' = \frac{5}{2}^+$  and  $\frac{1}{2}^+$ ) for  $E_d \leq 12$  MeV: see table 17.21 in (Aj 71) and (An 72j;  $E_d = 3.10, 3.40$  and 3.66 MeV). For polarization measurements see  $^{18}\text{F}$  in (Aj 83). See also (Hu 80c) and (Aj 77).

$$19. \ ^{16}\text{O}(^3\text{He}, \text{d})^{17}\text{F} \quad Q_m = -4.8932$$

At  $E(^3\text{He}) = 18$  MeV, angular distributions of the deuterons to  $^{17}\text{F}^*(0, 0.50, 3.104 \pm 0.003, 3.857 \pm 0.004)$  have been measured. The spectroscopic factors for  $^{17}\text{F}^*(0, 0.50)$  are 0.94 and 0.83. Two-step processes appear to be involved in the excitation of  $^{17}\text{F}^*(3.10, 3.86)$  (Fo 75j). Angular distributions have also been measured at  $E(^3\text{He}) = 30$  MeV (Ko 77k; to  $^{17}\text{F}^*(5.1, 5.7)$  and at  $E(^3\text{He}) = 33$  MeV (Lu 80a, Ro 81c;  $d_0, d_1$ ). See also  $^{19}\text{Ne}$  in (Aj 83) and (Co 77a; theor.).

$$20. \ ^{16}\text{O}(\alpha, \text{t})^{17}\text{F} \quad Q_m = -19.2137$$

See (Aj 71, Aj 77).

$$21. \ ^{16}\text{O}(^7\text{Li}, ^6\text{He})^{17}\text{F} \quad Q_m = -9.374$$

The angular distribution involving  $^{17}\text{F}_{g.s.}$  has been measured at  $E(^7\text{Li}) = 36$  MeV (Sc 73m).

$$\begin{aligned} 22. \text{ (a)} \ ^{16}\text{O}(^{10}\text{B}, ^9\text{Be})^{17}\text{F} \quad Q_m &= -5.985 \\ \text{(b)} \ ^{16}\text{O}(^{11}\text{B}, ^{10}\text{Be})^{17}\text{F} \quad Q_m &= -10.628 \end{aligned}$$

Angular distributions have been measured at  $E(^{10}\text{B}) = 100$  MeV for the transitions to  $^{17}\text{F}^*(0+0.50, 5.1, 8.1)$  (Na 75j), and at  $E(^{11}\text{B}) = 115$  MeV for the transition to  $^{17}\text{F}_{g.s.}$  (Ra 79c).

$$\begin{aligned} 23. \text{ (a)} \ ^{16}\text{O}(^{12}\text{C}, ^{11}\text{B})^{17}\text{F} \quad Q_m &= -15.3566 \\ \text{(b)} \ ^{16}\text{O}(^{13}\text{C}, ^{12}\text{B})^{17}\text{F} \quad Q_m &= -16.933 \end{aligned}$$

**<sup>17</sup>F**

114

F. AJZENBERG-SELOVE

Angular distributions have been measured at  $E(^{12}\text{C}) = 76.8 \text{ MeV}$  (Mo 77y, Mo 79z;  $^{17}\text{F}^*(0, 0.5)$ ) and at  $E(^{13}\text{C}) = 105 \text{ MeV}$  (Ra 79c;  $^{17}\text{F}_{\text{g.s.}}$ ).



Angular distributions involving  $^{17}\text{F}^*(0, 0.5)$  have been measured at  $E(^{14}\text{N}) = 76.2 \text{ MeV}$  (Mo 77y, Mo 79z) and 155 MeV (Na 75j, Na 76h).



Angular distributions involving  $^{17}\text{F}^*(0+0.5)$  have been measured at  $E(^{16}\text{O}) = 95.2 \text{ MeV}$  (Mo 77y, Mo 79z).



Angular distributions of the  $n_0$  and  $n_1$  groups have been obtained for  $E_p = 6.95$  to 13.50 MeV ( $n_0$ ) and 6.95 to 12.45 MeV ( $n_1$ ). There appears to be collective enhancement in the  $L = 2$  transition to  $^{17}\text{F}^*(0.5)$ . A large spin-flip term in the effective two-body force is necessary to account for the strength of the ground state transition (An 69a). See also  $^{18}\text{F}$  in (Aj 83).



Angular distributions have been studied for  $t_0$  and  $t_1$  at  $E(^3\text{He}) = 17.3 \text{ MeV}$  [see (Aj 77)] and for  $t_0$  at  $E(^3\overline{\text{He}}) = 33 \text{ MeV}$  (Ba 81c).



See  $^{17}\text{Ne}$ .



See  $^{19}\text{F}$  in (Aj 83).



Angular distributions have been measured for the  $t_0 \rightarrow t_3$  groups at  $E_p = 22.8$ , 42.4 and 45 MeV: see (Aj 77).



See (Gr 77d, Ko 80w) and (Aj 77).



At  $E(^3\text{He}) = 75$  and 88 MeV  $^{17}\text{F}^*(0, 0.5)$  have been populated in both reactions (Ke 78c).

$^{17}\text{Ne}$ 

(Figs. 8 and 9)

GENERAL<sup>†</sup>*Theory and reviews:* (Be 75cc, Ce 77c, Gu 78d, Wo 78f, Be 79a)*Other topics:* (Gr 81b)

**Mass of  $^{17}\text{Ne}$ :** The mass excess adopted by (Wa 77) is  $16.478 \pm 0.026$  MeV, based on unpublished data. We retain the mass excess  $16.48 \pm 0.05$  MeV based on the evidence reviewed in (Aj 77).

TABLE 17.22  
Energy levels of  $^{17}\text{Ne}^*$ )

$E_x$ in $^{17}\text{Ne}$ (MeV)	$J''; T$	$\tau_{1/2}$ (msec)	Decay	Reaction
0	$\frac{1}{2}^-; \frac{3}{2}^-$	$109.0 \pm 1.0$	$\beta^+ b)$	1

) The evidence for excited states of  $^{17}\text{Ne}$  has not been published: see (Aj 77).

b) See also tables 17.3 and 17.23.

1. (a)  $^{17}\text{Ne}(\beta^+)^{17}\text{F}^* \rightarrow {}^{16}\text{O} + p$        $Q_m = 13.93$   
 (b)  $^{17}\text{Ne}(\beta^+)^{17}\text{F}$        $Q_m = 14.53$

The half-life of  $^{17}\text{Ne}$  is  $109.0 \pm 1.0$  msec (Ha 71b). Earlier values (see (Aj 71)) gave a mean value of  $108.0 \pm 2.7$  msec. The decay is primarily to the proton unstable states of  $^{17}\text{F}$  at 4.70, 5.52 and 6.04 MeV with  $J'' = \frac{3}{2}^-, \frac{3}{2}^-$  and  $\frac{1}{2}^-$ : see table 17.23. The super-allowed decay to the analog state [ $^{17}\text{F}^*(11.20)$ ] has  $\log ft = 3.29^{+0.04}_{-0.07}$ . The character of the decay leads to  $J'' = \frac{1}{2}^-$  for  $^{17}\text{Ne}_{g.s.}$  (Ha 71b). See table 17.3 for a comparison of the mirror  $^{17}\text{N}$  and  $^{17}\text{Ne}$  decays and table 17.11 for the decay of  $^{17}\text{F}^*(11.20)$ . See also (Ra 78f).

2.  $^{20}\text{Ne}(^3\text{He}, ^6\text{He})^{17}\text{Ne}$        $Q_m = -26.19$

See (Aj 77).

 $^{17}\text{Na}$ 

(Not illustrated)

$^{17}\text{Na}$  has not been observed: its mass excess is predicted to be 35.61 MeV by (Ke 66c). It is then unbound with respect to breakup into  ${}^{16}\text{Ne} + p$  by 4.3 MeV and with respect to breakup into  ${}^{14}\text{O} + 3p$  by 5.7 MeV.

<sup>†</sup> See also (Aj 77).

TABLE 17.23  
 $\beta^+$  decay of <sup>17</sup>Ne <sup>a)</sup>

Decay to <sup>17</sup> F* (MeV $\pm$ keV)	$J^\pi$	Branching (%)	$\log ft$ <sup>b)</sup>	Decay to <sup>16</sup> O* (MeV)	Decay (%)
0	$\frac{5}{2}^+$	$0.5 \pm 0.2$ <sup>c)</sup>	$6.95 \pm 0.13$		
0.50	$\frac{1}{2}^+$	$1.1 \pm 0.5$ <sup>c)</sup>	$6.55 \pm 0.21$		
3.084 $\pm$ 30	$\frac{1}{2}^-$	$0.48 \pm 0.07$	$6.44 \pm 0.06$	0	100
4.609 $\pm$ 15	$\frac{3}{2}^-$	$16.2 \pm 0.7$	$4.59 \pm 0.02$	0	100
5.480 $\pm$ 10	$\frac{3}{2}^-$	$54.0 \pm 0.7$	$3.86 \pm 0.01$	0	100
6.037 $\pm$ 10	$\frac{1}{2}^+$	$10.6 \pm 0.2$	$4.42 \pm 0.01$	0	100
6.406 $\pm$ 30		$0.35 \pm 0.10$	$5.80 \pm 0.13$	0	100
7.708 $\pm$ 30	$\frac{1}{2}^+$	$0.18 \pm 0.05$	$5.67 \pm 0.12$	0	$>95$
				6.05	<5
8.075 $\pm$ 10	$\frac{5}{2}^+$	$6.83 \pm 0.11$	$3.96 \pm 0.01$	0	99.5
				6.05	$0.49 \pm 0.02$
8.436 $\pm$ 10		$6.51 \pm 0.26$	$3.85 \pm 0.02$	0	94.3
				6.05	$5.7 \pm 0.5$
8.825 $\pm$ 25		$1.90 \pm 0.06$	$4.23 \pm 0.02$	0	92.4
				6.05	$7.6 \pm 1.1$
11.19 <sup>c)</sup>	$\frac{1}{2}^-$ ; $T = \frac{3}{2}$	$0.71^{+0.10}_{-0.05}$	$3.29^{+0.04}_{-0.07}$	0	$10 \pm 2$
				6.13	$22 \pm 2$
				6.92	$24 \pm 6$
				7.12	$44 \pm 4$
<sup>d)</sup>					

<sup>a)</sup> (Ha 71b). See also table 17.23 in (Aj 71).

<sup>b)</sup>  $\log ft$  values calculated by (Ha 71b) using an atomic mass excess of  $16.517 \pm 0.026$  MeV [and  $\tau_{1/2} = 109.0 \pm 1.0$  msec] rather than the presently adopted  $16.48 \pm 0.05$  MeV. Since this energy difference leads to quite small changes, the original calculations are quoted here. However, table 17.3 (which compares the analog decays) shows corrected  $ft$  values.

<sup>c)</sup> Calculated branchings, based on the mirror <sup>17</sup>N decay.

<sup>d)</sup> A proton group with  $E_{c.m.} = 2.83$  MeV has been observed: the level in <sup>17</sup>F to which it corresponds is not known.

<sup>e)</sup> See also table 17.11.

## References

(Closed July 1, 1981)

- (i) Proc. of the Int. Conf. on nuclear structure – contributed papers – Tokyo, September 5–10, 1977, published by the Organizing Committee; referred to herein as “TOKYO”.
- (ii) Conf. Proc. of the 7th Int. Conf. on high energy physics and nuclear structure, Zurich, August 29–September 2, 1977; abstract volume published by the Swiss Institute for Nuclear Research, 5234 Villigen, Switzerland; referred to herein as “SIN”.
- (iii) Conf. Proc. of the 8th Int. Conf. on high energy physics and nuclear structure, Vancouver, August 13–17, 1979; abstract volume of contributed papers; referred to herein as “TRIUMF”.
- (iv) Abstracts of Contributed Papers to Int. Conf. on nuclear physics with electromagnetic interactions, Mainz, June 5–9, 1979; referred to herein as “MAINZ”.

(v) Abstracts of contributed papers to the Int. Conf. on nuclear physics, Berkeley, California, USA, August 24-30, 1980, published as LBL-11118 (1980); referred to herein as "BERKELEY".

(vi) Abstracts of contributed papers to the Int. Conf. on high energy physics and nuclear structure, Versailles, France, July 6-10, 1981, published by the Organizing Committee; referred to herein as "VERSAILLES".

## ERRATA<sup>†</sup>

Energy levels of light nuclei,  $A = 13-15$ , Nucl. Phys. A360 (1981) 1

p. 144 Table 15.18

The  $J^\pi$  of  $^{15}\text{O}^*(9.527)$  should be listed as  $(\frac{3}{2})^+$  and reaction 16 should be added.

The level listed at 9.72 MeV should be entirely deleted.

p. 152 Table 15.23

The same changes should be made here also.

- |       |  |
|-------|--|
| AB77  | ABUL-MAGD, KHALIL AND SHALABY, J. OF PHYS. G3 (1977) 381   |
| AB77A | ABGRALL, MORAND, CAURIER AND GRAMMATICOIS, PHYS. REV. LETT. 39 (1977) 922                            |
| AB77E | ABE, RIFP-296 (1977) (KYOTO U.)  |
| AB78A | ABE, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 132                                     |
| AB78B | ABUL-MAGD AND HUFNER, NUCLEAR PHYSICS A308 (1978) 429  |
| AB79A | ABDI AND WEIGEL, PHYS. REV. C19 (1979) 1553  |
| AB79C | ABELA ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 1825                                     |
| AB79D | ABDUL-JALIL AND JACKSON, J. OF PHYS. G 5 (1979) 1699   |
| AB79E | ABDI AND WEIGEL, NUOVO CIM. A 54 (1979) 11   |
| AB80B | ABLEEV ET AL, P. 71, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980) |
| AB80D | ABE, KONDO AND MATSUSE, SUPPL. PROGR. THEOR. PHYS. 68 (1980) 303                                     |
| AB81A | ABE, RIFP-421, 428   |
| AC81  | ACQUADRO, HUSSEIN, PEREIRA AND SALA, PHYS. LETT. 100B (1981) 381                                     |
| AD69  | ADELBERGER, MC DONALD AND BARNES, NUCLEAR PHYSICS A124 (1969) 49                                     |
| AD70A | ADELBERGER, NERO AND MC DONALD, NUCLEAR PHYSICS A143 (1970) 97                                       |
| AD73B | ADELBERGER ET AL, PHYS. REV. C7 (1973) 889   |
| AD78B | ADACHI AND YOSHIDA, NUCLEAR PHYSICS A306 (1978) 53   |
| AD79  | ADHIKARI, PHYS. REV. C19 (1979) 325  |
| AD79B | ADAMS ET AL, PHYS. REV. LETT. 43 (1979) 421  |
| AF77  | AFANAS'EV, DOBROMYSLOV, PHUNG AND SHILOV, BULL. ACAD. SCI. USSR, PHYS. SER. 41 (1977) 98             |
| AG80  | AGRELLO, AGUILERA-NAVARRO AND MAKI, LETT. NUOVO CIM. 28 (1980) 310                                   |

<sup>†</sup> I am very indebted to Dr. E. K. Warburton for pointing out these errors to me.

- AG81 AGRELLO AND PORTILHO, PHYS. REV. C23 (1981) 1898  
 AH70A AHRENS ET AL, PHYS. LETT. 31B (1970) 570  
 AH75B AHRENS ET AL, NUCLEAR PHYSICS A251 (1975) 479  
 AJ59 AJZENBERG-SELOVE AND LAURITSEN, NUCLEAR PHYSICS 11 (1959) 1  
 AJ68 AJZENBERG-SELOVE AND LAURITSEN, NUCLEAR PHYSICS A114 (1968) 1  
 AJ71 AJZENBERG-SELOVE, NUCLEAR PHYSICS A166 (1971) 1  
 AJ76 AJZENBERG-SELOVE, NUCLEAR PHYSICS A268 (1976) 1  
 AJ76B AJZENBERG-SELOVE, BETTS AND CROZIER, PHYS. REV. C14 (1976) 357  
 AJ77 AJZENBERG-SELOVE, NUCLEAR PHYSICS A281 (1977) 1  
 AJ78 AJZENBERG-SELOVE, NUCLEAR PHYSICS A300 (1978) 1  
 AJ79 AJZENBERG-SELOVE, NUCLEAR PHYSICS A320 (1979) 1  
 AJ80 AJZENBERG-SELOVE AND BUSCH, NUCL. PHYS. A336 (1980) 1  
 AJ81 AJZENBERG-SELOVE, NUCL. PHYS. A360 (1981) 1  
 AJ83 AJZENBERG-SELOVE, NUCLEAR PHYSICS (1983)  
 AK77 AKAISHI, KASAHARA, NAGATA AND YAMAGUCHI, IN TOKYO (1977), CONTRIBUTED PAPERS P. 52  
 AK77A AKBAROV, UZAKOVA AND UMIRBEKOV, SOVIET ATOMIC ENERGY 42 (1977) 468  
 AK78 AKSELSSON AND JOHANSSON, BULL. AM. PHYS. SOC. 23 (1978) 1035  
 AK80B AKSINENKO ET AL, NUCLEAR PHYSICS A348 (1980) 518  
 AL58C ALBURGER, PHYS. REV. 111 (1958) 1586  
 AL59F ALBURGER, GALLMANN AND WILKINSON, PHYS. REV. 116 (1959) 939  
 AL64G ALEXANDER, BROUDE AND LITHERLAND, NUCLEAR PHYSICS 53 (1964) 593  
 AL66G ALBURGER ET AL, PHYS. REV. 148 (1966) 1050  
 AL71C ALLEN AND MACKLIN, PHYS. REV. C3 (1971) 1737  
 AL72M ALBURGER AND WILKINSON, PHYS. REV. C6 (1972) 2019  
 AL76 ALBURGER AND WILKINSON, PHYS. REV. C13 (1976) 835  
 AL76D ALBURGER, NUCL. INSTR. METH. 136 (1976) 323  
 AL76K ALI, PROC. 7TH INTERN. CONF. ON FEW BODY PROBLEMS IN NUCLEAR AND PARTICLE PHYSICS, DELHI 1975 (AMSTERDAM, NORTH-HOLLAND 1976), P. 438, AND PHYS. ABSTRACTS 18501 (1977)  
 AL77AA ALDER ET AL, IN PROC., ZURICH - SIN, 1977, P. 45  
 AL77D ALKHAZOV, NUCLEAR PHYSICS A280 (1977) 330  
 AL77F ALDER ET AL, HELV. PHYS. ACTA 50 (1977) 197  
 AL77G ALBURGER, PHYS. REV. C16 (1977) 889  
 AL77I ALBERT ET AL, PHYS. REV. C16 (1977) 503  
 AL78A ALBANESE ET AL, PHYS. LETT. B73 (1978) 119  
 AL78D ALBURGER, PHYS. REV. C18 (1978) 576  
 AL78F ALCARAS AND DA SILVEIRA, NUCLEAR PHYSICS A298 (1978) 228  
 AL78I ALKHAZOV, BELOSTOTSKY AND VOROB'YOV, PHYS. REP. 42C (1978) 89  
 AL78K ALKHAZOV ET AL, YADERNAYA FIZ. 27 (1978) 333  
 AL78Q ALKHAZOV ET AL, PIS'MA V ZH. EKSP. & TEOR. FIZ. 28 (1978) 713  
 AL78R ALEXANDER AND FORSTER, IN ADVANCES IN NUCLEAR PHYSICS, VOL. 10, ED. BY BARANGER AND VOGT (PLENUM PRESS, NEW YORK, 1978), P. 197  
 AL79K ALSTER AND WARSZAWSKI, PHYS. REPORTS 52 (1979) 87  
 AL79N ALHASSID, LEVINE, KARP AND STEADMAN, PHYS. REV. C20 (1979) 1789  
 AL80B ALKHAZOV ET AL, PHYS. LETT. 90B (1980) 364

- AL80E ALMA-ATA, DUBNA AND TBILISI COLLABORATION, P. 519  
 PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR  
 PHYSICS, BERKELEY (1980)
- AL80G ALBANESE ET AL, NUCLEAR PHYSICS A350 (1980) 301
- AL80H ALBURGER, TRANSACTIONS OF THE N. Y. ACAD. OF SCIENCES 40  
 (1980) 1
- AL81D ALAMANOS ET AL, NUCL. PHYS. A363 (1981) 477
- AM76B AMOS AND GERAMB, LETT. NUOVO CIMENTO 16 (1976) 165
- AM78A AMES, BULL. AM. PHYS. SOC. 23 (1978) 540
- AM78B AMAKAWA, MORI AND YAZAKI, PHYS. LETT. B76 (1978) 157
- AM78C AMSEL ET AL, NUCL. INSTR. METH. 149 (1978) 705, 713
- AM79 AMES, BULL. AM. PHYS. SOC. 24 (1979) 612
- AM79A AMES, THESIS, U. WISCONSIN (1979)
- AM79B AMAKAWA AND KUBO, PHYS. REV. C19 (1979) 2403
- AM79E AMANN ET AL, BULL. AM. PHYS. SOC. 24 (1979) 819
- AM80A AMAKAWA, MORI AND YAZAKI, NUCL. PHYS. A340 (1980) 125
- AM80E AMADO, LENZ, MC NEIL AND SPARROW, PHYS. REV. C22  
 (1980) 2094
- AM81B AMES, TO BE PUBLISHED
- AN69A ANDERSON ET AL, PHYS. REV. 177 (1969) 1416
- AN72J ANDERSON, VELKLEY AND WILLARD, PHYS. REV. C6 (1972) 1513
- AN76C ANTONOV AND GARISTOV, BULG. J. PHYS. (BULGARIA) 3  
 (1976) 20; PHYS. ABS. 89932 (1976)
- AN76D ANTONOV, GARISTOV AND PETKOV, C. R. ACAD. BULG. SCI.  
 (BULGARIA) 29 (1976) 967, AND PHYS. ABSTRACTS 93027  
 (1976)
- AN77A ANDO, BANDO AND NAGATA, PROG. THEOR. PHYS. 57 (1977) 1303
- AN77B ANAGNOSTATOS, ATOMKERNENERGIE 29 (1977) 207
- AN77F ANGELI AND CSATLOS, NUCLEAR PHYSICS A288 (1977) 480
- AN78 ANDO, IKEDA AND TOHSAKI-SUZUKI, PROG. THEOR. PHYS. 59  
 (1978) 306
- AN78A ANDO, IKEDA AND TOHSAKI-SUZUKI, PROG. THEOR. PHYS. 59  
 (1978) 2168
- AN78B ANGELI AND CSATLOS, ATOMKI KOZLEM. 20 (1978) 1
- AN78F ANASTASIO ET AL, PHYS. REV. C18 (1978) 2416
- AN78H ANDRADE, ROBLEDO AND DORANTES, BULL. AM. PHYS. SOC. 23  
 (1978) 1064
- AN79C ANNI, RENNA AND TAFFARA, LETT. NUOVO CIM. 25 (1979) 121
- AN79D ANIOL ET AL, IN HOUSTON, AIP CONF. PROCEEDINGS 54  
 (1979), P. 276
- AN79G ANDO, IKEDA AND TOHSAKI-SUZUKI, PROG. THEOR. PHYS. 61  
 (1979) 101
- AN79L ANIOL ET AL, BULL. AM. PHYS. SOC. 24 (1979) 821
- AN79O ANDERSON, JARMIE, BARRETT AND ARTHUR, PHYS. REV. C20 (1979)  
 897
- AN79R ANDREEV AND TOMOZOVS, UKR. FIZ. ZH. (USSR) 24 (1979) 1888,  
 AND PHYS. ABSTRACTS 32288(1980)
- AN80F ANANTARAMAN, FULBRIGHT AND STWERTKA, PHYS. REV. C22  
 (1980) 501
- AN80G ANDERSON ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 729
- AN80J ANANTARAMAN, PHYS. REV. C22 (1980) 945
- AN80K ANDO AND BANDO, P. 5,6, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY  
 (1980)

- AN80L ANAGNOSTATOS, TOULIATOS, KYRITISIS AND YAPITZAKIS,  
ATOMKERNERNERG. KERNTECH (GERMANY) 35 (1980) 60
- AN80N ANDO, IKEDA AND TOHSAKI-SUZUKI, PROG. THEOR. PHYS. 64 (1980)  
1608
- AN81 ANDERSON ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 535
- AN81C ANIKINA ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 447
- A079 AOKI ET AL, NUCLEAR PHYSICS A322 (1979) 117
- AP76 APAGYI, MAGY. FIZ. FOLY. (HUNGARY) 24 (1976) 2
- AP76A APAGYI, FAI AND NEMETH, NUCLEAR PHYSICS A272 (1976) 303
- AP76B APAGYI, FAI AND NEMETH, NUCLEAR PHYSICS A272 (1976) 317
- AP77 APAGYI AND FAI, J. OF PHYS. G3 (1977) L163
- AP78 APAGYI, J. OF PHYS. G4 (1978) 1859
- AP80 APAGYI AND VERTSE, PHYS. REV. C21 (1980) 779
- AR76G ARNOULD AND HOWARD, NUCLEAR PHYSICS A274 (1976) 295
- AR76H ARTEMOV ET AL, SOVIET J. NUCL. PHYS. 23 (1976) 257
- AR76I ARTEMOV ET AL, YADERNAYA FIZ. 24 (1976) 3
- AR77A ARNETT, ASTROPHYS. J. (SUPPL. SERIES) 35 (1977) 145
- AR77B ARTUKH, GRIDNEV, MIKHEEV AND VOLKOV, NUCLEAR PHYSICS A283  
(1977) 350
- AR78A ARIMA, HYPERFINE INTERACTIONS 4 (1978) 151
- AR78B ARIMA, HORIKAWA, HYUGA AND SUZUKI, PHYS. REV. LETT. 40  
(1978) 1001, AND IBID. 42 (1979) 1186
- AR78G ARIMA, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47  
(1978) 1
- AR78H ARTEMOV ET AL, SOVIET J. NUCL. PHYS. 27 (1978) 1
- AR78J ARTEMOV ET AL, YADERNAYA FIZ. 28 (1978) 288
- AR78L ARTYUKH ET AL, IZV. AKAD NAUK SSSR (SER. FIZ.) 42  
(1978) 2234
- AR79E ARTEMOV ET AL, NUCLEAR PHYSICS A320 (1979) 479
- AR79K ARVIEUX ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
PAPER 1E18
- AR79N ARIMA, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108 (1979) P. 9
- AR80 ARVIEUX, NUCL. PHYS. A335 (1980) 353
- AR80D ARIMA AND STROTTMAN, PHYSICAL LETTERS B96 (1980) 23
- AR81 ARIMA, NUCL. PHYS. A354 (1981) 19c
- AS75A ASHER ET AL, J. OF PHYS. G1 (1975) 415
- AS80 ASHERY, NUCL. PHYS. A335 (1980) 385
- AS81A ASHERY, NUCL. PHYS. A354 (1981) 555c
- AT78 ATARASHI, HIRA AND NARUMI, PROG. THEOR. PHYS. 60 (1978)  
209
- AT81 ATKIN AND DUMBRAJS, PROCEEDINGS VERSAILLES CONF. (1981)  
p. 247
- AU75 AUDI, DETRAZ, LANGEVIN AND POUGHEON, NUCLEAR PHYSICS A237  
(1975) 300
- AU77D AUERBACH, VAN GIAI AND LEE, PHYS. LETT. B68 (1977) 225
- AU77F AUSTIN, MSUCL-254 (1977)
- AU77H AUDOUZE, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D.  
REIDEL, HOLLAND, 1977), P. 3
- AU77I AUDOUZE, LEQUEUX, ROCCA-VOLMERANGE AND VIGROUX, IN CNO  
ISOTOPES IN ASTROPHYSICS, J. AUDOUZE ED. (D. REIDEL,  
HOLLAND, 1977), P. 155
- AU77J AUER AND DEMARQUE, IN CNO ISOTOPES IN ASTROPHYS., J.  
AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 25
- AU78B AUGER AND LOMBARD, PHYS. LETT. B77 (1978) 165, AND J.  
OF PHYS. G4 (1978) L261

- AU80D AUGER AND LOMBARD, NUCL. PHYS. A350 (1980) 332  
AU80E AUDOUZE, CHIEZE AND VANGIONI-FLAM, ASTRONOMY AND ASTROPHYSICS 91 (1980) 49  
AU81 AUERBACH, DOVER AND KAHANA, PHYS. REV. LETT. 46 (1981) 702  
AU81A AULD, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 93  
AV78 AVISHAI, Z. PHYS. A285 (1978) 333  
AV81B AVER'YANOV, GOLUBEV AND SADOVOI, YAD. FIZ. 33 (1981) 66  
AY79 AYOUB AND MAVROMATIS, NUCL. PHYS. A323 (1979) 125  
AY80 AYTURK AND SVENNE, CAN. J. PHYS. 58 (1980) 1026  
AZ77 AZUELOS, KITCHING AND RAMAVATARAM, PHYS. REV. C15 (1977) 1847  
AZ77A AZUELOS AND KITCHING, NUCLEAR PHYSICS A285 (1977) 19  
BA61C BARROS, FORSYTH, JAFFE AND TAYLOR, PROC. PHYS. SOC. 77 (1961) 853  
BA69N BAHR, BECKER, JAHR AND KUHLMANN, NUCLEAR PHYSICS A129 (1969) 388  
BA70V BARKER, COCKBURN, SEILER AND MARMIER, PHYS. REV. LETT. 25 (1970) 1350  
BA70W BARKER ET AL, NUCLEAR PHYSICS A155 (1970) 401  
BA71K BASSANI, PAPPALARDO, SAUNIER AND TRAORE, PHYS. LETT. 34B (1971) 612  
BA76CC BARKER, PHYS. LETT. B64 (1976) 251  
BA76DD BAYE, NUCLEAR PHYSICS A272 (1976) 445  
BA76GG BANDO AND KRENCIGLOWA, NUCLEAR PHYSICS A273 (1976) 95  
BA76II BANGERT, BERG, WIENHARD AND WOLF, Z. PHYS. A278 (1976) 295  
BA76KK BARZ ET AL, SOVIET J. NUCL. PHYS. 24 (1976) 264  
BA77 BARZ, ROTTÉR AND HOHN, NUCLEAR PHYSICS A275 (1977) 111  
BA77A BARKER  
BA77AB BARNES, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 783  
BA77AJ BATTY ET AL, IN PROC., ZURICH - SIN, 1977, P. 301  
BA77AL BATIST, BERLOVICH, MEZILEV AND NOVIKOV, SOVIET J. NUCL. PHYS. 25 (1977) 605  
BA77AO BADAWY, ABDO AND TAWFIK, ACTA PHYS. ACAD. SCI. HUNGARY 43 (1977) 269  
BA77B BAYE AND HEENEN, NUCLEAR PHYSICS A276 (1977) 354  
BA77F BARKER  
BA77KK BAJAJ AND NOGAMI, ANN. OF PHYS. 103 (1977) 141  
BA77L BAZ ET AL, LETT. NUOVO CIM. 18 (1977) 227  
BA77M BACHELIER ET AL, PHYS. REV. C15 (1977) 2139  
BA77MM BAERI AND STRAZZERI, LETT. NUOVO CIM. 20 (1977) 41  
BA77OO BAYE, DEENEN AND SALMON, NUCLEAR PHYSICS A289 (1977) 511  
BA77QQ BALAMUTH, LIND, YOUNG AND ZURMUHLE, NUCL. PHYS. A290 (1977) 65  
BA77R BAYE AND HEENEN, NUCLEAR PHYSICS A283 (1977) 176  
BA77RR BAZ', GOL'DBERG, GRIDNEV AND SEMENOV, YADERNAYA FIZ. 25 (1977) 759  
BA77T BAUR AND LENSKE, NUCLEAR PHYSICS A282 (1977) 201  
BA77V BARROSO, NUCLEAR PHYSICS A281 (1977) 267  
BA77W BASS, PHYS. REV. LETT. 39 (1977) 265  
BA77WW BATUSOV AND ERAMZHIAN, FIZ. ELEM. CHASTITS & AT. YADRA (USSR) 8 (1977) 229  
BA77XX BAER, IN "HIGH ENERGY PHYSICS AND NUCLEAR STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 245  
BA77YY BACKENSTOSS, ACTA PHYS. AUSTRIACA SUPPL. 18 (1977) 593

- BA78 BARRETTE ET AL, PHYS. REV. LETT. 40 (1978) 445  
 BA78AA BARZ, ROTTER AND HOHN, NUCLEAR PHYS. A307 (1978) 285  
 BA78E BADING AND MAXSON, BULL. AM. PHYS. SOC. 23 (1978) 525  
 BA78F BARKER, AUSTRALIAN J. PHYS. 31 (1978) 27  
 BA78FF BALANDIN ET AL, YADERNAYA FIZ. 28 (1978) 582  
 BA78J BAUHOFF AND WILDERMUTH, IN PROCEEDINGS, CLUSTER,  
       WINNIPEG 1978, A9  
 BA78L BASSICHIS, STRAYER, READING AND SCHEERBAUM, PHYS. REV.  
       C18 (1978) 632  
 BA78P BACK ET AL, BULL. AM. PHYS. SOC. 23 (1978) 958  
 BA78R BAXTER, IKOSSI, MC DONALD AND KUEHNER, NUCLEAR PHYSICS  
       A305 (1978) 213  
 BA78W BALASHOV AND WUNSCH, J. OF PHYS. G4 (1978) L239  
 BA78Y BAYE, HEENEN AND LIBERT-HEINEMANN, NUCLEAR PHYSICS A308  
       (1978) 229  
 BA79AA BALL ET AL, NUCL. PHYS. A325 (1979) 305  
 BA79DD BASSICHIS AND STRAYER, PHYS. REV. C20 (1979) 915  
 BA79EE BARRETTE ET AL, PHYS. REV. C20 (1979) 1759  
 BA79GG BALTZ, GLENDENNING, KAUFFMANN AND PRUESS, NUCL. PHYS.  
       A327 (1979) 221  
 BA79HH BATTY ET AL, NUCL. PHYS. A329 (1979) 407  
 BA79I BALASHOV AND WUNSCH, IN 1978 RPI SYMPOSIUM "PHOTOPION  
       NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 125  
 BA79J BARGHOLTZ, PHYS. LETT. B81 (1979) 286  
 BA79KK BAYE AND SALMON, NUCL. PHYS. A331 (1979) 254  
 BA79MM BAUMGARDT AND SCHOPPER, J. OF PHYS. G 5 (1979) L231  
 BA80A BARTKE, NUCL. PHYS. A335 (1980) 481  
 BA80AA BACKENSTOSS ET AL, PHYS. LETT. B95 (1980) 212  
 BA80F BANERJEE AND WALLACE, PHYS. REV. C21 (1980) 1996  
 BA80H BARKER, ET AL, IN ATOMIC MASSES AND FUNDAMENTAL CONSTANTS,  
       PLENUM (1980) p. 233  
 BA80JJ BARTEL AND FELDMEIER, Z. PHYS. A297 (1980) 333  
 BA80K BARNES, DAP - 593 (JULY 1980)  
 BA80KK BARASHENCOV AND MUSULMANBECOV, Z. PHYS. A296 (1980) 371  
 BA80L BARFOOT ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 787  
 BA80O BAR-TOUV AND MOALEM, P. 782, PROCEEDINGS OF THE  
       INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY  
       (1980)  
 BA80OO BAYE, PHYS. LETT. 97B (1980) 17  
 BA80T BARREAU ET AL, P. 125, PROCEEDINGS OF THE INTERNATIONAL  
       CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 BA80V BALASHOV, BARZ, ROTTER AND WUNSCH, P. 143, PROCEEDINGS  
       OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS,  
       BERKELEY (1980)  
 BA80Y BASSALLECK ET AL, NUCL. PHYS. A343 (1980) 365  
 BA81 BALDOCK, ROBSON AND BARRETT, NUCL. PHYS. A351 (1981) 157  
 BA81A BAR-TOUV AND MOALEM, NUCL. PHYS. A351 (1981) 285  
 BA81C BASAK ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE  
       PROC. 69 (1981) p. 593  
 BA81K BASON ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 293  
 BA81L BAUER ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 321  
 BA81R BANAIGS ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 568  
 BA81U BARKER, SMITH, MORRISON AND AMOS, J. OF PHYS. G7 (1981) 657  
 BA81W BARREAU ET AL, NUCL. PHYS. A358 (1981) C371  
 BE64F BECKER AND WILKINSON, PHYS. REV. 134 (1964) B1200

- BE66AA BERG AND KASHY, NUCL. INSTR. METH. 39 (1966) 169  
 BE67J BECKER, OLNESS AND WILKINSON, PHYS. REV. 155 (1967) 1089  
 BE73GG BERGSTROM AND AUER, NUCLEAR PHYSICS A215 (1973) 232  
 BE75BB BELLOTTI ET AL, NUOVO CIM. A29 (1975) 106  
 BE75CC BENENSON, NUKLEONIKA (POLAND) 20 (1975) 775  
 BE75EE BERNARD AND TAYLOR, NUCLEAR CROSS SECTIONS AND TECHNOLOGY, PT. II, WASHINGTON, D.C., 1975 (WASH. D.C., NAT. BUR. STANDARDS 1975), P. 938, AND PHYS. ABSTRACTS 23093 (1977)  
 BE76BB BERTRAND, ANN. REV. NUCL. SCI. 26 (1976) 457  
 BE76V BEREZOVNI, UKR. FIZ. ZH. (USSR) 21 (1976) 391  
 BE76W BERMAN, PREPRINT UCRL-78482 (1976)  
 BE77AA BERNSTEIN ET AL, IN PROC., ZURICH - SIN, 1977, P. 252  
 BE77BB BERNHEIM ET AL, IN PROC., ZURICH - SIN, 1977, P. 267  
 BE77GG BERDNIKOV, GISMATULIN, LANTSEV AND OSTROUMOV, SOVIET J. NUCL. PHYS. 25 (1977) 499  
 BE77HH BERDNIKOV ET AL, SOVIET J. NUCL. PHYS. 25 (1977) 610  
 BE77N BEHRMAN, DAS GUPTA AND DE TAKACSY, NUCLEAR PHYSICS A289 (1977) 397  
 BE78A BERTRAND ET AL, BULL. AM. PHYS. SOC. 23 (1978) 80, 81  
 BE78B BECK, MULLER AND KOHLER, PHYS. REV. LETT. 40 (1978) 837  
 BE78CC BEAR AND HODGSON, J. OF PHYS. G4 (1978) L287  
 BE78D BENENSON AND WILLIAMS, BULL. AM. PHYS. SOC. 23 (1978) 636  
 BE78FF BERNHARDT, BOHN, EBERHARD AND VANDENBOSCH, PROC. INT. CONF. ON RESONANCES IN HEAVY ION REACTIONS, HVAR, 1977 (NORTH-HOLLAND, NETHERLANDS, 1978), P. 367; PHYSICS ABSTRACTS 13335 (1979)  
 BE78J BECCHETTI ET AL, IN PROCEEDINGS, CLUSTER, WINNIPEG 1978, H1  
 BE78P BERTSCH AND RISKA, PHYS. REV. C18 (1978) 317 (1978) 22  
 BE78T BECCHETTI, FLYNN, HANSON AND SUNIER, NUCLEAR PHYSICS A305 (1978) 293  
 BE78U BECCHETTI, JANECKE AND THORN, NUCLEAR PHYSICS A305 (1978) 313  
 BE78V BEHERA, GURU AND SATPATHY, CZECH. J. PHYS. B28 (1978) 615  
 BE78W BECCHETTI, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 308  
 BE78Z BERDNIKOV ET AL, SOVIET J. NUCL. PHYS. 27 (1978) 320  
 BE79A BENENSON AND KASHY, REV. MOD. PHYS. 51 (1979) 527  
 BE79D BECCHETTI, JANECKE, OVERWAY AND JACOBS, BULL. AM. PHYS. SOC. 24 (1979) 572  
 BE79DD BELYAK, SOVIET J. NUCL. PHYS. 29 (1979) 24  
 BE79EE BERTRAND, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 417.  
 BE79K BENKOULA ET AL, IN PROC. THIRD SYMP. ON NEUTRON CAPTURE GAMMA RAYS, 1978 (PLENUM PRESS, NEW YORK, 1979), P. 371  
 BE79Q BEHERA AND SATPATHY, J. OF PHYS. G5 (1979) 85  
 BE79T BEHERA, PANDA AND SATPATHY, PHYS. REV. C20 (1979) 683  
 BE79U BECCHETTI ET AL, BULL. AM. PHYS. SOC. 24 (1979) 852  
 BE79Z BEVELACQUA AND PREWETT, CAN. J. PHYS. 57 (1979) 1484  
 BE80A BECCHETTI ET AL, PHYS. REV. C21 (1980) 444  
 BE80B BETTS, HINDI, DI CENZO AND PARKER, PHYS. REV. C21 (1980) 175

- BE80C BEENE, DE VRIES, SHAPIRA AND CLOVER, PHYS. REV. C21 (1980) 167
- BE80D BEHERA, PANDA AND SATPATHY, PHYS. REV. C21 (1980) 1883
- BE80F BECCHETTI ET AL, NUCL. PHYS. A339 (1980) 132
- BE80G BEHERA, PANDA AND SATPATHY, PHYS. REV. C22 (1980) 148
- BE80O BECCHETTI, OVERWAY, JANECKE AND JACOBS, NUCL. PHYS. A344 (1980) 336
- BE80Q BETZ AND KERMAN, NUCL. PHYS. A345 (1980) 493
- BE80S BERDNIKOV AND GISMATULLIN, YAD. FIZ. 31 (1980) 1139
- BE80T BERTSCH, PHYS. LETT. 95B (1980) 157
- BE80U BERDNIKOV, GISMATULLIN AND MELENT'EV, YAD. FIZ. (USSR) 31 (1980) 1433
- BE81E BEVELACQUA AND PREWETT, CAN. J. PHYS. 59 (1981) 231
- BE81U BERTINI ET AL, NUCL. PHYS. A360 (1981) 315
- BH76B BHALERAO AND DHAVALE, PROG. THEOR. PHYS. 56 (1976) 1497
- BH78B BHOWMIK ET AL, PHYS. LETT. B80 (1978) 41
- BH78C BHAKAR AND VAN OERS, IN FEW BODY SYSTEMS AND NUCLEAR FORCES, GRAZ 1978, SPRINGER LECTURE NOTES 87 (1978), P. 364
- BH80 BHALERAO, LIU AND SHAKIN, PHYS. REV. C21 (1980) 2103
- BH80B BHALERAO AND SHAKIN, P. 705, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- BH81 BHOWMIK ET AL, NUCL. PHYS. A363 (1981) 516
- BI73A BINGHAM, FORTUNE, GARRETT AND MIDDLETON, PHYS. REV. C7 (1973) 57
- BI73Q BIRK, SOKOLOWSKI AND WOLFSON, NUCLEAR PHYSICS A216 (1973) 217
- BI76E BILLOIRE AND CHEMTOB, PHYS. LETT. B64 (1976) 389
- BI77D BIZZETI, NUCL. INSTR. METH. 146 (1977) 285
- BI77E BING, ACTA PHYS. SIN. (CHINA) 26 (1977) 333, AND PHYS. ABSTRACTS 8834 (1978)
- BI78 BIRKELUND AND HUIZENGA, PHYS. REV. C17 (1978) 126
- BI78A BILLEREY ET AL, Z. PHYS. A284 (1978) 319
- BI78J BILANIUK, FORTUNE AND MIDDLETON, NUCLEAR PHYSICS A305 (1978) 63
- BI78L BISHOP, HOWES, IRVINE AND MODARRES, J. OF PHYS. G4 (1978) 1709
- BI78M BIRBRAIR, GLEZER, LAPINA AND SADOVNIKOVA, SOVIET J. NUCL. PHYS. 27 (1978) 45
- BI78N BIRBRAIR ET AL, SOVIET J. NUCL. PHYS. 28 (1978) 321
- BI79 BILLEN, PHYS. REV. C20 (1979) 1648
- BI79B BIRBRAIR, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2242
- BI81C BICE, SHOTTER, STAHEL AND CERNY, PHYS. LETT. 101B (1981) 27
- BL75I BLOCK, FUJITA, KOBAYASHI AND OOSAKI, J. NUCL. SCI. TECHNOL. 12 (1975) 1
- BL77B BLAIZOT AND GOGNY, NUCLEAR PHYSICS A284 (1977) 429
- BL79C BLOCKI AND FLOCARD, PHYS. LETT. B85 (1979) 163
- BL79D BLATT ET AL, BULL. AM. PHYS. SOC. 24 (1979) 843
- BL79E BLECHER ET AL, PHYS. REV. C20 (1979) 1884
- BL79F BLESZYNSKI AND SANDER, NUCL. PHYS. A326 (1979) 525
- BL81 BLAIZOT AND GRAMMATICOIS, NUCL. PHYS. A355 (1981) 115
- B070B BONNER, PHYS. REV. C1 (1970) 1659
- B070C BONNER, PHYS. REV. C1 (1970) 1710
- B071B BOHNE ET AL, NUCLEAR PHYSICS A160 (1971) 257

- B073HH BOHNE ET AL, PHYS. LETT. B47 (1973) 342  
 B073O BOHNE ET AL, NUCLEAR PHYSICS A196 (1973) 41  
 B074A BOWMAN, POSKANZER, KORTELING AND BUTLER, PHYS. REV. C9 (1974) 836  
 B076CC BOUVIER AND WEIBEL, ARCH. SCI. (GENEVA) 29 (1976) 163  
 B076Y BOUSSY AND HUFNER, PHYS. LETT. B64 (1976) 276  
 B076Z BOX, GABRIC, LASSEY AND MC KELLAR, J. OF PHYS. G2 (1976) L107  
 B077D BORKOWSKI ET AL, PHYS. REV. LETT. 38 (1977) 742  
 B077O BOTTINO, CIOCCHETTI AND KIM, PHYS. REV. C16 (1977) 1120  
 B077S BOSCHITZ, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 133  
 B077U BOUSSY, NUCLEAR PHYSICS A290 (1977) 324  
 B078 BOWLES ET AL, PHYS. REV. LETT. 40 (1978) 97  
 B078D BONCHE, GRAMMATICOS AND KOONIN, PHYS. REV. C17 (1978) 1700  
 B078E BOHLEN ET AL, Z. PHYS. A285 (1978) 379  
 B078L BORDERS AND HARRIS, NUCL. INSTR. METH. 149 (1978) 279  
 B078R BOIKO, GISMATULLIN AND LANTSEV, SOVIET J. NUCL. PHYS. 27 (1978) 759  
 B078S BOIKO ET AL, SOVIET J. NUCL. PHYS. 27 (1978) 765  
 B079BB BODEK ET AL, HELV. PHYS. ACTA 52 (1979) 88  
 B079D BOFFI AND CAPUZZI, INT. CONF. NUCL. PHYS. WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 1.3  
 B079EE BOOTH ET AL, PHYS. REV. C20 (1979) 1603  
 B079G BONCHE, DOUBRE AND QUENTIN, PHYS. LETT. B82 (1979) 5  
 B079GG BOGATIN, LOZHGIN AND YAKOVLEV, NUCL. PHYS. A326 (1979) 508  
 B079J BOOTH, IN 1978 RPI SYMPOSIUM "PHOTOPION NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 129  
 B079K BOUSSY, PHYS. LETT. B84 (1979) 41  
 B079R BOLGER ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 1E19  
 B079Y BOHIGAS, LANE AND MORTORELL, PHYS. REPORTS 51 (1979) 267  
 B079Z BONCHE ET AL, PHYS. REV. C20 (1979) 641  
 B080A BOSTED ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 559  
 B080B BOHANNON, ZAMICK AND MOYA DE GUERRA, NUCL. PHYS. A334 (1980) 278  
 B080F BOUSSY, PHYS. LETT. 91B (1980) 15  
 B080L BORTIGNON, P. 183, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 B080N BOHIGAS, P. 116, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 B080O BOHIGAS, VAN GIAI AND VAUTHERIN, P. 225, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 B080Q BOSTED ET AL, PHYS. REV. LETT. 45 (1980) 1544  
 B081 BOWLES ET AL, PHYS. REV. C 23 (1981) 439  
 B081A BOUDARD ET AL, PHYS. REV. LETT. 46 (1981) 218  
 B081H BOFFI, GIUSTI AND PACATI, NUCL. PHYS. A359 (1981) 91  
 B081J BOUSSY, PHYS. LETT. 99B (1981) 373  
 B081K BOAL, GREEN, KORTELING AND SOROUSHIAN, PHYS. REV. C23 (1981) 2788  
 B081M BOFFI, GIUSTI AND PACATI, PROCEEDINGS VERSAILLES CONF. (1981) p. 116  
 B081P BOFFI, GIUSTI AND PACATI, NUCL. PHYS. A358 (1981) C271

- BO81R BOHIGAS, VAN GIAI AND VAUTHERIN, PHYS. LETT. 102B (1981) 105  
 BR64F BROWNE AND MICHAEL, PHYS. REV. 134 (1964) B133  
 BR73M BROUDE ET AL, NUCLEAR PHYSICS A215 (1973) 617  
 BR75S BRINK ET AL, IN JUL-CONF-16 (1975), P. 11  
 BR76D BROCHARD ET AL, PHYS. REV. C13 (1976) 967  
 BR76Q BRUCKNER ET AL, PHYS. LETT. B62 (1976) 481  
 BR77 BROWN, ARIMA AND MC GRORY, NUCLEAR PHYSICS A277 (1977) 77  
 BR77A BROCKMANN AND DILLIG, PHYS. REV. C15 (1977) 361  
 BR77C BRINK AND TAKIGAWA, NUCLEAR PHYSICS A279 (1977) 159  
 BR77E BRAUN-MUNZINGER ET AL, PHYS. REV. LETT. 38 (1977) 944  
 BR77GG BRISSAUD, J. OF PHYS. G3 (1977) 1497  
 BR77K BROCKMANN AND WEISE, PHYS. LETT. B69 (1977) 167  
 BR77Q BROCKMANN AND WEISE, PHYS. REV. C16 (1977) 1282  
 BR77S BRAY, FRAWLEY, OPHEL AND BARKER, NUCLEAR PHYSICS A288  
     (1977) 334  
 BR77W BRANFORD, NAGORCKA AND NEWTON, J. OF PHYS. G3 (1977) 1565  
 BR77Z BRUGE, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE,"  
     ZURICH 1977, BIRKHAUSER VERLAG, P. 69  
 BR78A BRAITHWAITE ET AL, BULL. AM. PHYS. SOC. 23 (1978) 48  
 BR78C BRENNER AND JACKSON, IN PROCEEDINGS, CLUSTER, WINNIPEG  
     1978, D2  
 BR78D BROWN, CHUNG AND WILDENTHAL, PHYS. REV. LETT. 40 (1978) 1631  
 BR78J BRINK AND STANCU, NUCLEAR PHYSICS A299 (1978) 321  
 BR78K BRIEVA AND ROOK, NUCLEAR PHYSICS A297 (1978) 206  
 BR78M BROCKMANN, PHYS. REV. C18 (1978) 1510  
 BR78Q BRIEVA AND ROOK, NUCLEAR PHYSICS A307 (1978) 493  
 BR79A BRADLOW ET AL, NUCLEAR PHYSICS A314 (1979) 207  
 BR79B BRADLOW ET AL, NUCLEAR PHYSICS A314 (1979) 171  
 BR79E BRAUN-MUNZINGER ET AL, BULL. AM. PHYS. SOC. 24 (1979)  
     571  
 BR79H BROWN, JENNINGS AND ROCKMORE, IN HOUSTON, AIP CONF.  
     PROCEEDINGS 54 (1979), P. 614  
 BR79I BROWN, MASSEN AND HODGSON, PHYS. LETT. B85 (1979) 167  
 BR79J BROWN, MASSEN AND HODGSON, J. OF PHYS. G 5 (1979) 1655  
 BR80 BREUNLICH, NUCL. PHYS. A335 (1980) 137  
 BR80A BROWN AND WILDENTHAL  
 BR80I BROWN, CHUNG AND WILDENTHAL, PHYS. REV. C22 (1980) 774  
 BR80J BREUER ET AL, PHYS. LETT. B96 (1980) 35  
 BR80N BROWN, RICHTER AND GUDWIN, PHYS. REV. LETT. 45 (1980) 1681  
 BR80Q BRAGIN ET AL, SOVIET J. NUCL. PHYS. 31 (1980) 14  
 BR81B BRANDAN AND MENCHACA-ROCHA, PHYS. REV. C23 (1981) 1272  
 BR81D BROCKMANN AND WEISE, PROCEEDINGS VERSAILLES CONF. (1981)  
     p. 376  
 BU75D BUENERD, ISN 75-24 (1975)  
 BU76E BUENERD, MARTIN, DE SAINTIGNON AND LOISEAUX, PHYS. REV.  
     C14 (1976) 1316  
 BU76I BURGOV ET AL, JETP LETT. 23 (1976) 272  
 BU76J BUKI, SHEVCHENKO AND MITROFANOVA, SOVIET J. NUCL. PHYS.  
     24 (1976) 237  
 BU76K BURTEBAEV ET AL, SOVIET J. NUCL. PHYS. 24 (1976) 457  
 BU77 BUCK, FRIEDRICH AND WHEATLEY, NUCLEAR PHYSICS A275  
     (1977) 246  
 BU77A BUENERD ET AL, J. PHYSIQUE LETT. 38 (1977) L-53  
 BU77L BUSSO, GALLINO AND MASANI, ASTROPHYS. AND SPACE SCI. 52  
     (1977) 479

- BU77S BURSAK ET AL, SOVIET J. NUCL. PHYS. 26 (1977) 260  
 BU78E BURMAN ET AL, PHYS. REV. C17 (1978) 1774  
 BU78G BUTLER, NEWMAN AND TALBOT, SCIENCE 201 (1978) 522  
 BU78J BUDZANOWSKI ET AL, NUOVO CIM. A48 (1978) 100  
 BU78K BUFFINGTON, ORTH AND MAST, ASTROPHYS. J. 226 (1978) 355  
 BU79 BUCHE ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
 PAPER 1E1  
 BU79E BUGAEV, BISNOVATYI-KOGAN, RUDZSKY AND SEIDOV, NUCL.  
 PHYS. A324 (1979) 350  
 BU79J Burov and Lukyanov, NUCLEAR INTERACTION, CANBERRA,  
 AUSTRALIA (SPRINGER-VERLAG 1979) P. 385, AND PHYS.  
 ABSTRACTS 9260 (1980)  
 BU80A BUTI ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 574  
 BU80B BURLESON ET AL, PHYS. REV. C22 (1980) 1180  
 BU81A BUTI ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 590  
 BU81C Burov, Dostovalov, Kaschiev and Shitikova, J. OF PHYS. G7  
 (1981) 137  
 BU81E BUCKLEY ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 340  
 BY77 BYRD, LISOWSKI, TORNOW AND WALTER, BULL. AM. PHYS. SOC.  
 22 (1977) 587  
 BY78A BYRNAK, NUOVO CIM. A44 (1978) 1  
 BY79A BYRD, BULL. AM. PHYS. SOC. 24 (1979) 814  
 BY80 BYRD ET AL, BULL. AM. PHYS. SOC. 25 (1980) 592  
 BY81 BYRD ET AL, NUCL. PHYS. A351 (1981) 189  
 BY81B BYRD AND WALTER, IN POLARIZATION, SANTA FE 1980, AIP  
 CONFERENCE PROC. 69 (1981) p. 1475  
 CA74L CARACCIOLI ET AL, LETT. NUOVO CIM. 11 (1974) 33  
 CA76R CALVI ET AL, LETT. NUOVO CIM. 17 (1976) 467  
 CA76S CASANOVA AND CASANOVA, AN. FIS. (SPAIN) 72 (1976) 174  
 CA77 CATARA ET AL, NUCLEAR PHYSICS A276 (1977) 433  
 CA77B CALVI ET AL, LETT. NUOVO CIM. 18 (1977) 77  
 CA77C CAURIER AND GRAMMATICOS, NUCLEAR PHYSICS A279 (1977) 333  
 CA77D CANTO, NUCLEAR PHYSICS A279 (1977) 97  
 CA77J CARTER ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 188  
 CA77L CALARCO ET AL, PHYS. REV. LETT. 39 (1977) 925  
 CA77N CASSE AND GORET, OAP-499 (1977)  
 CA77Q CAUGHLAN, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D.  
 REIDEL, HOLLAND, 1977), P. 121  
 CA77U CATARA, CONSOLI AND INSOLIA, NUOVO CIM. A41 (1977) 648  
 CA78F CARTER, CLARKSON, HNIZDO AND SELLSCHOP, NUCLEAR PHYSICS  
 A297 (1978) 520  
 CA79E CAMERON ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
 PAPER 4A15  
 CA79J CALARCO, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108  
 (1979) P. 114  
 CA80E CASTEL, CUSSON AND ZAMICK, NUCL. PHYS. A341 (1980) 93  
 CA80H CALARCO, BULL. AMER. PHYS. SOC. 25 (1980) 737  
 CA80K CANAL, ISERN AND LABAY, ASTROPHYS. J. 241 (1980) L33  
 CA80M CARLOS, J. PHYS. COLLOQ. 41 (1980) C3/149  
 CA81 CAREY, ROOS, CHANT, NADASEN AND CHEN, PHYS. REV. C 23  
 (1981) 576  
 CA81B CARDMAN, NUCL. PHYS. A354 (1981) 173c  
 CA81C CARTER ET AL, S. AFRICAN J. OF PHYS. 4 (1981) 13  
 CA81D CAVINATO, MARAGNONI AND SARUIS, PROCEEDINGS VERSAILLES CONF.  
 (1981) p. 81

- CE64B CERNY, PEHL AND GARVEY, PHYS. LETT. 12 (1964) 234  
 CE75A CEULENEER ET AL, PHYS. REV. C11 (1975) 631  
 CE77 CELENZA, HUFNER AND SANDER, NUCLEAR PHYSICS A276 (1977)  
 509  
 CE77A CECIL, SPARROW, SHEPARD AND BLOK, PHYS. LETT. B66 (1977) 127  
 CE77C CERNY AND HARDY, ANN. REV. NUCL. SCI. 27 (1977) 333  
 CE80 CECIL ET AL, PHYS. REV. C21 (1980) 2471  
 CE80A CERRUTI ET AL, P. 583, PROCEEDINGS OF THE INTERNATIONAL  
 CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 CH61A CHIBA, PHYS. REV. 123 (1961) 1316  
 CH76 CHANG, DIENER AND VENTURA, NUCLEAR PHYSICS A258 (1976) 91  
 CH76N CHEVARIER ET AL, NUCLEAR PHYSICS A274 (1976) 223  
 CH76P CHUA, BECCHETTI, JANECKE AND MILDNER, NUCLEAR PHYSICS  
 A273 (1976) 243  
 CH76R CHYI, BULL. AM. PHYS. SOC. 21 (1976) 1331  
 CH76S CHERDANTSEV, IN PROC. INTERN. SCHOOL-SEMINAR ON  
 REACTIONS OF HEAVY IONS WITH NUCLEI AND SYNTHESIS OF  
 NEW ELEMENTS (DUBNA, 1975), P. 181, DUBNA D7-9734 (1976)  
 CH76U CHATTERJEE, TOBIAS AND LYMAN, IN SPALLATION NUCLEAR  
 REACTIONS AND THEIR APPLICATIONS, EDS. SHEN AND MERKER  
 (D. REIDEL PUB. CO., THE NETHERLANDS, 1976), P. 169  
 CH77A CHANT AND ROOS, PHYS. REV. C15 (1977) 57  
 CH77B CHEW, NUCLEAR PHYSICS A283 (1977) 445  
 CH77E CHAIT AND SINCLAIR, NUCLEAR PHYSICS A279 (1977) 517  
 CH77G CHRISTENSEN AND SWITKOWSKI, NUCLEAR PHYSICS A280 (1977)  
 205  
 CH77J CHEW, LOWE, NELSON AND BARNETT, NUCLEAR PHYSICS A286  
 (1977) 451  
 CH77M CHECHIK, FRAENKEL, STOCKER AND EYAL, NUCLEAR PHYSICS A287  
 (1977) 353  
 CH77Q CHACHKUNASHVILI, KOPALEISHVILI AND MACHABELI, IN PROC.,  
 ZURICH - SIN, 1977, P. 25  
 CH78B CHANT, ROOS AND WANG, PHYS. REV. C17 (1978) 8  
 CH78E CHEUNG, HIGH AND CUJEC, NUCLEAR PHYSICS A296 (1978) 333  
 CH78F CHECHIK, EYAL, STOCKER AND FRAENKEL, NUCLEAR PHYSICS A296  
 (1978) 307  
 CH78K CHEW, LOWE, NELSON AND BARNETT, NUCLEAR PHYSICS A298  
 (1978) 19  
 CH78L CHAN ET AL, NUCLEAR PHYSICS A303 (1978) 500  
 CH78M CHECHIK, FRAENKEL, EYAL AND STOCKER, NUCLEAR PHYSICS A304  
 (1978) 243  
 CH78O CHANDRA AND MOSEL, NUCLEAR PHYSICS A298 (1978) 151  
 CH78P CHEW AND LOWE, NUCLEAR PHYSICS A306 (1978) 125  
 CH78R CHEN ET AL, BULL. AM. PHYS. SOC. 23 (1978) 953  
 CH78S CHANT, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47  
 (1978) 415  
 CH78T CHAUDHRI AND REEN, BULL. AM. PHYS. SOC. 23 (1978) 1062  
 CH78Y CHAUMEAUX, LAYLY AND SCHAEFFER, ANN. OF PHYS.  
 116 (1978) 247  
 CH78Z CHUNG, VAN HIENEN, WILDENTHAL AND BENNETT, PHYS. LETT. B79  
 (1978) 381  
 CH79A CHALMERS AND LONGLEY, BULL. AM. PHYS. SOC. 24 (1979) 62  
 CH79C CHAN ET AL, PHYS. REV. LETT. 42 (1979) 687, AND PHYS.  
 REV. LETT. 43 (1979) 1625

- CH79F CHRISTIANSSON, DUBOIS AND ROTH, PHYS. SCRIPTA 19 (1979) 245  
 CH79H CHEW ET AL, INT. CONF. NUCL. PHYS. WITH ELECTROMAGNETIC  
 INTERACTIONS, MAINZ (1979), 5.1  
 CH79I CHANCE AND HARRIS, ASTRON. & ASTROPHYS. 74 (1979) 247  
 CH79J CHAN AND SHELDON, BULL. AM. PHYS. SOC. 24 (1979) 760  
 CH79K CHENG ET AL, NUCLEAR PHYSICS A322 (1979) 168  
 CH79P CHANT, RUOS AND KITCHING, IN CONF. PROC. TRIUMF, VANCOUVER  
 (1979), PAPER 4C26  
 CH79U CHIANG AND HUFNER, PHYS. LETT. B84 (1979) 393  
 CH80B CHENG, LURAZO AND GOULARD, PHYS. REV. C21 (1980) 374  
 CH80C CHAPURAN ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 591  
 CH80H CHEW ET AL, P. 224, PROCEEDINGS OF THE INTERNATIONAL  
 CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 CH80M CHRISTILLIN, ROSA-CLOT AND SERVADIO, NUCL. PHYS. A345  
 (1980) 317  
 CH80N CHRISTILLIN, LIPPARINI, STRINGARI AND TRAINI, PHYS.  
 LETT. B95 (1980) 344  
 CH80O CHUNG AND SHIN, NUOVO CIM. A60 (1980) 27  
 CH81 CHENG AND GOULARD, PHYS. REV. C23 (1981) 869  
 CH81B CHEN ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 579  
 CH81C CHAN ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 554  
 CH81E CHRIEN ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 384  
 CH81G CHERKASOV, IZV. AKAD. NAUK SSSR, SER. FIZ. 45 (1981) 138  
 CI76 CIERJACKS, IN LOWELL CONF., CONF-760715-P2 (1976), P. 893  
 CI78 CINDRO ET AL, FIZIKA (YUGOSLAVIA) 10, SUPPL. (1978) 1,  
 AND PHYS. ABSTRACTS 74384 (1978)  
 CI80 CIERJACKS ET AL, NUCL. INSTR. METH. 169 (1980) 185  
 CI81 CIRIT AND YAZICI, PHYS. REV. C23 (1981) 2627  
 CL77B CLOVER AND ZOHNI, BULL. AM. PHYS. SOC. 22 (1977) 644  
 CL77C CLAYTON, DWEK AND WOOSLEY, ASTROPHYS. J. 214 (1977) 300  
 CL77I CLAYTON, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED.  
 (D. REIDEL, HOLLAND, 1977), P. 15  
 CL78F CLARK, KEMPER AND FOX, PHYS. REV. C18 (1978) 1262  
 CL78G CLAYTON, ANN. REV. NUCL. & PART. SCI. 28 (1978) 501  
 CL79D CLARK, JOHNSTON AND OPHEL, AUSTRALIAN J. PHYS. 32  
 (1979) 283  
 CL80B CLEMENT ET AL, POLARIZATION - SANTA FE (1980) 2.65  
 CL80C CLAYTON, NUKLEONIKA 25 (1980) 1477  
 C073U COLE ET AL, NUCLEAR PHYSICS A213 (1973) 107  
 C075P COELHO AND DAS, REV. BRAS. FIS. (BRAZIL) 5 (1975) 335  
 C076C COBERN, PISANO AND PARKER, PHYS. REV. C14 (1976) 491  
 C076U COTANCH AND VINCENT, PHYS. REV. C14 (1976) 1739  
 C077A COBERN AND FORTUNE, BULL. AM. PHYS. SOC. 22 (1977) 28  
 C077H COWAN AND ROSE, ASTROPHYS. J. 212 (1977) 149  
 C077M CORMIER ET AL, PHYS. REV. C16 (1977) 215  
 C077P CONJEAUD ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 663  
 C077R COLE, LONGQUEUE AND CAVAIGNAC, J. PHYSIQUE 38 (1977) 1043  
 C077V COOPER, JOHNSON AND WEST, NUCLEAR PHYSICS A292 (1977) 350  
 C077Z COWAN AND ROSE, ASTROPHYS. J. 217 (1977) 51  
 C078E CORMIER ET AL, PHYS. REV. LETT. 40 (1978) 924  
 C078H COTANCH, PHYS. LETT. B76 (1978) 19  
 C078O COTANCH, PHYS. REV. C18 (1978) 1941  
 C078P COTANCH, NUCLEAR PHYSICS A308 (1978) 253  
 C079C CONZE AND MANAKOS, J. OF PHYS. G5 (1979) 671

- C0790 COWLEY, MILLS AND HEYMANN, S. AFRICAN J. PHYS. 2 (1979) 85  
 C080A COTTINGAME AND HOLT Kamp, BULL. AMER. PHYS. SOC. 25  
       (1980) 503  
 C080Q COWAN, CAMERON AND TRURAN, ASTROPHYS. J. 241 (1980) 1090  
 C080S COOK, STONE AND VOGL, ASTROPHYS. J. 238 (1980) L97  
 C080T COLE ET AL, NUCL. PHYS. A341 (1980) 284  
 C081 CONZETT, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC.  
       69 (1981) p. 1195  
 C081G CORNELIUS, MOSS AND YAMAYA, PHYS. REV. C23 (1981) 1364  
 C081M COUVERT AND DILLIG, PROCEEDINGS VERSAILLES CONF. (1981)  
       p. 119  
 CR75A CROZIER AND FORTUNE, PHYS. REV. C11 (1975) 308  
 CR75E CRINEAN, HUDSON, WILD AND SPICER, NUCLEAR PHYSICS A244  
       (1975) 77  
 CR76D CRAMER ET AL, PHYSICAL REVIEW C14 (1976) 2158  
 CR79B CROWLEY, ANN. OF PHYS. 117 (1979) 323  
 CR79C CRAWFORD ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
       (1979), PAPER 4A1  
 CR80 CRAMER AND DE VRIES, PHYS. REV. C22 (1980) 91  
 CR80A CRAWFORD ET AL, PHYS. REV. C22 (1980) 1184  
 CU76F CUSSON, HILKO AND KOLB, NUCLEAR PHYSICS A270 (1976) 437  
 CU77 CUNSOLO ET AL, NUOVO CIM. A40 (1977) 293  
 CU78B CUZZOCREA ET AL, LETT. NUOVO CIM. 22 (1978) 257  
 CU78C CUNSOLO ET AL, PHYS. REV. C18 (1978) 856  
 CU78E CUSSON, GOMEZ DEL CAMPO AND MELDNER, IN CLUSTERING,  
       WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 35  
 CU78F CUSSON, MARUHN AND MELDNER, PHYS. REV. C18 (1978) 2589  
 CU78G CUNSOLO ET AL, LETT. NUOVO CIM. 23 (1978) 371  
 CU80 CUJEC, WU AND BARNES, LAP - 166 (1979)  
 CU80A CUNSOLO ET AL, PHYS. REV. C21 (1980) 952  
 CU80B CUNSOLO ET AL, PHYS. REV. C21 (1980) 2345  
 CU80D CUZZOCREA ET AL, LETT. NUOVO CIM. 28 (1980) 515  
 CU81B CUSSON AND CASTEL, PHYS. REV. C23 (1981) 1279  
 CU81F CUSSON AND CASTEL, PROCEEDINGS VERSAILLES CONF. (1981) p. 547  
 DA68 DAVIDS, NUCLEAR PHYSICS A110 (1968) 619  
 DA74I DAHLGREN, GRAFSTROM, HOISTAD AND ASBERG, NUCLEAR PHYSICS  
       A227 (1974) 24  
 DA76N DAVISON, FALK, SOURKES AND VOSNIAKOS, BULL. AM. PHYS.  
       SOC. 21 (1976) 1295  
 DA76O DALITZ AND GAL, PHYS. LETT. B64 (1976) 154  
 DA76P DALKHAZHAV ET AL, SOVIET J. NUCL. PHYS. 23 (1976) 643  
 DA76Q DADAYAN, SOVIET J. NUCL. PHYS. 24 (1976) 216  
 DA77E DAVYDOV, NAUMOV AND SHCHERBACHENKO, INDC(CCP)-104/LN  
       (1977)  
 DA77H DARWISCH, GRIDNEV, HEFTER AND SEMJONOV, NUOVO CIM. A42  
       (1977) 303  
 DA78N DAREMA-ROGERS AND TRUE, PHYS. REV. C18 (1978) 2355  
 DA78P DAVIES, FELDMEIER, FLOCARD AND WEISS, PHYS. REV. C18  
       (1978) 2631  
 DA78S DALITZ AND GAL, ANN. OF PHYS. 116 (1978) 167  
 DA79F DARDEN AND FOSTER, PHYS. REV. C19 (1979) 2396  
 DA80B DAVIES, FLOCARD, KRIEGER AND WEISS, NUCL. PHYS. A342 (1980)  
       111  
 DA81C DALITZ, NUCL. PHYS. A354 (1981) 101c  
 DE68G DENISOV, KOMAR AND KULCHITSKY, NUCLEAR PHYSICS A113  
       (1968) 289

- DE69C DETRAZ AND DUHM, PHYS. LETT. 29B (1969) 29  
 DE70E DETRAZ, DUHM AND HAFNER, NUCLEAR PHYSICS A147 (1970) 488  
 DE76L DEDONDER AND SCHMIT, PHYS. LETT. B65 (1976) 131  
 DE77D DEHESA, KREWALD, SPETH AND FAESSLER, PHYS. REV. C15  
     (1977) 1858  
 DE77F DELLAFFIORE, NUCLEAR PHYSICS A282 (1977) 493  
 DE77P DEARBORN, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D.  
     REIDEL, HOLLAND, 1977), P. 39  
 DE78G DEHNHARD, SHKOLNIK AND FRANEY, PHYS. REV. LETT. 40  
     (1978) 1549, AND IBID. 42 (1979) 1574  
 DE78H DESPLANQUES AND MISSIMER, NUCLEAR PHYSICS A300 (1978)  
     286  
 DE78N DEY AND ANSARI, PRAMANA 10 (1978) 213, AND PHYS.  
     ABSTRACTS 62491 (1978)  
 DE78S DEMENTII, OGURTSOV AND AFANAS'EV, YADERNAYA FIZ. 28  
     (1978) 865  
 DE79F DEUTCHMAN, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979)  
     P. 363  
 DE79K DEVANATHAN, GIRIJA AND PRASAD, IN CONF. PROC. TRIUMF,  
     VANCOUVER (1979), PAPER 5A12  
 DE79M DEINEKO ET AL, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43  
     (1979) 155  
 DE79N DE BOER, ASTROPHYS. J. 229 (1979) 132  
 DE79O DESPLANQUES AND MISSIMER, PHYS. LETT. B84 (1979) 363  
 DE79T DE VRIES, NUKLEONIKA 24 (1979) 75, 87  
 DE79W DEVRIES, IN BNL-51115 (1979), P. 513  
 DE79Z DEVRIES AND PENG, PHYS. REV. LETT. 43 (1979) 1373  
 DE80 DESPLANQUES, NUCL. PHYS. A335 (1980) 147  
 DE80AA DEUTCHMAN AND TOWNSEND, PHYS. REV. LETT. 45 (1980) 1622  
 DE80B DE VRIES ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 506  
 DE80BB DEVANATHAN, GIRIJA AND SIVARAMA PRASAD, CAN. J. PHYS.  
     58 (1980) 1151  
 DE80D DENNIS, FRAWLEY, KEMPER AND SZYMAKOWSKI, BULL. AMER.  
     PHYS. SOC. 25 (1980) 523  
 DE80FF DE SWINIARSKI AND DINH-LIEN PHAM, Z. PHYS. A296 (1980) 367  
 DE80GG DEMENTII, OGURTSOV, OMELAENKO AND AFANAS'EV, UKR. FIZ. ZH. 25  
     (1980) 628  
 DE80L DEVRIES AND PENG, PHYS. REV. C22 (1980) 1055  
 DE80M DEDONDER, MAILLET AND SCHMIT, ANN. OF PHYS. 127 (1980) 1  
 DE80P DEHNHARD, BULL. AMER. PHYS. SOC. 25 (1980) 732  
 DE80V DE LEO ET AL, P. 203, PROCEEDINGS OF THE INTERNATIONAL  
     CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 DE80W DEVRIES, DIGIACOMO AND PENG, P. 418, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 DE80Z DELLAFFIORE AND TRAINI, NUCL. PHYS. A344 (1980) 509  
 DE81G DETHIER AND STANCU, PHYS. REV. C23 (1981) 1503  
 DE81H DENNIS ET AL, NUCL. PHYS. A357 (1981) 521  
 DE81I DENNIS, ROY, FRAWLEY AND KEMPER, NUCL. PHYS. A359 (1981) 455  
 DE81K DEVI, DHAR AND STRAYER, PHYS. REV. C23 (1981) 2062  
 DE81N DE SWINIARSKI AND PHAM, PROCEEDINGS VERSAILLES CONF. (1981)  
     p. 149  
 DE81P DEMENTIJ, FARTUSHNYJ AND AFANAS'EV, PROCEEDINGS VERSAILLES  
     CONF. (1981) p. 127  
 DE81R DE YOUNG ET AL, PHYS. REV. C24 (1981) 166  
 DH79 DHAR AND NILSSON, NUCLEAR PHYSICS A315 (1979) 445

- DI77E DICKMAN, LANGER, MCCUTCHEON AND SHUTER, IN CNO ISOTOPES IN ASTROPHYSICS, J. AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 95
- DI78D DIETRICH AND SIMPSON, ASTROPHYS. J. 225 (1978) L41
- DI78F DI NAPOLI ET AL, J. INORG. & NUCL. CHEM. (GB) 40 (1978) 1619, AND PHYS. ABSTRACTS 22268 (1979)
- DI79B DILLIG ET AL, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 306
- DI80 DIEUMEGARD, MAUREL AND AMSEL, NUCL. INSTR. METH. 168 (1980) 93
- DI80A DIGREGORIO ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 523
- DM77 DMITRIEVA, LEBEDEV AND MATUSEVICH, IZV. AKAD. NAUK SSSR (SER. FIZ.) 41 (1977) 143
- DM80 DMITRIEV, KOSMACH, MOLCHANOV AND OSTROUMOV, SOVIET J. NUCL. PHYS. 31 (1980) 62
- DO56B DOUGLAS, GASTEN AND MUKERJI, CAN. J. PHYS. 34 (1956) 1097
- DO71A DOVER ET AL, NUCLEAR PHYSICS A166 (1971) 461
- DO77A DOGOTAR, ERAMZYAN, KISSENER AND SAKAEV, NUCL. PHYS. A282 (1977) 474
- DO78 DOBBRE ET AL, PHYS. REV. C17 (1978) 131
- DO79 DOLL ET AL, PHYS. REV. LETT. 42 (1979) 366
- DO79E DOLLHOPF, PERDRISAT, KITCHING AND OLSEN, NUCLEAR PHYSICS A316 (1979) 350
- DO79O DOGOTAR ET AL, J. OF PHYS. G 5 (1979) L221
- DO80D DO DANG AND RAMAVATARAM, J. PHYS. G 6 (1980) 47
- DO80E DOVER AND GAL, P. 795, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- DO80G DOVER, LUDEKING AND WALKER, PHYS. REV. C 22 (1980) 2073
- DO81 DOZIER AND CHALMERS, PHYS. REV. C 23 (1981) 399
- DO81A DOZIER AND CHALMERS, PHYS. REV. C 23 (1981) 563
- DR71B DRACOULIS AND LEGGE, NUCLEAR PHYSICS A177 (1971) 241
- DR80D DRAGUN AND UBERALL, PHYS. LETT. 94B (1980) 24
- DR81B DRAKE ET AL, PHYS. LETT. 100B (1981) 236
- DU70 DURAY AND BROWNE, PHYS. REV. C1 (1970) 776
- DU74C DUHAMEL ET AL, NUCLEAR PHYSICS A231 (1974) 349
- DU76B DURELL ET AL, NUCLEAR PHYSICS A269 (1976) 443
- DU77B DURELL, HARTER AND PHILLIPS, PHYS. LETT. B70 (1977) 405
- DU78A DUDEK-ELLIS ET AL, PHYS. REV. C18 (1978) 158
- DU78C DUDEK-ELLIS, SHKOLNIK AND DEHNHARD, PHYS. REV. C18 (1978) 1039
- DU78H DUBNA-WARSAW-LENINGRAD COLLABORATION, YADERNAYA FIZ. 27 (1978) 1246
- DU79A DUTTA, BHADURI, SRIVASTAVA AND VALLIERES, PHYS. LETT. B84 (1979) 17
- DU80B DUPLAINE AND GOULARD, CAN. J. PHYS. 58 (1980) 1099
- DU80D DUBOVOI, ISZV. AKAD. NAUK, SER. FIZ. 44 (1980) 2341, 2434
- DU81A DUMBRAJS AND MARTINIS, PROCEEDINGS VERSAILLES CONF. (1981) p. 240
- DU81B DUBOVIK AND TOSUNYAN, PROCEEDINGS VERSAILLES CONF. (1981) p. 471
- DU81C DUNPHY ET AL, ASTROPHYS. J. 244 (1981) 1081
- DW80 DWORZECKA, KAN AND GRIFFIN, BULL. AMER. PHYS. SOC. 25 (1980) 574
- DY74 DYER AND BARNES, NUCLEAR PHYSICS A233 (1974) 495

- DY77C DYUMIN AND LEBEDEV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 41 (1977) 2025
- DY79 DYER ET AL, PHYS. REV. LETT. 42 (1979) 560
- DY79A DYTMAN ET AL, PHYS. REV. C19 (1979) 971
- DY81 DYER ET AL, PHYS. REV. C23 (1981) 1865
- DY81A DYMARZ, MOLINA AND SHITIKOVA, Z. PHYS. A299 (1981) 245
- EC77 ECK, GRAY AND GARDNER, PHYS. REV. C16 (1977) 1873
- EC81 ECK ET AL, PHYS. REV. C23 (1981) 2068
- EG81B EGELHAAF ET AL, PHYS. REV. LETT. 46 (1981) 813
- EI77 EISEN ET AL, NUCLEAR PHYSICS A291 (1977) 459
- EI81 EISENBERG, NUCL. PHYS. A355 (1981) 269
- EL78A EL NAGGAR AND EL AZAB, J. OF PHYS. G4 (1978) 269
- EL81C EL-NADI ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 456
- EM78 EMILIANI, HUDSON, SHINN AND GEORGE, SCIENCE 202 (1978) 627
- EM81 EMRICH, NUCL. PHYS. A351 (1981) 397
- EM81A EMRICH AND ZABOLITZKY, NUCL. PHYS. A351 (1981) 439
- EN76 ENGELHARDT, LEWIS AND ULLRICH, NUCLEAR PHYSICS A258 (1976) 480
- EN77 ENGLAND ET AL, NUCLEAR PHYSICS A284 (1977) 29
- EN77A ENCRENAZ, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 79
- EN78C ENGELBRECHT, HAHNE AND HEISS, SOUTH AFRICAN J. PHYS. 1 (1978) 41
- EN79A ENDT, ATOMIC & NUCL. DATA TABLES 23 (1979) 3
- EP77A EPSTEIN, THOMPSON AND YAPP, SCIENCE 198 (1977) 1209
- EP79A EPONESHNIKOV AND KRECHETOV, PIS'MA V ZH. EKSP. & TEOR. FIZ. 29 (1979) 442
- ER75B EROKHIN, ORLOVA, SMIRNOV AND SHITIKOVA, SOVIET J. NUCL. PHYS. 22 (1975) 574
- ER77D ERAMZYAN, GMITRO, SAKAEV AND TOSUNJAN, NUCLEAR PHYSICS A290 (1977) 294
- ER78B ERICSON AND MYHRER, PHYS. LETT. B74 (1978) 163
- ER78C ERAMZYAN, GMITRO AND TOSUNJAN, J. OF PHYS. G4 (1978) L233
- ER79 ERNST, PHYS. REV. C19 (1979) 896
- ER79C ERAMZYAN, GMITRO AND TOSUNJAN, CZECH. J. PHYS. B29 (1979) 370
- ER80B ERAMZYAN, GMITRO AND KISSENER, NUCL. PHYS. A338 (1980) 436
- ER80C ERB, IN NUCLEI FAR FROM STABILITY, J. H. HAMILTON ET AL. EDS., NORTH-HOLLAND (1980) P. 311
- ER81A ERAMZYAN ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 503
- FA75I FAI, MAGY. FIZ. FOLY. (HUNGARY) 23 (1975) 393
- FA76F FAESSLER, IN PROC. DUBNA CONF. JINR D-9920, VOL. II, P. 242 (1976)
- FA77A FAYANS, SAPERSTEIN AND TOLOKONNIKOV, J. OF PHYS. G3 (1977) L51
- FA78D FARINE, PEARSON AND ROUBEN, NUCLEAR PHYSICS A304 (1978) 317
- FA79B FAESSLER, MILLENER, PAUL AND STROTTMAN, NUCL. PHYS. A330 (1979) 333
- FA80 FABRICI ET AL, PHYS. REV. C21 (1980) 830
- FA80A FABRICI ET AL, PHYS. REV. C21 (1980) 844
- FA80B FAESSLER, KUO AND MUTHER, Z. PHYS. A 294 (1980) 95
- FA80D FARHAN, UBERALL, DRAGUN AND MAQUEDA, P. 507, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)

- FA80E FAESSLER, MUTHER, SHIMIZU AND WADIA, NUCL. PHYS. A333 (1980) 428
- FE76E FEOFILOV, DENISOV, KOLALIS AND SADKOVSKII, IZV. AKAD. NAUK SSSR (SER. FIZ.) 40 (1976) 2210
- FE77E FESHBACH, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 291
- FE78B FERNANDEZ ET AL, NUCLEAR PHYSICS A306 (1978) 259
- FE78C FERNANDEZ DIAZ AND SIROTKIN, NUCLEAR PHYSICS A312 (1978) 17
- FE78E FELDMAN, ASTRON. & ASTROPHYS. 70 (1978) 547
- FE80A FENG ET AL, PHYS. REV. LETT. 44 (1980) 1037
- FI74F FICK, NUKLEONIKA 19 (1974) 693
- FI75E FIRK ET AL, NUCLEAR CROSS SECTIONS AND TECHNOLOGY, PT. II, WASHINGTON, D.C. 1975 (WASH., D.C., NAT. BUR. STANDARDS 1975), P. 875, AND PHYS. ABSTRACTS 23088 (1977)
- FI77B FINDLAY AND OWENS, NUCLEAR PHYSICS A279 (1977) 385
- FI78 FINDLAY ET AL, PHYS. LETT. B74 (1978) 305
- FI78A FICK, AIP CONF. PROC. 47 (1978) 326
- FI78C FINDLAY, GIBSON, OWENS AND MATTHEWS, PHYS. LETT. B79 (1978) 356
- FI79D FILIPOV ET AL, YADERNAYA FIZ. 29 (1979) 332
- FL74A FLYNN AND GARRETT, PHYS. REV. C10 (1974) 409
- FL76B FLIESSBACH, Z. PHYS. A278 (1976) 353
- FL77 FLECKNER AND MOSEL, NUCLEAR PHYSICS A277 (1977) 170
- FL78C FLOCARD, KOONIN AND WEISS, PHYS. REV. C17 (1978) 1682
- FL79 FLIESSBACH, NUCLEAR PHYSICS A315 (1979) 109
- FL80 FLOCARD, HEENEN AND VAUTHERIN, NUCL. PHYS. A339 (1980) 336
- FL80B FLODKVIST ET AL, P. 74, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- F073G FOWLER, JOHNSON AND FEEZEL, PHYS. REV. C8 (1973) 545
- F075I FORTUNE AND MEDSKER, PHYS. REV. C12 (1975) 1364
- F075J FORTUNE, MEDSKER, GARRETT AND BINGHAM, PHYS. REV. C12 (1975) 1723
- F076J FOWLER, IN SCIENTIFIC & INDUSTRIAL APPLICATIONS OF SMALL ACCELERATORS, FOURTH CONFERENCE - 1976 (INST. FOR ELEC. & ELECTRONIC ENGRS., INC.), P. 11
- F077B FONTES, PHYS. REV. C15 (1977) 2159
- F077D FORTUNE ET AL, PHYS. LETT. B70 (1977) 408
- F077G FORTUNE, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 99
- F078C FORTUNE, PHYS. REV. C17 (1978) 861
- F078D FORTUNE, COBERN AND MOORE, PHYS. REV. C17 (1978) 888
- F078F FORTUNE ET AL, PHYS. REV. LETT. 40 (1978) 1236
- F078G FORTUNE, J. OF PHYS. G4 (1978) L103
- F078L FORTUNE, MIDDLETON AND BILANIUK, J. OF PHYS. G4 (1978) L181
- F078O FORTUNE, BISHOP, MEDSKER AND WILDENTHAL, PHYS. REV. LETT. 41 (1978) 527
- F078R FORTUNE, MIDDLETON AND BILANIUK, PHYS. REV. C18 (1978) 1920
- F079 FORTUNE, BILANIUK, STEPHANS AND MIDDLETON, PHYS. REV. C19 (1979) 5
- F079F FORTUNE, RICHTER, SIEMSEN AND YNTEMA, PHYS. REV. C20 (1979) 648
- F079J FORTUNE ET AL, PHYS. REV. C20 (1979) 1228
- FR72B FREIESLEBEN AND WEIBEZAHN, NUCLEAR PHYSICS A184 (1972) 273
- FR76D FROHLICH ET AL, PHYS. LETT. B64 (1976) 408
- FR76E FRIAR, ANN. OF PHYS. 98 (1976) 490

- FR77B FRANCO AND TEKOU, PHYS. REV. C16 (1977) 658  
 FR77E FROBRICH, LIPPERHEIDE AND THOMA, ANN. OF PHYS.  
 104 (1977) 478  
 FR77F FRIEDRICH AND CANTO, NUCLEAR PHYSICS A291 (1977) 249  
 FR77G FREEDMAN, SCHRAMM AND TUBBS, ANN. REV. NUCL. SCI. 27  
 (1977) 167  
 FR78B FRAWLEY, BRAY AND OPHEL, NUCLEAR PHYSICS A294 (1978) 161  
 FR78C FRAWLEY, ROY, MATEJA AND FLETCHER, BULL. AM. PHYS. SOC.  
 23 (1978) 539  
 FR78F FREEMAN AND HAAS, PHYS. REV. LETT. 40 (1978) 927  
 FR78G FRANCO AND NUTT, PHYS. REV. C17 (1978) 1347  
 FR78I FRANCO AND VARMA, PHYS. REV. C18 (1978) 349  
 FR78J FRANEY, BAYMAN, LILLEY AND PHILLIPS, PHYS. REV. LETT. 41  
 (1978) 837  
 FR78N FRAHN AND REHM, PHYS. REPORTS 37C (1978) 1  
 FR78O FRAHN, S. AFRICAN J. PHYS. 1 (1978) 133  
 FR79A FRANEY, SHKOLNIK AND DEHNHARD, PHYS. LETT. B81 (1979) 132  
 FR79B FREEMAN, PIEL, WILSCHUT AND PAUL, BULL. AM. PHYS. SOC.  
 24 (1979) 665  
 FR79C FREEDMAN ET AL, PHYS. REV. C19 (1979) 1907  
 FR79G FRISCHKNECHT ET AL, IN CONF. PROC. TRIUMF, VANCOUVER  
 (1979), PAPER 5B11  
 FR79K FRIEDMAN, MCVOY AND NEMES, PHYS. LETT. B87 (1979) 179  
 FR80A FREEMAN ET AL, PHYS. REV. LETT. 45 (1980) 1479  
 FR80C FRAWLEY, ROY AND FLETCHER, PHYS. REV. LETT. 44 (1980) 1377  
 FR80E FRAHN, NUCL. PHYS. A337 (1980) 324, 351  
 FR80G FRAWLEY, FLETCHER, DENNIS AND ABDO, BULL. AMER. PHYS.  
 25 (1980) 727  
 FR80H FREEDMAN, MILLER AND HENLEY, BULL. AMER. PHYS. SOC.  
 25 (1980) 725  
 FR80J FRANZ AND HAMANN, P. 168, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 FR80K FRANZ ET AL, P. 890, PROCEEDINGS OF THE INTERNATIONAL  
 CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 FR81E FREEDMAN, MILLER AND HENLEY, PROCEEDINGS VERSAILLES CONF.  
 (1981) p. 422  
 FR81F FRIEDMAN, PROCEEDINGS VERSAILLES CONF. (1981) p. 349  
 FR81G FRIEDMAN, PHYS. REV. C24 (1981) 125  
 FR81H FRAWLEY, ROY, MATEJA AND FLETCHER, NUCL. PHYS. A363 (1981)  
 280  
 FU71A FUCHS, HOMEYER, LORENZ AND OESCHLER, PHYS. LETT. 37B  
 (1971) 285  
 FU72C FUCHS ET AL, NUCLEAR PHYSICS A196 (1972) 286  
 FU75C FUJII ET AL, J. PHYS. SOC. JAPAN 38 (1975) 1563  
 FU76G FUJITA, ISHIDA AND HIRATA, SUPPL. PROG. THEOR. PHYS. 60  
 (1976) 73  
 FU76H FULLER, J. PHYS. & CHEM. REF. DATA 5 (1976) 835  
 FU77L FULLER, PHYS. REV. C16 (1977) 1865  
 FU78C FURUI, NUCLEAR PHYSICS A300 (1978) 385  
 FU78G FURUI, NUCLEAR PHYSICS A312 (1978) 311  
 FU79G FURUNO ET AL, NUCLEAR PHYSICS A321 (1979) 250  
 FU80C FULTON, CORMIER AND HERMAN, P. 235, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 FU80E FUJITA AND HUFNER, NUCL. PHYS. A343 (1980) 493  
 FU80G FUJIWARA AND HORIUCHI, PROG. THEOR. PHYS. 63 (1980) 895

- FU80H FURUTANI ET AL, SUPPL. PROGR. THEOR. PHYS. 68 (1980) 193  
 FU80I FUJIWARA ET AL, SUPPL. PROGR. THEOR. PHYS. 68 (1980) 29  
 FU81B FULMER ET AL, PHYS. LETT. 100B (1981) 305  
 GA66D GALLMANN, FINTZ, NELSON AND ALBURGER, PHYS. REV. 147 (1966) 753  
 GA69E GALLMANN, JUNDT, ASLANIDES AND ALBURGER, PHYS. REV. 179 (1969) 921  
 GA70 GALLMANN ET AL, CAN. J. PHYS. 48 (1970) 1595  
 GA73N GALE AND ECK, PHYSICAL REVIEW C7 (1973) 1950  
 GA76F GARBER AND KINSEY, BNL 325 (1976)  
 GA76K GARI, HYUGA AND ZABOLITZKY, NUCLEAR PHYSICS A271 (1976) 365  
 GA77E GAREEV, GIZZATKULOV AND REVAI, NUCLEAR PHYSICS A286 (1977) 512  
 GA77G GADIOLI AND ERBA, NUCL. INSTR. METH. 146 (1977) 265  
 GA77H GABRIC AND AMOS, AUSTRALIAN J. PHYS. 30 (1977) 287  
 GA77L GALIN, GUERREAU AND BABINET, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 683  
 GA77N GASTEBOIS AND CHARLES, BULL. INF. SCI. & TECH. (FRANCE) 224-225 (1977) 141, AND PHYS. ABSTRACTS 3166 (1979)  
 GA78 GARCILAZO AND GIBBS, BULL. AM. PHYS. SOC. 23 (1978) 553  
 GA78B GARI AND HEBACH, PHYS. REV. C18 (1978) 1071  
 GA78C GAL, SOPER AND DALITZ, ANN. OF PHYS. 113 (1978) 79  
 GA78D GARI, HEBACH, SOMMER AND ZABOLITZKY, PHYS. REV. LETT. 41 (1978) 1288  
 GA78E GAI ET AL, BULL. AM. PHYS. SOC. 23 (1978) 933  
 GA78H GASTEBOIS, PROC. INT. CONF. ON RESONANCES IN HEAVY ION REACTIONS, HVAR, 1977 (NORTH-HOLLAND, AMSTERDAM, 1978), P. 61; PHYSICS ABSTRACTS 9616 (1979)  
 GA79 GAIDOS ET AL, PHYS. REV. LETT. 42 (1979) 82  
 GA79A GAI ET AL, BULL. AM. PHYS. SOC. 24 (1979) 570  
 GA79C GAZIS ET AL, BULL. AM. PHYS. SOC. 24 (1979) 695  
 GA79E GAL, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 680  
 GA79G GAI, BULL. AM. PHYS. SOC. 24 (1979) 843  
 GA79K GAL, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 485.  
 GA80 GAL AND DOVER, PHYS. REV. LETT. 44 (1980) 379  
 GA80B GAI, BULL. AMER. PHYS. SOC. 25 (1980) 592  
 GA80C GAI ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 592  
 GA80I GAI, P. 424, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 GA81B GARCIA-MUNOZ, GUZIK, SIMPSON AND WEFEL, BULL. AMER. PHYS. SOC. 26 (1981) 557  
 GA81C GAI, BULL. AMER. PHYS. SOC. 26 (1981) 612  
 GA81E GASPARIAN, PROCEEDINGS VERSAILLES CONF. (1981) p. 514  
 GE750 GERAMB ET AL, IN JUL-CONF-16 (1975), P. 6  
 GE78B GELBKE ET AL, PHYS. REPORTS 42C (1978) 311  
 GE78C GEESAMAN ET AL, PHYS. REV. C18 (1978) 284  
 GE78F GELBKE ET AL, PHYS. REV. LETT. 41 (1978) 1778  
 GE78I GEIGER AND VANDERZWAN, NUCL. INSTR. METH. 157 (1978) 199  
 GE79B GELBKE, IN BNL-51115 (1979), P. 1  
 GI63E GIVENS ET AL, NUCLEAR PHYSICS 46 (1963) 504  
 GI76E GIANNONI ET AL, PHYS. LETT. B65 (1976) 305  
 GI77A GIBBS, GIBSON AND STEPHENSON, PHYS. REV. LETT. 39 (1977) 1316  
 GI77C GISMATULLIN, LANTSEV AND OSTROUMOV, YADERNAYA FIZ. 26 (1977) 243

- GI78B GIBBS, GIBSON AND STEPHENSON, PHYS. REV. C18 (1978) 2782  
 GI78C GISMATULLIN, KOSMACH, MARTIROSOVA AND OSTROUMOV,  
*YADERNAYA FIZ.* 27 (1978) 37  
 GI79 GIBSON, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 691  
 GI79C GIBBS, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 595.  
 GI80A GIANNONI AND QUENTIN, PHYS. REV. C21 (1980) 2076  
 GI81B GIRIJA AND DEVANATHAN, PROCEEDINGS VERSAILLES CONF. (1981)  
 p. 120  
 GI81C GIFFON ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 500  
 GL77 GLENDINNING ET AL, BULL. AM. PHYS. SOC. 22 (1977) 631  
 GL78B GLOECKLER ET AL, BULL. AM. PHYS. SOC. 23 (1978) 510  
 GL79A GLOECKLER AND HOVESTADT, BULL. AM. PHYS. SOC. 24 (1979) 659  
 GL79C GLASHAUSSER ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
 PAPER 4A2  
 GL79E GLAGOLA, BREUER, GOKMEN AND VIOLA, BULL. AM. PHYS. SOC.  
 24 (1979) 846  
 GL79K GLASHAUSSER, ANN. REV. NUCL. & PART. SCI. 29 (1979) 33  
 GM77 GMITRO, TINKOVA, RIMINI AND WEBER, NUOVO CIM. LETT. 20  
 (1977) 449  
 GM79 GMITRO, TINKOVA, RIMINI AND WEBER, CZECH. J. PHYS.  
 B29 (1979) 155; IBID B29 (1979) 1308  
 GM79A GMITRO AND OVCHINNIKOVA, SOVIET J. NUCL. PHYS. 30 (1979) 171  
 G071T GOVE AND MARTIN, NUCLEAR DATA TABLES 10 (1971) 205  
 G075CC GOGNY AND MAIRE, PROC. INTERN. CONF. ON NUCLEAR  
 SELF-CONSISTENT FIELDS, ITALY, 1975 (AMSTERDAM -  
 NORTH-HOLLAND 1975), P. 149; PHYS. ABS. 82569 (1976)  
 G076K GOEKE, MOSZKOWSKI AND KREWALD, PHYS. LETT. B65 (1976) 113  
 G076N GOLOVANOVA ET AL, SOVIET J. NUCL. PHYS. 23 (1976) 33  
 G076Q GOLDBERG ET AL, HYPERFINE INTERACTIONS 2 (1976) 390  
 G077 GOEKE, PHYS. REV. LETT. 38 (1977) 212  
 G077N GOLUBEV, AVERYANOV AND SADOVOI, IZV. AKAD. NAUK SSSR  
 (SER. FIZ.) 41 (1977) 2096  
 G077P GONCHAROVA, MULEI AND SHITIKOVA, YADERNAYA FIZ. 26  
 (1977) 1183  
 G078C GOLDEN ET AL, APPL. PHYS. LETT. (TO BE PUBLISHED)  
 G078F GOLOVANOVA, IBRAEVA AND NEUDATCHIN, PROG. THEOR. PHYS.  
 59 (1978) 127  
 G078K GOLIN, PHYS. LETT. B74 (1978) 23  
 G078N GOLDBERG ET AL, HYPERFINE INTERACTIONS 4 (1978) 262  
 G078P GOLIKOV, ZHUKOV, LOSHKAKOV AND OSTROUMOV, SOVIET J. NUCL.  
 PHYS. 27 (1978) 3  
 G079C GOEKE AND CASTEL, PHYS. REV. C19 (1979) 201  
 G079D GORGEN ET AL, NUCLEAR PHYSICS A320 (1979) 296  
 G079I GOMEZ DEL CAMPO ET AL, PHYS. REV. LETT. 43 (1979) 26  
 G079J GOMEZ DEL CAMPO ET AL, PHYS. REV. C19 (1979) 2170  
 G079N GOMEZ DEL CAMPO, BULL. AM. PHYS. SOC. 24 (1979) 814  
 G079T GOTO AND HORIUCHI, PROGR. THEOR. PHYS. 62 (1979) 662  
 G079U GORBATOV, KRYLOV AND SOLOVEI, SOVIET J. NUCL. PHYS. 30  
 (1979) 490  
 G080E GOLDFARB AND NAGEL, NUCL. PHYS. A341 (1980) 494  
 G080F GOEKE, CASTEL AND REINHARD, NUCL. PHYS. A339 (1980) 377  
 G080J GOLDFARB, MANAGOLI AND SINGLETON, P. 485, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 G080L GORINGER AND SCHOCH, PHYS. LETT. B97 (1980) 41  
 G080N GORBATOV, KRYLOV AND SOLOVEI, YAD. FIZ. 32 (1980) 636

- GO81B GOKMEN ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 580  
 GR77A GROSSIORD ET AL, PHYS. REV. C15 (1977) 843  
 GR77B GRASSAU, GARI AND HEBACH, NUCLEAR PHYSICS A282 (1977) 66  
 GR77D GRUHLE AND KOBER, NUCLEAR PHYSICS A286 (1977) 523  
 GR77G GREBEN AND LEVIN, PHYS. LETT. B71 (1977) 252  
 GR77H GROSS AND TALMI, NUCLEAR PHYSICS A286 (1977) 211  
 GR77I GRUHLE AND MOBIUS, Z. PHYS. A283 (1977) 97  
 GR77J GRAY AND THOMPSON, NATURE 270, NO. 5639 (1977) 708  
 GR77K GRECKSCH, DILLIG AND HUBER, PHYS. LETT. B72 (1977) 11  
 GR77M GRECHUKHIN AND SOLDATOV, ZH. EKSP. TEOR. FIZ. 72 (1977) 390  
 GR78G GRIGORYAN AND SHAKHBAZYAN, YADERNAYA FIZ. 28 (1978) 1372  
 GR78H GRIDNEV ET AL, IZV. AKAD. NAUK SSSR (SER. FIZ.) 42 (1978) 2361  
 GR79C GREENWOOD, IN PROC. THIRD SYMP. ON NEUTRON CAPTURE GAMMA RAYS, 1978 (PLENUM PRESS, NEW YORK, 1979), P. 441  
 GR79M GREBEN AND LEVIN, NUCL. PHYS. A325 (1979) 145  
 GR79Q GRIDNEV, SEMIJDONOV, SUBBOTIN AND HETTER, MICROSCOPIC OPTICAL POTENTIALS, HAMBURG, GERMANY (SPRINGER-VERLAG 1979) P. 277, AND PHYS. ABSTRACTS 97715 (1979)  
 GR80 GRAVES ET AL, CAN. J. PHYS. 58 (1980) 48  
 GR80B GREENE ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 504  
 GR80D GREEN AND KORTELING, PHYS. REV. C22 (1980) 1594  
 GR80F GREENE ET AL, P. 687, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 GR80J GRABMAYR, RAPAPORT AND FINLAY, NUCL. PHYS. A350 (1980) 167  
 GR81A GREENE ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 582  
 GR81B GROTOWSKI ET AL, PHYS. REV. C23 (1981) 2513  
 GR81C GREENE ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 314  
 GU76D GUILLAUME, RASTEGAR, FINTZ AND GALLMANN, NUCLEAR PHYSICS A272 (1976) 338  
 GU78A GUPTA AND KAILAS, PHYS. LETT. B73 (1978) 398  
 GU78C GUICHON, GIFFON AND SAMOUR, PHYS. LETT. B74 (1978) 15  
 GU78D GUL, J. PHYS. SOC. JAPAN 44 (1978) 353  
 GU78F GUICHON ET AL, Z. PHYS. A285 (1978) 183  
 GU79 GUICHON ET AL, PHYS. REV. C19 (1979) 987  
 GU79C GUICHON AND SAMOUR, PHYS. LETT. B82 (1979) 28  
 GU79G GUARDIOLA, NUCL. PHYS. A328 (1979) 490  
 GU80B GUARDIOLA AND POLLS, NUCL. PHYS. A342 (1980) 385  
 GU81 GURBANOVICH AND ZELENSKAYA, PROCEEDINGS VERSAILLES CONF. (1981) p. 142  
 GU81B GUZIK, ASTROPHYS. J. 244 (1981) 695  
 GY78 GYULASSY AND KAUFFMANN, PHYS. REV. LETT. 40 (1978) 298  
 HA65C HART, NORBECK AND CARLSON, PHYS. REV. 137 (1965) B17  
 HA69S HATTIG, HUNCHEN, ROTH AND WAFFLER, NUCLEAR PHYSICS A137 (1969) 144  
 HA71B HARDY, ESTERL, SEXTRO AND CERNY, PHYS. REV. C3 (1971) 700  
 HA75DD HANNA ET AL, IN JUL-CONF-16 (1975), P. 8  
 HA75EE HANNA ET AL, IN JUL-CONF-16 (1975), P. 9  
 HA75G HARAKEH, PAUL AND SNOVER, PHYS. REV. C11 (1975) 998  
 HA75H HARAKEH, PAUL AND GORODETZKY, PHYS. REV. C11 (1975) 1008  
 HA76N HARAKEH, COMFORT AND VAN DER WOUDE, PHYS. LETT. B62 (1976) 155  
 HA76O HARAKEH ET AL, NUCLEAR PHYSICS A265 (1976) 189  
 HA76T HANNA, IN PROC. DUBNA CONF. JINR D-9920, VOL. II, P. 195 (1976)

- HA77AA HANNA, IN PROC. HAIFA INT. WORKSHOP (1976), (BRISTOL, ENGLAND, ADAM HILGER 1977), P. 577, AND PHYS. ABSTRACTS 70448 (1978)
- HA77H HARRIS ET AL, PHYS. REV. LETT. 38 (1977) 1460
- HA77I HAGEN, FISHER AND ORMES, ASTROPHYS. J. 212 (1977) 262
- HA77N HARTMANN, Z. PHYS. A282 (1977) 293
- HA77V HAEBERLI, PROC. INT. CONF. NUCL. STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 435
- HA77W HARVEY, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 127
- HA78A HAMM AND NAGATANI, PHYS. REV. C17 (1978) 586, 2258
- HA78AA HAAS ET AL, PROC. INT. CONF. ON RESONANCES IN HEAVY ION REACTIONS, HVAR, 1977 (NORTH-HOLLAND, AMSTERDAM, 1978), P. 349; PHYSICS ABSTRACTS 9626 (1979)
- HA78F HARRINGTON AND VARMA, PHYS. LETT. B74 (1978) 316
- HA78M HAKANSSON, BERGGREN AND BENGTSSON, NUCLEAR PHYSICS A306 (1978) 406
- HA78N HAXTON, NUCLEAR PHYSICS A306 (1978) 429
- HA78O HARRINGTON AND VARMA, NUCLEAR PHYSICS A306 (1978) 477
- HA78W HALKIA, ELLIOTT AND SANDERSON, FIZIKA (YUGOSLAVIA) 10, SUPPL. (1978) 39, AND PHYS. ABSTRACTS 74319 (1978)
- HA78Z HASSAN, MOHAMMAD AND ABDELMONEIM, ATOMKERNENERGIE 32 (1978) 193
- HA79H HARWOOD AND CRAWLEY, BULL. AM. PHYS. SOC. 24 (1979) 817
- HA79K HANNA, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108 (1979) P. 288
- HA79V HASSAN, MOHAMMAD, OSMAN AND ABDEL MONEM, ACTA PHYS. POLONICA B10 (1979) 845
- HA80A HARAKEH, PHYS. LETT. 90B (1980) 13
- HA80D HATANAKA ET AL, NUCL. PHYS. A340 (1980) 93
- HA80P HALDERSON AND PHILPOTT, NUCL. PHYS. A345 (1980) 141
- HA81D HAMILL ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 579
- HA81G HARAKEH AND DIEPERINK, PHYS. REV. C23 (1981) 2329
- HA81K HAAS AND ABE, PHYS. REV. LETT. 46 (1981) 1667
- HA81L HANNA, NUCL. PHYS. A358 (1981) C229
- HE66A HEWKA, HOLBROW AND MIDDLETON, NUCLEAR PHYSICS 88 (1966) 561
- HE68D HENSLEY, NETTLES AND BARNES, PHYS. LETT. 26B (1968) 435
- HE76J HEENEN, NUCLEAR PHYSICS A272 (1976) 399
- HE76K HEIL AND STOCK, PHYS. LETT. B65 (1976) 412
- HE76M HEUSCH, IN 1974 GAMMA RAY TRANSITION PROBABILITIES (U. OF DELHI PRESS, 1976), P. 92
- HE77B HERMANS, THESIS, U. UTRECHT (1977)
- HE77E HERMANS ET AL, NUCLEAR PHYSICS A284 (1977) 307
- HE77H HEISS, ENGELBRECHT AND HAHNE, NUCLEAR PHYSICS A289 (1977) 386
- HE77I HECHT, PHYS. REV. C16 (1977) 2401
- HE77K HERSCOVITZ, MARIS AND TEODORO, PHYS. LETT. B69 (1977) 33
- HE77L HENLEY, PROC. INT. CONF. NUCL. STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 812
- HE78A HECHT AND BRAUNSCHWEIG, NUCLEAR PHYSICS A295 (1978) 34
- HE78B HECKMAN ET AL, PHYS. REV. C17 (1978) 1651
- HE78C HECKMAN, GREINER, LINDSTROM AND SHWE, PHYS. REV. C17 (1978) 1735
- HE78D HENNINGSSEN, SCHATZLER AND KUHN, ATOMKERNENERGIE 31 (1978) 131
- HE78G HEFTER, GERAMB, OSTERFELD AND UDAGAWA, Z. PHYS. A284 (1978) 413

- HE78H HENDRIE, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 402
- HE78J HERTZ ET AL, PHYS. REV. C18 (1978) 2780
- HE78K HEITZMANN, ATOMKERNENERGIE 31 (1978) 262
- HE78M HERING ET AL, S. AFR. J. PHYS. 1 (1978) 169, AND PHYS. ABSTRACTS 18284 (1980)
- HE79 HECHT AND ZAHN, NUCLEAR PHYSICS A313 (1979) 77
- HE79A HERMAN, CORMIER AND FULTON, BULL. AM. PHYS. SOC. 24 (1979) 664
- HE79D HEENEN AND BAYE, PHYS. LETT. B81 (1979) 295
- HE79E HEFTER, LETT. NUOVO CIM. 24 (1979) 425
- HE79G HENDERSON ET AL, AUSTRALIAN J. PHYS. 32 (1979) 411
- HE79H HENNING, IN BNL-51115 (1979), P. 445
- HE79I HERING ET AL, IN BNL-51115 (1979), P. 759
- HE80D HEARST, BULL. AMER. PHYS. SOC. 25 (1980) 793
- HE80O HEIL AND MANAKOS, Z. PHYS. A294 (1980) 331
- HE81 HEENEN, PHYS. LETT. 99B (1981) 298
- HI70B HILDENBRAND, GUTBROD, VON OERTZEN AND BOCK, NUCL. PHYS. A157 (1970) 297
- HI76D HINTERBERGER ET AL, NUCLEAR PHYSICS A263 (1976) 460
- HI77 HIGH AND CUJEC, NUCLEAR PHYSICS A278 (1977) 149
- HI77C HIGH AND CUJEC, NUCLEAR PHYSICS A282 (1977) 181
- HI77F HIRATA, KOCH, LENZ AND MONIZ, PHYS. LETT. B70 (1977) 281
- HI78C HIRVONEN AND LUCKE, NUCL. INSTR. METH. 149 (1978) 295
- HI79C HIRATA, KOCH, LENZ AND MONIZ, ANN. OF PHYS. 120 (1979) 205
- HI80C HIRSEKORN, Z. PHYS. A297 (1980) 141
- HI81A HINTERBERGER ET AL, NUCL. PHYS. A352 (1981) 93
- HO69H HOFMANN ET AL, Z. PHYS. 221 (1969) 417
- HO75M HODGSON, NATURE 257, NO. 5526 (1975) 446
- HO76F HOFMANN ET AL, IN PROC. DUBNA CONF. JINR D-9920, VOL. II, P. 370 (1976)
- HO76H HORIKAWA, HOSHINO AND ARIMA, PHYS. LETT. B63 (1976) 134
- HO76J HORIUCHI, PROG. THEOR. PHYS. 55 (1976) 1448
- HO77 HORIKAWA, HOSHINO AND ARIMA, NUCLEAR PHYSICS A278 (1977) 297
- HO77B HOLINDE, MACHLEIDT, FAESSLER AND MUTHER, PHYS. REV. C15 (1977) 1432
- HO77C HOLT, SPECHT, JACKSON AND LASZEWSKI, NUCL. INSTR. METH. 141 (1977) 125
- HO77F HOLT ET AL, PHYS. LETT. B69 (1977) 55
- HO77H HOODBHOY AND NEGELE, NUCLEAR PHYSICS A288 (1977) 23
- HO77K HORIUCHI, PROG. THEOR. PHYS. 58 (1977) 204
- HO77L HORIUCHI, INVITED TALK AT INTERN. SYMP. NUCL. COLLISIONS AND THEIR MICROSCOPIC DESCRIPTION, BLED (YUGOSLAVIA), 1977
- HO77M HORIUCHI, PROC. INT. CONF. NUCL. STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 85
- HO77O HO ET AL, Z. PHYS. A283 (1977) 235
- HO77P HOISTAD, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 215
- HO77R HORIUCHI, PROG. THEOR. PHYS. SUPPL. 62 (1977) 90
- HO77S HOHN, BARZ AND ROTTER, NUKLEONIKA 22 (1977) 1175
- HO78F HORLACHER AND ARENHOVEL, NUCLEAR PHYSICS A300 (1978) 348
- HO78H HORN AND FERGUSON, PHYS. REV. LETT. 41 (1978) 1529
- HO78I HOLT ET AL, PHYS. REV. C18 (1978) 1962
- HO78K HORN, FERGUSON AND HAUSSER, NUCLEAR PHYSICS A311 (1978) 238
- HO78N HODGSON, RIV. NUOVO CIM. 1, NO. 1 (1978) 1

- H079D HOLSTEIN AND KIM, PHYS. REV. C19 (1979) 1433  
 H079I HOISTAD ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
 PAPER 4B13  
 H080C HOLTKAMP ET AL, PHYS. REV. LETT. 45 (1980) 420  
 H080D HONDA, KUDO AND HORIE, PROGR. THEOR. PHYS. 63 (1980) 872  
 H080K HOLTKAMP AND COTTINGAME, P. 699, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 H080L HORIKAWA, THIES AND LENZ, NUCL. PHYS. A345 (1980) 386  
 H081 HOSONO ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE  
 PROC. 69 (1981) p. 629  
 HU76H HUMBLET, DYER AND ZIMMERMAN, NUCLEAR PHYSICS A271  
 (1976) 210  
 HU77B HUSKEN, NUCLEAR PHYSICS A291 (1977) 206  
 HU79 HUSSEIN, PHYS. REV. C19 (1979) 807  
 HU79B HUFNER AND THIES, PHYS. REV. C20 (1979) 273  
 HU79C HUBER AND KLINGENBECK, NUKLEONIKA 24 (1979) 95  
 HU79D HUIZENGA ET AL, IN BNL-51115 (1979), P. 235  
 HU80 HUSSEIN, CANTO AND DONANGELO, PHYS. REV. C21 (1980) 772  
 HU80C HUGG AND HANNA, POLARIZATION - SANTA FE (1980) 2.94  
 HU80E HUGI ET AL, P. 508, PROCEEDINGS OF THE INTERNATIONAL  
 CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 HU80G HULKI, ROLFS AND TRAUTVETTER, Z. PHYS. A297 (1980) 161  
 HU81A HUNGERFORD ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 606  
 HU81C HUFNER AND NEMES, PHYS. REV. C23 (1981) 2538  
 HY79 HYNES ET AL, PHYS. REV. LETT. 42 (1979) 1444  
 HY80 HYNES ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 575  
 HY80A HYUGA, ARIMA AND SHIMIZU, NUCL. PHYS. A336 (1980) 363  
 HY81 HYDE ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 27  
 IB77A IBEN, ASTROPHYS. J. 217 (1977) 788  
 IB78 IBEN, ASTROPHYS. J. 219 (1978) 213  
 IB78A IBEN, ASTROPHYS. J. 226 (1978) 996  
 IK77 IKEDA, IN TOKYO (1977), CONTRIBUTED PAPERS P. 218  
 IK77B IKEDA AND TAMAGAKI, PROG. THEOR. PHYS. SUPPL. 62 (1977) 1  
 IK80 IKOSSI, SNOVER, ADELBERGER AND LESKO, BULL. AMER. PHYS.  
 25 (1980) 577  
 IK80A IKEDA, HORIUCHI AND SAITO, SUPPL. PROGR. THEOR. PHYS., NO. 68  
 (1980) 1  
 IM63 IMHOFF, GRENCH AND JOHNSON, NUCLEAR PHYSICS 49 (1963) 503  
 IM77B IMANISHI, TANIMURA AND OHNISHI, PROC. INT. CONF. NUCL.  
 STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978)  
 SUPPL., P. 257  
 IM79B IMADA, KURITA, BANDO AND SHONO, PROG. THEOR. PHYS.  
 62 (1979) 1422  
 IN76D INOUE AND TANAKA, J. INORG. & NUCL. CHEM. (GB) 38 (1976) 1425  
 IN78 INGRAM ET AL, PHYS. LETT. B76 (1978) 173  
 IN79B INOPIN, KINCHAKOV, LUKYANOV AND POL', ANN. OF PHYS. 118  
 (1979) 307  
 IN79C INGRAM, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 455  
 IN79D INOPIN, INOPIN, KOPANETS AND KORDA, IZV. AKAD. NAUK  
 SSSR (SER. FIZ.) 43 (1979) 2308  
 IO78A IOANNIDES AND JACKSON, NUCLEAR PHYSICS A308 (1978) 305, 317  
 IO81 ION AND ION-MIHAI, NUCL. PHYS. A360 (1981) 400  
 IS78A ISMAIL, ATOMKERNENERGIE 31 (1978) 44  
 IS80B ISHKHANOV ET AL, P. 861, PROCEEDINGS OF THE INTERNATIONAL  
 CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 IS80E ISHKHANOV ET AL, YAD. FIZ. 31 (1980) 1407

- IT78 ITOH AND KOHYAMA, NUCLEAR PHYSICS A306 (1978) 527  
 IZ77 IZUMOTO, NUCLEAR PHYSICS A275 (1977) 342  
 IZ77B IZUMOTO, INS REPORT 300 (1977)  
 IZ79 IZUMOTO AND MORI, PHYS. LETT. B82 (1979) 163  
 IZ80 IZUMOTO, KREWALD AND FAESSLER, P. 502, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 IZ80A IZUMOTO, P. 566, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 JA68H JAYARAMAN AND HOLMGREN, PHYS. REV. 172 (1968) 1015  
 JA76N JAHN ET AL, PHYS. LETT. B65 (1976) 339  
 JA76Q JACOBSSON AND KULLBERG, PHYS. SCRIPTA 13 (1976) 327  
 JA77 JAKOBSSON, KULLBERG AND OTTERLUND, NUCLEAR PHYSICS A276 (1977) 523  
 JA77C JANG AND BANG, IN TOKYO (1977), CONTRIBUTED PAPERS P. 586  
 JA77J JAUSEL-HUSKEN AND FREIESLEBEN, Z. PHYS. A283 (1977) 363  
 JA78A JACHCINSKI ET AL, PHYS. REV. C17 (1978) 1263  
 JA78F JACKLE, PILKUHN AND SCHLAILE, PHYS. LETT. B76 (1978) 177  
 JA78G JAMES AND FLETCHER, PHYS. REV. C17 (1978) 2248  
 JA78I JAHN ET AL, PHYS. REV. C18 (1978) 9  
 JA78J JAKOBSSON ET AL, NUCLEAR PHYSICS A300 (1978) 397  
 JA78K JANSEN ET AL, PHYS. LETT. B77 (1978) 359  
 JA78M JAROS ET AL, PHYS. REV. C18 (1978) 2273  
 JA78O JAKOBSSON, PHYS. SCRIPTA 17 (1978) 491  
 JA79 JAIN AND SHAstry, PHYS. REV. C19 (1979) 848  
 JA79A JAIN, NUCLEAR PHYSICS A314 (1979) 51  
 JA79D JAIN, J. OF PHYS. G5 (1979) 663  
 JA79F JAKOBSSON, BONDORF AND FAI, PHYS. LETT. B82 (1979) 35  
 JA79G JAIN AND SARMA, NUCLEAR PHYSICS A321 (1979) 429  
 JA79H JACKSON AND BRENNER (TO BE PUBLISHED)  
 JA79J JACKSON, IOANNIDES AND THOMAS, NUCL. PHYS. A322 (1979) 493  
 JA79M JADID AND MAVROMATIS, NUCLEAR PHYSICS A317 (1979) 399  
 JA79T JARCZYK ET AL, NUCL. PHYS. A325 (1979) 510  
 JA79U JAMINON, MAHAUX AND ROCHUS, PHYS. REV. LETT. 43 (1979) 1097  
 JA79V JAKOBSSON, JONSSON AND KRISTIANSSON, ASTROPHYS. & SPACE SCI. 63 (1979) 215  
 JA79W JACHCINSKI ET AL, PHYS. LETT. B87 (1979) 354  
 JA80B JACHCINSKI ET AL, PHYS. REV. C22 (1980) 101  
 JA80E JAMINON, MAHAUX AND ROCHUS, PHYS. REV. C 22 (1980) 2027  
 JE75 JELLEY, CERNY, STAHEL AND WILCOX, PHYS. REV. C11 (1975) 2049  
 JE76B JEUKENNE, LEJEUNE AND MAHAUX, IN LOWELL CONF. CONF-760715-P1 (1976), P. 451  
 JE77A JEUKENNE, LEJEUNE AND MAHAUX, PHYS. REV. C16 (1977) 80  
 JE79A JENNEWIN AND SCHOCH, Z. PHYS. A292 (1979) 27  
 JE80C JENKINS ET AL, P. 685, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 JO73 JOHNSON, PHYS. REV. C7 (1973) 561; IBID. C8 (1973) 851  
 JO76C JOGLEKAR, ANN. OF PHYS. 101 (1976) 433  
 JO77 JOHNSON, NILSSON AND LINDGREN, NUCLEAR PHYSICS A278 (1977) 365  
 JO77C JOSS, NATURE 270, NO. 5635 (1977) 310  
 JO79 JOHNSON ET AL, PHYS. REV. C20 (1979) 27  
 JO79A JONES, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 116  
 JO79D JOHNSON, FOWLER, HILL AND ORTOLF, BULL. AM. PHYS. SOC. 24 (1979) 886

- J079G JOHNSON ET AL, PHYS. REV. LETT. 43 (1979) 844  
 J079H JORDAN, MAHER AND PENG, PHYS. LETT. B87 (1979) 38  
 J079I JOHN, KRAMER AND HETZEL, SEVENTH INTERNATIONAL COLLOQUIUM AND INTEGRATIVE CONFERENCE ON GROUP THEORY AND MATHEMATICAL PHYSICS, AUSTIN, TEXAS, USA (SPRINGER-VERLAG 1979) P. 439, AND PHYS. ABSTRACTS 9276 (1980)  
 JU77C JURIC, IN PROC., ZURICH - SIN, 1977, P. 296  
 JU80 JURY ET AL, PHYS. REV. C21 (1980) 503  
 KA61C KAUFMANN AND WAFFLER, NUCLEAR PHYSICS 24 (1961) 62  
 KA76Q KABACHNIK AND RAZUNAEV, BULL. ACAD. SCI. USSR, PHYS. SER. 40 (1976) 148  
 KA77 KALISH, HASS, BRENNAN AND KING, NUCL. PHYS. A276 (1977) 339  
 KA77BB KALINSKY, MELNIK, SMILANSKY AND TRAUTNER, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 263  
 KA77C KAZAKS, BULL. AM. PHYS. SOC. 22 (1977) 616  
 KA77CC KAPTARI, LUKYANOV AND TITOV, IN PROC., ZURICH - SIN, 1977, P. 175  
 KA77EE KAMUNTAVICIUS, SOV. PHYS.-COLLECT. (USA) 17, NO. 3 (1977) 1, AND PHYS. ABSTRACTS 63610 (1979)  
 KA77FF KAMUNTAVICIUS, SOV. PHYS.-COLLECT. (USA) 17, NO. 4 (1977) 28, AND PHYS. ABSTRACTS 63582 (1979)  
 KA77K KAWAI, IMANISHI AND ICHIMURA, IN TOKYO (1977), CONTRIBUTED PAPERS, P. 526  
 KA77V KALINSKY ET AL, NUCLEAR PHYSICS A289 (1977) 205  
 KA78A KATORI, FURUNO AND OOI, PHYS. REV. LETT. 40 (1978) 1489, AND IBID. 41 (1978) 138  
 KA78I KATO AND BANDO, PROG. THEOR. PHYS. 59 (1978) 774  
 KA78R KAMUNTAVICIUS, SOVIET J. NUCL. PHYS. 27 (1978) 52  
 KA78S KALER, IBEN AND BECKER, ASTROPHYS. J. 224 (1978) L63  
 KA78T KAPTAR' AND TITOV, YADERNAYA FIZ. 28 (1978) 647  
 KA78U KAWAI, KAMIMURA, MITO AND TAKESAKO, PROG. THEOR. PHYS. 59 (1978) 676  
 KA78W KAMUNTAVICIUS, LITOV. FIZ. SB. (USSR) 18 (1978) 693, AND PHYS. ABSTRACTS 42551 (1980)  
 KA79A KASSIS AND KHALIL, NUCLEAR PHYSICS A315 (1979) 381  
 KA79C KALER, ASTROPHYS. J. 228 (1979) 163  
 KA79D KAMPFER AND ROTTER, J. OF PHYS. G5 (1979) 747  
 KA79M KAHANA, KIM AND MERMAZ, PHYS. REV. C20 (1979) 2124  
 KA79N KAMBLE AND KHADIKAR, PRAMANA 13 (1979) 475  
 KA79Q KARBAN, MICROSCOPIC OPTICAL POTENTIALS, HAMBURG, GERMANY (SPRINGER-VERLAG 1979) P. 264, AND PHYS. ABSTRACTS 97712 (1979)  
 KA79S KASCHIEV AND SHITIKOVA, YAÖ. FIZ. (USSR) 30 (1979) 1479  
 KA80H KARAOGLU, KARAPIPERIS AND MONIZ, PHYS. REV. C22 (1980) 1806  
 KA80M KASATKIN, GONCHAR AND INOPIN, UKR. FIZ. ZH. 25 (1980) 705  
 KA80N KAN, PHYS. REV. C 22 (1980) 2228  
 KA80P KALER, ASTROPHYS. J. 239 (1980) 78  
 KA81C KARBAN, BASAK AND ROMAN, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 706  
 KA81D KARBAN ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 703  
 KA81E KASSIS AND ARAFAH, J. OF PHYS. G7 (1981) 165  
 KA81F KAWAZOE AND TSUKAMOTO, PHYS. REV. C23 (1981) 2364  
 KA81G KALETKA ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 300

- KE66C KELSON AND GARVEY, PHYS. LETT. 23 (1966) 689  
 KE71E KERNEL, MASON AND VON WIMMERSPERG, NUCLEAR PHYSICS A167 (1971) 352  
 KE78B KEMPER, PARKS, MATEJA AND ROY, BULL. AM. PHYS. SOC. 23 (1978) 615  
 KE78C KEKELIS ET AL, PHYS. REV. C17 (1978) 1929  
 KE80 KENNY, BIRD AND CLAYTON, NUCL. INSTR. METH. 168 (1980) 115  
 KE80C KELLY ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 575  
 KE80G KELLY ET AL, PHYS. REV. LETT. 45 (1980) 2012  
 KE80H KEMPER AND OPHEL, AUSTRALIAN J. PHYS. 33 (1980) 197  
 KH78A KHANNA, TOWNER AND LEE, NUCLEAR PHYSICS A305 (1978) 349  
 KH78B KHYR ET AL, PHYS. LETT. B76 (1978) 273  
 KH79B KHAN, LOWE, NELSON AND BARNETT, NUCLEAR PHYSICS A322 (1979) 33  
 KH80 KHODYACHIKH, VATSET, DOGYUST AND KIRICHENKO, UKR. FIZ. ZH. (USSR) 25 (1980) 229, AND PHYS. ABSTRACTS 47465 (1980)  
 KH80A KHALLAF, REV. ROU. PHYS. 25 (1980) 741  
 KH81 KHAIROZZAMAN, CURRENT SCIENCE 50 (1981) 21  
 KH81B KHARITONOV, SMIRNOV AND SLIV, IZV. AKAD. NAUK SSSR, SER. FIZ. 45 (1981) 66  
 KI76K KIRK, IN LOWELL CONF., CONF-760715-P1 (1976), P. 389  
 KI76M KITCHING ET AL, PHYS. REV. LETT. 37 (1976) 1600  
 KI77 KIDD ET AL, BULL. AM. PHYS. SOC. 22 (1977) 567  
 KI77H KISSLINGER AND VAN GIAI, PHYS. LETT. B72 (1977) 19  
 KI77J KILIAN, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 91  
 KI77M KIRPICHNIKOV ET AL, IN PROC., ZURICH - SIN, 1977, P. 206  
 KI78A KIM ET AL, NUCLEAR PHYSICS A297 (1978) 301  
 KI79E KIRSON, NUCL. PHYS. A317 (1979) 388  
 KI79G KILIAN, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 666  
 KI79K KIRCHBACH AND JAGER, YADERNAYA FIZ. 29 (1979) 1191  
 KI79M KIM AND PRIMAKOFF, IN "MESONS IN NUCLEI," EDS. RHO AND WILKINSON (NORTH-HOLLAND, AMSTERDAM, 1979), P. 69  
 KI79N KIRICHENKO, INOPIN AND GONCHAR, UKR. FIZ. ZH. 24 (1979) 388  
 KI80A KITCHING ET AL, NUCL. PHYS. A340 (1980) 423  
 KI80E KIRICHENKO, VATSET, DOGYUST AND KHODYACHIKH, UKR. FIZ. ZH. 25 (1980) 798  
 KI81A KIRPICHNIKOV ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 324  
 KL79E KLINGENBECK, PHYS. LETT. B85 (1979) 21  
 KL79F KLEIN ET AL, NUCL. PHYS. A329 (1979) 339  
 KN70A KNUDSON AND YOUNG, NUCLEAR PHYSICS A149 (1970) 323  
 KN75D KNEISSL ET AL, NUCL. INSTR. METH. 127 (1975) 1  
 KN76G KNUPFER AND HUBER, PHYS. REV. C14 (1976) 2254  
 KN77 KNUPFER, GRECKSCH AND HUBER, Z. PHYS. A280 (1977) 175  
 KN77C KNOPFLE ET AL, BULL. AM. PHYS. SOC. 22 (1977) 541  
 KN78A KNOPFLE ET AL, PHYS. LETT. B74 (1978) 191  
 KN79 KNUPFER, ENDER AND HUBER, INT. CONF. NUCL. PHYS. WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 1.21  
 KN79C KNUPFER, ENDER AND HUBER, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 5A13  
 KN79I KNOPFLE, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108 (1979) P. 311  
 KN79J KNOPFLE ET AL, NUCLEAR INTERACTIONS, CANBERRA, AUSTRALIA (SPRINGER-VERLAG 1979) P. 445, AND PHYS. ABS. 13440 (1980)

- KN79K KNOPFLE ET AL, NUCLEAR INTERACTIONS, CANBERRA,  
AUSTRALIA (SPRINGER-VERLAG 1979) P. 442, AND PHYS.  
ABSTRACTS 13440 (1980)
- K074 KOONIN, TOMBRELLO AND FOX, NUCLEAR PHYSICS A220 (1974) 221
- K074V KOZLOWSKI AND ZGLINSKI, NUKLEONIKA (POLAND) 19 (1974) 721
- K077CC KOLTUN AND MYHRER, Z. PHYS. A283 (1977) 397
- K077D KOONIN ET AL, PHYS. REV. C15 (1977) 1359
- K077G KOSTIN, KOPANETS AND KOVAL, IZV. AKAD. NAUK SSSR (SER.  
FIZ.) 41 (1977) 153
- K077H KOENIG ET AL, NUOVO CIM. 39 (1977) 9
- K077J KOLATA ET AL, PHYS. REV. C16 (1977) 891
- K077JJ KOLYBASOV AND SMORODINSKAYA, JETP LETT. 26 (1977) 109
- K077K KOYAMA, NAKANISHI AND OHNUMA, IN TOKYO (1977), CONTRIBUTED  
PAPERS, P. 546
- K077MM KOIKE, RES. REP. NAGAOKA TECH. COLL. (JAPAN) 13 (1977) 145;  
PHYS. ABS. 37355 (1978)
- K077W KONOPKA, GARI AND ZABOLITZKY, NUCLEAR PHYSICS A290 (1977)  
360, AND IBID. A306 (1978) 536
- K078A KOVAR ET AL, PHYS. REV. C17 (1978) 83
- K078H KOONIN, FLANDERS, FLOCARD AND WEISS, PHYS. LETT. B77  
(1978) 13
- K078J KOONIN ET AL, BULL. AM. PHYS. SOC. 23 (1978) 955
- K078N KO AND RISKA, NUCLEAR PHYSICS A312 (1978) 217
- K079B KOLATA ET AL, PHYS. REV. C19 (1979) 408
- K0798B KOSHIGIRI, OHTSUBO AND MORITA, PROGR. THEOR. PHYS. 62  
(1979) 706
- K079E KOBOS AND MACKINTOSH, J. OF PHYS. G5 (1979) 97
- K079FF KOLYBASOV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2033
- K079GG KOO AND TASSIE, AUSTRALIAN J. OF PHYS. 32 (1979) 427
- K079H KOSHEL AND GRIFFIN, BULL. AM. PHYS. SOC. 24 (1979) 611
- K079KK KOBOS AND MACKINTOSH, ANN. PHYS. 123 (1979) 296
- K079LL KOHNO, NAGATA AND YAMAGUCHI, PROG. THEOR. PHYS. SUPPL. 65  
(1979) 200
- K079M KOLATA ET AL, PHYS. REV. C19 (1979) 2237
- K079O KOHNO AND ANDO, PROG. THEOR. PHYS. 61 (1979) 1065
- K079P KOCH AND MONIZ, PHYS. REV. C20 (1979) 235
- K079R KOZLOWSKI, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
PAPER 5B6
- K079S KOSICE AND LENINGRAD COLLABORATION, YADERNAYA FIZ. 29  
(1979) 117
- K079T KOLATA ET AL, BULL. AM. PHYS. SOC. 24 (1979) 825
- K079W KOESTER, KNOPF AND WASCHKOWSKI, Z. PHYS. A292 (1979) 95
- K079Y KOVAR ET AL, PHYS. REV. C20 (1979) 1305
- K080 KOLATA ET AL, PHYS. REV. C21 (1980) 579
- K080B KOVAR ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 524
- K080C KOX, COLE AND OST, PHYS. REV. LETT. 44 (1980) 1204
- K080CC KOPTEV, MAEV, MAKAROV AND KHANZADEEV, YAD. FIZ. 31 (1980)  
1501
- K080E KO, PHYS. REV. C21 (1980) 2672
- K080H KONDO, BROMLEY AND ABE, PROGR. THEOR. PHYS. 63 (1980) 722
- K080K KONDO, BROMLEY AND ABE, PHYS. REV. C22 (1980) 1068
- K080S KOZLOWSKI ET AL, P. 229, PROCEEDINGS OF THE INTERNATIONAL  
CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- K080U KOLATA, PHYS. LETT. B95 (1980) 215
- K080W KOBER, GRUHLE AND HILLEN, NUCL. INSTRUM. & METHODS 174  
(1980) 173

- K081A KOLATA ET AL, PHYS. REV. C23 (1981) 1056  
 K081G KOCH AND MONIZ, PROCEEDINGS VERSAILLES CONF. (1981) p. 420  
 KR75F KREWALD ET AL, PROC. INTERN. CONF. ON NUCLEAR  
     SELF-CONSISTENT FIELDS, ITALY, 1975 (AMSTERDAM -  
     NORTH-HOLLAND 1975), P. 267; PHYS. ABS. 86092 (1976)  
 KR75G KREWALD, BIRKHOLZ, FAESSLER AND SPETH, IN JUL-CONF-16  
     (1975), P. 10  
 KR77 KREWALD, KLEMT, SPETH AND FAESSLER, NUCLEAR PHYSICS  
     A281 (1977) 166  
 KR78 KRENCIGLOWA AND BANDO, NUCLEAR PHYSICS A294 (1978) 191  
 KR78C KRATSCHMER, NUCLEAR PHYSICS A298 (1978) 477  
 KR78D KRIEGER AND DAVIES, PHYS. REV. C18 (1978) 2567  
 KR78F KRANE, ATOMIC DATA & NUCL. DATA TABLES 22 (1978) 269  
 KR79A KRIVINE, TREINER AND BOHIGAS, INT. CONF. NUCL. PHYS.  
     WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 1.22  
 KR79B KRIEGER AND DAVIES, PHYS. REV. C20 (1979) 167  
 KR79C KRAPPE, NIX AND SIERK, PHYS. REV. C20 (1979) 992  
 KR80 KRETSCHMER, PROBSTLE AND STACH, NUCL. PHYS. A333 (1980) 13  
 KR80H KRETSCHMER ET AL, P. 848, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 KR80I KRASNOPOL'SKY AND KUKULIN, PHYS. LETT. B96 (1980) 4  
 KR81 KRETSCHMER, PROBSTLE AND STACH, IN POLARIZATION, SANTA FE  
     1980, AIP CONFERENCE PROC. 69 (1981) p. 724  
 KR81A KRETSCHMER AND RENNER, IN POLARIZATION, SANTA FE 1980, AIP  
     CONFERENCE PROC. 69 (1981) p. 1329  
 KU78D KUBO AND AMAKAWA, PHYS. REV. C17 (1978) 1271  
 KU78E KULLBERG, OSKARSSON AND OTTERLUND, PHYS. REV. LETT. 40  
     (1978) 289  
 KU78H KULLBERG AND OSKARSSON, Z. PHYS. A288 (1978) 283  
 KU79 KUBONO ET AL, NUCLEAR PHYSICS A313 (1979) 434  
 KU79B KUBONO, BOND AND THORN, PHYS. B81 (1979) 140  
 KU79G KUDO AND MANO, LETT. NUOVO CIM. 24 (1979) 289  
 KU79J KUBONO, BOND, HORN AND THORN, PHYS. LETT. B84 (1979) 408  
 KU79L KUMAR, CONF. STRUCTURE MEDIUM-HEAVY NUCLEI, RHODOS (1979)  
 KU80D KUMAR, IN NUCLEI FAR FROM STABILITY, J. H. HAMILTON ET  
     AL EDS., NORTH-HOLLAND (1980) P. 265  
 KU80G KUBONO, BOND AND HORN, P. 542, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 KW78 KWON AND TABAKIN, PHYS. REV. C18 (1978) 932  
 LA76J LACANNA, GLAVISH, CALARCO AND HANNA, BULL. AM. PHYS.  
     SOC. 21 (1976) 1294  
 LA77A LANGWORTHY, LAMERS AND UBERALL, NUCLEAR PHYSICS A280  
     (1977) 351  
 LA77F LANCMAN, VAN'T WESTENDE AND GRABER, NUCLEAR PHYSICS  
     A291 (1977) 293  
 LA77J LASSILA AND LARY, IN PROC., ZURICH - SIN, 1977, P. 180  
 LA78 LAZZARINI ET AL, BULL. AM. PHYS. SOC. 23 (1978) 15  
 LA78A LATTIMER, SCHRAMM AND GROSSMAN, ASTROPHYS. J. 219  
     (1978) 230  
 LA79 LANDOWNE, PHYS. REV. LETT. 42 (1979) 633  
 LA79C LAL ET AL, BULL. AM. PHYS. SOC. 24 (1979) 564  
 LA79F LAZAREFF, AUDOUZE, STARRFIELD AND TRURAN, ASTROPHYS. J. 228  
     (1979) 875  
 LA79J LANGANKE, LEUTENANTSMEYER, STINGL AND WEIGUNY, Z. PHYS.  
     A291 (1979) 267

- LA79O LARSON, HARVEY AND HILL, BULL. AM. PHYS. SOC. 24 (1979) 862  
 LA79P LAZZARINI, DOUBRE, SEAMSTER AND VANDENBOSCH, BULL. AM.  
       PHYS. SOC. 24 (1979) 852  
 LA80A LANDAU, NUCL. PHYS. A335 (1980) 289  
 LA80B LARSON, HETRICK AND HARVEY, BULL. AM. PHYS. SOC. 25 (1980)  
       543  
 LA80F LANDOWNE ET AL, PHYS. LETT. 90B (1980) 389  
 LA80G LARRY AND BERES, PHYS. REV. C22 (1980) 1145  
 LA80I LAUL, BULL. AMER. PHYS. SOC. 25 (1980) 770  
 LA80J LANDOWNE, WOLTER AND SCHLICHER, P. 648, PROCEEDINGS OF THE  
       INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 LA80K LANGANKE, LEUTENANTSMEIER, STADEMANN AND WEIGUNY,  
       P. 498, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE  
       ON NUCLEAR PHYSICS, BERKELEY (1980)  
 LA80L LARRY AND BERES, P. 496, PROCEEDINGS OF THE  
       INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 LA81C LAZZARINI ET AL, PHYS. REV. C24 (1981) 309  
 LE63B LEE, MO AND WU, PHYS. REV. LETT. 10 (1963) 258  
 LE72C LEMAIRE, MERMAZ AND SETH, PHYS. REV. C5 (1972) 328  
 LE76P LE MERE, TANG AND THOMPSON, PHYS. REV. C14 (1976) 1715  
 LE76Q L'ECUYER ET AL, BULL. AM. PHYS. SOC. 21 (1976) 1344  
 LE76R LEE, KHANNA, LONE AND MC DONALD, PHYS. LETT. B65 (1976) 201  
 LE76T LEFORT, REP. PROG. PHYS. 39 (1976) 129  
 LE77C LEMERE, BROWN, TANG AND THOMPSON, PHYS. REV. C15 (1977) 1191  
 LE77P LEE, KURATH AND ZEIDMAN, PHYS. REV. LETT. 39 (1977) 1307  
 LE78C LEJEUNE AND HODGSON, NUCLEAR PHYSICS A295 (1978) 301  
 LE78F LE MERE AND TANG, IN PROCEEDINGS, CLUSTER, WINNIPEG 1978, A21  
 LE78G LEFORT AND NGO, ANN. DE PHYSIQUE 3 (1978) 5  
 LE78K LEMERE AND TANG, PHYS. REV. C18 (1978) 1114  
 LE78N LEDERER AND SHIRLEY, TABLE OF ISOTOPES, JOHN WILEY PUBS.  
       (1978)  
 LE78Q LEE, TAKIGAWA AND MARTY, NUCLEAR PHYSICS A308 (1978) 161, 189  
 LE78R LEVINE, STEADMAN, KARP AND ALHASSID, PHYS. REV. LETT. 41  
       (1978) 1537  
 LE78S LEE, NUCLEAR PHYSICS A311 (1978) 518  
 LE78T LEE, ASTROPHYS. J. 224 (1978) 217  
 LE79A LE MERE AND TANG, PHYS. REV. C19 (1979) 391  
 LE79B LEE ET AL, PHYS. REV. LETT. 42 (1979) 429  
 LE79D LEMERE, STUBEDA, HORIUCHI AND TANG, NUCLEAR PHYSICS  
       A320 (1979) 449  
 LE79E LEWIS AND PERKINS, IEEE TRANSACTIONS ON NUCL. SCI. 26  
       (1979) 1818  
 LE79G LEE, IN PROC. THIRD SYMP. ON NEUTRON CAPTURE GAMMA  
       RAYS, 1978 (PLENUM PRESS, NEW YORK, 1979), P. 353  
 LE79H LEITCH ET AL, IN HOUSTON, AIP CONF. PROCEEDINGS 54  
       (1979), P. 357  
 LE79M LEE, SCHRAMM, WEFEL AND BLAKE, ASTROPHYS. J. 232 (1979) 854  
 LE79O LEMERE AND TANG, PHYS. REV. C20 (1979) 2003  
 LE80C LEE AND KURATH, PHYS. REV. C21 (1980) 293  
 LE80G LEE, MATSUSE AND ARIMA, PHYS. REV. LETTERS 45  
       (1980) 165  
 LE80R LEMERE, TANG, KANELLOPOULOS AND SUNKEL, NUCL. PHYS.  
       A348 (1980) 321  
 LE80T LEEB AND SCHMID, Z. PHYS. A298 (1980) 113  
 LE80U LEVKOVSKII AND DREIMAN, SOVIET J. NUCL. PHYS. 31 (1980) 295

- LE81C LEE ET AL, PHYS. LETT. 98B (1981) 418  
 LE81E LEVIN AND EISENBERG, NUCL. PHYS. A355 (1981) 277  
 LE81F LEWIS ET AL, IN POLARIZATION, SANTA FE, 1980, AIP CONFERENCE PROC. 69 (1981) p. 590  
 LI76 LI, DEHNHARD, BROWN AND ELLIS, PHYS. REV. C13 (1976) 55  
 LI76R LIU AND SHAKIN, PHYS. REV. C14 (1976) 1885  
 LI76S LIU AND VAN GIAI, PHYS. LETT. B65 (1976) 23  
 LI76T LIPPARINI, STRINGARI AND TRAINI, NUOVO CIM. A34 (1976) 48  
 LI77E LIPPARINI, ORLANDINI AND LEONARDI, PHYS. REV. C16 (1977) 812  
 LI77G LIPPARINI ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 134  
 LI77I LINDAHL AND SHULTIS, NUCL. INSTR. METH. 144 (1977) 317  
 LI77J LISKIEN AND PAULSEN, ATOMKERNENERGIE 30 (1977) 59  
 LI77K LISBONA, NUCL. INSTR. METH. 146 (1977) 243  
 LI77L LIPPARINI, ORLANDINI, STRINGARI AND TRAINI, NUOVO CIM. A42 (1977) 296  
 LI78D LIU AND SHAKIN, PHYS. LETT. B78 (1978) 389  
 LI79A LIU AND SHAKIN, PHYS. REV. C19 (1979) 129  
 LI79G LIEB ET AL, PHYS. REV. C19 (1979) 2405  
 LI79H LINDGREN ET AL, PHYS. REV. LETT. 42 (1979) 1524  
 LI79I LIPPERHEIDE, FIEDELDEY, HABERZETTL AND NAIDOO, PHYS. LETT. B82 (1979) 39  
 LI79N LIU AND SHAKIN, NUOVO CIM. 53A (1979) 142  
 LI80B LINCK, KRAUS AND BLATT, PHYS. REV. C21 (1980) 791  
 LI80F LIBERT-HEINEMANN, BAYE AND HEENEN, NUCL. PHYS. A339 (1980) 429  
 LI80J LIU, WE, HWANG AND LI, P. 688, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 LI80K LIEB ET AL, P. 715, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 LI80N LIU ET AL, PHYS. ENERG. FORTIS & PHYS. NUCL. 4 (1980) 394; PHYS. ABS. 102211 (1980)  
 LI81B LI, LI AND LEE, PHYS. LETT. B99 (1981) 200  
 LI81D LIPPARINI AND STRINGARI, PHYS. LETT. B99 (1981) 183  
 LI81H LIU AND LI, PROCEEDINGS VERSAILLES CONF. (1981) p. 286  
 LI81I LILJESTRAND ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 544  
 LI81J LIN AND ZAMICK, PROCEEDINGS VERSAILLES CONF. (1981) p. 467  
 L072B LOWE AND BARNETT, NUCLEAR PHYSICS A187 (1972) 323  
 L076F LOMNITZ-ADLER AND BRINK, PHYS. LETT. B65 (1976) 106  
 L076G LODHI AND WAAK, ANN. OF PHYS. 101 (1976) 1  
 L077H LOMBAARD AND MEYER, NUCL. INSTR. METH. 145 (1977) 347  
 L079B LONDERGAN AND NIXON, PHYS. REV. C19 (1979) 998  
 L080 LOZANO AND MADURGA, PHYS. LETTERS B90 (1980) 50  
 LU69A LU, ZISMAN AND HARVEY, PHYS. REV. 186 (1969) 1086  
 LU78 LUCK, ASTROPHYS. J. 219 (1978) 148  
 LU79A LUHN ET AL, BULL. AM. PHYS. SOC. 24 (1979) 841  
 LU80 LUI ET AL, NUCL. PHYS. A333 (1980) 205  
 LU80A LUI ET AL, NUCL. PHYS. A333 (1980) 221  
 MA74C MACDONALD ET AL, PHYS. REV. C9 (1974) 1694  
 MA77 MATTHEWS ET AL, PHYS. REV. LETT. 38 (1977) 8  
 MA77E MASUDA AND UCHIYAMA, PHYS. REV. C15 (1977) 972  
 MA77F MASUDA AND UCHIYAMA, PHYS. REV. C15 (1977) 1598  
 MA77G MARTZ ET AL, BULL. AM. PHYS. SOC. 22 (1977) 634  
 MA77GG MANDELZWEIG, NUCLEAR PHYSICS A292 (1977) 333  
 MA77H MAUSNER ET AL, BULL. AM. PHYS. SOC. 22 (1977) 547

- MA77K MAIRLE ET AL, NUCLEAR PHYSICS A280 (1977) 97  
 MA77L MARTZ ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 177  
 MA77OO MARANGONI, OTTAVIANI AND SRRUIS, IN PROC., ZURICH - SIN,  
 1977, P. 241  
 MA77QQ MARIS AND SCHNEIDER, IN PROC., ZURICH - SIN, 1977, P. 224  
 MA77U MACKINTOSH AND CORDERO, PHYS. LETT. B68 (1977) 213  
 MA77UU MAYER-BORICKE, NUKLEONIKA 22 (1977) 1131  
 MA77WW MADEY ET AL, INT. J. RADIAT. ONCOLOGY BIOL. PHYS. (GB)  
 3 (1977) 325  
 MA77Y MAEDA, OKADA AND KATO, IN TOKYO (1977), CONTRIBUTED  
 PAPERS P. 562  
 MA78A MARQUARDT, SPRENGEL, BUTTLAR AND HOPPE, Z. PHYS. A285  
 (1978) 89  
 MA78B MARTZ ET AL, BULL. AM. PHYS. SOC. 23 (1978) 501  
 MA78C MALMIN, KOLATA, GRANT AND RISING, BULL. AM. PHYS. SOC.  
 23 (1978) 502  
 MA78G MALBROUGH ET AL, PHYS. REV. C17 (1978) 1395  
 MA78H MATSUSE, ABE AND KONDO, PROG. THEOR. PHYS. 59 (1978) 1037  
 MA78HH MALECKI, NAMYSLOWSKI, REALE AND MINETTI, RIV. NUOVO  
 CIM. 1, NO. 10 (1978) 1  
 MA78I MAKOWSKA-RZESZUTKO ET AL, PHYS. LETT. B74 (1978) 187  
 MA78II MATSUSE, PROC. INT. CONF. ON RESONANCES IN HEAVY ION  
 REACTIONS, HVAR, 1977 (NORTH-HOLLAND, AMSTERDAM, 1978),  
 P. 247; PHYSICS ABSTRACTS 9621 (1979)  
 MA78J MALMIN, KAHN AND PAUL, PHYS. REV. C17 (1978) 2097  
 MA78K MACKINTOSH AND KOBOS, J. OF PHYS. G4 (1978) L135  
 MA78L MATSUSE, ABE AND KONDO, PROG. THEOR. PHYS. 59 (1978) 1904  
 MA78N MALMIN, HARRIS AND PAUL, PHYS. REV. C18 (1978) 163  
 MA78P MAIRLE ET AL, NUCLEAR PHYSICS A299 (1978) 39  
 MA78V MATEJA ET AL, PHYS. REV. C18 (1978) 2622  
 MA79AA MAKOWSKA-RZESZUTKO, NUKLEONIKA 24 (1979) 495  
 MA79BB MAVROMATIS AND SKOURAS, NUCL. PHYS. A326 (1979) 83  
 MA79C MASON, GLOECKLER AND HOVESTADT, BULL. AM. PHYS. SOC. 24  
 (1979) 693  
 MA79CC MALFIET AND KARANT, PHYS. LETT. B86 (1979) 251  
 MA79E MARANGONI, OTTAVIANI AND SARUIS, INT. CONF. NUCL. PHYS.  
 WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 1.26  
 MA79F MAC GREGOR, PHYS. REV. LETT. 42 (1979) 1724  
 MA79FF MATTHEWS, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108  
 (1979) P. 369  
 MA79GG MARRS, POLLICK AND JACOBS, PHYS. REV. C20 (1979) 2308  
 MA79HH MARRS AND POLLICK, PHYS. REV. C20 (1979) 2446  
 MA79II MAHAUX, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979)  
 P. 265.  
 MA79J MAILLET, DEDONDER AND SCHMIT, NUCLEAR PHYSICS A316  
 (1979) 267  
 MA79KK MAYER-BORICKE, MAGY. FIZ. FOLY. 27 (1979) 465, AND  
 PHYS. ABSTRACTS 52642 (1980)  
 MA79LL MASSEN, INT. J. THEOR. PHYS. 18 (1979) 695  
 MA79MM MANAENKOV, JETP LETT. (USSR) 30 (1979) 334  
 MA79P MARIS, TEODORO AND VASCONCELLOS, NUCL. PHYS. A322  
 (1979) 461  
 MA79Q MADEY ET AL, IN CONF. PROC. TRIUMF, VANCOUVER (1979),  
 PAPER 1E22  
 MA80CC MARIS, TEODORO AND VEIT, PHYS. LETT. 94B (1980) 6

- MA80E MAIRLE ET AL, NUCL. PHYS. A339 (1980) 61  
 MA80EE MANAENKOV, SOVIET J. NUCL. PHYS. 31 (1980) 189  
 MA80J MAHER ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 723  
 MA80Y MADURGA, LOZANO AND JADRAQUE, PHYS. LETT. B95 (1980) 358  
 MA81 MAIRLE ET AL, NUCL. PHYS. (TO BE PUBLISHED) (1981)  
 MA81C MAIRLE ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE  
 PROC. 69 (1981) p. 739  
 MA81D MAIRLE ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE  
 PROC. 69 (1981) p. 685  
 MA81E MADEY ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 634  
 MA81G MASUDA, INOUE AND ITO, PHYS. REV. C23 (1981) 1543  
 MA81H MASUTANI AND YAZAKI, PROCEEDINGS VERSAILLES CONF. (1981)  
 p. 279  
 MA81K MADEY ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 538  
 MC66A MC GRATH, PHYS. REV. 145 (1966) 802  
 MC70B MC DONALD ET AL, PHYS. LETT. 31B (1970) 119  
 MC76I MC DONALD, ALEXANDER AND HAUSSER, NUCLEAR PHYSICS A273  
 (1976) 464  
 MC76M MCGRORY, IN 1974 GAMMA RAY TRANSITION PROBABILITIES (U. 3F  
 DELHI PRESS, 1976), P. 469  
 MC77 MC VOY, NUCLEAR PHYSICS A276 (1977) 491  
 MC77E MC DONALD ET AL, NUCLEAR PHYSICS A281 (1977) 325  
 MC77H MC DONALD ET AL, NUCLEAR PHYSICS A287 (1977) 189  
 MC78A MC GUIRE, VON ROSENVINGE AND MC DONALD, BULL. AM. PHYS.  
 SOC. 23 (1978) 509  
 MC78D MC DONALD AND ADELBERGER, PHYS. REV. LETT. 40 (1978) 1692  
 MC79B MC CALL, ASTROPHYS. J. 229 (1979) 962  
 MC79D MC VOY AND NEMES, BULL. AM. PHYS. SOC. 24 (1979) 816  
 MC80 McDONALD, NUCL. PHYS. A335 (1980) 463  
 MC80A MCGRORY, BULL. AMER. PHYS. SOC. 25 (1980) 573  
 MC80D MCCUTCHEON, DICKMAN, SHUTER AND ROGER, ASTROPHYS.  
 J. 237 (1980) 9  
 MC80F MCKLVEEN, BULL. AMER. PHYS. SOC. 25 (1980) 771  
 ME70 MENDELSON, HARDY AND CERNY, PHYS. LETT. 31B (1970) 126  
 ME78D MEYER, NAHABETIAN AND ELBAZ, IN PROCEEDINGS, CLUSTER,  
 WINNIPEG 1978, G4  
 ME78F MEYER, NATURE 272, NO. 5655 (1978) 675  
 ME78I MEYER, NAHABETIAN AND ELBAZ, LETT. NUOVO CIM. 22 (1978)  
 355  
 ME78J MEYER-TER-VEHN, Z. PHYS. A287 (1978) 241  
 ME79 MERMAZ ET AL, PHYS. REV. C19 (1979) 794  
 ME79Q MECKING, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108  
 (1979) P. 382  
 ME79R MEKJIAN, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979)  
 P. 149.  
 ME80 MEWALDT, SPALDING, STONE AND VOGT, BULL. AM. PHYS. SOC.  
 25 (1980) 546  
 ME80B MEAD AND CLARK, PHYS. LETT. 90B (1980) 331  
 ME80C MEKJIAN AND ZAMICK, PHYS. LETT. 92B (1980) 249  
 ME81 MERMAZ, PHYS. REV. C23 (1981) 755  
 ME81C MEYER AND ELBAZ, IN POLARIZATION, SANTA FE 1980, AIP  
 CONFERENCE PROC. 69 (1981) p. 1091  
 MI74L MINAMISONO, NOJIRI, MIZOBUCHI AND SUGIMOTO, NUCLEAR PHYSICS  
 A236 (1974) 416  
 MI75 MICHELETTI, PIGNANELLI AND GUZZONI, PHYS. REV. C11 (1975) 64

- MI75I MISKA ET AL, PHYS. LETT. B58 (1975) 155  
 MI75L MISKA ET AL, PHYS. LETT. B59 (1975) 441  
 MI76I MITO AND KAMIMURA, PROG. THEOR. PHYS. 56 (1976) 583  
 MI77C MILLER, PHYS. REV. C16 (1977) 2325  
 MI77E MILDNER ET AL, PHYS. LETT. B72 (1977) 159  
 MI78E MIGDAL, REV. MOD. PHYS. 50 (1978) 107  
 MI79D MISKA ET AL, PHYS. LETT. B83 (1979) 165  
 MI79I MILLER, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979)  
     P. 513.  
 MI80A MIKUMO ET AL, PHYS. REV. C21 (1980) 620  
 MI80B MIKUMO, UTTAC-28, 1979, AND NSSRP-26  
 MI80D MISCHKE ET AL, PHYS. REV. LETT. 44 (1980) 1197  
 MI80F MILLER AND NOBLE, PHYS. REV. C22 (1980) 1211  
 MI80H MILLENER, PHYS. REV. C22 (1980) 1355  
 MI81A MILLER, NUCL. PHYS. A353 (1981) 157c  
 MO71L MOSS AND COMITER, NUCLEAR PHYSICS A178 (1971) 241  
 MO73H MO, BLUE AND WELLER, NUCLEAR PHYSICS A197 (1973) 290  
 MO74C MOALEM, BENENSON AND CRAWLEY, NUCLEAR PHYSICS A236  
     (1974) 307  
 MO76 MOTOBAYASHI ET AL, PHYS. REV. LETT. 36 (1976) 390  
 MO76M MOLLER AND NIX, NUCLEAR PHYSICS A272 (1976) 502  
 MO77A MOLLER, HUF SCHMIDT AND KAMKE, NUCL. INSTR. METH. 140  
     (1977) 157  
 MO77K MOYLE, GLAGOLA, MATHEWS AND VIOLA, BULL. AM. PHYS. SOC. 22  
     (1977) 998  
 MO77P MOFFA, DOVER AND VARY, PHYS. REV. C16 (1977) 1857  
 MO77V MOSKALEV, SOVIET J. NUCL. PHYS. 25 (1977) 630  
 MO77W MORS, HERSCOVITZ AND JACOB, REV. BRAS. FIS. (BRAZIL) 7  
     (1977) 283, AND PHYS. ABSTRACTS 54762 (1978)  
 MO77Y MOTOBAYASHI, REP. INST. PHYS. & CHEM. RES. (JAPAN) 53 (1977)  
     81  
 MO78B MORRIS ET AL, PHYS. REV. C17 (1978) 227  
 MO78G MOORE ET AL, PHYS. LETT. B76 (1978) 192  
 MO78M MOSS, CORNELIUS AND BROWN, PHYS. REV. LETT. 41 (1978) 930  
 MO79E MOBIUS AND GRUHLE, Z. PHYS. A290 (1979) 81  
 MO79G MONTGOMERY, ERTEL AND OBERALL, INT. CONF. NUCL. PHYS.  
     WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 1.28  
 MO79P MONTGOMERY, ERTEL AND UBERALL, IN CONF. PROC. TRIUMF,  
     VANCOUVER (1979), PAPER 5C3  
 MO79V MORRISSEY ET AL, PHYS. REV. LETT. 43 (1979) 1139  
 MO79Y MOUGEY, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108  
     (1979) P. 124  
 MO79Z MOTOBAYASHI, KOHNO, OOI AND NAKAJIMA, NUCL. PHYS. A331  
     (1979) 193  
 MO80 MOUGEY, NUCL. PHYS. A355 (1980) 35  
 MO80N MORGAN, ASTROPHYS. J. 238 (1980) 674  
 MO80O MOLINARI, PHYS. REP. 64 (1980) 283  
 MO81 MOSS, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC.  
     69 (1981) p. 334  
 MO81E MONIZ, NUCL. PHYS. A354 (1981) 535c  
 MO81F MORRISON AND AMOS, PHYS. REV. C23 (1981) 1679  
 MO81K MOUGEY, NUCL. PHYS. A358 (1981) C293  
 MU73I MUGHABGHAB AND GARBER, BNL 325, THIRD EDITION, VOL. 1  
     (1973)  
 MU76B MUTHER, WAGHMARE AND FAESSLER, PHYS. LETT. B64 (1976) 125

- MU77 MUKHERJEE, SHYAM, PAL AND GANGULY, PHYS. REV. C15 (1977) 1238  
 MU77B MUKHOPADHYAY, PHYS. REPORTS 30C (1977) 1  
 MU78E MUTHER, NUKLEONIKA 23 (1978) 1141  
 MU79B MUKHOPADHYAY, TOKI AND WEISE, IN CONF. PROC. TRIUMF,  
       VANCOUVER (1979), PAPER 5B13  
 MU79C MUTHER, NUCL. PHYS. A328 (1979) 429  
 MU79D MUKHERJEE, AGARWAL, MAHESHWARI AND SOOD, NUCLEAR  
       INTERACTIONS, CANBERRA, AUSTRALIA (SPRINGER-VERLAG 1979)  
       p. 439, AND PHYS. ABSTRACTS 13504 (1980)  
 MU80 MUTHER, WAGHMARE AND FAESSLER, Z. PHYS. A 294 (1980) 87  
 MU80G MUTHER, FAESSLER, SHIMIZU AND WADIA, P. 13, PROCEEDINGS  
       OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS,  
       BERKELEY (1980)  
 MU81 MURPHY, BYRD, GUSS, FLOYD AND WALTER, NUCL. PHYS. A355  
       (1981) 1  
 MU81B MURPHY ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE  
       PROC. 69 (1981) p. 1478  
 MY80 MYERS AND SWIATECKI, NUCL. PHYS. A336 (1980) 267  
 NA75J NAIR ET AL, PHYS. REV. C12 (1975) 1575  
 NA76H NAIR, NAGATANI AND VOIT, PHYS. REV. LETT. 36 (1976) 1293  
 NA76P NAGORNYI AND INOPIN, Ukr. Fiz. Zh. 21 (1976) 1895; PHYS. ABS.  
       14090 (1977)  
 NA77 NATOWITZ ET AL, NUCLEAR PHYSICS A277 (1977) 477  
 NA77A NAGL AND UBERALL, BULL. AM. PHYS. SOC. 22 (1977) 617  
 NA77C NANN ET AL, PHYS. REV. C16 (1977) 1684  
 NA77D NAKANO, CAN. J. PHYS. 55 (1977) 134  
 NA77F NAGORCKA, SYMONS, TREACY AND MACLEAN, AUSTRALIAN J.  
       PHYS. 30 (1977) 149  
 NA77R NAGL AND UBERALL, IN PROC., ZURICH - SIN, 1977, P. 263  
 NA77S NADEZHDIN, PETROV AND SATAROV, JETP LETT. 26 (1977) 115  
 NA78 NAGL AND UBERALL, BULL. AM. PHYS. SOC. 23 (1978) 552  
 NA78E NAYAK AND SATPATHY, NUCLEAR PHYSICS A304 (1978) 64  
 NA78I NAGL, CANNATA AND UBERALL, ACTA PHYS. AUSTRIACA 48  
       (1978) 267  
 NA78K NADEZHDIN, PETROV AND SATAROV, YADERNAYA FIZ. 27 (1978) 1164  
 NA79 NAGATANI AND PENG, PHYS. REV. C19 (1979) 747  
 NA79C NAGL AND UBERALL, IN 1978 RPI SYMPOSIUM "PHOTOPION  
       NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 155  
 NA79E NATH, FIRK AND CHIU, IN PROC. THIRD SYMP. ON NEUTRON  
       CAPTURE GAMMA RAYS, 1978 (PLENUM PRESS, NEW YORK, 1979),  
       P. 701  
 NA79Q NAGL AND UBERALL, NUCLEAR INTERACTIONS, CANBERRA,  
       AUSTRALIA (SPRINGER-VERLAG 1979) p. 494, AND PHYS.  
       ABSTRACTS 18292 (1980)  
 NA80A NAGL AND UBERALL, BULL. AM. PHYS. SOC. 25 (1980) 606  
 NA80I NAVON ET AL, PHYS. LETT. B95 (1980) 365  
 NE64F NELSON, HUDSPETH AND BERNSTEIN, PHYS. REV. 136 (1964) B71  
 NE74D NEUBECK, SCHOBERT AND WAFFLER, PHYS. REV. C10 (1974) 320  
 NE77C NEGELE, BULL. AM. PHYS. SOC. 22 (1977) 1009  
 NE77D NEGELE, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE,"  
       ZURICH 1977, BIRKHAUSER VERLAG, P. 17  
 NE78 NEEDHAM ET AL, BULL. AM. PHYS. SOC. 23 (1978) 14  
 NE78A NEGELE AND RISKA, PHYS. REV. LETT. 40 (1978) 1005  
 NE78B NEEDHAM ET AL, BULL. AM. PHYS. SOC. 23 (1978) 943  
 NG80 NGO AND NGO, NUCL. PHYS. A348 (1980) 140

- NI78 NIILER AND BIRKMIRE, NUCL. INSTR. METH. 149 (1978) 301  
 NI80 NIILER AND BIRKMIRE, NUCL. INSTR. METH. 168 (1980) 105  
 NI80D NISHIOKA AND JOHNSON, PHYS. REV. C 22 (1980) 2457  
 NO57A NORBECK AND LITTLEJOHN, PHYS. REV. 108 (1957) 754  
 NO59A NORBECK, BLAIR, PINSONNEAULT AND GERBRACHT, PHYS. REV.  
     116 (1959) 1560  
 NO76C NOLEN AND AUSTIN, PHYS. REV. C13 (1976) 1773  
 NO76L NORGAARD AND FRICKE, ASTRON. & ASTROPHYS. 49 (1976) 337  
 NO77 NORGAARD, ASTROPHYS. J. 215 (1977) 200  
 NO77B NORUM, BERGSTROM AND CAPLAN, NUCLEAR PHYSICS A289 (1977) 275  
 NO77C NOLEN ET AL, PHYS. LETT. B71 (1977) 314  
 NO77D NORMAND, NUCLEAR PHYSICS A291 (1977) 126  
 NO78A NOVOTNY ET AL, NUCLEAR PHYSICS A294 (1978) 255  
 NO78E NORUM ET AL, BULL. AM. PHYS. SOC. 23 (1978) 960  
 NO78G NORDBORG, NILSSON, CONDE AND STROMBERG, NUCL. SCI. & ENG. 66  
     (1978) 75  
 NO79C NOMURA, PHYS. LETT. B85 (1979) 187  
 NO81 NORTHCLIFFE, CORNELIUS, YORK AND HIEBERT, IN POLARIZATION,  
     SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 1335  
 OB77 O'BRIEN, SALADIN, BAKTASH AND ALESSI, NUCLEAR PHYSICS  
     A291 (1977) 510  
 OB78A OBENSHAIN ET AL, PHYS. REV. C18 (1978) 764  
 OB80A OBENSHAIN ET AL, P. 701, PROCEEDINGS OF THE INTERNATIONAL  
     CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 OB81 OBENSHAIN ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 581  
 OB81A OBENSHAIN ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 290  
 OC78A O'CONNELL AND HANNA, PHYS. REV. C17 (1978) 892  
 OE79A OELERT ET AL, PHYS. REV. C20 (1979) 459  
 OF76 OFFERMANN, EY AND KUMMEL, NUCLEAR PHYSICS A273 (1976) 349  
 OG76 OGOLOBLIN, IN PROC. DUBNA CONF. JINR D-9920, VOL. II, P.  
     31 (1976)  
 OH76E OHM, RUDOLPH AND KRATZ, NUCLEAR PHYSICS A274 (1976) 45  
 OH77 OHNUMA, SUEHIRO, TANABE AND YAMADA, J. PHYS. SOC. JAPAN 42  
     (1977) 382  
 OH77C OHKUBO, KONDO AND NAGATA, PROG. THEOR. PHYS. 57 (1977) 82  
 OH78 OHTSUKA AND OHTSUBO, NUCLEAR PHYSICS A306 (1978) 513  
 OH79 OHASHI ET AL, IN 1978 RPI SYMPOSIUM "PHOTOPION NUCLEAR  
     PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 193  
 OH80 OHTA AND OKAI, PROGR. THEOR. PHYS. 63 (1980) 881  
 OK80D OKHRIMENKO AND STESHENKO, YAD. FIZ. (USSR) 32 (1980) 381  
 OL78A OLMER ET AL, BULL. AM. PHYS. SOC. 23 (1978) 941  
 OL80 OLSON ET AL, P. 200, PROCEEDINGS OF THE INTERNATIONAL  
     CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 ON78 ONISHI AND NEGELE, NUCLEAR PHYSICS A301 (1978) 336  
 OP76 OPHEL, FRAWLEY, TREACY AND BRAY, NUCLEAR PHYSICS A273  
     (1976) 397  
 OR78 ORTH, BUFFINGTON, SMOOT AND MAST, ASTROPHYS. J. 226  
     (1978) 1147  
 OR79 ORTIZ ET AL, PHYS. LETT. B84 (1979) 63  
 OR81 ORLANDINI, TRAINI AND DELLAGIACOMA, PROCEEDINGS VERSAILLES  
     CONF. (1981) pp. 117 AND 118  
 OS77A OSTERFELD, HNIZDO AND TOEPFFER, PHYS. LETT. B68 (1977) 319  
 OS77B OSMAN, INTER. J. THEOR. PHYS. 16 (1977) 575  
 OS79 OSET AND RHO, PHYS. REV. LETT. 42 (1979) 47  
 OS79C OSET, WEISE AND BROCKMANN, PHYS. LETT. B82 (1979) 344

- OS79I OSET AND WEISE, NUCL. PHYS. A329 (1979) 365  
 OS80C OSET AND WEISE, PHYS. LETT. 94B (1980) 19  
 OS81A OSET AND STROTTMAN, NUCL. PHYS. A355 (1981) 437  
 OS81G OST ET AL., NUCL. PHYS. A361 (1981) 453  
 OT76A OTSUBO ET AL., NUCLEAR PHYSICS A259 (1976) 452  
 OV80 OVERWAY, BECCHETTI AND JANECKE, P. 420, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 OV81 OVERWAY ET AL.  
 PA71F PALFFY, LEHMANN, GRENACS AND SUBOTOWICZ, NUOVO CIM. A3 (1971) 505  
 PA75A PALFFY ET AL., PHYS. REV. LETT. 34 (1975) 212  
 PA75O PAUL ET AL., IN JUL-CONF-16 (1975), P. 2  
 PA76I PARTRIDGE, TANG, THOMPSON AND BROWN, NUCLEAR PHYSICS A273 (1976) 341  
 PA77C PARKINSON, ASTRON. AND ASTROPHYS. 57 (1977) 185  
 PA77J PAPINEAU, NUCL. INSTR. METH. 146 (1977) 253  
 PA78 PARK, GREINER AND SCHEID, BULL. AM. PHYS. SOC. 23 (1978) 538  
 PA78A PAL, SRIVASTAVA AND GANGULY, IN PROCEEDINGS, CLUSTER, WINNIPEG 1978, B17  
 PA78B PANTIS, CAN. J. PHYS. 56 (1978) 659  
 PA78H PARTHASARATHY, ACTA PHYS. POLONICA B9 (1978) 483  
 PA79 PARK, SCHEID AND GREINER, BULL. AM. PHYS. SOC. 24 (1979) 665  
 PA79A PANTIS AND SPRUNG, CAN. J. PHYS. 57 (1979) 132  
 PA80 PARK, GREINER AND SCHEID, PHYS. REV. C21 (1980) 958  
 PA80A PARKS, STANLEY, COURTNEY AND KEMPER, PHYS. REV. C21 (1980) 217  
 PA80C PAUL ET AL., PHYS. REV. C21 (1980) 1802  
 PA80E PAL, CHATTERJEE AND GHOSE, NUCL. INSTR. METH. 171 (1980) 347  
 PA80G PARK AND SCHEID, P. 636, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 PE65 PEHL AND CERNY, PHYS. LETT. 14 (1965) 137  
 PE77 PETERSEN, DEHNHARD AND BAYMAN, PHYS. REV. C15 (1977) 1719  
 PE77B PERRIN ET AL., PHYS. LETT. B68 (1977) 55  
 PE77I PETRAUSKAS AND SABALYauskas, SOV. PHYS.-COLLECT. (USA) 17 (1977) 1, AND PHYS. ABSTRACTS 63627 (1979)  
 PE78A PETROVICH, STANLEY, PARKS AND NAGEL, PHYS. REV. C17 (1978) 1642  
 PE78C PETERSEN, SEQLIE AND ASCUITTO, BULL. AM. PHYS. SOC. 23 (1978) 947  
 PE78D PENG ET AL., PHYS. REV. C18 (1978) 2179  
 PE78I PETERSON, ASTROPHYS. J. 224 (1978) 595  
 PE79A PERROUD, IN 1978 RPI SYMPOSIUM "PHOTOPION NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 69  
 PE79B PEARSON, ROUBEN, SAUNIER AND BRUT, NUCLEAR PHYSICS A317 (1979) 447  
 PE79C PENG ET AL., BULL. AM. PHYS. SOC. 24 (1979) 842  
 PE79D PETERSEN, ASCUITTO AND SEQLIE, BULL. AM. PHYS. SOC. 24 (1979) 816  
 PE79E PENZIAS, REV. MOD. PHYS. 51 (1979) 425  
 PE80D PENZIAS, SCIENCE 208 (1980) 663  
 PE80J PETERSON AND HECHT, NUCL. PHYS. A344 (1980) 361  
 PE81 PENG, DEVRIES AND DIGIACOMO, PHYS. LETT. 98B (1981) 244  
 PE81B PETROVITCH AND LOVE, NUCL. PHYS. A354 (1981) 499c  
 PH77B PHILPOTT, NUCLEAR PHYSICS A289 (1977) 109

- PH77C PHAM AND DE SWINIARSKI, NUOVO CIM. A41 (1977) 543  
 PH78 PHILPOTT, MUKHOPADHYAY AND PURCELL, NUCLEAR PHYSICS A294 (1978) 6  
 PH79 PHILLIPS, ERB, BROMLEY AND WENESER, PHYS. REV. LETT. 42 (1979) 566  
 PH79C PHAM AND DE SWINIARSKI, Z. PHYS. A291 (1979) 327  
 PH79D PHILLIPS AND JOHNSON, PHYS. REV. C20 (1979) 1689  
 PH81A PHILPOTT, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 1144  
 PI73D PIGNANELLI ET AL, PHYS. REV. C8 (1973) 2120  
 PI74B PIGNANELLI ET AL, PHYS. REV. C10 (1974) 445  
 PI75F PILT, NUKLEONIKA (POLAND) 20 (1975) 801  
 PI77D PINKSTON, NUCLEAR PHYSICS A291 (1977) 342  
 PI78A PICKLESIMER AND WALKER, PHYS. REV. C17 (1978) 237  
 PI79F PICKAR ET AL, BULL. AM. PHYS. SOC. 24 (1979) 819  
 PI79G PINKSTON AND IANO, NUCL. PHYS. A330 (1979) 91  
 PL77 PLAGNOL, DOBBRE AND MARTY, PHYS. LETT. B67 (1977) 377  
 PL77A PLAVKO, IZV. AKAD. NAUK SSSR (SER. FIZ.) 41 (1977) 1293  
 PO73D POLETTI AND PRONKO, PHYS. REV. C8 (1973) 1285  
 PO76A POLING, NORBECK AND CARLSON, PHYS. REV. C13 (1976) 648  
 PO77C POVES, CEDILLO AND GOMEZ, NUCLEAR PHYSICS A293 (1977) 397  
 PO77E POVH, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 775  
 PO78 POOTH ET AL, NUCLEAR PHYSICS A294 (1978) 435  
 PO78A POVH, ANN. REV. NUCL. & PART. SCI. 28 (1978) 1  
 PO78B PODOSEK, ANN. REV. ASTRON. & ASTROPHYS. 16 (1978) 293  
 PO78C POSSNERT ET AL, PHYS. SCRIPTA 18 (1978) 353  
 PO79 POVH, BULL. AM. PHYS. SOC. 24 (1979) 69  
 PO79A POLLACCO, GARRETT AND WEISSER, PHYS. REV. C19 (1979) 2073  
 PO80 POVH, NUCL. PHYS. A335. (1980) 233  
 PR76A PRINCE, IN LOWELL CONF., CONF-760715-P2 (1976), P. 1083  
 PR77 PRUESS, NUCLEAR PHYSICS A278 (1977) 124  
 PR77A PRICE, STEVENSON AND FRANKEL, PHYS. REV. LETT. 39 (1977) 177  
 PR77G PREEDOM, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 119  
 PR77H PRIALNIK, SHARA AND SHAVIV, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 63  
 PR79A PROKOPENKO, SKLYARENKO, CHERNIEVSKII AND SHUSTOV, YADERNAYA FIZ. 29 (1979) 1223  
 PR79C PRIMAKOFF, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 3.  
 PR80B PRAKASH, SHLOMO, NILSSON AND BONDORF, P. 635, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 PR80C PROUDFOOT ET AL, NUCL. PHYS. A345 (1980) 278  
 PR81 PREEDOM ET AL, PHYS. REV. C23 (1981) 1134  
 PU81 PUGH ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 606  
 QA78 QAIM AND WOLFLE, NUCLEAR PHYSICS A295 (1978) 150  
 QU76A QUERCI AND QUERCI, ASTRON. & ASTROPHYS. 49 (1976) 443  
 QU80 QUENTIN, IN ATOMIC MASSES AND FUND. CONSTANTS, NOLEN AND BENENSON EDS., PLENUM (1980) P. 327  
 RA73C RANDOLPH ET AL, PHYS. LETT. B44 (1973) 36  
 RA76G RAMSTROM AND WIEDLING, NUCLEAR PHYSICS A272 (1976) 259  
 RA76H RADOMSKI AND RISKA, NUCLEAR PHYSICS A274 (1976) 428

- RA77 RAMSTROM AND WIEDLING, ASTROPHYS. J. 211 (1977) 223
- RA78F RAMAN, Houser, Walkiewicz and Towner, ATOMIC DATA & NUCL. DATA TABLES 21 (1978) 567, AND IBID. 22 (1978) 369
- RA78I RAGNARSSON, NILSSON AND SHELTON, PHYS. REPORTS 45 (1978) 1
- RA79C RAE ET AL., NUCLEAR PHYSICS A319 (1979) 239
- RA79F RAO, GUPTA, SINGH AND PRAKASH, J. PHYS. SOC. JAPAN 46 (1979) 1683
- RA79H RAMATY, KOZLOVSKY AND LINGENFELTER, ASTROPHYS. J., SUPPL. SERIES 40 (1979) 487
- RA79I RASCHER, MULLER AND LIEB, PHYS. REV. C20 (1979) 1028
- RA79K RAY, PHYS. REV. C20 (1979) 1857
- RA80 RAY, NUCL. PHYS. A335 (1980) 443
- RA80E RAE ET AL., PHYS. REV. LETT. 45 (1980) 884
- RA80F RACCA ET AL., BULL. AMER. PHYS. SOC. 25 (1980) 733
- RA81B RAY, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 295
- RA81F RAYNAL, PHYS. REV. C23 (1981) 2571
- RE75D REISDORF, LAU AND VANDENBOSCH, NUCLEAR PHYSICS A253 (1975) 490
- RE77 REHM, ESSEL AND KORNER, BULL. AM. PHYS. SOC. 22 (1977) 39
- RE77B REISS ET AL., NUCLEAR PHYSICS A283 (1977) 149
- RE78B REHM, HENNING, ERSKINE AND KOVAR, PHYS. REV. LETT. 40 (1978) 1479
- RE78E REMLER AND SATHE, PHYS. REV. C18 (1978) 2293
- RE79A REDWINE, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 501
- RE79C RENNER ET AL., BULL. AM. PHYS. SOC. 24 (1979) 842
- RE79D RENNER ET AL., BULL. AM. PHYS. SOC. 24 (1979) 843
- RE80 REAMES, VON ROSENVINGE AND MC GUIRE, BULL. AMER. PHYS. 25 (1980) 597
- RH79 RHO, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 105.
- RH80 RHOADES-BROWN, MACFARLANE AND PIEPER, PHYS. REV. C21 (1980) 2436
- RI78 RISKA AND NEGELE, BULL. AM. PHYS. SOC. 23 (1978) 29
- RO73AA ROLFS, NUCLEAR PHYSICS A217 (1973) 29
- RO74J ROGERS ET AL., NUCLEAR PHYSICS A226 (1974) 424
- RO74K ROGERS ET AL., NUCLEAR PHYSICS A226 (1974) 445
- RO74S ROLFS AND RODNEY, NUCLEAR PHYSICS A235 (1974) 450
- RO76R ROGERS, NUCLEAR PHYSICS A271 (1976) 447
- RO76T ROLFS, RODNEY AND WINKLER, IN ATOMIC MASSES AND FUNDAMENTAL CONSTANTS, ED. BY SANDERS AND WAPSTRA (PLENUM PRESS, 1976), P. 60
- RO76U ROTTER, BARZ AND HOHN, SOVIET J. NUCL. PHYS. 24 (1976) 266
- RO77AA ROLFS, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 133
- RO77I ROHWER, MORIKE AND STAUDT, Z. PHYS. A281 (1977) 107
- RO77K ROTTER, PHYS. LETT. B67 (1977) 385
- RO77U ROSENFELDER, NUCLEAR PHYSICS A290 (1977) 315
- RO77W ROESEL, TRAUTMANN AND VIOLIER, NUCLEAR PHYSICS A292 (1977) 523
- RO78B ROLFS AND TRAUTVETTER, ANN. REV. NUCL. & PART. SCI. 28 (1978) 115
- RO78E ROZAK, MONTESTRUQUE AND DARDEN, BULL. AM. PHYS. SOC. 23 (1978) 501
- RO78H ROWE AND VOGT, NUCLEAR PHYSICS A302 (1978) 511

- R078J ROTTER, BARZ AND HOHN, NUCLEAR PHYSICS A297 (1978) 237  
 R078L ROSENFELDER, NUCLEAR PHYSICS A298 (1978) 397  
 R078O ROLFS, AIP CONF. PROC. 47 (1978) 197  
 R078P ROBSON, NUCLEAR PHYSICS A308 (1978) 381  
 R079B ROMERO ET AL, BULL. AM. PHYS. SOC. 24 (1979) 656  
 R079C ROTTER, J. OF PHYS. G5 (1979) 251  
 R079D ROTTER, J. OF PHYS. G5 (1979) 685  
 R079E ROSENFELDER, IN 1978 RPI SYMPOSIUM "PHOTOPION NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 367  
 R079J ROY, KEMPER AND FRAWLEY, BULL. AM. PHYS. SOC. 24 (1979) 851  
 R079L ROTTER, J. OF PHYS. G5 (1979) 1575  
 R079M ROBERT, MERLIVAT AND JAVOY, NATURE 282, NO. 5741 (1979) 785  
 R079P ROBSON, PHYS. REV. LETT. 42 (1979) 876  
 R080G ROTBERG AND MITTIG, PHYS. REV. C22 (1980) 1574  
 R081C ROMAN, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 282  
 R081J ROSENFELDER, PROCEEDINGS VERSAILLES CONF. (1981) p. 246  
 R081L ROSENSTEIN AND ROWE, PHYS. REV. LETT. 47 (1981) 223  
 R081M ROCKMORE AND KAMAL, PHYS. REV. C24 (1981) 236  
 SA75I SANCHEZ, CASANOVA AND CASANOVA, AN. FIS. (SPAIN) 71 (1975) 119  
 SA76I SATO, NUCLEAR PHYSICS A269 (1976) 378  
 SA76M SAYED AND SHALABY, ATOMKERNENERGIE 28 (1976) 217  
 SA77A SAGE AND CUSSON, BULL. AM. PHYS. SOC. 22 (1977) 644  
 SA77AA SADOVOI, SOVIET J. NUCL. PHYS. 25 (1977) 27  
 SA77B SACKMANN, ASTROPHYS. J. 212 (1977) 159  
 SA77C SANDERS, MARTZ AND PARKER, IN TOKYO (1977), CONTRIBUTED PAPERS P. 625  
 SA77CC SAITO, PROG. THEOR. PHYS. SUPPL. 62 (1977) 11  
 SA77DD SAPERSHTAIN, TOLOKONNIKOV AND FAYANS, SOVIET J. NUCL. PHYS. 25 (1977) 510  
 SA77EE SANCHEZ-DEHESA, REPORT JUL-1425 (1977), AND PHYS. ABSTRACTS 538 (1978)  
 SA77G SALONER AND TOEPFFER, NUCLEAR PHYSICS A283 (1977) 108  
 SA77GG SADIKOVSKII, FEOFILOV, DENISOV AND KOLALIS, YADERNAYA FIZ. 26 (1977) 1200  
 SA77H SALONER, TOEPFFER AND FINK, NUCLEAR PHYSICS A283 (1977) 131  
 SA77J SANDERS, PH.D. DISSERTATION, YALE U. (1977)  
 SA77P SATO, IN TOKYO (1977), CONTRIBUTED PAPERS P. 589  
 SA77V SANTANA, COELHO AND DAS, PHYS. REV. C16 (1977) 1785  
 SA78C SANDORFI, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 185  
 SA78I SATCHLER ET AL, NUCLEAR PHYSICS A298 (1978) 313  
 SA78K SAKAGUCHI ET AL, FIZIKA (YUGOSLAVIA) 10, SUPPL. (1978) 53  
 SA78N SAPERSHTAIN AND TOLOKONNIKOV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 42 (1978) 1890  
 SA79C SACKMANN, OAP-551 (1979)  
 SA79I SANDORFI, CALARCO, RAND AND SCHWETTMAN, BULL. AM. PHYS. SOC. 24 (1979) 851  
 SA79K SANDERS, MARTZ AND PARKER, PHYS. REV. C20 (1979) 1743  
 SA79L SATCHLER, NUCL. PHYS. A329 (1979) 233  
 SA79M SAINT-LAURENT ET AL, NUCL. PHYS. A327 (1979) 517  
 SA79N SANDORFI, IN BNL-51115 (1979), P. 405  
 SA79P SAKAGUCHI ET AL, PHYS. LETT. 89B (1979) 40  
 SA79U SAVUSHKIN, YAD. FIZ. (USSR) 30 (1979) 660

- SA80B SAMANTA ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 593  
 SA80C SARTOR AND MAHAUX, PHYS. REV. C21 (1980) 2613  
 SA80H SANDHYA DEVI ET AL, P. 415, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SA80I SASAGASE ET AL, P. 527, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SA80J SAMANTA ET AL, P. 878, PROCEEDINGS OF THE INTERNATIONAL  
     CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SA80M SATO ET AL, P. 536, PROCEEDINGS OF THE INTERNATIONAL  
     CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SA81A SAMADDAR, SPERBER AND ZIELINSKA-PFABE, PHYS. LETT. 98B  
     (1981) 340  
 SA81C SAKAGUCHI ET AL, IN POLARIZATION, SANTA FE 1980, AIP  
     CONFERENCE PROC. 69 (1981) p. 445  
 SA81E SANDORFI ET AL, PHYS. REV. LETT. 46 (1981) 884  
 SA81F SARTOR, FAESSLER, KHADKIKAR AND KREWALD, NUCL. PHYS. A359  
     (1981) 467  
 SA81G SAHARIA, WOLOSHYN AND KISSLINGER, PHYS. REV. C23 (1981) 2140  
 SA81I SARTOR AND STANCU, PROCEEDINGS VERSAILLES CONF. (1981) p. 143  
 SC69U SCHAEFER, KLEMM AND HARRIS, PHYS. REV. 181 (1969) 137  
 SC72C SCHLÖTHAUER-VOOS ET AL, NUCLEAR PHYSICS A186 (1972) 225  
 SC73M SCHUMACHER ET AL, NUCLEAR PHYSICS A212 (1973) 573  
 SC75Q SCHNEIDER, KOHLMAYER, PUHLHOFER AND BOCK, NUCLEAR  
     PHYSICS A251 (1975) 331  
 SC77D SCHOCH ET AL, BULL. AM. PHYS. SOC. 22 (1977) 591  
 SC77E SCHIMIZU, FUJII AND TAKIMOTO, PROG. THEOR. PHYS. 57  
     (1977) 680  
 SC77I SCALO, OAP-498 (1977)  
 SC77O SCHRODER AND HUIZENGA, ANN. REV. NUCL. SCI. 27 (1977) 465  
 SC78A SCHIMIZU ET AL, J. PHYS. SOC. JAPAN 44 (1978) 7  
 SC78B SCHIFFER, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO,  
     1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 9  
 SC78C SCHULTHEIS ET AL, IN PROCEEDINGS, CLUSTER, WINNIPEG 1978, C3  
 SC78E SCHOCH, PHYS. REV. LETT. 41 (1978) 80  
 SC78G SCHNEIDER, NUCLEAR PHYSICS A300 (1978) 313  
 SC78I SCHWESINGER, Z. PHYS. A287 (1978) 373  
 SC78L SCHULTHEIS AND SCHULTHEIS, PHYS. REV. LETT. 41 (1978) 1020  
 SC78M SCHEID AND GREINER, IN CLUSTERING, WINNIPEG 1978, AIP  
     CONF. PROC. 47 (1978) 374  
 SC79B SCHMITT, BORKOWSKI, SIMON AND WALTHER, INT. CONF. NUCL.  
     PHYS. WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 6.9  
 SC79C SCHWEIZER ET AL, PHYS. REV. C19 (1979) 1408  
 SC79D SCHEURER ET AL, NUCLEAR PHYSICS A319 (1979) 274  
 SC80 SCHIFFER, NUCL. PHYS. A335 (1980) 339  
 SC80D SCHOCH AND GORINGER, P. 682, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SC80E SCHULTHEIS AND SCHULTHEIS, P. 267, PROCEEDINGS OF THE  
     INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SC80F SCOTT, LECTURES PRESENTED AT THE INTERNATIONAL SCHOOL ON  
     NUCLEAR PHYSICS, ALUSHTA, CRIMEA, USSR, 14-25 APRIL 1980  
 SC80I SCHOCH ET AL, PHYS. REV. C 22 (1980) 2630  
 SC80J SCHWANNER ET AL, PHYS. LETT. B96 (1980) 268  
 SC81B SCHLOEMER ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 610  
 SC81D SCHOBBERT, GERSCH AND WIEBICKE, J. OF PHYS. G7 (1981) L73  
 SC81G SCHOCH, NUCL. PHYS. A358 (1981) C307

- SE63E SEGEL, SINGH, ALLAS AND HANNA, PHYS. REV. LETT. 10 (1963) 345  
 SE77 SERCELY, SHEPARD, SPARROW AND CECIL, BULL. AM. PHYS. SOC. 22 (1977) 552  
 SE78A SERCELY, PETERSON AND FLYNN, PHYS. REV. C17 (1978) 1919  
 SE78D SEGLIE, ASCUITTO AND PETERSEN, BULL. AM. PHYS. SOC. 23 (1978) 947  
 SE78F SENS, REFAEI AND PAPE, PHYS. REV. C18 (1978) 2007  
 SE79 SEILER AND ROSEN, NUCLEAR PHYSICS A315 (1979) 45  
 SE80E SEKI, MASUTANI, OKA AND YAZAKI, PHYS. LETT. B97 (1980) 200  
 SE81C SEYA, KOHNO AND NAGATA, PROG. THEOR. PHYS. 65 (1981) 204  
 SH75H SHERA, PHYS. REV. C12 (1975) 1003  
 SH77I SHERR, PHYS. REV. C16 (1977) 1159  
 SH77K SHAPIRA ET AL, PHYS. LETT. B71 (1977) 293  
 SH77M SHURPIN, MUTHER, KUO AND FAESSLER, NUCLEAR PHYSICS A293 (1977) 61  
 SH77N SHODA ET AL, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 482  
 SH77O SHIBATA ET AL, IN PROC., ZURICH - SIN, 1977, P. 166  
 SH78A SHAPIRA ET AL, PHYS. REV. LETT. 40 (1978) 371  
 SH78D SHLOMO AND WAGNER, Z. PHYS. A285 (1978) 283  
 SH78E SHIMIZU AND FAESSLER, PHYS. REV. C17 (1978) 1891  
 SH78F SHKOLNIK ET AL, PHYS. LETT. B74 (1978) 195  
 SH78G SHIMIZU AND FAESSLER, Z. PHYS. A287 (1978) 299  
 SH78I SHAPIRA ET AL, BULL. AM. PHYS. SOC. 23 (1978) 949  
 SH78J SHIMIZU AND FAESSLER, NUCLEAR PHYSICS A306 (1978) 311  
 SH78K SHEPARD AND KING, BULL. AM. PHYS. SOC. 23 (1978) 953  
 SH78O SHALABY AND KHALIL, J. OF PHYS. G4 (1978) L293  
 SH78P SHABRATOVA ET AL, ACTA PHYS. SLOV. 28 (1978) 132  
 SH78Q SHLOMO, REP. PROG. PHYS. (GB) 41 (1978) 957  
 SH79G SHAKIN, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 550  
 SH79J SHODA, NAHARA, OHASHI AND YAMAZAKI, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 5A4  
 SH79T SHIRIKOVA AND SHITIKOVA, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2396  
 SH80C SHKOLNIK, DEHNHARD AND FRANEY, BULL. AMER. PHYS. SOC. 25 (1980) 592  
 SH80E SHAPIRA, FORD, GOMEZ DEL CAMPO AND STELSON, PHYS. REV. C21 (1980) 1824  
 SH80F SHU ET AL, PHYS. REV. C21 (1980) 1828  
 SH80L SHKOLNIK, FRANEY AND DEHNHARD, P. 588, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SH80N SHITICOVA, NUCL. PHYS. A331 (1979) 365  
 SH81E SHEPARD AND ROST, PHYS. REV. LETT. 46 (1981) 1544  
 SH81H SHIMIZU, POLLS, MUTHER AND FAESSLER, PROCEEDINGS VERSAILLES CONF. (1981) p. 428  
 SI75F SIMONIUS, PROC. INTERN. SYMP. ON INTERACTION STUDIES IN NUCLEI, GERMANY, 1975 (AMSTERDAM - NORTH-HOLLAND 1975), P. 3; PHYS. ABS. 89907 (1976)  
 SI76G SILBERBERG, TSAO AND SHAPIRO, IN SPALLATION NUCLEAR REACTIONS AND THEIR APPLICATIONS, EDS. SHEN AND MERKER (D. REIDEL PUB. CO., THE NETHERLANDS, 1976), P. 49  
 SI76I SINGH AND SCHWANDT, NUKLEONIKA 21 (1976) 451  
 SI77 SILBAR, GINOCCHIO AND STERNHEIM, PHYS. REV. C15 (1977) 371  
 SI77D SILBERBERG AND TSAO, ASTROPHYS. J. (SUPPL. SERIES) 35 (1977) 129

- SI77E SILBERBERG AND TSAO, ASTROPHYS. J. 35 (1977) 137  
 SI78B SINHA AND MOSzkowski, NUCLEAR PHYSICS A302 (1978) 237  
 SI78D SINGH, INDIAN J. PHYS. A52 (1978) 389  
 SI79C SINHA, PHYS. REV. LETT. 42 (1979) 690  
 SI79F SINHA AND MOSzkowski, PHYS. LETT. B81 (1979) 289  
 SI80C SILVER AND AUSTERN, PHYS. REV. C21 (1980) 272  
 SI80D SINHA, PHYS. REV. LETT. 44 (1980) 1207  
 SI80G SINGLETON, MANAGOLI AND GOLDFARB, NUCL. PHYS. A342 (1980) 312  
 SI80I SINHA, P. 596, 598, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SI81 SIE, NEWTON, LEIGH AND DIAMOND, PHYS. REV. LETT. 46 (1981) 405  
 SI81B SICK, NUCL. PHYS. A354 (1981) 37c  
 SI81E SINHA, PROCEEDINGS VERSAILLES CONF. (1981) p. 465  
 SJ80A SJOREEN ET AL, P. 684, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 SJ81 SJOREEN ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 193  
 SK74 SKWIERSKY, BAGLIN AND PARKER, PHYS. REV. C9 (1974) 910  
 SK75A SKOPIK, SHIN AND MURPHY, CAN. J. PHYS. 53 (1975) 1398  
 SL76A SLAUS, IN LOWELL CONF., CONF-760715-P1 (1976), P. 272  
 SL78 SLAUS, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 57  
 SL79A SLOBODRIAN ET AL, Z. PHYS. A291 (1979) 261  
 SM77 SMIRNOV AND TCHUVIL'SKY, PHYS. REV. C15 (1977) 84  
 SM77C SMITH ET AL, PHYS. LETT. B72 (1977) 176  
 SM78D SMIRNOV AND FILIPPOV, SOVIET J. NUCL. PHYS. 27 (1978) 39  
 SN69B SNYDER AND WAGGONER, PHYS. REV. 186 (1969) 999  
 SN74 SNOVER, ADELBERGER AND BROWN, PHYS. REV. LETT. 32 (1974) 1061  
 SN79 SNOVER, IKOSSI AND TRAINOR, PHYS. REV. LETT. 43 (1979) 117  
 SN79A SNOVER, IN PROC. THIRD SYMP. ON NEUTRON CAPTURE GAMMA RAYS, 1978 (PLENUM PRESS, NEW YORK, 1979), P. 319  
 SN81 SNOVER, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 321  
 SO77A SOTONA, CZECH. J. PHYS. B27 (1977) 386, AND PHYS. ABSTRACTS 41877 (1977)  
 SO78B SOGA ET AL, PHYS. REV. C18 (1978) 2457  
 SO78C SOTONA AND ZOFKA, CZECH. J. PHYS. B28 (1978) 593  
 SO78D SOKOL, IZV. AKAD. NAUK SSSR (SER. FIZ.) 42 (1978) 1829  
 SO79A SOGA ET AL, BULL. AM. PHYS. SOC. 24 (1979) 614  
 SO79C SOLIN, NEMLIK, KUZMIN AND ZHEREBTSOVA, YADERNAYA FIZ. 29 (1979) 289  
 SP77D SPENCER, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 153  
 SP78 SPARROW, BULL. AM. PHYS. SOC. 23 (1978) 591  
 SP79A SPERGEL, LAZARETH AND LEVY, BULL. AM. PHYS. SOC. 24 (1979) 660  
 SP80 SPARROW, PHYS. REV. LETT. 44 (1980) 625  
 SP80C SPETH, KLEMT, WAMBACH AND BROWN, NUCL. PHYS. A343 (1980) 382  
 SP80D SPETH AND WAMBACH, NUCL. PHYS. A347 (1980) 389  
 SP81A SPETH AND SUZUKI, NUCL. PHYS. A358 (1981) C139  
 SR79A SRINIVASA RAO AND SUSILA, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 430  
 ST76G STANCU AND BRINK, NUCLEAR PHYSICS A270 (1976) 236

- ST76I STOCK ET AL, PHYS. REV. C14 (1976) 1824  
 ST76L STROBEL, NUCLEAR PHYSICS A271 (1976) 162  
 ST76N STRAZZERI, NUOVO CIM. A34 (1976) 229, AND IBID. A41 (1977)  
     592  
 ST77 STABLER, BUCK, STAUDT AND OESCHLER, NUCLEAR PHYSICS  
     A275 (1977) 269  
 ST77E STEVENSON, PRICE AND FRANKEL, PHYS. REV. LETT. 38  
     (1977) 1125  
 ST77N STRAYER AND WERBY, J. OF PHYS. G3 (1977) L179  
 ST77O STEPHENSON, GIBBS AND GIBSON, BULL. AM. PHYS. SOC. 22  
     (1977) 1029  
 ST77P STAHEL ET AL, PHYS. REV. C16 (1977) 1456  
 ST78A STECK, PHYS. REV. C17 (1978) 1034  
 ST78G STANCU, J. PHYSIQUE 39 (1978) 799  
 ST78I STARRFIELD, TRURAN AND SPARKS, ASTROPHYS. J. 226 (1978) 186  
 ST78J STRINGARI ET AL, NUCLEAR PHYSICS A309 (1978) 189  
 ST78L STOKSTAD, S. AFRICAN J. PHYS. 1 (1978) 191  
 ST79 STRICKER, MC MANUS AND CARR, PHYS. REV. C19 (1979) 929  
 ST79D STANLEY, PHILPOTT, HALDERSON AND PETROVICH, BULL. AM.  
     PHYS. SOC. 24 (1979) 592  
 ST79J STANLEY, GLOVER AND PETROVICH, BULL. AM. PHYS. SOC. 24  
     (1979) 816  
 ST79L STEHLING ET AL, HELV. PHYS. ACTA 52 (1979) 93  
 ST79M STOCK, NUKLEONIKA 24 (1979) 165  
 ST79Q STRINGARI AND VAUTHERIN, PHYS. LETT. B88 (1979) 1  
 ST80D STOLER ET AL, PHYS. REV. C22 (1980) 911  
 ST80G STRICKER, CARR AND MC MANUS, PHYS. REV. C 22 (1980) 2043  
 ST80I STRIKMAN AND FRANKFURT, YAD. FIZ. 32 (1980) 1403  
 ST81B STOKER ET AL, NUCL. INSTR. & METH. 180 (1981) 515  
 SU76E SUZUKI, PROG. THEOR. PHYS. 56 (1976) 111  
 SU77E SUZUKI, HYUGA AND HORIKAWA, IN TOKYO (1977), CONTRIBUTED  
     PAPERS P. 176  
 SU77F SUZUKI, IN TOKYO (1977), CONTRIBUTED PAPERS P. 180  
 SU77G SUZUKI AND ROWE, NUCLEAR PHYSICS A286 (1977) 307  
 SU77H SUNG ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 802  
 SU77K SUZUKI, ANDO AND IMANISHI, INS-REPORT-299 (1977)  
 SU78 SUBRAMANIAN ET AL, BULL. AM. PHYS. SOC. 23 (1978) 12  
 SU78C SUNKEL, IN PROCEEDINGS, CLUSTER, WINNIPEG 1978, A20  
 SU78D SUZUKI AND IMANISHI, IN PROCEEDINGS, CLUSTER, WINNIPEG  
     1978, A2  
 SU78E SUBRAMANIAN ET AL, BULL. AM. PHYS. SOC. 23 (1978) 943  
 SU78F SUZUKI AND IMANISHI, INS REPORT 324 (1978)  
 SU79A SUZUKI, HYUGA, ARIMA AND HORIKAWA, INT. CONF. NUCL. PHYS.  
     WITH ELECTRO-MAGNETIC INTERACTIONS, MAINZ (1979), 7.8  
 SU79D SUBRAMANIAN AND DEVANATHAN, PHYS. REV. C20 (1979) 1615  
 SU79G SUZUKI AND KUBO, IN TSUKUBA SYMPOSIUM ON  
     POLARIZATION PHENOMENA, NOV. 1979, ED. K. YAGI, P. 211  
 SU80C SUZUKI ET AL, PHYS. LETT. B95 (1980) 202  
 SU80D SUZUKI, PROG. THEOR. PHYS. 64 (1980) 2041  
 SU81A SUZUKI AND IMANISHI, PHYS. REV. C23 (1981) 2414  
 SW70 SWANN, NUCLEAR PHYSICS A150 (1970) 300  
 SW74A SWITKOWSKI, WIELAND AND WINther, PHYS. REV. LETT 33 (1974)  
     840  
 SW77 SWITKOWSKI, WU, OVERLEY AND BARNES, NUCLEAR PHYSICS A289  
     (1977) 236

- SW77B SWITKOWSKI, STOKSTAD AND WIELAND, NUCLEAR PHYSICS A279 (1977) 502
- SW79 SWEIGART AND MENDEL, ASTROPHYS. J. 229 (1979) 624
- TA69B TAUBER, HOFMANN, MAUSBERG AND ROSSLE, Z. PHYS. 227 (1969) 71
- TA77 TARAS ET AL., PHYS. REV. C15 (1977) 834
- TA78B TABOR ET AL., PHYS. REV. C17 (1978) 2136
- TA78C TAKIGAWA, LEE AND MARTY, PHYS. LETT. B76 (1978) 187
- TA78E TARAS, RAO AND AZUELOS, PHYS. REV. LETT. 41 (1978) 840
- TA78F TAKEMASA AND YOSHIDA, NUCLEAR PHYSICS A304 (1978) 229
- TA78J TAKEMASA AND TAMURA, PHYS. REV. C18 (1978) 1282
- TA78M TARAS, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 234
- TA78S TAAM AND PICKLUM, ASTROPHYS. J. 224 (1978) 210
- TA78U TANG, LEMERE AND THOMPSON, PHYS. REPORTS 47 (1978) 167
- TA79C TANIMURA AND TAZAWA, PHYS. LETT. B83 (1979) 22
- TA79E TANIMURA AND TAZAWA, NUCLEAR PHYSICS A321 (1979) 490
- TA79F TANIMURA AND TAZAWA, PHYS. REV. C20 (1979) 183
- TA79H TANAKA, ODA, PETROVICH AND SHELINE, PHYS. LETT. B83 (1979) 279
- TA80A TABOR, DENNIS, NEUSCHAEFER AND THEISEN, BULL. AM. PHYS. SOC. 25 (1980) 524
- TA80C TANIMURA, PHYS. LETT. 90B (1980) 204
- TA80D TAVARES ET AL., LETT. NUOVO CIM. 27 (1980) 358
- TA80E TANIMURA AND TAZAWA, PHYS. REPORTS 61 (1980) 253
- TA80K TAKIGAWA, PLAGNOL AND MARTY, P. 427, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- TA80N TANIMURA AND TAZAWA, NUCL. PHYS. A346 (1980) 266
- TA80P TADIC, REP. PROG. PHYS. 43 (1980) 67
- TA81 TAKADA ET AL., PHYS. REV. C23 (1981) 772
- TA81A TAZAWA AND TANIMURA, PHYS. REV. LETT. 46 (1981) 408
- TA81G TAKAKI, PROG. THEOR. PHYS. 65 (1981) 241
- TA81I TAM ET AL., NUCL. PHYS. A361 (1981) 412
- TA81J TANIMURA AND MOSEL, PHYS. REV. C24 (1981) 321
- TE77B TERRIEN, IN TOKYO (1977), CONTRIBUTED PAPERS P. 843
- TE78A TERENETSKI AND GARRETT, PHYS. REV. C18 (1978) 1944
- TE79 TERLECKI ET AL., MICROSCOPIC OPTICAL POTENTIALS, HAMBURG, GERMANY (SPRINGER-VERLAG 1979) P. 410, AND PHYS. ABSTRACTS 97730 (1979)
- TH69A THROOP, PHYS. REV. 179 (1969) 1011
- TH74 THIBAULT AND KLAISCH, PHYS. REV. C9 (1974) 793
- TH77C THOMPSON AND GRAY, INTERN. J. APPL. RAD. AND ISOTOPES 28 (1977) 411
- TH77D THOMPSON, LEMERE AND TANG, NUCLEAR PHYSICS A286 (1977) 53
- TH77F THIES, PHYS. LETT. B70 (1977) 401
- TH78D THOMAS, PIJOLAT AND FALLAVIER, REV. PHYS. APPL. 13 (1978) 433
- TH78E THOMPSON, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC. 47 (1978) 69
- TH80 THIESSEN, NUCL. PHYS. A335 (1980) 329
- TH80A THOMAS AND LANDAU, PHYS. REPORTS 58 (1980) 121
- TH81A THORNTON, PARKS, CORDELL AND WIEDNER, BULL. AMER. PHYS. SOC. 26 (1981) 611
- TH81B THOMAS, NUCL. PHYS. A354 (1981) 51c
- TH81D THIES, PROCEEDINGS VERSAILLES CONF. (1981) p. 426
- TI79A TINSLEY, ASTROPHYS. J. 229 (1979) 1046

- T071H TOWNER, WARBURTON AND GARVEY, ANN. OF PHYS. 66 (1971) 674  
 T077 TOWNSLEY ET AL, PHYS. REV. C15 (1977) 281, 1607  
 T077D TORNOW, REINHARD AND DRECHSEL, Z. PHYS. A280 (1977) 253  
 T077G TOMBRELLO, OAP-501/LIAP (1977)  
 T077I TOYAMA, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 450  
 T077J TONEEV AND GUDIMA, IN PROC., ZURICH - SIN, 1977, P. 181  
 T078C TOHSAKI-SUZUKI, PROG. THEOR. PHYS. 60 (1978) 1013  
 T078D TOMKIN AND LAMBERT, ASTROPHYS. J. 223 (1978) 937  
 T079 TOWNER AND KHANNA, PHYS. REV. LETT. 42 (1979) 51  
 T079D TOKI AND WEISE, Z. PHYS. A 292 (1979) 389  
 T080D TOHSAKI-SUZUKI, KAMIMURA AND IKEDA, P. 411, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCL. PHYS., BERKELEY (1980)  
 T080E TORNOW ET AL, NUCL. PHYS. A348 (1980) 157  
 T080F TOHSAKI-SUZUKI, KAMIMURA AND IKEDA, SUPPL. PROGR. THEOR. PHYS. 68 (1980) 359  
 T081B TOWNSEND AND DEUTCHMAN, NUCL. PHYS. A355 (1981) 505  
 T081E TOWNER AND KHANNA, PROCEEDINGS VERSAILLES CONF. (1981) p. 400  
 TR77A TRIPATHI, SANDERSON AND ELLIOTT, NUCLEAR PHYSICS A278 (1977) 78  
 TR77C TRABER, TRAUTMANN AND ALDER, HELV. PHYS. ACTA 50 (1977) 195  
 TR77G TRABER, TRAUTMANN AND ROSEL, NUCLEAR PHYSICS A291 (1977) 221  
 TR77I TRURAN, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D. REIDEL, HOLLAND, 1977), P. 145  
 TR78 TRURAN AND CAMERON, ASTROPHYS. J. 219 (1978) 226  
 TR78D TRAINI, PHYS. REV. LETT. 41 (1978) 1535  
 TR78H TROMBKA, JOHNSON AND EVANS, BULL. AM. PHYS. SOC. 23 (1978) 1024  
 TR78I TRURAN, COWAN AND CAMERON, ASTROPHYS. J. 222 (1978) L63  
 TR79D TRUEX AND BANERJEE, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 263  
 TR79H TRUOL, PROC. MAINZ, 1979, SPRINGER LEC. NOTES 108 (1979) P. 351  
 TR80C TROST ET AL, NUCL. PHYS. A337 (1980) 377  
 TR80D TRIBBLE, IN ATOMIC MASSES AND FUND. CONSTANTS, NOLEN AND BENENSON EDS., PLenum (1980) P. 13  
 TR80F TROST, SCHWARZ, LEZOCH AND STROHBUSCH, P. 443  
 PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS BERKELEY (1980)  
 TR80H TRIPATHI, FAESSLER AND SHIMIZU, Z. PHYS. A297 (1980) 275  
 TR80J TRURAN, NUKLEONIKA 25 (1980) 1463  
 TR81A TRAUTMANN ET AL, PHYS. REV. LETT. 46 (1981) 1188  
 TS78A TSERRUYA ET AL, PHYS. REV. C18 (1978) 1688  
 TU76A TURCHINETZ, AUSTRALIAN J. PHYS. 29 (1976) 573  
 TU77A TURCK ET AL, IN TOKYO (1977), CONTRIBUTED PAPERS P. 152  
 TU81 TUTUNJIAN AND HARRINGTON, BULL. AMER. PHYS. SOC. 26 (1981) 622  
 UB81 UBERALL, FARHAN, DRAGUN AND MAQUEDA, NUCL. PHYS. A362 (1981) 241  
 UC81 UCHIYAMA, PHYS. REV. C23 (1981) 935  
 UD77 UDAGAWA, PROC. INT. CONF. NUCLEAR STRUCTURE, TOKYO, 1977, J. PHYS. SOC. JAPAN 44 (1978) SUPPL., P. 667  
 UL78 ULEHLA AND NGUYEN, CZECH. J. PHYS. 28 (1978) 508  
 UL79 ULLRICH, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 154  
 UN79 UNGRICHT ET AL, NUCLEAR PHYSICS A313 (1979) 376

- US79 USTIMENKOV ET AL, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2189
- UZ79 UZUREAU, FIENI, COCU AND MICHAUDON, 5TH SESS. D'ETUDES BIENNALE DE PHYS. NUCL. - AUSSOIS (FRANCE) (1979)
- VA75E VANDER MOLEN, BECCHETTI, JANECKE AND CHUA, PHYS. REV. C11 (1975) 734
- VA76H VALEK, VERTSE, SCHLENK AND HUNYADI, NUCLEAR PHYSICS A270 (1976) 200
- VA76M VAN PARADIJS AND KESTER, ASTRON. AND ASTROPHYS. 53 (1976) 1
- VA77 VAN BIBBER ET AL, PHYS. REV. LETT. 38 (1977) 334
- VA77D VAGRADOV, GAREEV AND BANG, NUCLEAR PHYSICS A278 (1977) 319
- VA77E VARMA AND ZAMICK, PHYS. REV. C16 (1977) 308, AND IBID. C18 (1978) 1081
- VA78A VARMA, PHYS. REV. C17 (1978) 267
- VA78G VAZ AND ALEXANDER, PHYS. REV. C18 (1978) 2152
- VA79A VAN ORDEN, BANERJEE AND TRUEX, BULL. AM. PHYS. SOC. 24 (1979) 677
- VA79B VAAGEN, NILSSON, BANG AND IBARRA, NUCLEAR PHYSICS A319 (1979) 143
- VA79H VAN BIBBER ET AL, PHYS. REV. LETT. 43 (1979) 840, AND IBID. 43 (1979) 1274
- VA79K VAN BIBBER, IN BNL-51115 (1979), P. 365
- VA79L VARY, BELEHRAD AND DALTON, NUCL. PHYS. A328 (1979) 526
- VA80A VARY, BELEHRAD AND MCCARTHY, PHYS. REV. C21 (1980) 1626
- VA80B VAN ORDEN, TRUEX AND BANERJEE, PHYS. REV. C21 (1980) 2628
- VA80C VANDENBOSCH, NUCL. PHYS. A339 (1980) 167
- VA81B VANDENBOSCH AND LAZZARINI, PHYS. REV. C23 (1981) 1074
- VA81C VARY AND MCCARTHY, BULL. AMER. PHYS. SOC. 26 (1981) 567
- VA81E VAN GIAI, AUERBACH AND MEKIJIAN, PHYS. REV. LETT. 46 (1981) 1444
- VA81F VAZ, ALEXANDER AND SATCHLER, PHYS. REP. 69 (1981) 373
- VA81G VAN GIAI AND AUERBACH, PROCEEDINGS VERSAILLES CONF. (1981) p. 498
- VD79 VDOVIN ET AL, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 148
- VE79 VENTURA ET AL, PHYS. REV. C19 (1979) 1705
- VE79A VERGADOS, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 5D14
- VE79C VERSHININ AND CHERNOV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 1047
- VE79E VERSHININ, IZV. VUZ. FIZ. (USSR) 22 (1979) 44
- VI77A VILAITHONG ET AL, BULL. AM. PHYS. SOC. 22 (1977) 1007
- VI79 VIDAL-QUADRAS AND ORTEGA, NUOVO CIM. A49 (1979) 235
- VI79C VIGDOR ET AL, PHYS. REV. C20 (1979) 2147
- VI80A VICHNIAC, P. 556, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- VI81 VINAS, LOZANO AND MADURGA, PHYS. REV. C23 (1981) 780
- VL79 VLADIMIROV, GONCHAROVA AND ZHIVOPISTSEV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2311
- VO75I DE VOIGT ET AL, IN JUL-CONF-16 (1975), P. 5
- VO77 VON ROSENVINGE, MC DONALD, TRAINOR AND WEBBER, BULL. AM. PHYS. SOC. 22 (1977) 567
- VO77B VOLLLIER AND WALECKA, ACTA PHYS. POLONICA B8 (1977) 25
- VO77E VOLANT, CONJEAUD, HARAR AND DA SILVEIRA, J. PHYSIQUE 38 (1977) 1179

- V078 VOLKOV, PHYS. REPORTS 44 (1978) 93  
 V078A VOLKOV, IN CLUSTERING, WINNIPEG 1978, AIP CONF. PROC.  
 47 (1978) 352  
 V078B VON CHARZEWSKI, HNIZDO AND TOEPFFER, NUCLEAR PHYSICS  
 A307 (1978) 309  
 V078C VOOK, PICRAUX AND DOYLE, BULL. AM. PHYS. SOC. 23 (1978)  
 1038, 1061  
 V078F VON CHARZEWSKI, HNIZDO AND TOEPFFER, S. AFRICAN J.  
 PHYS. 1 (1978) 222  
 V079A VOSNIAKOS ET AL, NUCL. PHYS. A332 (1979) 157  
 V080E VON OERTZEN, NUKLEONIKA 25 (1980) 939  
 V081 VOLANT AND GARY, J. PHYSIQUE 42 (1981) 27  
 V081B VOIT ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 612  
 WA64G WACHTER, PHYS. REV. 135 (1964) B1180  
 WA68N WARBURTON, HARRIS AND ALBURGER, PHYS. REV. 175 (1968) 1275  
 WA75H WATSON, BARDIN, BECKER AND FISHER, PHYS. REV. LETT. 35  
 (1975) 1333  
 WA76K WALL, CRAIG AND EZROW, NUCLEAR PHYSICS A268 (1976) 459  
 WA76P WALTER AND LISOWSKI, IN LOWELL CONF., CONF-760715-P2  
 (1976), P. 1061  
 WA77 WAPSTRA AND BOS, ATOMIC DATA AND NUCLEAR DATA TABLES 19  
 (1977) 175  
 WA77E WADA, SCHIMIZU AND TAKIMOTO, PHYS. REV. LETT. 38 (1977) 1341  
 WA77K WAGHMARE, PRAMANA 9 (1977) 7  
 WA77L WAGNER ET AL, PHYS. REV. C16 (1977) 1271  
 WA77O WALECKA, AIP CONF. PROC. NO. 37 (INDIANA U.), WEAK  
 INTERACTION PHYSICS - 1977, ED. LICHTENBERG (AIP, NEW  
 YORK, 1977), P. 125  
 WA77P WALTER, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE,"  
 ZURICH 1977, BIRKHAUSER VERLAG, P. 225  
 WA77R WALKER, IN "MOMENTUM WAVE FUNCTIONS - 1976" (AIP CONF.  
 PROC. NO. 36), ED. DEVINS (AIP, NEW YORK, 1977), P. 21  
 WA77T WANNIER, IN CNO ISOTOPES IN ASTROPHYS., J. AUDOUZE ED. (D.  
 REIDEL, HOLLAND, 1977), P. 71  
 WA79B WALKER, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 206  
 WA79C WAMBACH, OSTERFELD, SPETH AND MADSEN, NUCL. PHYS. A324  
 (1979) 77  
 WA79G WALKER, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979)  
 P. 351.  
 WA79J WANG ET AL, ANNU. REP. INST. PHYS. ACAD. SIN. 9 (1979) 71;  
 PHYS. ABS. 5785 (1981)  
 WA80B WANG ET AL, PHYS. REV. C21 (1980) 1705  
 WA80E WANG, LI AND ZHANG, PHYS. ENERG. FORTIS & PHYS. NUCL.  
 4 (1980) 63, AND PHYS. ABSTRACTS 47407 (1980)  
 WA80F WADA ET AL, PHYS. REV. C22 (1980) 557  
 WA80I WATSON, SMITHSON, FORTUNE AND MIDDLETON, P. 564, PROCEEDINGS  
 OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY  
 (1980)  
 WA80J WAROQUIER, HEYDE AND PROVOOST, P. 738, PROCEEDINGS OF THE  
 INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)  
 WA80K WARBURTON, ALBURGER AND MILLENER, PHYS. REV. C22 (1980) 2330  
 WA80L WALTER AND BYRD, IN THE (P,N) REACTION AND THE NUCLEON-  
 NUCLEON FORCE (1980) P. 469, PLENUM PUBS. (1980) 469  
 WA80N WANNIER, ANN. REV. ASTRON. & ASTROPHYS. 18 (1980) 399  
 WA81 WALTHAM ET AL

- WA81A WALTER ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 1326
- WA81B WALTER, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 344
- WE71Q WEIBEZAHN, FREIESLEBEN, PUHLHOFER AND BOCK, NUCLEAR PHYSICS A176 (1971) 645
- WE72A WELLER, BLUE AND BLATT, PHYS. REV. C5 (1972) 648
- WE76J WESTFALL AND ZAIDI, PHYS. REV. C14 (1976) 610
- WE77G WELSH, IN "HIGH ENERGY PHYSICS AND NUCL. STRUCTURE," ZURICH 1977, BIRKHAUSER VERLAG, P. 95
- WE77K WERBY, NAGARAJAN, GOLDFINGER AND TOBOCMAN, FIZIKA (YUGOSLAVIA) 9, SUPPL. 2 (1977) 46, AND PHYS. ABSTRACTS 627 (1978)
- WE77L WEISS, FIZIKA (YUGOSLAVIA) 9, SUPPL. 3 (1977) 315, AND PHYS. ABSTRACTS 54781 (1978)
- WE78D WERNITZ, PHYS. REV. LETT. 41 (1978) 1030
- WE78H WEISE, IN FEW BODY SYSTEMS AND NUCLEAR FORCES, GRAZ 1978, SPRINGER LECTURE NOTES 87 (1978), P. 177
- WE78I WEST AND SHERWOOD, REPORT AERE/R9195, UKAEA (1978) AND PHYS. ABSTRACTS 6651 (1979)
- WE79 WEBBER, KISH AND YUSHAK, BULL. AM. PHYS. SOC. 24 (1979) 692
- WE79E WEISE, IN HOUSTON, AIP CONF. PROCEEDINGS 54 (1979), P. 576
- WE80 WERBY, STRAYER AND NAGARAJAN, PHYS. REV. C21 (1980) 2235
- WE80A WELLS, BALAMUTH, BYBELL AND MULLIGAN, BULL. AMER. PHYS. 25 (1980) 738
- WE80D WELLER AND ROBERTSON, REV. OF MODERN PHYS. 52 (1980) 699
- WE80E WERBY, BASSICHIS AND STRAYER, NUOVO CIMENTO A58 (1980) 219
- WE81E WEISE, NUCL. PHYS. A358 (1981) C163
- WH80A WHITTEN, NUCL. PHYS. A335 (1980) 419
- WH81 WHARTON AND KEISTER, PHYS. REV. C23 (1981) 1141
- WH81B WHARTON ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 327
- WI56 WILKINSON, TOPPEL AND ALBURGER, PHYS. REV. 101 (1956) 673
- WI72B WILKINSON AND ALBURGER, PHYS. REV. C5 (1972) 719
- WI78A WISSINK, CALARCO, FISHER AND KURJAN, BULL. AM. PHYS. SOC. 23 (1978) 62
- WI79 WILKERSON, CLEGG AND LUDWIG, BULL. AM. PHYS. SOC. 24 (1979) 107
- WI79B WILKERSON ET AL, BULL. AM. PHYS. SOC. 24 (1979) 612
- WI79C WISSINK, CALARCO AND HANNA, BULL. AM. PHYS. SOC. 24 (1979) 646
- WI79D WINHOLD ET AL, INT. CONF. NUCL. PHYS. WITH ELECTROMAGNETIC INTERACTIONS, MAINZ (1979), 6.16
- WI79E WINHOLD, IN 1978 RPI SYMPOSIUM "PHOTOPION NUCLEAR PHYSICS," (PLENUM PRESS, NEW YORK, 1979), P. 19
- WI79M WILLETT, LING, MAHONEY AND JACOBSON, ASTROPHYS. J. 234 (1979) 753
- WI79O WILDERMUTH AND KANELLOPOULOS, REP. PROG. PHYS. 42 (1979) 1719
- WI80B WILSCHUT ET AL, BULL. AMER. PHYS. SOC. 25 (1980) 591
- WI80E WIELECZKO, HARAR, CONJEAUD AND SAINT LAURENT, PHYS. LETTERS B93 (1980) 35
- WI80H WILSCHUT, BRAUN-MUNZINGER, FREIFELDER AND RENNER, BULL. AMER. PHYS. SOC. 25 (1980) 726
- WI81B WIEBICKE AND ZHUKOV, NUCL. PHYS. A351 (1981) 321
- WI81C WIEDENBECK AND GREINER, PHYS. REV. LETT. 46 (1981) 682

- WI81D WIENHARD ET AL, IN POLARIZATION, SANTA FE 1980, AIP CONFERENCE PROC. 69 (1981) p. 747
- WI81F WILSCHUT ET AL, BULL. AMER. PHYS. SOC. 26 (1981) 610
- WI81I WILSON, LANGER AND GOLDSMITH, ASTROPHYS. J. 243 (1981) L47
- WO76C WOZNIAK, STAHEL, CERNY AND JELLEY, PHYSICAL REVIEW C14 (1976) 815
- WO78F WOOSLEY, FOWLER, HOLMES AND ZIMMERMAN, ATOMIC DATA AND NUCLEAR DATA TABLES 22 (1978) 371
- WO79B WONG, IN BNL-51115 (1979), P. 329
- WO80A WOLF, MOSEL AND PIEPER, Z. PHYS. A 294 (1980) 261
- WO80F WONG AND DAVIES, PHYS. LETT. B96 (1980) 258
- WR78 WRIGHT, STORER AND MC CARTHY, PHYS. REV. C17 (1978) 473
- WU78B WU AND CHANG, PHYS. REV. C17 (1978) 1540
- WU78C WU, OVERLEY, BARNES AND SWITKOWSKI, NUCLEAR PHYSICS A312 (1978) 177
- WU79D WULLSCHLEGER AND SHECK, NUCL. PHYS. A326 (1979) 325
- WU80 WUNSCH, NUCL. PHYS. A336 (1980) 446
- WU80A WU, PHYS. LETT. 91B (1980) 169
- YA76B YAMAMOTO, HINO, ITAGAKI AND SUGIYAMA, J. NUCL. SCI. & TECHNOL. (JAPAN) 15 (1978) 797
- YA80A YASUE ET AL, P. 259, PROCEEDINGS OF THE INTERNATIONAL CONFERENCE ON NUCLEAR PHYSICS, BERKELEY (1980)
- YA81 YAMAZAKI ET AL, PROCEEDINGS VERSAILLES CONF. (1981) p. 115
- YO66M YOUNG ET AL, NUCL. INSTR. METH. 44 (1966) 109
- YO77D YORK AND CARPENTER, NUCLEAR PHYSICS A282 (1977) 351
- YO77K YOUNG, GOLDEN AND KAPETANAKOS, REV. SCI. INSTRUM. 48 (1977) 432
- YO78C YOUNG AND GIBBS, PHYS. REV. C17 (1978) 837
- YO79 YOUNG, ZURMUHLE, LIND AND BALAMUTH, NUCLEAR PHYS. A330 (1979) 452
- YU78A YU, TZU AND WU, PHYS. ENERG. FORTIS & PHYS. NUCL. (CHINA) 2 (1978) 280, AND PHYS. ABSTRACTS 82271 (1978)
- ZA65 ZAFIRATOS, AJZENBERG-SELOVE AND DIETRICH, PHYS. REV. 137 (1965) B1479
- ZA77 ZAMICK, GOLIN AND MOSzkowski, PHYS. LETT. B66 (1977) 116
- ZA78 ZAMICK, PHYS. REV. LETT. 40 (1978) 381
- ZA78A ZABOLITZKY AND EY, PHYS. LETT. B76 (1978) 527
- ZA78B ZALM, VAN HIENEN AND GLAUDEMANS, Z. PHYS. A287 (1978) 255
- ZA79 ZAMICK, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 5C1
- ZA79A ZAMICK, IN NATO INSTITUTE VOL. 45 (PLENUM PRESS, 1979) P. 397.
- ZA79B ZABOLITZKY AND EY, NUCL. PHYS. A328 (1979) 507
- ZA81 ZANELLI ET AL, PHYS. REV. C23 (1981) 1015
- ZA81A ZAMICK, PROCEEDINGS VERSAILLES CONF. (1981) p. 89
- ZE71 ZEITNITZ ET AL, NUCLEAR PHYSICS A166 (1971) 443
- ZE76 ZELLER ET AL, PHYS. REV. C14 (1976) 2162
- ZE77B ZELENSKAYA, TEPLOV AND YUSHCHENKO, IZV. AKAD. NAUK SSSR (SER. FIZ.) 41 (1977) 1691
- ZE77C ZELENSKAYA AND TEPLOV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 41 (1977) 1711
- ZE77D ZEIDMAN, NUCL. INSTR. METH. 146 (1977) 199
- ZE78C ZELENSKAYA, LEBEDEV AND YUSHCHENKO, SOVIET J. NUCL. PHYS. 28 (1978) 44
- ZE79A ZELENSKAYA AND TEPLOV, IZV. AKAD. NAUK SSSR (SER. FIZ.) 43 (1979) 2375

- ZH77 ZHANG, ZHAO AND LI, ACTA PHYS. SIN. (CHINA) 26 (1977) 54;  
PHYS. ABSTRACTS 57614 (1977)
- ZI75C ZIMMERMAN, FOWLER AND CAUGHLAN, OAP-399 (1975)
- ZI77C ZINT, Z. PHYS. A281 (1977) 373
- ZI80A ZISMAN ET AL, PHYS. REV. C21 (1980) 2398
- Z079 ZOFKA, IN CONF. PROC. TRIUMF, VANCOUVER (1979), PAPER 5E7
- Z080 ZOFKA, CZECH. J. PHYS. B30 (1980) 95
- ZY79A ZYSKIND AND PARKER, NUCLEAR PHYSICS A320 (1979) 404