

## Post Hoc Comparisons

```
> ### Descriptive Statistics
```

```
> by(Outcome, Factor, sd)
```

```
> mean(Outcome)
```

```
[1] 5
```

```
> tapply(Outcome, Factor, function(x) c(length(x), mean(x), sd(x)))
```

```
$`1`
```

```
[1] 4.00000 2.00000 2.44949
```

```
$`2`
```

```
[1] 4.00000 6.00000 2.44949
```

```
$`3`
```

```
[1] 4.00000 7.00000 2.44949
```

```
> ### Inferential Statistics
```

```
> TukeyHSD(Results)
```

```
Tukey multiple comparisons of means
95% family-wise confidence level
```

```
Fit: aov(formula = Outcome ~ Factor)
```

```
$Factor
```

	diff	lwr	upr	p-adj
2-1	4	-0.8358956	8.835896	0.1055254
3-1	5	0.1641044	9.835896	0.0431300
3-2	1	-3.8358956	5.835896	0.8352889

These values of the group statistics are calculated separately for each group. They are not identical to the values obtained from analyzing the variable as a whole.

A grand mean can be determined by taking the weighted average of all of the group means:

$$M_{TOTAL} = \frac{\sum n(M_{GROUP})}{N} = \frac{4(2) + 4(6) + 4(7)}{4 + 4 + 4} = 5.000$$

The "Standard Errors" are for the difference between the two group means. The values are a function of the  $MS_{WITHIN}$  (from the ANOVA) and the sample sizes:

$$SE_{DIFF} = \sqrt{\left(\frac{MS_{WITHIN}}{n_{GROUP}}\right) + \left(\frac{MS_{WITHIN}}{n_{GROUP}}\right)} = \sqrt{\left(\frac{6}{4}\right) + \left(\frac{6}{4}\right)} = 1.732$$

In this case, because all groups are of the same size, the standard error for each comparison is the same.

An HSD value is conceptually similar to a  $t$  statistic in that it is a function of the "Diff" and the "SE". For the first comparison in the example:

$$HSD = \frac{M_2 - M_1}{SE_{DIFF}} = \frac{4.000}{1.732} = 2.309$$

The "p adj" column provides the probability of the HSD statistic. An HSD of -2.309 (with 2  $df_{BETWEEN}$  and 9  $df_{WITHIN}$  like in the ANOVA source table) has a two-tailed probability ( $p$ ) of .106, a finding that is not statistically significant.

This section provides confidence intervals around (centered on) the Mean Differences. Calculation requires the appropriate critical value. Specifically, the HSD statistic (with 2  $df_{BETWEEN}$  and 9  $df_{WITHIN}$ ) that has a probability of .05 equals 3.068. For the first comparison in the example:

$$CI_{DIFF} = M_{DIFF} \pm (HSD_{CRITICAL})(SE_{DIFF})$$

$$CI_{DIFF} = 4.000 \pm (2.792)(1.732)$$

Thus, the estimates that the true population mean difference is somewhere between -8.836 and 0.836 (knowing that the estimate could be incorrect).