

RESEARCH STATEMENT


MILTON LIN

INTRODUCTION

My areas of interest in mathematics are:

- (1) The **Langlands correspondence**, which is now a huge web of conjectures, from special values of L functions to conformal field theory. My particular interest is in various incarnations of the **Casselman–Shalika formula**, [A](#), and **relative aspects of the ℓ -adic geometric local Langlands correspondence**, [C](#).
- (2) **Stable homotopy theory**, through the lens of **categorical deformation theory**, [F](#).
- (3) **Motivic phenomena**, where I hope to explore the motivic version of Casselman–Shalika formula, [D](#), and a stacky approach to periods, [E](#).

Majority of the research presented here originates from my study of the Casselman–Shalika formula in the mixed characteristic setting, as outlined in [Section 1](#). These areas of research are interconnected, as shown in the following diagram.



The priority of research is listed in the following order¹,

$$\mathbf{A}=\mathbf{C}=\mathbf{F}>\mathbf{D}>\mathbf{E}>\mathbf{B}.$$

Date: October 26, 2024.

¹The alphabet links to the goal rather than the section.

Notations. Theorems stated have full proof written by either me or joint with collaborators, *unless* it is annotated with: **conjecture** – no proofs have been written down but is believed to hold, or **in progress** – where we have partial progress. We use freely the language of higher categories and higher algebra, [Lur09], [Lur18].

1. MIXED CHARACTERISTIC GEOMETRY AND THE CASSELMAN–SHALIKA FORMULA

Let G be a connected reductive group over a nonarchimedean local field with residue characteristic $p \neq \ell$, and $\Lambda := \overline{\mathbb{Q}}_\ell$. In [FS24], Fargues and Scholze have formulated the geometric Langlands conjecture for the Fargues–Fontaine curve, similar to the function field correspondence of Beilinson–Drinfeld [AG15], [Gai14]. They constructed two actions

$$\begin{array}{ccc} & \text{IndPerf}(\text{LS}_{\hat{G}}) & \\ \circlearrowleft & & \circlearrowright \\ \text{IndCoh}_{\text{Nilp}}^{qc}(\text{LS}_{\hat{G}}) & & D_{\text{lis}}(\text{Bun}_G) \end{array}$$

where $\text{LS}_{\hat{G}}$ be the moduli stack of L -parameters of \hat{G} , and Bun_G the moduli stack of G -bundles on the Fargues–Fontaine curve. The left-hand action in the diagram is induced from the natural tensor structure, while the right-hand is the so-called *spectral action* [NY19]. For a fixed choice of Whittaker datum, [FS24, I.10.2] conjectured a $\text{IndPerf}(\text{LS}_{\hat{G}})$ -linear equivalence

$$\mathbb{L}_G : D_{\text{lis}}(\text{Bun}_G, \Lambda) \simeq \text{IndCoh}_{\text{Nilp}}^{qc}(\text{LS}_{\hat{G}})$$

which can be thought of as a generalization of the classical local Langlands.

One fundamental aspect of the program is to understand the *Whittaker Fourier coefficient functor*,

$$\text{coeff} : D_{\text{lis}}(\text{Bun}_G, \Lambda) \rightarrow D(\Lambda)$$

and its various properties, see [FR22, Ch. 5] for definitions and theorems. In the classical setting, this corresponds to finding the Fourier coefficients of automorphic functions.

Example 1.1. Let $G = \text{PGL}_2$ be the projective linear group over \mathbb{Q} . A modular function, f , has an adelic formulation, \tilde{f} on $G(\mathbb{A}_{\mathbb{Q}})$. Up to a *normalizing factor*, the Fourier coefficients coincide with the Whittaker coefficients (Jacquet period),

$$a_m \sim \int_{N(\mathbb{Q}) \backslash N(\mathbb{A}_{\mathbb{Q}})} \tilde{f}(n\alpha_m)\psi(-n) \, dn \quad \text{for } m \geq 1$$

where $\alpha_m \in T(\mathbb{A}_{\mathbb{Q}}^{\text{fin}})$ is m considered as a finite idèle and ψ is a standard character on $N(\mathbb{A}_{\mathbb{Q}}) = \mathbb{A}_{\mathbb{Q}}$, where N is the radical of standard Borel. For more details, see [Gel75, Ch.3].

The first fundamental result in this context is the *global Casselman–Shalika formula*, as proven in [FGV01], which we aim to replicate in the mixed characteristic setting. In joint work with Ashwin Iyengar (American Mathematical Society) and Konrad Zou (Bonn University) [ILZ24], we proved a variation of this problem: the *geometric Casselman–Shalika formula* over the Witt vector affine Grassmannian Gr_G , analogous to the equi-characteristic geometrization carried out in [NP01].

Theorem 1.2. [ILZ24] *The geometric Casselman–Shalika formula holds over the Witt vector Grassmannian.*

At the same time, we have established basic properties of averaging functors (cf. [FR22, Ch. 7], [BBM21]), yielding Iwahori–Satake equivalence, [Bez+19] without using nearby cycles.

Theorem 1.3 (I.-Lin-Z., in progress). *The Iwahori-Whittaker category is equivalent to the spherical Hecke category in mixed characteristics.*²

As of writing, D. Hansen, L. Hamman, and L. Mann have announced significant advancements to the global version of the formula. Building on all these progresses, I have developed a strong interest in exploring the geometrization of the Casselman–Shalika formula for covering groups [GGW18].

Research Goal A. Geometrization of Metaplectic/Quasi-split Casselman–Shalika. *We propose two explorations of the Casselman–Shalika formula over equal-characteristic local fields. First, a geometric metaplectic Casselman–Shalika formula, building on the works of Gaitsgory and Lysenko [GL22], McNamara [McN16], and Brubaker et al. [Bru+20; Bru+24]. Second, a geometric Casselman–Shalika formula for quasi-split groups, following [GK20].*

Lastly, we would like to highlight that in our recent work [ILZ24], we have studied the smoothness of perfect Schubert subvarieties, $\mathrm{Gr}_{G, \leq \mu}$, for minuscule and quasi-minuscule μ . In the equal characteristic case over $\mathbb{C}((t))$, this was first examined by Evens and Mirkovic and later extended by Haines [HR20], with applications to classifying Shimura varieties with good or semi-stable reductions.

Research Goal B. Geometry of general perfect Schubert variety *Prove the results of Pappas and Zou, [PZ24] in perfect geometry. Associated to an absolutely special vertex in the Bruhat-Tits building of G , we have an associated group scheme \mathcal{G} over \mathcal{O} .*

Conjecture 1.4. The smooth locus of $\mathrm{Gr}_{\mathcal{G}, \leq s}$ is $\mathrm{Gr}_{\mathcal{G}, s}$ in perfect geometry, in the sense of [Zhu17].

2. RELATIVE LANGLANDS ON THE FARGUES FONTAINE CURVE

Joint with Yuta Takaya (Tokyo University), we explicitly compare the period sheaves on \mathcal{A} -side (automorphic) and L -sheaves on \mathcal{B} -side (Galois) under the relative Langlands conjectures of Ben-Zvi-Sakellaridis-Venkatesh. [BSV]. In *op. cit.*, one interprets Langlands correspondence as a form of arithmetic 4d quantum field theory, which has genesis in the Kapustin-Witten interpretation, [KW07], and the Knots and Primes promoted by B. Mazur, M. Kim, [Kim16]. The Fargues-Fontaine curve should be a *global object* of dimension 2 under the dictionary.

The theory suggests a duality extending that Langlands dual group: (G, X) with (\hat{G}, \hat{X}) of hyper-spherical varieties. We considered the *Iwasawa-Tate case*: $G = \mathbb{G}_{m, F}$ and $X = \mathbb{A}_F$ with dual pair $\hat{G} = \mathbb{G}_{m, \Lambda}$ and $\hat{X} = \mathbb{A}_{\Lambda}$. We constructed two maps

$$\pi : \mathrm{Bun}_G^X \rightarrow \mathrm{Bun}_G, \quad \hat{\pi} : \mathrm{LS}_{\hat{G}}^{\hat{X}} \rightarrow \mathrm{LS}_{\hat{G}}$$

yielding the *period sheaf*, $\mathcal{P}_X := \pi_! \Lambda$, and *L-sheaf*, $\mathcal{L}_{\hat{X}} := \pi_* \omega_{\mathrm{Loc}_{\hat{G}}^{\hat{X}}}$. Bun_G has a Hardar-Narasimhan stratification by locally closed substacks Bun_G^b indexed by the Kottwitz set $B(G)$. In our case, $G = \mathbb{G}_m$, $\mathrm{Bun}_{\mathbb{G}_m}$ is stratified by $\mathrm{Bun}_{\mathbb{G}_m}^n$ for $n \in \mathbb{Z} = B(T)$. The following theorem explicitly computes the period sheaf.

Proposition 2.1 (Lin-T.). *Let $n > 0$. Let $\mathcal{BC}(n)$ be the Banach-Colmez space of the line bundle $\mathcal{O}(n)$. The relative stack $\mathrm{Bun}_G^{X, n} := \pi^{-1}(\mathrm{Bun}_G^n)$ is a $\mathcal{BC}(n)$ -torsor over Bun_G^n . $\mathcal{P}_X^n := \mathcal{P}_X|_{\mathrm{Bun}_G^n} \simeq \Lambda[-2n]$.*

On the other hand, we expect that one can study the spectral side using derived Fourier vector bundles (cf. [FW24, Ch. 6]).

²In the set up of [Bez+19], the equivalence was applied to modular representation theory, as the Iwahori-Whittaker category has a much simpler categorical structure.

Conjecture 2.2. [Lin-T., in progress] Under the geometric local Langlands correspondence, \mathbb{L}_G , (appropriately normalized) \mathcal{P}_X is sent to³ $\mathcal{L}_{\hat{X}}$.

Research Goal C. Relative Langlands on the Fargues Fontaine curve. *Complete Conjecture 2.2 as a first step and then the Hecke case, which classically corresponds to Hecke's integral representation of standard L -function for GL_2 . Lastly, one can ask whether on the \mathcal{B} -side, the same constructions of [BSV, Ch. 11] works for the p -adic (Emerton-Gee) L -parameter stacks, which potentially give new interpretations to p -adic L -functions.*

3. MOTIVIC PHENOMENA

In this section S denotes a base scheme satisfying the Beilinson-Soulé vanishing conjecture, such as $\text{Spec } \mathbb{Z}$, finite field. R denotes a regular coherent coefficient ring. I discuss one side project I have done, and one area of research, [D](#), which extends my joint work [\[ILZ24\]](#).

In Grothendieck's quest towards the Weil conjectures, he reduced to problems of motives. Beilinson proposed an extension of the notion of pure motives to mixed motives, which was realized through the works of Voevodsky, Levine for a field, and extended by Ayoub and Cisikinki-Dégliše, [\[CD19\]](#). For a finite type S -scheme $X \in \text{Sch}_S^{\text{ft}}$, we can construct the derived categories of motives over X , $\text{DM}(X, R)$. If X is smooth then its Ext-groups,

$$\pi_{-m} \text{Map}_{\text{DM}(X, R)}(1_X, 1_X(n)) \simeq \text{CH}^n(X, 2n - m)_R$$

are Bloch's higher Chow groups. Motivic categories and the Chow groups are difficult to explicitly work with due to the lack of motivic t -structures. One method is due to the work of Beauville [\[Bea83\]](#), using Fourier transform. In my joint work, [\[Has+24\]](#) we extended the work of Beauville,

Theorem 3.1 (Lin et al.). *Let $X \rightarrow Y$ be an abelian scheme, whose base Y is smooth and quasi-projective over a field. There is an explicit N , such that one obtains a Beauville decomposition*

$$\text{CH}^i(X)_{\mathbb{Z}[1/N]} \simeq \bigoplus_s \text{CH}_{(s)}^i(X)_{\mathbb{Z}[1/N]}$$

where $\text{CH}_{(s)}^i(X)_R := \{x \in \text{CH}^i(X)_R : [n]_X^* x = n^{2i-s} x \quad \forall n \in \mathbb{Z}\}$ and $[n]$ is the multiplication of an abelian scheme.

This extends to a \mathfrak{sl}_2 action, which we discussed in *op.cit*; and if S were an algebraically closed field, this implies various structural results. The key ingredient was using G. Pappas' version of integral Grothendieck–Riemann–Roch.

Returning to motivic t -structures, it was shown by Levine that t -structures exist on a nice subcategory $\text{DTM}(X) \hookrightarrow \text{DM}(X)$ of *mixed Tate motives* for nice schemes X . This was extended to schemes with cellular Whitney-Tate stratification by Soergel and Wendt, and to prestacks in [\[RS20\]](#).

Research Goal D. Motivic Whittaker categories. *Define a Whittaker category in motivic setting, obtaining a similar equivalence at [\[FGV01\]](#). To begin, we can prove the same statement in [\[NP01\]](#) in the category of mixed Tate motives, in which all the relevant objects are well defined. The difficulty, however, is that the original argument requires characteristic 0 coefficients, whilst in positive characteristic, there are problems of extensions.*

³One has to take into account shearing, twisting and tensoring. There is also an additional $\mathbb{G}_m := \mathbb{G}_{gr}$ action on X which we do not discuss.

4. STACKY APPROACHES AND PERIODS

Various cohomology theories – crystalline cohomology, syntomic cohomology, and Dolbeault cohomology – admit a factorization to the category of stacks over some affine scheme $\mathrm{Spec} R$,

$$\mathrm{Sch}_{\mathbb{Z}}^{\mathrm{sep}, \mathrm{ft}} \rightarrow \mathrm{Stk}_R \rightarrow D(R)$$

$$X \mapsto X^? \mapsto \Gamma(X^?, \mathcal{O}_{X^?})$$

For instance, the *de Rham stack* X^{dR} over \mathbb{Q} , has points given by $X^{\mathrm{dR}}(A) := X(A_{\mathrm{red}})$ for any \mathbb{Q} -algebra A (cf. [GR14]). This is often referred to as a *stacky approach* [Dri22] or *transmutation* [Bha23], which allows one to use six functor formalism and geometric techniques. I explored this concept in basic constructions of F. Brown's work on *motivic periods*, [Bro14]. If X were a smooth variety over \mathbb{Q} , the matrix coefficient from Grothendieck's comparison theorem

$$H_{\mathrm{dR}}^*(X) \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\sim} H_{\mathrm{Betti}}^*(X) \otimes_{\mathbb{Q}} \mathbb{C}$$

with respect to the \mathbb{Q} structure of de-Rham and Betti cohomology (of $X(\mathbb{C})$) are *periods associated to X* . These periods along with their enhancements through Hodge structures, has a natural action of "Galois group"⁴ which should govern the arithmetic structure of periods.

Let $X := \mathbb{P}^1 \setminus \{0, 1, \infty\}$ be the projective space minus three points over \mathbb{Q} . The period associated is studied through Chen's de Rham comparison theorem, [Che76], which relates to iterated integrals [Che73] and multiple zeta values [Bro14]. To give a "stacky" perspective of this period one uses unipotent types of stacks, which is defined in Toën, [Toë06], and recently in [MR23]. This is an endofunctor \mathbf{U} on stacks, sending a stack X to its unipotent homotopy type. My first result is:

Theorem 4.1. (*Lin*) *Unipotent de Rham fundamental group, $\pi_1^{u, \mathrm{dR}}(X, x)$ is isomorphic to $\pi_1(\mathbf{U}(X^{\mathrm{dR}}))$.*

Research Goal E. A stacky approach to motivic periods. *This research aims to use geometric techniques in the study of periods. As a proof of concept, we will recover de Rham comparison theorem through the Riemann-Hilbert correspondence between the analytic Betti stack and the de Rham stack. $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, is not proper, which requires us to incorporate log structures. I expect to prove:*

Conjecture 4.2. There exists X^{Betti} such that the unipotent Betti group $\pi_1^{u, \mathrm{Betti}}(X(\mathbb{C}), x)$ is isomorphic to $\pi_1(\mathbf{U}(X^{\mathrm{Betti}}))$. A logarithmic Riemann-Hilbert comparison should induce Chen's comparison theorem [Hai01, Thm 3.1]

$$\pi_1^{u, \mathrm{dR}}(X, x) \simeq \pi_1^{u, \mathrm{Betti}}(X(\mathbb{C}), x) \otimes \mathbb{C}$$

*By similar techniques of [Bha23], we should recover Haine's theorem: the pro-unipotent completion of de Rham fundamental group admits a mixed Hodge structure.*⁵ *We hope that such work can spark new techniques and new phenomena, such as those used in p -adic integration theory, [Vol01].*

5. DEFORMATION THEORY AND THE SPHERE SPECTRUM \mathbb{S}

Let \mathcal{S} denote the ∞ -category of ∞ -groupoids/anima. The *stabilization* of \mathcal{S} is Sp , the ∞ -category of spectra. This is the natural category to study cohomological invariants. Within Sp , lies the universal cohomology theory, \mathbb{S} , the *sphere spectrum* By Chevalley's works, connected reductive groups over \mathbb{C} have a canonical split \mathbb{Z} -form $G_{\mathbb{Z}}$, see [Con15]. One can analogously ask: *is there a \mathbb{S} -form for algebraic groups?* A first approximation is the existence of an algebraic category $\mathrm{Rep}_{\mathbb{S}}(G_{\mathbb{S}})$, which

⁴For instance, in the approach of Deligne, he defined a *systems of realizations* [Del89]

⁵This is important in Brown's approach, where he reduced his study of motivic periods to mixed Hodge periods [Bro17, p. 3].

deforms to $\text{Rep}_{\mathbb{Z}}(G_{\mathbb{Z}})$. To study, we begin with *formal deformation of categories*, which we briefly recall.

Let \mathcal{C} be a symmetric monoidal category. There is a natural hierarchy of commutativity fitting in the diagram

$$\cdots \rightarrow \mathbb{E}_n(\mathcal{C}) \rightarrow \mathbb{E}_{n-1}(\mathcal{C}) \rightarrow \cdots \rightarrow \mathbb{E}_1(\mathcal{C})$$

where $\text{CAlg}(\mathcal{C}) := \mathbb{E}_{\infty}(\mathcal{C}) := \varprojlim \mathbb{E}_n(\mathcal{C})$ of symmetric algebra objects can be identified with the limit. One can formalize the notion of \mathbb{E}_n -algebra objects via *disk operads*, or *little cubes operads*.

Example 5.1. Let $\mathcal{C} = (\mathcal{S}, \times)$. Let $Y \in \mathcal{S}_*$ is a pointed ∞ -groupoid, its k -fold based loop spaces, $\Omega_*^k Y$ is a classical example of \mathbb{E}_k algebra object in (\mathcal{S}, \times) .

Let $R \in \mathbb{E}_n(\text{Sp})$ be an \mathbb{E}_n ring, and consider LMod_R , the derived category of left R -modules, as an \mathbb{E}_1 object in Pr^{st} , the category of presentable stable categories. This defines $\text{RMod}_{\text{LMod}_R}(\text{Pr}^{\text{st}})$, the category of presentable stable (right) R -linear categories, [Lur18, Appendix D]. Set $\text{Pr}_R^{\text{st}, \text{cg}}$, as the full subcategory spanned by those whose underlying category is compactly generated.⁶ For G a connected reductive group over a field k , $D^b(\text{Rep}_k^{\text{fd}}(G))$, the bounded derived category of finite dimensional algebraic representations with k coefficients lies in $\text{Pr}_R^{\text{st}, \text{cg}}$.

From now on, $k = \mathbb{C}$. Let $\text{Art}_k^{(n)}$ denote the category of \mathbb{E}_n *artinian* ring spectrum, for $n \geq 0$ over k . We refer to [Lur18, Ch. 15] for definition. This extends the classical definition of Artinian local ring, in particular, $R \in \text{Art}_k^{(n)}$ admits an augmentation map $\epsilon : R \rightarrow k$. One defines the \mathbb{E}_{n+2} -formal moduli problem,

$$\begin{aligned} \text{CatDef}^{(n)}(\mathcal{C}) : \text{Art}_k^{(n+2)} &\rightarrow \hat{\mathcal{S}} \\ R &\mapsto |\{\mathcal{C}\} \times_{\text{Pr}_R^{\text{st}, \text{cg}}} \text{Pr}_k^{\text{st}, \text{cg}}| \end{aligned}$$

where $|\cdot|$ is the underlying Kan complex of the ∞ -category. An object consists of: a \mathcal{C}_B right stable R -linear category, and an equivalence $\mathcal{C}_B \otimes_{\text{LMod}_B} \text{LMod}_k \simeq \mathcal{C}$. Our \mathbb{E}_4 -moduli problem is when $n = 2$ and $\mathcal{C} = D^b(\text{Rep}_k^{\text{fd}}(G))$. The geometric Casselman–Shalika [FGV01], which is the \mathbb{E}_2 -algebra equivalence of \mathcal{C} with the *Whittaker sheaves* on the *affine Grassmanian* $\text{Gr}_{\hat{G}}$, describes this moduli problem. Consider moduli of functor of \mathbb{G}_m -gerbes over X

$$\text{Ge}_{\mathbb{G}_m}(X) : R \mapsto \text{Map}_{\mathbb{E}_2(\mathcal{S})}(X, B^2 R^{\times}) \quad R \in \text{Art}_k^{(4)}$$

where $R^{\times} \subset \Omega^{\infty} R$ are the invertible elements of the underlying space of R ⁷ and B^2 is the second deloop. It was stated without proof in [Lur10]

Theorem 5.2 (Lurie). *There is an equivalence of formal \mathbb{E}_4 -moduli problems*

$$\widehat{\text{Ge}_{\mathbb{G}_m}}(\text{Gr}_{\hat{G}}) \xrightarrow{\sim} \text{CatDef}^{(2)}(\text{Rep}_k^{\text{fd}} G)$$

where $\widehat{}$ is the formal completion of the moduli functor at a base point.

Research Goal F. Categorical deformations for the sphere spectrum. We will first document carefully Lurie’s theorem, [Theorem 5.2](#). Then, we will explore deformations of representation of Lusztig’s small quantum group, see *op.cit.* Remark 10.12, using recent advances in quantum geometric Langlands. Ultimately, this research aims to contribute to the foundations of categorical deformations.

⁶The compact generation is only a smallness condition for our version.

⁷is the union of the connected components of invertibles in the $\pi_0 R$ of the 0th space of R and is equivalent to the n th loop space of some space, $R^{\times} \simeq \Omega^n Z$, $n \geq 4$, hence the deloop $B^2 R^{\times} \in \mathbb{E}_2(\mathcal{S})$

REFERENCES

- [AG15] Arinkin, D. and Gaitsgory, D. “Singular support of coherent sheaves and the geometric Langlands conjecture”. In: *Selecta Math. (N.S.)* 21.1 (2015), pp. 1–199. ISSN: 1022-1824,1420-9020. URL: <https://doi.org/10.1007/s00029-014-0167-5> (cit. on p. 2).
- [BBM21] Bezrukavnikov, Roman, Braverman, Alexander, and Mirkovic, Ivan. *Some results about the geometric Whittaker model*. 2021. arXiv: [math/0210250](https://arxiv.org/abs/math/0210250) [math.AG] (cit. on p. 2).
- [Bea83] Beauville, A. “Quelques remarques sur la transformation de Fourier dans l’anneau de Chow d’une variété abélienne”. In: *Algebraic geometry (Tokyo/Kyoto, 1982)*. Vol. 1016. Lecture Notes in Math. Springer, Berlin, 1983, pp. 238–260. ISBN: 3-540-12685-6. URL: <https://doi.org/10.1007/BFb0099965> (cit. on p. 4).
- [Bez+19] Bezrukavnikov, Roman, Gaitsgory, Dennis, Mirković, Ivan, Riche, Simon, and Rider, Laura. “An Iwahori-Whittaker model for the Satake category”. In: *J. Éc. polytech. Math.* 6 (2019), pp. 707–735. ISSN: 2429-7100,2270-518X. URL: <https://doi.org/10.5802/jep.104> (cit. on pp. 2, 3).
- [Bha23] Bhatt, Bhargav. *Prismatic F -gauges*. 2023 (cit. on p. 5).
- [Bro14] Brown, Francis. *Motivic periods and the projective line minus three points*. 2014. arXiv: [1407.5165](https://arxiv.org/abs/1407.5165) [math.NT]. URL: <https://arxiv.org/abs/1407.5165> (cit. on p. 5).
- [Bro17] Brown, Francis. “Notes on motivic periods”. In: *Commun. Number Theory Phys.* 11.3 (2017), pp. 557–655. ISSN: 1931-4523,1931-4531. URL: <https://doi.org/10.4310/CNTP.2017.v11.n3.a2> (cit. on p. 5).
- [Bru+20] Brubaker, Ben, Buciumas, Valentin, Bump, Daniel, and Gustafsson, Henrik P. A. *Vertex operators, solvable lattice models and metaplectic Whittaker functions*. 2020. arXiv: [1806.07776](https://arxiv.org/abs/1806.07776) [math.RT]. URL: <https://arxiv.org/abs/1806.07776> (cit. on p. 3).
- [Bru+24] Brubaker, Ben, Buciumas, Valentin, Bump, Daniel, and Gustafsson, Henrik P. A. *Metaplectic Iwahori Whittaker functions and supersymmetric lattice models*. 2024. arXiv: [2012.15778](https://arxiv.org/abs/2012.15778) [math.RT]. URL: <https://arxiv.org/abs/2012.15778> (cit. on p. 3).
- [BSV] Ben-Zvi, David, Sakellaridis, Yiannis, and Venkatesh, Akshay. “Relative Langlands duality”. In: () (cit. on pp. 3, 4).
- [CD19] Cisinski, Denis-Charles and Déglise, Frédéric. *Triangulated categories of mixed motives*. Springer Monographs in Mathematics. Springer, Cham, [2019] ©2019, pp. xlii+406. ISBN: 978-3-030-33241-9; 978-3-030-33242-6. URL: <https://doi.org/10.1007/978-3-030-33242-6> (cit. on p. 4).
- [Che73] Chen, Kuo-tsai. “Iterated integrals of differential forms and loop space homology”. In: *Ann. of Math. (2)* 97 (1973), pp. 217–246. ISSN: 0003-486X. URL: <https://doi.org/10.2307/1970846> (cit. on p. 5).
- [Che76] Chen, Kuo Tsai. “Reduced bar constructions on de Rham complexes”. In: *Algebra, topology, and category theory (a collection of papers in honor of Samuel Eilenberg)*. Academic Press, New York-London, 1976, pp. 19–32 (cit. on p. 5).
- [Con15] Conrad, Brian. “Non-split reductive groups over \mathbf{Z} ”. In: *Autours des schémas en groupes. Vol. II*. Vol. 46. Panor. Synthèses. Soc. Math. France, Paris, 2015, pp. 193–253. ISBN: 978-2-85629-819-0. URL: <https://doi.org/10.1017/CB09781316092439> (cit. on p. 5).
- [Del89] Deligne, Par P. “Le groupe fondamental de la droite projective moins trois points”. In: *Galois Groups over \mathbb{Q} Proceedings of a Workshop Held March 23–27, 1987*. Springer, 1989, pp. 79–297 (cit. on p. 5).
- [Dri22] Drinfeld, Vladimir. *A stacky approach to crystals*. 2022. arXiv: [1810.11853](https://arxiv.org/abs/1810.11853) [math.AG]. URL: <https://arxiv.org/abs/1810.11853> (cit. on p. 5).
- [FGV01] Frenkel, E., Gaitsgory, D., and Vilonen, K. “Whittaker patterns in the geometry of moduli spaces of bundles on curves”. In: *Ann. of Math. (2)* 153.3 (2001), pp. 699–748.

- ISSN: 0003-486X, 1939-8980. URL: <https://doi.org/10.2307/2661366> (cit. on pp. 2, 4, 6).
- [FR22] Faergeman, Joakim and Raskin, Sam. *Non-vanishing of geometric Whittaker coefficients for reductive groups*. 2022. arXiv: [2207.02955](https://arxiv.org/abs/2207.02955) [math.RT]. URL: <https://arxiv.org/abs/2207.02955> (cit. on p. 2).
- [FS24] Fargues, Laurent and Scholze, Peter. “Geometrization of the local Langlands correspondence”. In: *arXiv e-prints*, arXiv:2102.13459 (Feb. 2024), arXiv:2102.13459. arXiv: [2102.13459](https://arxiv.org/abs/2102.13459) [math.RT] (cit. on p. 2).
- [FW24] Feng, Tony and Wang, Jonathan. *Geometric Langlands duality for periods*. 2024. arXiv: [2402.00180](https://arxiv.org/abs/2402.00180) [math.NT]. URL: <https://arxiv.org/abs/2402.00180> (cit. on p. 3).
- [Gai14] Gaitsgory, Dennis. *Outline of the proof of the geometric Langlands conjecture for $GL(2)$* . 2014. arXiv: [1302.2506](https://arxiv.org/abs/1302.2506) [math.AG]. URL: <https://arxiv.org/abs/1302.2506> (cit. on p. 2).
- [Gel75] Gelbart, Stephen S. *Automorphic forms on adèle groups*. Vol. No. 83. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1975, pp. x+267 (cit. on p. 2).
- [GGW18] Gan, Wee Teck, Gao, Fan, and Weissman, Martin H. “L-groups and the Langlands program for covering groups: a historical introduction”. In: 398. L-groups and the Langlands program for covering groups. 2018, pp. 1–31. ISBN: 978-2-85629-845-9 (cit. on p. 3).
- [GK20] Gurevich, Nadya and Karasiewicz, Edmund. *The Twisted Satake Transform and the Casselman-Shalika Formula for Quasi-Split Groups*. 2020. arXiv: [2012.09893](https://arxiv.org/abs/2012.09893) [math.RT]. URL: <https://arxiv.org/abs/2012.09893> (cit. on p. 3).
- [GL22] Gaitsgory, D. and Lysenko, S. *Parameters and duality for the metaplectic geometric Langlands theory*. 2022. arXiv: [1608.00284](https://arxiv.org/abs/1608.00284) [math.AG]. URL: <https://arxiv.org/abs/1608.00284> (cit. on p. 3).
- [GR14] Gaitsgory, Dennis and Rozenblyum, Nick. *Crystals and D-modules*. 2014. arXiv: [1111.2087](https://arxiv.org/abs/1111.2087) [math.AG]. URL: <https://arxiv.org/abs/1111.2087> (cit. on p. 5).
- [Hai01] Hain, Richard. *Iterated Integrals and Algebraic Cycles: Examples and Prospects*. 2001. arXiv: [math/0109204](https://arxiv.org/abs/math/0109204) [math.AG]. URL: <https://arxiv.org/abs/math/0109204> (cit. on p. 5).
- [Has+24] Hasan, Junaid, Hassan, Hazem, Lin, Milton, Manivel, Marcella, McBeath, Lily, and Moonen, Ben. *Integral aspects of Fourier duality for abelian varieties*. 2024. arXiv: [2407.06184](https://arxiv.org/abs/2407.06184) [math.AG]. URL: <https://arxiv.org/abs/2407.06184> (cit. on p. 4).
- [HR20] Haines, Thomas J. and Richarz, Timo. “Smoothness of Schubert varieties in twisted affine Grassmannians”. In: *Duke Mathematical Journal* 169.17 (Nov. 2020). ISSN: 0012-7094. URL: [http://dx.doi.org/10.1215/00127094-2020-0025](https://doi.org/10.1215/00127094-2020-0025) (cit. on p. 3).
- [ILZ24] Iyengar, Ashwin, Lin, Milton, and Zou, Konrad. *Geometric Casselman-Shalika in mixed characteristic*. 2024. arXiv: [2408.07953](https://arxiv.org/abs/2408.07953) [math.AG]. URL: <https://arxiv.org/abs/2408.07953> (cit. on pp. 2–4).
- [Kim16] Kim, Minhyong. *Arithmetic Chern-Simons Theory I*. 2016. arXiv: [1510.05818](https://arxiv.org/abs/1510.05818) [math.NT]. URL: <https://arxiv.org/abs/1510.05818> (cit. on p. 3).
- [KW07] Kapustin, Anton and Witten, Edward. *Electric-Magnetic Duality And The Geometric Langlands Program*. 2007. arXiv: [hep-th/0604151](https://arxiv.org/abs/hep-th/0604151) [hep-th]. URL: <https://arxiv.org/abs/hep-th/0604151> (cit. on p. 3).
- [Lur09] Lurie, Jacob. *Higher topos theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925. ISBN: 978-0-691-14049-0; 0-691-14049-9. URL: <https://doi.org/10.1515/9781400830558> (cit. on p. 2).
- [Lur10] Lurie, Jacob. “Moduli problems for ring spectra”. In: *Proceedings of the International Congress of Mathematicians. Volume II*. Hindustan Book Agency, New Delhi, 2010, pp. 1099–1125. ISBN: 978-81-85931-08-3; 978-981-4324-32-8; 981-4324-32-9 (cit. on p. 6).

- [Lur18] Lurie, Jacob. “Spectral Algebraic Geometry”. In: (2018) (cit. on pp. 2, 6).
- [McN16] McNamara, Peter J. “The metaplectic Casselman-Shalika formula”. In: *Trans. Amer. Math. Soc.* 368.4 (2016), pp. 2913–2937. ISSN: 0002-9947,1088-6850. URL: <https://doi.org/10.1090/tran/6597> (cit. on p. 3).
- [MR23] Mondal, Shubhodip and Reinecke, Emanuel. *Unipotent homotopy theory of schemes*. 2023. arXiv: [2302.10703](https://arxiv.org/abs/2302.10703) [math.AG] (cit. on p. 5).
- [NP01] Ngô, B. C. and Polo, P. “Résolutions de Demazure affines et formule de Casselman-Shalika géométrique”. In: *J. Algebraic Geom.* 10.3 (2001), pp. 515–547. ISSN: 1056-3911,1534-7486 (cit. on pp. 2, 4).
- [NY19] Nadler, David and Yun, Zhiwei. “Spectral action in Betti geometric Langlands”. In: *Israel J. Math.* 232.1 (2019), pp. 299–349. ISSN: 0021-2172,1565-8511. URL: <https://doi.org/10.1007/s11856-019-1871-9> (cit. on p. 2).
- [PZ24] Pappas, Georgios and Zhou, Rong. *On the smooth locus of affine Schubert varieties*. 2024. arXiv: [2312.14827](https://arxiv.org/abs/2312.14827) [math.AG]. URL: <https://arxiv.org/abs/2312.14827> (cit. on p. 3).
- [RS20] Richarz, Timo and Scholbach, Jakob. “The intersection motive of the moduli stack of Shtuka”. In: *Forum of Mathematics, Sigma* 8 (2020). ISSN: 2050-5094 (cit. on p. 4).
- [Toë06] Toën, Bertrand. “Champs affines”. In: *Selecta Math. (N.S.)* 12.1 (2006), pp. 39–135. ISSN: 1022-1824,1420-9020. URL: <https://doi.org/10.1007/s00029-006-0019-z> (cit. on p. 5).
- [Vol01] Vologodsky, Vadim. *Hodge structure on the fundamental group and its application to p-adic integration*. 2001. arXiv: [math/0108109](https://arxiv.org/abs/math/0108109) [math.AG] (cit. on p. 5).
- [Zhu17] Zhu, Xinwen. “Affine Grassmannians and the geometric Satake in mixed characteristic”. In: *Ann. of Math. (2)* 185.2 (2017), pp. 403–492. ISSN: 0003-486X,1939-8980. URL: <https://doi.org/10.4007/annals.2017.185.2.2> (cit. on p. 3).