

Derived smooth induction with applications

Peter Schneider, Claus Sorensen

Abstract

In natural characteristic, smooth induction from an open subgroup does not always give an exact functor. In this article we initiate a study of the right derived functors, and we give applications to the non-existence of projective representations and duality.

1 Introduction

Let G be a profinite group, and let k be a field of characteristic p . The category of smooth G -representations on k -vector spaces $\text{Mod}_k(G)$ has nonzero projective objects if and only if p has finite exponent in the pro-order $|G|$. See [CK23, Thm. 3.1] for example (or [DK23, Rk. 2.20, p. 20] for a less precise result). In this paper we study the question about non-existence of projectives for *locally* profinite groups. More precisely for p -adic Lie groups G . We approach the problem via the right derived functors of smooth induction Ind_K^G from a compact open subgroup K . This is the *right* adjoint to the restriction functor and, in contrast to compact induction, the functor Ind_K^G is provably not exact in general for non-compact G .

As a sample result, suppose G is a p -adic Lie group with a non-discrete center. We show in Proposition 4.2 that the category $\text{Mod}_k(G)$ has no nonzero projective objects.

For general p -adic reductive groups (with no restriction on the center) we prove the following result, which has been a folklore expectation for some time:

Theorem 1.1. *Let $G = \mathbf{G}(\mathfrak{F})$ for a nontrivial connected reductive group \mathbf{G} defined over a finite extension $\mathfrak{F}/\mathbb{Q}_p$. Then $\text{Mod}_k(G)$ has no nonzero projective objects.*

We deduce 1.1 from the vanishing of $R^d \text{Ind}_K^G(k)$ for certain principal congruence subgroups K . More precisely, we fix a special vertex x_0 in the Bruhat-Tits building of G and consider the associated group scheme $\mathbf{G}_{x_0}^\circ$ over \mathfrak{O} (the valuation ring in \mathfrak{F}). The congruence subgroup

$$K_m := \ker(\mathbf{G}_{x_0}^\circ(\mathfrak{O}) \longrightarrow \mathbf{G}_{x_0}^\circ(\mathfrak{O}/\pi^m \mathfrak{O}))$$

is a uniform pro- p group for $m \in e\mathbb{N}$ (with the extra assumption that $m > e$ if $p = 2$). Here π is a choice of uniformizer in \mathfrak{F} , and e denotes the ramification index of $\mathfrak{F}/\mathbb{Q}_p$.

The most technical part of our paper is finding the precise vanishing range for $R^i \text{Ind}_{K_m}^G(k)$. This range is given by the number

$$i_0 := \dim_{\mathbb{Q}_p}(G/P_{\min})$$

where P_{\min} denotes the group of \mathfrak{F} -points of a minimal parabolic \mathfrak{F} -subgroup of \mathbf{G} . We have:

Theorem 1.2. *$R^i \text{Ind}_{K_m}^G(k) = 0$ if and only if $i > i_0$.*

This answers a question in [Sor] about the higher smooth duals of the compactly induced representation $\text{ind}_{K_m}^G(k)$.

We introduce the functor $\underline{\text{Ind}}$ by taking the union of Ind_K^G as K varies. This takes a smooth G -representation to a smooth $G \times G$ -representation. At the derived level this gives a functor

$$R\underline{\text{Ind}} : D(G) \longrightarrow D(G \times G)$$

where $D(G) := D(\text{Mod}_k(G))$. In [SS] we studied the smooth duality functor $R\underline{\text{Hom}}(-, k)$ on this category. The complex $R\underline{\text{Ind}}(k)$ in some sense represents $R\underline{\text{Hom}}(-, k)$ on the subcategory $D(G)^c$ of compact objects. The precise statement is the following: For all compact V^\bullet there is an isomorphism

$$\tau_{V^\bullet} : R\underline{\text{Hom}}(V^\bullet, k) \xrightarrow{\sim} R\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V^\bullet, R\underline{\text{Ind}}(k))$$

in $D(G)$. Here, if V is a smooth G -representation and W is a smooth $G \times G$ -representation, we let $\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V, W)$ denote the space of k -linear maps $V \longrightarrow W$ which are G -equivariant on the right and smooth on the left (referring to the two G -factors in $G \times G$ acting on W).

When G is a p -adic reductive group, as in Theorems 1.1 and 1.2, we show that $R\underline{\text{Ind}}(k)$ only has cohomology in degree zero:

Theorem 1.3. *Keep the group $G = \mathbf{G}(\mathfrak{F})$ as in Theorem 1.1. There is an isomorphism*

$$\mathcal{C}^\infty(G, k)[0] \xrightarrow{\sim} R\underline{\text{Ind}}(k)$$

in $D(G \times G)$. (Here $\mathcal{C}^\infty(G, k)[0]$ is the space of k -valued functions on G , which are smooth on both sides, viewed as a complex concentrated in degree zero.)

2 The derived functors of smooth induction

For now G denotes an arbitrary locally profinite group, and k is any field. Let Mod_k and $\text{Mod}_k(G)$ be the category of k -vector spaces and of smooth G -representations on k -vector spaces, respectively. Fix a compact open subgroup $K \subseteq G$ and consider the restriction functor $\text{res}_K^G : \text{Mod}_k(G) \longrightarrow \text{Mod}_k(K)$. The compact induction functor ind_K^G is an exact left adjoint. The full smooth induction functor Ind_K^G is a right adjoint of res_K^G , but it is not exact in general when the characteristic of k divides the pro-order of G . The purpose of this paper is to understand the derived functors $R^i \text{Ind}_K^G$ better in that case.

Starting with an object V from $\text{Mod}_k(K)$ we will follow the convention in [Vig96, Ch. I, Sect. 5] and realize $\text{Ind}_K^G(V)$ as the space of all smooth functions $f : G \longrightarrow V$ satisfying the transformation property $f(\kappa x) = \kappa f(x)$ for $\kappa \in K$. Thus in this article our convention is that G acts by right translations.

Definition 2.1. *For an open subgroup $U \subset G$ the K -action on G/U gives rise to the following:*

- i. Let $K \curvearrowright G/U$ denote the groupoid with objects the elements $x \in G/U$ and morphisms $\text{Hom}(x, y) = \{\kappa \in K : \kappa x = y\}$ for $x, y \in G/U$;*
- ii. The representation V gives a functor $F_V : K \curvearrowright G/U \longrightarrow \text{Mod}_k$ sending $x \mapsto V^{K \cap x U x^{-1}}$, and if $\kappa x = y$ the k -linear map associated with κ is*

$$F_V(\kappa) : V^{K \cap x U x^{-1}} \xrightarrow{\sim} V^{K \cap y U y^{-1}} \\ v \longmapsto \kappa v.$$

We can think of the U -invariants $\text{Ind}_K^G(V)^U$ as the limit of F_V .

Lemma 2.2. $\text{Ind}_K^G(V)^U \simeq \varprojlim_{x \in G/U} V^{K \cap xUx^{-1}}$.

Proof. The space $\text{Ind}_K^G(V)^U$ consists of all K -equivariant functions $f : G/U \rightarrow V$. For such an f , as $x \in G/U$ varies the vectors $f(x) \in V^{K \cap xUx^{-1}}$ are compatible via the isomorphisms $F_V(\kappa)$. Vice versa, a compatible tuple of vectors arise from a unique U -invariant function. \square

Remark 2.3. The point of this categorical description of $\text{Ind}_K^G(V)^U$ is to avoid having to pick double coset representatives R for $K \backslash G/U$. With such a choice R one can of course describe the U -invariants in simpler terms as just a product $\prod_{x \in R} V^{K \cap xUx^{-1}}$. However, as U varies the transition maps become more cumbersome to work with.

There is a formula for $R^i \text{Ind}_K^G(V)$ of the same nature.

Proposition 2.4. $R^i \text{Ind}_K^G(V) \simeq \varinjlim_U \varprojlim_{x \in G/U} H^i(K \cap xUx^{-1}, V)$.

Proof. Let $V \rightarrow J^\bullet$ be an injective resolution of V in $\text{Mod}_k(K)$. This remains injective upon restriction to an open subgroup since compact induction is exact. Therefore we have

$$\begin{aligned} R^i \text{Ind}_K^G(V) &\simeq h^i(\text{Ind}_K^G(J^\bullet)) \\ &\simeq \varinjlim_U \varprojlim_{x \in G/U} h^i((J^\bullet)^{K \cap xUx^{-1}}) \\ &\simeq \varinjlim_U \varprojlim_{x \in G/U} H^i(K \cap xUx^{-1}, V). \end{aligned}$$

In the second isomorphism we moved h^i inside \varinjlim_U and $\varprojlim_{x \in G/U}$, which is justified by the fact that Mod_k satisfies AB5 and AB4* (filtered colimits and products are exact). Recall from Remark 2.3 that $\varprojlim_{x \in G/U}$ can be identified with a product $\prod_{x \in R}$. \square

Remark 2.5. For a fixed U the limit $\varprojlim_{x \in G/U} H^i(K \cap xUx^{-1}, V)$ coincides with the groupoid cohomology of $K \curvearrowright G/U$ as described in [Ron, Df. 6] for example. It is the limit of the functor F_V^i sending $x \mapsto H^i(K \cap xUx^{-1}, V)$. Concretely, an element of this limit is a function

$$c : G/U \rightarrow \bigoplus_{x \in G/U} H^i(K \cap xUx^{-1}, V)$$

with the following properties:

- i. $c_x := c(x) \in H^i(K \cap xUx^{-1}, V)$ for all $x \in G/U$;
- ii. If $\kappa x = y$ then $c_x \mapsto c_y$ via the isomorphism

$$\kappa_* = F_V^i(\kappa) : H^i(K \cap xUx^{-1}, V) \xrightarrow{\sim} H^i(K \cap yUy^{-1}, V).$$

With this description we can make the transition maps in the colimit \varinjlim_U explicit. Let $U' \subset U$ be an open subgroup. Then the transition map is

$$\begin{aligned} t_{U, U'}^i : \varprojlim_{x \in G/U} H^i(K \cap xUx^{-1}, V) &\rightarrow \varprojlim_{x' \in G/U'} H^i(K \cap x'U'x'^{-1}, V) \\ c &\mapsto \left(\text{res}_{K \cap x'U'x'^{-1}}^{K \cap x'U'x'^{-1}} c_{x'U} \right)_{x' \in G/U'}. \end{aligned}$$

This above formula for $t_{U, U'}^i$ will play a crucial role throughout this paper.

3 The connection to higher smooth duality

To motivate the ensuing discussion we establish a relation between the functors $R^i \text{Ind}_K^G$ and the higher smooth duality functors introduced in [Koh], and recast in [SS].

Lemma 3.1. *For V in $\text{Mod}_k(K)$ and W in $\text{Mod}_k(G)$ there are functorial isomorphisms*

$$\text{Ind}_K^G \underline{\text{Hom}}(V, W|_K) \simeq \underline{\text{Hom}}(\text{ind}_K^G V, W).$$

(Here $\underline{\text{Hom}}$ denotes the smooth k -linear maps, as defined in [SS, Sect. 1] for example.)

Proof. For any representation X in $\text{Mod}_k(G)$ we have functorial isomorphisms

$$\begin{aligned} \text{Hom}_{\text{Mod}_k(G)}(X, \text{Ind}_K^G \underline{\text{Hom}}(V, W|_K)) &\simeq \text{Hom}_{\text{Mod}_k(K)}(X|_K, \underline{\text{Hom}}(V, W|_K)) \\ &\simeq \text{Hom}_{\text{Mod}_k(K)}(X|_K \otimes_k V, W|_K) \\ &\simeq \text{Hom}_{\text{Mod}_k(G)}(\text{ind}_K^G(X|_K \otimes_k V), W) \\ &\simeq \text{Hom}_{\text{Mod}_k(G)}(X \otimes_k \text{ind}_K^G V, W) \\ &\simeq \text{Hom}_{\text{Mod}_k(G)}(X, \underline{\text{Hom}}(\text{ind}_K^G V, W)). \end{aligned}$$

The fourth isomorphism follows from [Vig96, p. 40]; part d) just prior to Section 5.3. The others use standard adjunction properties. The claim then follows from the Yoneda lemma. \square

This gives the following spectral sequence (with V and W as above).

Proposition 3.2. $E_2^{i,j} = R^i \text{Ind}_K^G \underline{\text{Ext}}^j(V, W|_K) \implies \underline{\text{Ext}}^{i+j}(\text{ind}_K^G V, W).$

Proof. Note that the functor $\underline{\text{Hom}}(V, -)$ preserves injective objects since $(-)\otimes_k V$ is an exact left adjoint. So does $(-)|_K$ as observed in the proof of Proposition 2.4. The Grothendieck spectral sequence for Ind_K^G composed with $\underline{\text{Hom}}(V, (-)|_K)$ takes the stated form by 3.1. \square

We emphasize the special case $W = k$ below.

Corollary 3.3. *Suppose V is a finite-dimensional object of $\text{Mod}_k(K)$ and let $V^* = \text{Hom}_k(V, k)$ denote its contragredient. Then there is an isomorphism of G -representations*

$$R^i \text{Ind}_K^G(V^*) \simeq \underline{\text{Ext}}^i(\text{ind}_K^G V, k).$$

Proof. When $W = k$ the spectral sequence in Proposition 3.2 becomes

$$E_2^{i,j} = R^i \text{Ind}_K^G \underline{\text{Ext}}^j(V, k) \implies \underline{\text{Ext}}^{i+j}(\text{ind}_K^G V, k).$$

When V is finite-dimensional $\underline{\text{Hom}}(V, -) = \text{Hom}_k(V, -)$ is exact, so $\underline{\text{Ext}}^j(V, k) = 0$ for $j > 0$ and $\underline{\text{Ext}}^0(V, k) = V^*$. In this case the spectral sequence degenerates into the isomorphisms in 3.3, and we are done. \square

4 The top-dimensional derived functor

In this section we take G to be a p -adic Lie group of dimension $d = \dim_{\mathbb{Q}_p}(G)$, and we assume $\text{char}(k) = p$.

Remark 4.1. In part i of Proposition 6.2 we will show that, for any V in $\text{Mod}_k(K)$,

$$R^i \text{Ind}_K^G(V) = 0 \quad \forall i > d.$$

A natural question is whether it is possible to compute the top-dimensional derived functors $R^d \text{Ind}_K^G(V)$. For the trivial representation $V = k$ we have the following.

Proposition 4.2. *Assume G has a non-discrete center. Then:*

- i. $R^d \text{Ind}_K^G(k) = 0$ for all compact open subgroups $K \subset G$;
- ii. The category $\text{Mod}_k(G)$ has no nonzero projective objects.

Proof. For the proof of part one let $U \subset K$ be any open Poincaré subgroup, and let c be as in Remark 2.5 with $V = k$. We must find an open subgroup $U' \subset U$ such that $t_{U,U'}^d(c) = 0$. The corestriction map

$$\text{cor}_{K \cap x'U'x'^{-1}}^{K \cap x'Ux'^{-1}} : H^d(K \cap x'U'x'^{-1}, k) \longrightarrow H^d(K \cap x'Ux'^{-1}, k)$$

is known to be an isomorphism (of one-dimensional spaces) for all U' . Its composition with the restriction map $\text{res}_{K \cap x'U'x'^{-1}}^{K \cap x'Ux'^{-1}}$ is multiplication by the index. Thus $\text{res}_{K \cap x'U'x'^{-1}}^{K \cap x'Ux'^{-1}} = 0$ if this index is > 1 . To summarize, we must find $U' \subsetneq U$ such that we have strict inclusions

$$K \cap gU'g^{-1} \subsetneq K \cap gUg^{-1}$$

for all $g \in G$. Let Z denote the center of G . Intersecting both sides above with Z shows it is enough to pick a U' such that $Z \cap U' \subsetneq Z \cap U$. For example, for the right-hand side we get

$$Z \cap (K \cap gUg^{-1}) = K \cap g(Z \cap U)g^{-1} = K \cap (Z \cap U) = Z \cap U.$$

Since we are assuming Z is non-discrete $Z \cap U$ contains a non-identity element z say. Pick an open neighborhood $z(Z \cap U') \subset Z \cap U$ not containing the identity. Any such U' works.

For part two let W be an arbitrary object of $\text{Mod}_k(G)$ and let V be an object of $\text{Mod}_k(K)$ as before. Note that Ind_K^G takes injective objects to injective objects (since restriction is an exact left adjoint) and we therefore have a Grothendieck spectral sequence of the form

$$E_2^{j,i} = \text{Ext}_{\text{Mod}_k(G)}^j(W, R^i \text{Ind}_K^G(V)) \implies \text{Ext}_{\text{Mod}_k(K)}^{i+j}(W, V)$$

coming from Frobenius reciprocity. See [Vig96, p. 42] for example. If $\text{Hom}_{\text{Mod}_k(G)}(W, -)$ is exact we find that $E_2^{j,i} = 0$ for all $j > 0$, from which we deduce an isomorphism

$$\text{Hom}_{\text{Mod}_k(G)}(W, R^i \text{Ind}_K^G(V)) \simeq \text{Ext}_{\text{Mod}_k(K)}^i(W, V).$$

We specialize to the case $V = k$ and $i = d$. As we have just shown, the left-hand side vanishes in this case. For part two K only plays an auxiliary role and we may take it to be Poincaré. We infer that

$$\text{Hom}_k(H^0(K, W), k) \simeq \text{Ext}_{\text{Mod}_k(K)}^d(W, k) = 0 \implies H^0(K, W) = 0 \implies W = 0$$

by duality for Poincaré groups. See the review in [SS, Sect. 1] for instance. \square

Remark 4.3. Both parts of the previous Proposition clearly fail if G is discrete (as $d = 0$ and smooth G -representations are the same as abstract $k[G]$ -modules). We do not know whether the Proposition holds if we only assume G itself is non-discrete.

Also, a natural question in the context of Proposition 4.2 is whether every homotopically projective complex of objects in $\text{Mod}_k(G)$ is necessarily contractible. An affirmative answer would vastly generalize part ii of Proposition 4.2.

Theorem 1.1 in the introduction gives a supplement to Proposition 4.2 for p -adic reductive groups G .

5 The case of p -adic reductive groups

5.1 Notation, conventions, and background

In this article $\mathbb{N} = \{1, 2, 3, \dots\}$ denotes the set of all positive integers.

We let $\mathfrak{F}/\mathbb{Q}_p$ be a finite extension with valuation ring \mathfrak{O} , and we choose a uniformizer π . Take $\text{val}_{\mathfrak{F}}$ to be the valuation on \mathfrak{F} satisfying $\text{val}_{\mathfrak{F}}(\pi) = 1$. As usual $q = p^f$ is the cardinality of the residue field $\mathfrak{O}/\pi\mathfrak{O}$, and $e = e(\mathfrak{F}/\mathbb{Q}_p)$ denotes the ramification index.

In this section \mathbf{G} is an arbitrary connected reductive group defined over \mathfrak{F} , and $G = \mathbf{G}(\mathfrak{F})$. We choose a maximal \mathfrak{F} -split subtorus $\mathbf{S} \subset \mathbf{G}$ and let \mathbf{Z} denote its centralizer. Following our earlier convention, $S = \mathbf{S}(\mathfrak{F})$ and $Z = \mathbf{Z}(\mathfrak{F})$ denote the p -adic Lie groups of \mathfrak{F} -rational points.

Let Φ denote the roots of \mathbf{G} relative to \mathbf{S} , and select a subset of positive roots Φ^+ once and for all (then $\Phi^- = -\Phi^+$ is the set of negative roots). The root system may not be reduced, and we let Φ_{red} be the subset of reduced roots ($\alpha \in \Phi$ such that $\frac{1}{2}\alpha \notin \Phi$). By Φ_{red}^+ and Φ_{red}^- we mean the subsets of positive and negative reduced roots respectively.

Furthermore, we pick a special vertex x_0 in the apartment associated with \mathbf{S} , and consider the Bruhat-Tits group scheme $\mathbf{G}_{x_0}^\circ$ over \mathfrak{O} . The special subgroup $K_0 = \mathbf{G}_{x_0}^\circ(\mathfrak{O})$ and its principal congruence subgroups

$$K_m = \ker (\mathbf{G}_{x_0}^\circ(\mathfrak{O}) \longrightarrow \mathbf{G}_{x_0}^\circ(\mathfrak{O}/\pi^m\mathfrak{O}))$$

play a pivotal role. The argument proving [OS19, Cor. 7.8] applies verbatim and shows K_m is a uniform pro- p group if $m \in e\mathbb{N}$ and $m > e$ if $p = 2$. We give the proof below and compute the lower p -series for such K_m .

Let \mathbf{A} be the maximal \mathfrak{F} -split subtorus of the center of \mathbf{G} . We can arrange for x_0 to be the origin of the apartment $X_*(\mathbf{S})/X_*(\mathbf{A}) \otimes \mathbb{R}$. In this case the action of $z \in Z$ on the apartment is translation by the image of $\nu(z)$ where

$$\nu : Z \longrightarrow X_*(\mathbf{S}) \otimes \mathbb{R}$$

is the homomorphism for which

$$\langle \nu(z), \chi|_{\mathbf{S}} \rangle = -\text{val}_{\mathfrak{F}} \chi(z)$$

for all $\chi \in X^*(\mathbf{Z})$. We refer to [SSt, Sect. I.1] for more details. Later on we consider the monoid Z^+ of all $z \in Z$ such that $\langle \nu(z), \alpha \rangle \leq 0$ for all $\alpha \in \Phi^+$.

For each $\alpha \in \Phi$ we have a root subgroup \mathcal{U}_α of \mathbf{G} with \mathfrak{F} -rational points U_α ; the latter is normalized by Z . According to [KP23, Thm. 9.6.5] the root datum of \mathbf{G} carries a valuation

attached to x_0 . By [KP23, Df. 6.1.2] this gives rise to a descending filtration of U_α by subgroups $(U_{\alpha,r})_{r \in \mathbb{R}}$ satisfying

$$(1) \quad zU_{\alpha,r}z^{-1} = U_{\alpha,r-\langle \nu(z), \alpha \rangle} \quad \text{for any } z \in Z.^1$$

This filtration needs to be modified. For any concave function f on $\Phi \cup \{0\}$ [KP23, Df. 7.3.3] introduces the subgroup

$$U_{\alpha,f} := U_{\alpha,x_0,f} := U_{\alpha,f(\alpha)} \cdot U_{2\alpha,f(2\alpha)}$$

of U_α . For any real number $r \geq 0$ the constant function f_r with value r is concave, and we abbreviate $\tilde{U}_{\alpha,r} := U_{\alpha,f_r}.$ ²

Before [KP23, Lem. 9.8.1] a descending filtration $(Z_r)_{r \geq 0}$ of Z is constructed. We point out that this filtration depends on the connected reductive group \mathbf{Z} and not the ambient group \mathbf{G} . Since the Bruhat-Tits building of Z is a single point ([KP23, Prop. 9.3.9]), namely x_0 , it follows from [KP23, Prop. 13.2.5, part (2)] that each Z_r is a normal subgroup of Z .

Using these filtrations [KP23, Def. 7.3.3] then introduces a descending filtration $(\mathcal{P}_{x_0,r})_{r \geq 0}$ of $K_0 = \mathbf{G}_{x_0}^\circ(\mathfrak{O})$. The crucial property of these filtrations is the Iwahori factorization ([KP23, Prop. 13.2.5, part (3)]): For any $r > 0$ the multiplication map defines a homeomorphism

$$(2) \quad \prod_{\alpha \in \Phi_{\text{red}}^-} \tilde{U}_{\alpha,r} \times Z_r \times \prod_{\alpha \in \Phi_{\text{red}}^+} \tilde{U}_{\alpha,r} \xrightarrow{\sim} \mathcal{P}_{x_0,r}.$$

(The factors in each product are ordered in a fixed but arbitrary way.)

Lemma 5.1. *For any $m > 0$ we have $K_m = \mathcal{P}_{x_0,m}$.*

Proof. By [KP23, Prop. 9.8.3] there is, for any $r \geq 0$, a smooth affine \mathfrak{O} -group scheme of finite type $\mathcal{G}_{x_0,r}$ such that $\mathcal{G}_{x_0,r}(\mathfrak{O}) = \mathcal{P}_{x_0,r}$. It comes by descend from the maximal unramified extension \mathfrak{F}^{ur} of \mathfrak{F} . By [KP23, Prop. 8.5.16, Df. A.5.12] and passing to $\text{Gal}(\mathfrak{F}^{\text{ur}}/\mathfrak{F})$ -invariants we have

$$\mathcal{G}_{x_0,r+m}(\mathfrak{O}) = \ker(\mathcal{G}_{x_0,r}(\mathfrak{O}) \longrightarrow \mathcal{G}_{x_0,r}(\mathfrak{O}/\pi^m \mathfrak{O})).$$

Now take $r = 0$ and observe that $\mathcal{G}_{x_0,0} = \mathbf{G}_{x_0}^\circ$ since $K_0 = \mathcal{P}_{x_0,0}$. □

As a consequence we obtain from (2) the Iwahori factorization

$$(3) \quad \prod_{\alpha \in \Phi_{\text{red}}^-} \tilde{U}_{\alpha,m} \times Z_m \times \prod_{\alpha \in \Phi_{\text{red}}^+} \tilde{U}_{\alpha,m} \xrightarrow{\sim} K_m \quad \text{for any } m > 0.$$

Remark 5.2. For any $m > 0$ and $\alpha \in \Phi_{\text{red}}$ the filtration subgroups $\tilde{U}_{\alpha,m}$ and Z_m are themselves principal congruence subgroups. For Z_m this is a special case of Lemma 5.1 since Z_m is the analog of $\mathcal{P}_{x_0,m}$ for the connected reductive \mathfrak{F} -group \mathbf{Z} .

¹The ν in [KP23, Df. 6.1.2 V6] is the composite of our ν and the canonical splitting of the natural monomorphism $X_*(\mathbf{S}') \otimes \mathbb{R} \hookrightarrow X_*(\mathbf{S}) \otimes \mathbb{R}$ where \mathbf{S}' is the maximal subtorus of \mathbf{S} contained in the derived subgroup of \mathbf{G} ([KP23, 4.1.4]). This canonical splitting exists since the difference between \mathbf{S} and \mathbf{S}' , up to isogeny, comes from the center of \mathbf{G} . Hence our number $\langle \nu(z), \alpha \rangle$, for a root $\alpha \in \Phi$, coincides with $\alpha(\nu(z))$ in [KP23, Df. 6.1.2 V6]. Note that [KP23] contains a sign mistake as explained in [Kal, 2nd item on p. 23].

²[KP23, Sect. 13.2] replaces f_r again by r in the notation, which we find too confusing. For simplicity we drop the point x_0 , which we fixed once and for all, from the notation.

To see that $\tilde{U}_{\alpha,m}$ is a principal congruence subgroup we argue as follows. By [KP23, Prop. 2.11.9, Prop. 9.8.3] there is, for $r \geq 0$, a unique closed subgroup scheme $\mathcal{U}_{\alpha,x_0,r} \subset \mathcal{G}_{x_0,r}$ such that $\mathcal{U}_{\alpha,x_0,r}(\mathfrak{O}) = \tilde{U}_{\alpha,r}$. Because $\mathcal{U}_{\alpha,x_0,r}$ is closed the map $\mathcal{U}_{\alpha,x_0,r}(\mathfrak{O}/\pi^m\mathfrak{O}) \hookrightarrow \mathcal{G}_{x_0,r}(\mathfrak{O}/\pi^m\mathfrak{O})$ is still an inclusion and therefore

$$\begin{aligned}\tilde{U}_{\alpha,m} &= \mathcal{P}_{x_0,m} \cap \tilde{U}_{\alpha,0} = K_m \cap \tilde{U}_{\alpha,0} \\ &= \ker(\mathbf{G}_{x_0}^\circ(\mathfrak{O}) \longrightarrow \mathbf{G}_{x_0}^\circ(\mathfrak{O}/\pi^m\mathfrak{O})) \cap \tilde{U}_{\alpha,0} \\ &= \ker(\mathcal{U}_{\alpha,x_0,0}(\mathfrak{O}) \longrightarrow \mathbf{G}_{x_0}^\circ(\mathfrak{O}/\pi^m\mathfrak{O})) \\ &= \ker(\mathcal{U}_{\alpha,x_0,0}(\mathfrak{O}) \longrightarrow \mathcal{U}_{\alpha,x_0,0}(\mathfrak{O}/\pi^m\mathfrak{O})).\end{aligned}$$

The first equality follows from [KP23, Prop. 13.2.5, part (3)], applied to $\mathcal{P}_{x_0,m}$, after comparing the components associated with the root α . The second equality is Lemma 5.1.

5.2 Some uniform subgroups and their cohomology

We begin with the following variant of [OS19, Cor. 7.7]. Let $\mathcal{G} = \text{Spec}(A)$ be a smooth affine \mathfrak{O} -group scheme, and $\hat{\mathcal{G}} = \text{Spf}(\hat{A}^{\mathfrak{p}})$ be its formal completion in the unit section. Upon choosing an isomorphism $\hat{A}^{\mathfrak{p}} \simeq \mathfrak{O}[[X_1, \dots, X_\delta]]$ we get a homeomorphism $\xi : \hat{\mathcal{G}}(\mathfrak{O}) \xrightarrow{\sim} (\pi\mathfrak{O})^\delta$. We let

$$\hat{\mathcal{G}}_m(\mathfrak{O}) := \xi^{-1}((\pi^m\mathfrak{O})^\delta) = \ker(\mathcal{G}(\mathfrak{O}) \longrightarrow \mathcal{G}(\mathfrak{O}/\pi^m\mathfrak{O})).$$

With this notation we have the following slight generalization of [OS19, Cor. 7.7].

Lemma 5.3. *Let $m \in e\mathbb{N}$, and assume $m > e$ if $p = 2$. Then the congruence subgroup $\hat{\mathcal{G}}_m(\mathfrak{O})$ is a uniform pro- p group. Furthermore*

$$\hat{\mathcal{G}}_{m+e}(\mathfrak{O}) = \hat{\mathcal{G}}_m(\mathfrak{O})^p = \{g^p : g \in \hat{\mathcal{G}}_m(\mathfrak{O})\}.$$

(This gives the lower p -series for $\hat{\mathcal{G}}_m(\mathfrak{O})$ by iteration.)

Proof. The fact that $\hat{\mathcal{G}}_m(\mathfrak{O})$ is uniform is precisely the content of [OS19, Cor. 7.7]. Here we compute its lower p -series. By considering $\mathcal{G}_0 = \text{Res}_{\mathfrak{O}/\mathbb{Z}_p}\mathcal{G}$, and noting that $\hat{\mathcal{G}}_{0,j}(\mathbb{Z}_p) = \hat{\mathcal{G}}_m(\mathfrak{O})$ if $m = ej$, we may (and will) assume $\mathfrak{O} = \mathbb{Z}_p$.

We first deal with the case $p > 2$. As explained in [OS19, 7.2.1] the group $\hat{\mathcal{G}}(\mathbb{Z}_p)$ is a standard group in the sense of [DdSMS, Df. 8.22]. The (last paragraph of the) proof of [DdSMS, Thm. 8.31] shows that $\hat{\mathcal{G}}(\mathbb{Z}_p)^{p^{m-1}} = \hat{\mathcal{G}}_m(\mathbb{Z}_p)$. (See also part (iii) of [DdSMS, Thm. 3.6] which gives the lower p -series of a uniform group.) Raising both sides to the p^{th} power gives the result.

For $p = 2$ we work with $\hat{\mathcal{G}}_2(\mathbb{Z}_2)$ which again is standard for the same reason. The proof of [DdSMS, Thm. 8.31] (with $\varepsilon = 1$) also shows that $\hat{\mathcal{G}}_2(\mathbb{Z}_2)^{2^{m-2}} = \hat{\mathcal{G}}_m(\mathbb{Z}_2)$ for $m \geq 2$. Taking squares gives the result. \square

In particular K_m as well as $\tilde{U}_{\alpha,m}$ and Z_m (by Remark 5.2) all are uniform pro- p groups if $m \in e\mathbb{N}$ and $m > e$ if $p = 2$; moreover

$$(4) \quad K_m^p = K_{m+e}, \quad Z_m^p = Z_{m+e}, \quad \text{and} \quad \tilde{U}_{\alpha,m}^p = \tilde{U}_{\alpha,m+e}.$$

Next we will see that in certain situations intersections of uniform subgroups are uniform. In the following we put $\wp := p$ if $p > 2$, and $\wp := 4$ if $p = 2$.

Lemma 5.4. *Let $(A, \|\cdot\|)$ be a finite dimensional normed \mathbb{Q}_p -algebra whose norm $\|\cdot\|$ is submultiplicative and let $A_0 := \{a \in A : \|a\| \leq \wp\}$. Fix a closed subgroup $\Gamma \leq 1 + A_0$. Let K, K' be uniform open subgroups of Γ , and $a \in A^\times$ an arbitrary unit. If $a^{-1}K'a \cap K$ is open in K then it is uniform.*

Proof. First of all note that A_0 is a open pro- p subgroup of the additive group A , and $1 + A_0$ is an open pro- p subgroup of the multiplicative group A^\times . It is well known (cf. [DdSMS, Prop. 6.22 and Cor. 6.25]) that the usual exponential and logarithm power series induce homeomorphisms

$$\exp : A_0 \longrightarrow 1 + A_0 \quad \text{and} \quad \log : 1 + A_0 \longrightarrow A_0 ,$$

which are inverse to each other.

We recall that a pro- p group H is uniform if and only if the following conditions are satisfied ([DdSMS, Thm. 4.5]):

- (1) H is (topologically) finitely generated;
- (2) H is torsion-free;
- (3) H is powerful – which means $H/\overline{H^\wp}$ is abelian.

Here H^n , for $n \in \mathbb{N}$, is a priori the subgroup generated by the set of n -powers. However, when H is uniform H^\wp is the same as the set of all \wp -powers by [DdSMS, Lemma 3.4] ; the same result shows that H^\wp is open.

Obviously the intersection is torsion-free since K is. Since $a^{-1}K'a \cap K$ is open in K by assumption it is finitely generated by [DdSMS, Prop. 1.7] . It remains to show it is powerful, so choose two elements $x, y \in a^{-1}K'a \cap K$ arbitrarily. It suffices to show the commutator $[x, y]$ is a \wp -power from $a^{-1}K'a \cap K$. Since K and $a^{-1}K'a \cong K'$ are uniform we can write

$$[x, y] = \kappa^\wp = (a^{-1}\kappa'a)^\wp$$

for suitable $\kappa \in K$ and $\kappa' \in K'$. We just need to argue that $\kappa = a^{-1}\kappa'a$. Taking log above, using [DdSMS, Cor. 6.25(iii)], yields $\log[x, y] = \wp \log \kappa$. Since $a^{-1}\kappa'a$ may not lie in $1 + A_0$ we have to note that nevertheless $\log(a^{-1}\kappa'a)$ makes sense. Indeed the series

$$\log(a^{-1}\kappa'a) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} (a^{-1}\kappa'a - 1)^n = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} a^{-1}(\kappa' - 1)^n a = a^{-1} \log(\kappa') a$$

still converges. Having checked this, taking log as above we deduce $\log[x, y] = \wp \log(a^{-1}\kappa'a)$. Comparing the two expressions for $\log[x, y] \in \wp A_0$ and dividing by \wp (in A) we infer the equality $\log(\kappa) = \log(a^{-1}\kappa'a)$ in A_0 . Finally take exp on both sides, again noting that

$$\exp(\log(a^{-1}\kappa'a)) = \exp(a^{-1} \log(\kappa') a) = a^{-1} \exp(\log(\kappa')) a = a^{-1} \kappa' a .$$

This gives the equality $\kappa = a^{-1}\kappa'a$ as desired. We conclude $a^{-1}K'a \cap K$ that is uniform. \square

Example 5.5. *Our main example is the matrix algebra $A = M_N(\mathbb{Q}_p)$ with the norm $\|a\| = \max_{i,j} |a_{ij}|$. The multiplicative group $1 + A_0$ is the congruence subgroup $\ker(\mathrm{GL}_N(\mathbb{Z}_p) \rightarrow \mathrm{GL}_N(\mathbb{Z}_p/\wp\mathbb{Z}_p))$. Then Lemma 5.4 implies the following: If K and K' are uniform closed subgroups of $\ker(\mathrm{GL}_N(\mathbb{Z}_p) \rightarrow \mathrm{GL}_N(\mathbb{Z}_p/\wp\mathbb{Z}_p))$ then so is $a^{-1}K'a \cap K$ for any $a \in \mathrm{GL}_N(\mathbb{Q}_p)$ such that $a^{-1}K'a \cap K$ is open in K .*

Proposition 5.6. *Let $m, n \in e\mathbb{N}$ assuming $m, n > e$ if $p = 2$; we then have:*

- a. $K_m \cap gK_n g^{-1}$ is uniform for all $g \in G$;
- b. $\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1}$ is uniform for all $\alpha \in \Phi$ and $z \in Z$,

Proof. As we noted before (4) all groups of which we take intersections are uniform. The assertion then follows from the above Example by using a faithful representation of the Weil restriction to \mathbb{Z}_p of $\mathbf{G}_{x_0}^\circ$ into some GL_N . \square

Throughout we fix a field k of characteristic p . A uniform pro- p group U admits a canonical p -valuation defined by the lower p -series. Thus U becomes equi- p -valued, and by [Laz, Ch. V, Prop. 2.5.7.1] the cup product gives an isomorphism of graded k -algebras

$$\bigwedge H^1(U, k) \xrightarrow{\sim} H^*(U, k).$$

Moreover $H^1(U, k) = \mathrm{Hom}_{\mathbb{F}_p}(U/U^p, k)$ is the dual Frattini quotient, which we will henceforth denote U_Φ^* . (Here the subscript Φ is standard notation for Frattini quotients, and should not be confused with the root system.) Our exterior products $\bigwedge = \bigwedge_k$ are always over k .

Proposition 5.7. *Consider arbitrary $m, n \in e\mathbb{N}$, both assumed to be $> e$ if $p = 2$, and $z \in Z$. Then $H^i(K_m \cap zK_n z^{-1}, k)$ decomposes as a direct sum*

$$\bigoplus \left[\bigotimes_{\alpha \in \Phi_{\mathrm{red}}^-}^{a_\alpha} \bigwedge (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* \otimes \bigwedge (Z_{\max\{m, n\}})_\Phi^* \otimes \bigotimes_{\alpha \in \Phi_{\mathrm{red}}^+}^{c_\alpha} \bigwedge (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* \right]$$

where $\{a_\alpha\}_{\alpha \in \Phi_{\mathrm{red}}^-}$, b , $\{c_\alpha\}_{\alpha \in \Phi_{\mathrm{red}}^+}$ in the sum \bigoplus range over non-negative integers with sum i .

Proof. Conjugating the Iwahori factorization of K_n by z , using that Z_n is normal in Z , and intersecting with K_m we find that

$$(5) \quad \left(\prod_{\alpha \in \Phi_{\mathrm{red}}^-} \tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1} \right) \times Z_{\max\{m, n\}} \times \left(\prod_{\alpha \in \Phi_{\mathrm{red}}^+} \tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1} \right) \xrightarrow{\sim} K_m \cap zK_n z^{-1}.$$

By the same argument as in the proof of [OS19, Cor. 7.11], which uses the uniformity of all groups involved, quotienting out p -powers induces an isomorphism on the level of Frattini quotients. Taking k -linear duals and forming the exterior algebra gives an isomorphism of graded k -algebras

$$\begin{aligned} & \bigwedge (K_m \cap zK_n z^{-1})_\Phi^* \xrightarrow{\sim} \\ & \bigwedge \left[\bigoplus_{\alpha \in \Phi_{\mathrm{red}}^-} (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* \oplus (Z_{\max\{m, n\}})_\Phi^* \oplus \bigoplus_{\alpha \in \Phi_{\mathrm{red}}^+} (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* \right] \simeq \\ & \bigotimes_{\alpha \in \Phi_{\mathrm{red}}^-} \bigwedge (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* \otimes \bigwedge (Z_{\max\{m, n\}})_\Phi^* \otimes \bigotimes_{\alpha \in \Phi_{\mathrm{red}}^+} \bigwedge (\tilde{U}_{\alpha, m} \cap z\tilde{U}_{\alpha, n} z^{-1})_\Phi^* . \end{aligned}$$

Comparing the degree i graded pieces gives the result. \square

5.3 On the vanishing of certain restriction maps

We fix an $m \in e\mathbb{N}$. If $p = 2$ we always assume $m > e$. For $n, n' \in e\mathbb{N}$ such that $n \leq n'$ we have restriction maps

$$\text{res}_{n,n'}^i(g) : H^i(K_m \cap gK_n g^{-1}, k) \longrightarrow H^i(K_m \cap gK_{n'} g^{-1}, k).$$

When $g = z \in Z$, this map is compatible with the decomposition in Prop. 5.7 in the obvious sense. (In general restriction commutes with cup products in group cohomology.)

Lemma 5.8. *Suppose $m \leq n < n'$ all lie in $e\mathbb{N}$ (and are $> e$ if $p = 2$). Then $\text{res}_{n,n'}^i(g) = 0$ for all $g \in G$ and $i > i_0 := \dim_{\mathbb{Q}_p}(G/P_{\min})$.*

Proof. By the Cartan decomposition $G = K_0 Z^+ K_0$, as described in [KP23, Thm. 5.2.1, part (1)] for example, we may write $g = hzh'$ for some $z \in Z^+$ and $h, h' \in K_0$. Since K_m and K_n are both normal in K_0 we note that

$$K_m \cap gK_n g^{-1} = h(K_m \cap zK_n z^{-1})h^{-1}.$$

Therefore $x \mapsto hx$ induces isomorphisms h_* on cohomology which fit in the following commutative diagram, where the horizontal maps are the restriction maps:

$$\begin{array}{ccc} H^i(K_m \cap gK_n g^{-1}, k) & \longrightarrow & H^i(K_m \cap gK_{n'} g^{-1}, k) \\ \uparrow h_* & & \uparrow h_* \\ H^i(K_m \cap zK_n z^{-1}, k) & \longrightarrow & H^i(K_m \cap zK_{n'} z^{-1}, k). \end{array}$$

Fix an $i > i_0$. Our claim that $\text{res}_{n,n'}^i(g) = 0$ is therefore equivalent to $\text{res}_{n,n'}^i(z) = 0$, which we now proceed to show using the decomposition in Proposition 5.7.

Consider $\text{res}_{n,n'}^i(z)$ restricted to the piece of $H^i(K_m \cap zK_n z^{-1}, k)$ indexed by $\{a_\alpha\}_{\alpha \in \Phi_{\text{red}}^-}$, b , $\{c_\alpha\}_{\alpha \in \Phi_{\text{red}}^+}$. Observe that

- $\bigwedge^b(Z_n)_\Phi^* \longrightarrow \bigwedge^b(Z_{n'})_\Phi^*$ vanishes for $b > 0$ since $Z_{n'} \subseteq Z_{n+e} = (Z_n)^p$ by (4).
- When $\alpha \in \Phi_{\text{red}}^+$ we have $\langle \nu(z), \alpha \rangle \leq 0$. In particular

$$z\tilde{U}_{\alpha,n}z^{-1} = U_{\alpha,n-\langle \nu(z), \alpha \rangle} \cdot U_{2\alpha,n-\langle \nu(z), 2\alpha \rangle} \subseteq \tilde{U}_{\alpha,n-\langle \nu(z), \alpha \rangle} \subseteq \tilde{U}_{\alpha,n} \subseteq \tilde{U}_{\alpha,m},$$

so that

$$\tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n}z^{-1} = z\tilde{U}_{\alpha,n}z^{-1}.$$

For such α the map in question is

$$\bigwedge^{c_\alpha} (z\tilde{U}_{\alpha,n}z^{-1})_\Phi^* \longrightarrow \bigwedge^{c_\alpha} (z\tilde{U}_{\alpha,n'}z^{-1})_\Phi^*$$

which vanishes for $c_\alpha > 0$ since

$$z\tilde{U}_{\alpha,n'}z^{-1} \subseteq z\tilde{U}_{\alpha,n+e}z^{-1} = (z\tilde{U}_{\alpha,n}z^{-1})^p,$$

again by (4).

If $\text{res}_{n,n'}^i(z) \neq 0$ there must be a piece of cohomology with $b = 0$ and all $c_\alpha = 0$ where $\text{res}_{n,n'}^i(z)$ is nonzero. Since $\sum_{\alpha \in \Phi_{\text{red}}^-} a_\alpha = i$ and all $a_\alpha \leq \dim_{\mathbb{Q}_p} U_\alpha (= \dim_{\mathbb{Q}_p} (\tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n}z^{-1}))$ we conclude that

$$i \leq \sum_{\alpha \in \Phi_{\text{red}}^-} \dim_{\mathbb{Q}_p} U_\alpha = \dim_{\mathbb{Q}_p} (G/P_{\min}) = i_0.$$

See [Bor, Sect. 21.11] for the first equality. \square

Next we show the above bound i_0 is sharp, as made precise in the result below.

Lemma 5.9. *Given $m \leq n \leq n'$ in $e\mathbb{N}$ (all $> e$ if $p = 2$) there exists $z \in Z^+$ satisfying*

$$-\langle \nu(z), \alpha \rangle \leq m - n', \quad \forall \alpha \in \Phi^-.$$

For such z , $\text{res}_{n,n'}^i(z) \neq 0$ for all $i \leq i_0$.

Proof. For the existence part, take $z := \mu(\pi) \in S \subseteq Z$ where $\mu \in X_*(\mathbf{S})^+$ is a dominant cocharacter satisfying $\langle \mu, \alpha \rangle \leq m - n'$ for all $\alpha \in \Phi^-$. We note that $X^*(\mathbf{Z}) \rightarrow X^*(\mathbf{S})$ has finite cokernel, so some multiple of α extends to an \mathfrak{F} -rational character of \mathbf{Z} , and therefore $-\langle \nu(\mu(\pi)), \alpha \rangle = \text{val}_{\mathfrak{F}} \alpha(\mu(\pi)) = \langle \mu, \alpha \rangle$ by the defining property of ν . We see that $z \in S \cap Z^+$.

For the non-vanishing part, we first work out the case $i = i_0$. Consider the contribution to $H^{i_0}(K_m \cap zK_n z^{-1}, k)$ with indices $a_\alpha = \dim_{\mathbb{Q}_p} U_\alpha$ for all $\alpha \in \Phi_{\text{red}}^-$ (and therefore $b = 0$ and $c_\alpha = 0$ for $\alpha \in \Phi_{\text{red}}^+$) which is the line

$$\mathcal{L}_{z,n} := \bigotimes_{\alpha \in \Phi_{\text{red}}^-}^{\text{top}} (\tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n}z^{-1})_{\Phi}^*.$$

We are assuming $z \in Z^+$ satisfies the inequalities $-\langle \nu(z), \alpha \rangle \leq m - n'$ for all $\alpha \in \Phi^-$ (even the non-reduced roots). Under this assumption we have

$$U_{\alpha,m} \subseteq zU_{\alpha,n'}z^{-1} \subseteq zU_{\alpha,n}z^{-1}$$

since $m \geq n' - \langle \nu(z), \alpha \rangle$, and similarly for 2α if it is a root. As a result

$$\tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n}z^{-1} = \tilde{U}_{\alpha,m} = \tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n'}z^{-1}$$

and consequently $\text{res}_{n,n'}^{i_0}(z)$ maps $\mathcal{L}_{z,n}$ isomorphically to the line $\mathcal{L}_{z,n'}$ in $H^{i_0}(K_m \cap zK_{n'}z^{-1}, k)$.

In particular $\text{res}_{n,n'}^{i_0}(z)$ is nonzero on $\mathcal{L}_{z,n}$ for all such z .

For $i \leq i_0$ we generalize the argument from the previous paragraph as follows. Once and for all we write $i = \sum_{\alpha \in \Phi_{\text{red}}^-} a_\alpha$ for a choice of integers $0 \leq a_\alpha \leq \dim_{\mathbb{Q}_p} U_\alpha$. One way of doing this is to list the roots, $\Phi_{\text{red}}^- = \{\alpha_1, \alpha_2, \dots\}$. Then let q be the largest index for which

$$\dim_{\mathbb{Q}_p} U_{\alpha_1} + \dots + \dim_{\mathbb{Q}_p} U_{\alpha_q} \leq i.$$

By convention $q := 0$ if $i < \dim_{\mathbb{Q}_p} U_{\alpha_1}$. Now let $a_{\alpha_j} := \dim_{\mathbb{Q}_p} U_{\alpha_j}$ for $j \leq q$. If $i < i_0$ we let

$$a_{\alpha_{q+1}} := i - (\dim_{\mathbb{Q}_p} U_{\alpha_1} + \dots + \dim_{\mathbb{Q}_p} U_{\alpha_q}).$$

If there are any remaining roots α_j with $j > q + 1$ we declare that $a_{\alpha_j} := 0$.

Having chosen this expansion $i = \sum_{\alpha \in \Phi_{\text{red}}^-} a_\alpha$, we introduce the following auxiliary subspace of $H^i(K_m \cap zK_n z^{-1}, k)$:

$$\mathcal{V}_{z,n} := \bigotimes_{\alpha \in \Phi_{\text{red}}^-} \bigwedge^{a_\alpha} (\tilde{U}_{\alpha,m} \cap z\tilde{U}_{\alpha,n} z^{-1})_\Phi^*.$$

The restriction map $\text{res}_{n,n'}^i(z)$ restricts to an isomorphism $\mathcal{V}_{z,n} \xrightarrow{\sim} \mathcal{V}_{z,n'}$ for z as above. In particular $\text{res}_{n,n'}^i(z) \neq 0$ as claimed. \square

5.4 The proof of Theorem 1.2

We can now prove our main result for p -adic reductive groups, which we recall here:

Theorem 5.10. *Fix an $m \in e\mathbb{N}$ ($> e$ if $p = 2$) and let $i_0 = \dim_{\mathbb{Q}_p}(G/P_{\min})$ as above. Then:*

- (a) $R^i \text{Ind}_{K_m}^G(k) = 0$ for all $i > i_0$;
- (b) $R^i \text{Ind}_{K_m}^G(k) \neq 0$ for all $i \leq i_0$.

Proof. To show the vanishing in part (a) suppose $i > i_0$. Let $n \in (e\mathbb{N})_{\geq m}$ be arbitrary and consider any

$$c \in \varprojlim_{g \in G/K_n} H^i(K_m \cap gK_n g^{-1}, k)$$

as in Remark 2.5. It suffices to show $t_{K_n, K_{n'}}^i(c) = 0$ for all $n' \in (e\mathbb{N})_{>n}$, which follows from Lemma 5.8 since

$$t_{K_n, K_{n'}}^i(c)_{gK_{n'}} = \text{res}_{n,n'}^i(g)(c_{gK_n}) = 0$$

for all $g \in G$. (We have used the formula for the transition maps $t_{K_n, K_{n'}}^i$ given in 2.5.)

For the non-vanishing in part (b) now suppose $i \leq i_0$ and pick some $n \in (e\mathbb{N})_{\geq m}$ once and for all. We will construct a nonzero class c which survives all the transition maps. That is, such that $t_{K_n, K_{n'}}^i(c) \neq 0$ for all $n' \in (e\mathbb{N})_{>n}$.

The class c will be prescribed on a set of representatives \mathcal{X} for Z^+/Z^0 . (For comparison Z^0 is shorthand notation for the subgroup denoted by $Z(\mathfrak{F})^0$ in [KP23, Df. 2.6.23].) We recall the Cartan decomposition $G = \bigsqcup_{z \in \mathcal{X}} K_0 z K_0$.

For every $z \in \mathcal{X}$ we once and for all select a nonzero vector $v_z \in \mathcal{V}_{z,n}$ (recall the notation introduced in the proof of Lemma 5.9). Corresponding to these data there is a unique groupoid cohomology class

$$c : G/K_n \longrightarrow \bigoplus_{g \in G/K_n} H^i(K_m \cap gK_n g^{-1}, k)$$

with the following properties:

- c is K_m -equivariant (in the sense described in Remark 2.5);
- c is supported on $K_m \mathcal{X} K_n / K_n$;
- $c_{zK_n} = v_z$ for all $z \in \mathcal{X}$.

The uniqueness of c is clear. For its existence let $g \in K_m \mathcal{X} K_n$ and write $g = hzh'$ accordingly. Thus $h \in K_m$, $h' \in K_n$, and $z \in \mathcal{X}$. By the Cartan decomposition the factor z is uniquely determined, and the left coset $h(K_m \cap zK_n z^{-1})$ is independent of the factorization of g . Now let $c_{gK_n} = h_*(v_z)$ which is therefore well-defined.

To see this c has the desired property let $n' \in (e\mathbb{N})_{>n}$. Our task is to verify the tuple $t_{K_n, K_{n'}}^i(c)$ has at least one nonzero component. Take $z \in \mathcal{X}$ to be an element such that $-\langle \nu(z), \alpha \rangle \leq m - n'$ for all $\alpha \in \Phi^-$. Then by (the proof of) Lemma 5.9 we know that $\text{res}_{n, n'}^i(z)$ restricts to an isomorphism $\mathcal{V}_{z, n} \xrightarrow{\sim} \mathcal{V}_{z, n'}$. In particular

$$t_{K_n, K_{n'}}^i(c)_{zK_{n'}} = \text{res}_{n, n'}^i(z)(c_{zK_n}) = \text{res}_{n, n'}^i(z)(v_z) \neq 0$$

which finishes the proof. \square

We finish this section by emphasizing an application to duality, which essentially reproves and strengthens one of the main results from [Sor].

Corollary 5.11. *Let $m \in e\mathbb{N}$ ($m > e$ if $p = 2$). Then $\underline{\text{Ext}}^i(\text{ind}_{K_m}^G k, k) = 0 \iff i > i_0$.*

Proof. Due to Corollary 3.3 this is just a restatement of Theorem 5.10. \square

5.5 The proof of Theorem 1.1

By assumption \mathbf{G} is nontrivial and connected. Hence the group Z has positive \mathbb{Q}_p -dimension. To show $\text{Mod}_k(G)$ has no nonzero projective objects, the proof of Proposition 4.2 applies, noting that $R^d \text{Ind}_{K_m}^G(k) = 0$. Indeed, by (3) for instance,

$$d = \sum_{\alpha \in \Phi_{\text{red}}^-} \dim_{\mathbb{Q}_p} U_\alpha + \dim_{\mathbb{Q}_p} Z + \sum_{\alpha \in \Phi_{\text{red}}^+} \dim_{\mathbb{Q}_p} U_\alpha = 2i_0 + \dim_{\mathbb{Q}_p} Z > i_0.$$

We immediately deduce from Theorem 5.10 that $R^d \text{Ind}_{K_m}^G(k) = 0$.

Remark 5.12. There is a quicker route to Theorem 1.1 by directly showing $R^d \text{Ind}_{K_m}^G(k) = 0$. This amounts to verifying the condition appearing in the proof of Proposition 4.2. Indeed, for all large enough n there is an $n' > n$ (for instance $n' = n + 1$) such that we have a *strict* inclusion

$$K_m \cap gK_{n'}g^{-1} \subsetneq K_m \cap gK_n g^{-1}$$

for all $g \in G$. Indeed, by the Cartan decomposition it suffices to check this for $g = z \in Z^+$. For such z the strict inclusion follows from the factorization (5) by noting that $Z_{n'} \subsetneq Z_n$.

6 The derived functor $R\underline{\text{Ind}}$

6.1 Preliminary remarks

We return to the general setup and let G be an arbitrary p -adic Lie group of \mathbb{Q}_p -dimension d , and k is still a field of characteristic p . Recall that $\text{Mod}_k(G)$ is the abelian category of smooth G -representations on k -vector spaces. We let $D(G)$ denote its (unbounded) derived category.

We remind the reader that our convention is that G acts on $\text{Ind}_K^G(V)$ by right translations. Thus, for any V in $\text{Mod}_k(K)$, the space $\text{Ind}_K^G(V)$ consists of all functions $f : G \rightarrow V$ such that

- $f(\kappa g) = \kappa f(g)$ for any $g \in G$ and $\kappa \in K$;
- $f(gu) = f(g)$ for any $g \in G$ and $u \in U$ for some open $U \leq G$ (depending on f).

The fact that Ind_K^G is right adjoint to restriction is a consequence of Frobenius reciprocity (see [Vig96, I.5.7(i)] for example) which is the isomorphism

$$(6) \quad \begin{aligned} \text{Hom}_{\text{Mod}_k(K)}(W, V) &\xrightarrow{\cong} \text{Hom}_{\text{Mod}_k(G)}(W, \text{Ind}_K^G(V)) \\ A &\longmapsto B(v)(g) := A(gv) \end{aligned}$$

for V in $\text{Mod}_k(K)$ and W in $\text{Mod}_k(G)$. The functor Ind_K^G is left exact and preserves injective objects (see [Vig96, I.5.9]). If $K' \leq K$ then $\text{Ind}_K^G(V) \subseteq \text{Ind}_{K'}^G(V)$. We may therefore take the union over K and introduce the left exact functor

$$\begin{aligned} \underline{\text{Ind}} : \text{Mod}_k(G) &\longrightarrow \text{Mod}_k(G) \\ V &\longmapsto \varinjlim_K \text{Ind}_K^G(V) . \end{aligned}$$

Later we will endow $\underline{\text{Ind}}(V)$ with a smooth $G \times G$ -action, provided V is in $\text{Mod}(G)$. We are interested in the derived functors $R^i \underline{\text{Ind}}$ and their relation to $R^i \text{Ind}_K^G$.

Lemma 6.1. $R^i \underline{\text{Ind}}(V) = \varinjlim_K R^i \text{Ind}_K^G(V)$.

Proof. This is immediate from the exactness of inductive limits and the fact that any injective object in $\text{Mod}_k(G)$ remains injective in any $\text{Mod}_k(K)$. (Indeed, the left adjoint of restriction is compact induction, which is exact since $k[G]$ is flat over $k[K]$.) \square

As a preliminary observation, we prove that $R^i \underline{\text{Ind}}$ vanishes for $i > d$:

Proposition 6.2. *The following holds:*

- i. *For any compact open subgroup $K \subseteq G$, the functor Ind_K^G has cohomological dimension at most d .*
- ii. *The functor $\underline{\text{Ind}}$ has cohomological dimension at most d .*

Proof. This is a consequence of Proposition 2.4, but we prefer to include the following direct argument (which also gives an alternate proof of the derived Mackey decomposition 2.4).

i. Consider any V in $\text{Mod}_k(K)$ and let $U \subseteq G$ be a compact open subgroup. We choose a set $R \subseteq G$ of representatives for the double cosets $K \backslash G / U$. The Mackey decomposition (see [Vig96, I.5.5] for example) is a natural isomorphism

$$\begin{aligned} \text{Ind}_K^G(V)^U &\xrightarrow{\cong} \prod_{x \in R} V^{K \cap x U x^{-1}} \\ f &\longmapsto (f(x))_{x \in R} . \end{aligned}$$

Let $V \xrightarrow{\text{qis}} \mathcal{J}^\bullet$ be a choice of injective resolution in $\text{Mod}_k(K)$. On the one hand it remains an injective resolution in any $\text{Mod}_k(K \cap x U x^{-1})$. Hence the complex $\prod_{x \in R} (\mathcal{J}^\bullet)^{K \cap x U x^{-1}}$ computes $\prod_{x \in R} H^*(K \cap x U x^{-1}, V)$. On the other hand the complex $\text{Ind}_K^G(\mathcal{J}^\bullet)$ computes $R^* \text{Ind}_K^G(V)$

and is a complex of injective objects in $\text{Mod}_k(G)$. Therefore the composed functor spectral sequence for the functors $(-)^U$ and $\text{Ind}_K^G(-)$ exists and reads

$$E_2^{r,s} = H^r(U, R^s \text{Ind}_K^G(V)) \implies \prod_{x \in R} H^{r+s}(K \cap xUx^{-1}, V) .$$

By passing to the limit with respect to U it degenerates into isomorphisms

$$(7) \quad R^s \text{Ind}_K^G(V) \cong \varinjlim_U \prod_{x \in R} H^s(K \cap xUx^{-1}, V) .$$

For this note that profinite group cohomology commutes with inductive limits and that, for any M in $\text{Mod}_k(K)$, we have

$$\varinjlim_U H^r(U, M) = \begin{cases} M & \text{if } r = 0, \\ 0 & \text{otherwise.} \end{cases}$$

In the limit (7) we may take U to run over Poincaré subgroups of G . However, with U the open subgroup $K \cap xUx^{-1}$ is also a Poincaré group of dimension d . Therefore all the cohomology groups on the right-hand side of (7) vanish for $s > d$.

ii. Because of Lemma 6.1 this follows from i. \square

Remark 6.3. If G is compact then the functors Ind_K^G and $\underline{\text{Ind}}$ are exact.

Proof. For compact G we have $\text{Ind}_K^G = \text{ind}_K^G = \text{compact induction}$, which is exact. \square

As a consequence of Prop. 6.2.ii, $\underline{\text{Ind}}$ has finite cohomological dimension, and we therefore have (by [Har, Cor. I.5.3(γ)]) the total derived functor between the unbounded derived categories:

$$R\underline{\text{Ind}} : D(G) \longrightarrow D(G) .$$

This functor has more structure, as we now explain. In the following we use the convention that for a $G \times G$ -action we write G_ℓ , resp. G_r , if we refer to the action of the left, resp. right, factor in the product $G \times G$.

Lemma 6.4. *For any two representations V and V' in $\text{Mod}_k(G)$ we have*

i. *For $f \in \text{Ind}_K^G(V')$ and $(g_1, g_2) \in G \times G$ the function*

$$(g_1, g_2) f(g) := g_1 f(g_1^{-1} g g_2)$$

lies in $\text{Ind}_{g_1 K g_1^{-1}}^G(V')$;

ii. *this defines a smooth $G \times G$ -action on $\underline{\text{Ind}}(V')$; more precisely, $\text{Ind}_K^G(V') = \underline{\text{Ind}}(V')^{K_\ell}$;*

iii. *the earlier G -action on $\underline{\text{Ind}}(V')$ is the G_r -action;*

iv. *the adjunction isomorphism*

$$(8) \quad \underline{\text{Hom}}(V, V') \xrightarrow{\cong} \underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V, \underline{\text{Ind}}(V')) := \varinjlim_K \text{Hom}_{\text{Mod}_k(G_r)}(V, \underline{\text{Ind}}(V'))^{K_\ell}$$

obtained by passing to the inductive limit with respect to K in (6) is G -equivariant where G acts on the target through the G_ℓ -action on $\underline{\text{Ind}}(V')$;

v. if the representation V is finitely generated then

$$\underline{\mathrm{Hom}}_{\mathrm{Mod}_k(G_r)}(V, \underline{\mathrm{Ind}}(V')) = \mathrm{Hom}_{\mathrm{Mod}_k(G_r)}(V, \underline{\mathrm{Ind}}(V')) .$$

Proof. i. Suppose f is fixed by right translation by U . Then $^{(g_1, g_2)}f$ is fixed by right translation by $g_2 U g_2^{-1}$. Furthermore, for $\kappa \in K$, we compute

$$^{(g_1, g_2)}f((g_1 \kappa g_1^{-1})g) = g_1 f(g_1^{-1}(g_1 \kappa g_1^{-1})g g_2) = g_1 \kappa f(g_1^{-1}g g_2) = (g_1 \kappa g_1^{-1}) \cdot ^{(g_1, g_2)}f(g) .$$

Thus $^{(g_1, g_2)}f \in \underline{\mathrm{Ind}}(V')^{g_1 K g_1^{-1} \times g_2 U g_2^{-1}}$. Now ii and iii are obvious.

iv. For a fixed K we have by (6) and ii:

$$\begin{aligned} \mathrm{Hom}_{\mathrm{Mod}_k(K)}(V, V') &\xrightarrow{\cong} \mathrm{Hom}_{\mathrm{Mod}_k(G)}(V, \mathrm{Ind}_K^G(V')) \\ &= \mathrm{Hom}_{\mathrm{Mod}_k(G)}(V, \underline{\mathrm{Ind}}(V')^{K_\ell}) \\ &= \mathrm{Hom}_{\mathrm{Mod}_k(G_r)}(V, \underline{\mathrm{Ind}}(V'))^{K_\ell} \end{aligned}$$

which in the limit with respect to K gives rise to the isomorphism (8). A straightforward computation shows its equivariance. (We refer to [SS, p. 32] for the definition of the smooth linear maps $\underline{\mathrm{Hom}}(V, V')$.)

v. Under the finiteness assumption, the image of any G -homomorphism from V into $\underline{\mathrm{Ind}}(V')$ lies in $\mathrm{Ind}_K^G(V')$ for some open $K \leq G$ (which depends on the homomorphism). \square

Hence we actually have a left exact functor $\underline{\mathrm{Ind}} : \mathrm{Mod}_k(G) \rightarrow \mathrm{Mod}_k(G \times G)$ which derives to a functor $R\underline{\mathrm{Ind}} : D(G) \rightarrow D(G \times G)$ computable by homotopically injective resolutions. Our next goal is to lift the adjunction (8) to the derived level, using [KS, Thm. 14.4.8]. For this we first have to discuss its right-hand side in more detail.

We begin by recalling some general nonsense about the adjunction between tensor products and the Hom-functor which for three k -vector spaces V_1 , V_2 , and V_3 is given by the linear isomorphism

$$(9) \quad \begin{aligned} \mathrm{Hom}_k(V_1 \otimes_k V_2, V_3) &\xrightarrow{\cong} \mathrm{Hom}_k(V_1, \mathrm{Hom}_k(V_2, V_3)) \\ A &\mapsto \lambda_A(v_1)(v_2) := A(v_1 \otimes v_2) . \end{aligned}$$

Suppose that all three vector spaces carry a left G -action. Then $\mathrm{Hom}_k(V_1 \otimes_k V_2, V_3)$ and $\mathrm{Hom}_k(V_1, \mathrm{Hom}_k(V_2, V_3))$ are equipped with the $G \times G \times G$ -action defined by

$$^{(g_1, g_2, g_3)}A(v_1 \otimes v_2) := g_3 A(g_1^{-1}v_1 \otimes g_2^{-1}v_2) \quad \text{and} \quad ^{(g_1, g_2, g_3)}\lambda(v_1)(v_2) := g_3(\lambda(g_1^{-1}v_1)(g_2^{-1}v_2)),$$

respectively. The above adjunction is equivariant for these two actions. If we restrict to the diagonal G -action, then the above adjunction induces the adjunction isomorphism

$$\mathrm{Hom}_{k[G]}(V_1 \otimes_k V_2, V_3) \xrightarrow{\cong} \mathrm{Hom}_{k[G]}(V_1, \mathrm{Hom}_k(V_2, V_3)) .$$

If the G -action on the V_i is smooth then this can also be written as an isomorphism

$$(10) \quad \mathrm{Hom}_{\mathrm{Mod}_k(G)}(V_1 \otimes_k V_2, V_3) \cong \mathrm{Hom}_{\mathrm{Mod}_k(G)}(V_1, \underline{\mathrm{Hom}}(V_2, V_3)) .$$

In the next paragraph we discuss a variant of this.

Now suppose that in the adjunction (9) the vector spaces V_1 and V_2 carry a G -action whereas V_3 carries a $G \times G$ -action. Then $V_1 \otimes_k V_2$ carries a $G \times G$ -action as well. Moreover, $\text{Hom}_{k[G_r]}(V_2, V_3)$ still carries a G -action through the G_ℓ -action on V_3 . The above adjunction induces the adjunction isomorphism

$$\text{Hom}_{k[G \times G]}(V_1 \otimes_k V_2, V_3) \xrightarrow{\cong} \text{Hom}_{k[G_\ell]}(V_1, \text{Hom}_{k[G_r]}(V_2, V_3)) .$$

Suppose in addition that the actions on the V_i are smooth. Then the G_ℓ -action on

$$\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V_2, V_3) := \varinjlim_K \text{Hom}_{\text{Mod}_k(G_r)}(V_2, V_3)^{K_\ell} ,$$

where K runs over the compact open subgroups, is smooth. Hence we may rewrite the latter isomorphism as

$$\text{Hom}_{\text{Mod}_k(G \times G)}(V_1 \otimes_k V_2, V_3) \cong \text{Hom}_{\text{Mod}_k(G_\ell)}(V_1, \underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V_2, V_3)) .$$

This works as well with V_1 and V_2 interchanged:

Remark 6.5. We could have defined the initial adjunction for vector spaces analogously by the linear isomorphism

$$\begin{aligned} \text{Hom}_k(V_1 \otimes_k V_2, V_3) &\xrightarrow{\cong} \text{Hom}_k(V_2, \text{Hom}_k(V_1, V_3)) \\ A &\longmapsto \mu_A(v_2)(v_1) := A(v_1 \otimes v_2) . \end{aligned}$$

This leads to the isomorphism

$$\text{Hom}_{\text{Mod}_k(G \times G)}(V_1 \otimes_k V_2, V_3) \cong \text{Hom}_{\text{Mod}_k(G_r)}(V_2, \underline{\text{Hom}}_{\text{Mod}_k(G_\ell)}(V_1, V_3)) .$$

We are now in a position to apply [KS, Thm. 14.4.8]. For our present context we conclude that the functor

$$\begin{aligned} \text{Mod}_k(G)^{op} \times \text{Mod}_k(G \times G) &\longrightarrow \text{Mod}_k(G) \\ (V, V') &\longmapsto \underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V, V') \end{aligned}$$

has the left adjoint functor

$$\begin{aligned} \text{Mod}_k(G) \times \text{Mod}_k(G) &\longrightarrow \text{Mod}_k(G \times G) \\ (V, V') &\longmapsto V \otimes_k V' \end{aligned}$$

This shows that the adjointness requirements in [KS, Thm. 14.4.8] are satisfied, so that we have the total derived functor

$$R\underline{\text{Hom}}_{\text{Mod}_k(G_r)} : D(G)^{op} \times D(G \times G) \longrightarrow D(G)$$

satisfying (14.4.6) in [KS]. Namely, for V_1^\bullet, V_2^\bullet in $D(G)$ and V_3^\bullet in $D(G \times G)$ we have isomorphisms

$$\begin{aligned} R \text{Hom}_{\text{Mod}_k(G \times G)}(V_1^\bullet \otimes_k V_2^\bullet, V_3^\bullet) &\cong R \text{Hom}_{\text{Mod}_k(G_\ell)}(V_1^\bullet, R\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V_2^\bullet, V_3^\bullet)) \\ &\cong R \text{Hom}_{\text{Mod}_k(G_\ell)}(V_2^\bullet, R\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(V_1^\bullet, V_3^\bullet)) . \end{aligned}$$

The derived functor is computable via homotopically injective resolutions, see part (ii) of [KS, Thm. 14.4.8]:

$$(11) \quad R\text{Hom}_{\text{Mod}_k(G_r)}(V^\bullet, V'^\bullet) = \text{tot}_\Pi \text{Hom}_{\text{Mod}_k(G_r)}(V^\bullet, J'^\bullet)$$

where $V'^\bullet \xrightarrow{\sim} J'^\bullet$ is a homotopically injective resolution in $\text{Mod}_k(G \times G)$. Here tot_Π is the direct product totalization of the double complex. We emphasize that Π refers to the direct product in $\text{Mod}_k(G)$, in other words the smooth vectors of the direct product of abstract $k[G]$ -modules.

6.2 Duality on compact objects

Next we relate the duality functor $R\text{Hom}(-, k)$ from [SS] to the object $R\text{Ind}(k)$. In this section we are primarily interested in the restriction of the duality functor to $D(G)^c$ (the subcategory of compact objects). Our main result here is Corollary 6.9 below.

First, let V_1^\bullet and V_2^\bullet be any two objects of $D(G)$ and fix a homotopically injective resolution $V_2^\bullet \xrightarrow{\sim} J^\bullet$. Then, by [SS, Prop. 3.1]:

$$(12) \quad R\text{Hom}(V_1^\bullet, V_2^\bullet) = \text{Hom}^\bullet(V_1^\bullet, J^\bullet) \quad \text{and} \quad R\text{Ind}(V_2^\bullet) = \text{Ind}(J^\bullet) .$$

By Lemma 6.4.iv. we have the G -equivariant adjunction isomorphism of actual complexes

$$(13) \quad \text{Hom}^\bullet(V_1^\bullet, J^\bullet) \xrightarrow{\sim} \text{tot}_\Pi \text{Hom}_{\text{Mod}_k(G_r)}(V_1^\bullet, \text{Ind}(J^\bullet)) .$$

Using (12), (13), and (11) we now consider the G -equivariant map

$$\begin{aligned} \tau_{V_1^\bullet, V_2^\bullet} : R\text{Hom}(V_1^\bullet, V_2^\bullet) &= \text{Hom}^\bullet(V_1^\bullet, J^\bullet) \\ &\cong \text{tot}_\Pi \text{Hom}_{\text{Mod}_k(G_r)}(V_1^\bullet, \text{Ind}(J^\bullet)) \\ &\rightarrow \text{tot}_\Pi \text{Hom}_{\text{Mod}_k(G_r)}(V_1^\bullet, J_{\text{Ind}(J^\bullet)}^\bullet) \\ &= R\text{Hom}_{\text{Mod}_k(G_r)}(V_1^\bullet, \text{Ind}(J^\bullet)) \\ &= R\text{Hom}_{\text{Mod}_k(G_r)}(V_1^\bullet, R\text{Ind}(V_2^\bullet)) \end{aligned}$$

where $\text{Ind}(J^\bullet) \xrightarrow{\sim} J_{\text{Ind}(J^\bullet)}^\bullet$ is a homotopically injective resolution in $\text{Mod}_k(G \times G)$.

Proposition 6.6. *Let $U \subseteq G$ be a torsionfree pro- p open subgroup. Viewing $\mathbf{X}_U := \text{ind}_U^G(k)$ as a complex (concentrated in degree zero) the above map $\tau_{\mathbf{X}_U, V_2^\bullet}$ is a quasi-isomorphism for any V_2^\bullet .*

Proof. We have to show that the map

$$\text{Hom}_{\text{Mod}_k(G_r)}(\mathbf{X}_U, \text{Ind}(J^\bullet)) \longrightarrow \text{Hom}_{\text{Mod}_k(G_r)}(\mathbf{X}_U, J_{\text{Ind}(J^\bullet)}^\bullet)$$

is a quasi-isomorphism. Frobenius reciprocity implies that

$$(14) \quad \text{Hom}_{\text{Mod}_k(G_r)}(\mathbf{X}_U, -) = \varinjlim_K \text{Hom}_{\text{Mod}_k(G_r)}(\mathbf{X}_U, -)^{K \times \{1\}} = \varinjlim_K (-)^{K \times U} = (-)^{\{1\} \times U} .$$

Hence the map in question is the map $\text{Ind}(J^\bullet)^{\{1\} \times U} \longrightarrow (J_{\text{Ind}(J^\bullet)}^\bullet)^{\{1\} \times U}$. Of course we have the isomorphism $H^*(U, \text{Ind}(J^\bullet)) \xrightarrow{\sim} H^*(U, J_{\text{Ind}(J^\bullet)}^\bullet)$ where, for simplicity, we write simply U instead of $\{1\} \times U$. We adopt this convention for the rest of this proof. Hence it is enough to verify the following:

- (a) $H^*(U, \underline{\text{Ind}}(J^\bullet)) = h^*(\underline{\text{Ind}}(J^\bullet)^U);$
- (b) $H^*(U, J_{\underline{\text{Ind}}(J^\bullet)}^\bullet) = h^*((J_{\underline{\text{Ind}}(J^\bullet)}^\bullet)^U).$

We first note ([Lur, 1.3.5], [ScSc, 3.1]) that we may assume that all our homotopically injective resolutions are even semi-injective, i.e., in addition each of their terms is an injective object.

• *Part (a)*

Each J^m , for $m \in \mathbb{Z}$, is injective in $\text{Mod}_k(G)$ and hence in $\text{Mod}_k(K)$. Then any $\text{Ind}_K^G(J^m)$ is injective in $\text{Mod}_k(G)$ and hence in $\text{Mod}_k(U)$. Since the cohomology of U commutes with inductive limits it follows that the complex $\underline{\text{Ind}}(J^\bullet)$ consists of $H^0(U, -)$ -acyclic objects. The functor $H^0(U, -)$ has finite cohomological dimension. Therefore a) holds by [Har, Cor. I.5.3(γ)] (and its proof).

• *Part (b)*

This is a similar argument since in the subsequent Lemma 6.7 we will show that any term of the complex $J_{\underline{\text{Ind}}(J^\bullet)}^\bullet$ being injective in $\text{Mod}_k(G \times G)$ is $H^0(U, -)$ -acyclic. \square

At the end of the previous proof we alluded to:

Lemma 6.7. *Let $U \subseteq G$ be an open subgroup. Then any injective object V in $\text{Mod}_k(G \times G)$ is $H^0(\{1\} \times U, -)$ -acyclic.*

Proof. First we allow an arbitrary object V in $\text{Mod}_k(G \times G)$. For any compact open subgroup $K \subset G$ we have the Hochschild-Serre spectral sequence (where we write U instead of $\{1\} \times U$ on the E_2 -page):

$$E_2^{rs} = H^r(K, H^s(U, V)) \implies H^{r+s}(K \times U, V) .$$

It is functorial with respect to the restriction to a smaller compact open subgroup $K' \subseteq K$ (see [NSW, II.4 Ex. 3]). Hence we may pass to the limit with respect to smaller $K' \subseteq K$ in this spectral sequence. As in the proof of Prop. 6.2.i the limit spectral sequence degenerates into isomorphisms

$$H^s(U, V) \cong \varinjlim_{K'} H^s(K' \times U, V) .$$

Now, if V is injective in $\text{Mod}_k(G \times G)$ then it is injective in each $\text{Mod}_k(K' \times U)$ so that the above right-hand side vanishes for $s > 0$. Therefore so does $H^s(U, V)$. \square

We easily deduce the following result.

Corollary 6.8. *Let $U \subseteq G$ be a torsionfree pro- p open subgroup; we then have:*

- i. *The functors $R\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(\mathbf{X}_U, -)$ and $RH^0(\{1\} \times U, -)$ from $D(G \times G)$ to $D(G)$ are naturally isomorphic;*
- ii. *the functors $R\underline{\text{Hom}}(\mathbf{X}_U, -)$ and $RH^0(\{1\} \times U, R\underline{\text{Ind}}(-))$ from $D(G)$ to $D(G)$ are naturally isomorphic;*
- iii. *if G is compact, then the functors $R\underline{\text{Hom}}(\mathbf{X}_U, -)$ and $RH^0(\{1\} \times U, \underline{\text{Ind}}(-))$ from $D(G)$ to $D(G)$ are naturally isomorphic.*

Proof. i. This follows from (14) and the above Lemma 6.7. ii. Combine i. and Prop. 6.6. iii. If G is compact then, by Remark 6.3, the functor $\underline{\text{Ind}}$ is exact so that $R\underline{\text{Ind}} = \underline{\text{Ind}}$. Now apply part ii. \square

We now specialize the map $\tau_{V_1^\bullet, V_2^\bullet}$ to the case where V_2^\bullet is the complex with the trivial representation k in degree zero, and obtain a natural transformation

$$(15) \quad \tau_- := \tau_{-,k} : R\underline{\text{Hom}}(-, k) \longrightarrow R\underline{\text{Hom}}_{\text{Mod}_k(G_r)}(-, R\underline{\text{Ind}}(k))$$

between exact functors from $D(G)$ to $D(G)$.

Recall from [DGA, Rk. 10] that the full subcategory $D(G)^c$ of all compact objects in $D(G)$ is the smallest strictly full triangulated subcategory closed under direct summands which contains \mathbf{X}_U for some (or equivalently any) open torsionfree pro- p subgroup $U \subseteq G$.

Corollary 6.9. τ_- restricted to $D(G)^c$ is a natural isomorphism.

Proof. The full subcategory of all objects V^\bullet in $D(G)$ for which τ_{V^\bullet} is an isomorphism is a strictly full triangulated subcategory closed under direct summands which contains \mathbf{X}_U by Prop. 6.6. \square

6.3 The complex $R\underline{\text{Ind}}(k)$ for reductive groups

In this section we again focus on p -adic reductive groups, and we put ourselves in the setup from Section 5. Thus $G = \mathbf{G}(\mathfrak{F})$ is the group of \mathfrak{F} -rational points of a connected reductive group \mathbf{G} defined over some finite extension $\mathfrak{F}/\mathbb{Q}_p$. The goal in this section is to establish the following vanishing result.

Theorem 6.10. $R^i \underline{\text{Ind}}(k) = 0$ for all $i > 0$.

The proof requires some preparation. We start with the following observation.

Lemma 6.11. Let $U' \subset U$ be two uniform pro- p groups; if $U' \subset U^p$, then the restriction map

$$H^i(U, k) \xrightarrow{\text{res}} H^i(U', k)$$

is the zero map for all $i > 0$.

Proof. For $i = 1$ the map in question is the natural map $\text{Hom}_k(U/U^p, k) \rightarrow \text{Hom}_k(U'/U'^p, k)$ which is the zero map by assumption. For uniform pro- p groups, and a coefficient field k of characteristic p , the cohomology is generated under the cup product in degree 1 (one can reduce to the case $k = \mathbb{F}_p$ which is [Laz, Prop. 2.5.7.1, p. 567]). Since restriction maps commute with cup products the assertion follows. \square

To apply Lemma 6.11 the key input is the following.

Lemma 6.12. Let $n \in e\mathbb{N}$ (and assume $n > e$ if $p = 2$). Then, for all $g \in G$ we have

$$(K_n \cap gK_n g^{-1})^p = K_{n+e} \cap gK_{n+e} g^{-1}.$$

Proof. By the Cartan decomposition $G = K_0 Z^+ K_0$ we may assume that $g = z \in Z^+$, noting that K_n and K_{n+e} are both normal subgroups of K_0 . For any $z \in Z$ we have, by (5) in the proof of Proposition 5.7, the homeomorphism

$$\left(\prod_{\alpha \in \Phi_{\text{red}}^-} \tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1} \right) \times Z_n \times \left(\prod_{\alpha \in \Phi_{\text{red}}^+} \tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1} \right) \xrightarrow{\sim} K_n \cap z K_n z^{-1}.$$

In that proof we also argued that by dividing out by the subgroups of p -powers (even if $p = 2$) we get an isomorphism. Hence the above map restricted to p -powers must still be a homeomorphism, i.e.,

$$(K_n \cap z K_n z^{-1})^p = \prod_{\alpha \in \Phi_{\text{red}}^-} (\tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1})^p \times Z_n^p \times \prod_{\alpha \in \Phi_{\text{red}}^+} (\tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1})^p.$$

Using (1) we compute, now for $z \in Z^+$,

$$\begin{aligned} z \tilde{U}_{\alpha,n} z^{-1} &= z U_{\alpha,n} z^{-1} \cdot z U_{2\alpha,n} z^{-1} \\ &= U_{\alpha,n - \langle \nu(z), \alpha \rangle} \cdot U_{2\alpha,n - \langle \nu(z), 2\alpha \rangle} \\ &\begin{cases} \supseteq U_{\alpha,n} \cdot U_{2\alpha,n} = \tilde{U}_{\alpha,n} & \text{if } \alpha \in \Phi^-, \\ \subseteq U_{\alpha,n} \cdot U_{2\alpha,n} = \tilde{U}_{\alpha,n} & \text{if } \alpha \in \Phi^+ \end{cases} \end{aligned}$$

and therefore

$$\tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1} = \begin{cases} \tilde{U}_{\alpha,n} & \text{if } \alpha \in \Phi^-, \\ z \tilde{U}_{\alpha,n} z^{-1} & \text{if } \alpha \in \Phi^+. \end{cases}$$

Using (4) it follows that

$$(\tilde{U}_{\alpha,n} \cap z \tilde{U}_{\alpha,n} z^{-1})^p = \tilde{U}_{\alpha,n}^p \cap z \tilde{U}_{\alpha,n}^p z^{-1} = \tilde{U}_{\alpha,n+e} \cap z \tilde{U}_{\alpha,n+e} z^{-1}$$

as well as $Z_n^p = Z_{n+e}$. We conclude that

$$\begin{aligned} (K_n \cap z K_n z^{-1})^p &= \prod_{\alpha \in \Phi_{\text{red}}^-} (\tilde{U}_{\alpha,n+e} \cap z \tilde{U}_{\alpha,n+e} z^{-1}) \times Z_{n+e} \times \prod_{\alpha \in \Phi_{\text{red}}^+} (\tilde{U}_{\alpha,n+e} \cap z \tilde{U}_{\alpha,n+e} z^{-1}) \\ &= K_{n+e} \cap z K_{n+e} z^{-1} \end{aligned}$$

as desired. \square

We can now prove our vanishing result for $R^i \underline{\text{Ind}}(k)$.

Proof. (Theorem 6.10.) Upon passing to the diagonal colimit $K = U$ when combining Proposition 2.4 and Lemma 6.1 we see that

$$\begin{aligned} R^i \underline{\text{Ind}}(k) &\simeq \varinjlim_K \varinjlim_U \varprojlim_{x \in G/U} H^i(K \cap x U x^{-1}, k) \\ &\simeq \varinjlim_U \varprojlim_{x \in G/U} H^i(U \cap x U x^{-1}, k) \\ &\simeq \varinjlim_{n \in \mathbb{N}} \varprojlim_{x \in G/K_n} H^i(K_n \cap x K_n x^{-1}, k). \end{aligned}$$

Now it is obvious from Lemmas 6.11 and 6.12 that the transition maps

$$\begin{aligned} \varprojlim_{x \in G/K_n} H^i(K_n \cap x K_n x^{-1}, k) &\longrightarrow \varprojlim_{x' \in G/K_{n'}} H^i(K_{n'} \cap x' K_{n'} x'^{-1}, k) \\ c &\longmapsto \left(\text{res}_{K_{n'} \cap x' K_{n'} x'^{-1}}^{K_n \cap x' K_n x'^{-1}} c_{x' K_n} \right)_{x' \in G/K_{n'}} \end{aligned}$$

are all zero for $i > 0$ and $n' = n + e$. Therefore $R^i \underline{\text{Ind}}(k) = 0$ for $i > 0$. \square

Since the complex $R\text{Ind}(k)$ is concentrated in non-negative degrees, and has zero cohomology in positive degrees by 6.10, there is a quasi-isomorphism

$$(16) \quad \text{Ind}(k)[0] \xrightarrow{\text{qis}} R\text{Ind}(k)$$

for p -adic reductive groups G . Note that $\text{Ind}(k)$ is simply the space $\mathcal{C}^\infty(G, k)$ of smooth vectors in the $G \times G$ -representation on the space of all k -valued functions on G .

Corollary 6.13. *The functors $R\text{Hom}(-, k)$ and $R\text{Hom}_{\text{Mod}_k(G_r)}(-, \mathcal{C}^\infty(G, k))$ restricted to $D(G)^c$ are isomorphic.*

Proof. Combine Corollary 6.9 with (16). □

References

- [Bor] Borel A.: *Linear algebraic groups*. Second edition. Graduate Texts in Mathematics, 126. Springer-Verlag, New York, 1991.
- [CK23] Chirvasitu A. and Kanda R.: *Projective discrete modules over profinite groups*. Journal of Pure and Applied Algebra 227 (2023) 107260.
- [DdSMS] Dixon J. D., du Sautoy M. P. F., Mann A., Segal D.: *Analytic pro- p Groups*. 2nd Ed., Cambridge Univ. Press 1999.
- [DK23] Dupré N. and Kohlhaase J.: *Model categories and pro- p Iwahori-Hecke modules*. J. Inst. Math. Jussieu (2023), 1–48.
- [Har] Hartshorne R.: *Residues and Duality*. Springer Lect. Notes Math. vol. 20, 1966.
- [KP23] Kaletha T. and Prasad, G.: *Bruhat-Tits theory – a new approach*. New Mathematical Monographs, 44. Cambridge University Press, Cambridge, 2023.
- [Kal] Kaletha T.: *Corrections and additions in the book “Bruhat-Tits theory: a new approach”*. <https://websites.umich.edu/~kaletha/v1errata.pdf>
- [KS] Kashiwara M., Shapira P.: *Categories and sheaves*. Springer Grundlehren vol. 332, 2006.
- [Koh] Kohlhaase J.: *Smooth duality in natural characteristic*. Adv. Math. 317, 1-49 (2017).
- [Laz] Lazard M.: *Groupes analytiques p -adiques*. Publ. Math. IHES 26, 389-603 (1965).
- [Lur] Lurie J.: *Higher Algebra*. Preprint 2012.
- [NSW] Neukirch J., Schmidt A., Wingberg K.: *Cohomology of Number Fields*. 2nd Ed., Springer Grundlehren vol. 323, 2008.
- [OS19] Ollivier R., Schneider P.: *The modular pro- p Iwahori-Hecke Ext-algebra*. Proc. Symp. Pure Math. 101, 255–308 (2019).
- [Ron] Ronchetti N.: *A Satake homomorphism for the mod p derived Hecke algebra*. Preprint, 2019. arXiv:1808.06512.

- [ScSc] Scherotzke S., Schneider P.: *Derived parabolic induction*. Bull. Lond. Math. Soc. 54 (2022), no. 1, 264–274.
- [DGA] Schneider P.: *Smooth representations and Hecke modules in characteristic p* . Pacific J. Math. 279, 447–464 (2015).
- [Sor] Sorensen C.: *A vanishing result for higher smooth duals*. Algebra Number Theory 13 (7) 1735–1763 (2019).
- [SS] Schneider P., Sorensen C.: *Duals and admissibility in natural characteristic*. Represent. Theory 27 (2023), 30–50.
- [SSt] Schneider P., Stuhler U.: *Representation theory and sheaves on the Bruhat-Tits building*. Publ. Math. IHES 85, 97–191 (1997).
- [Vig96] Vignéras M.-F.: *Représentations ℓ -modulaires d’un groupe réductif p -adique avec $\ell \neq p$* . Progress Math. vol. 131, Birkhäuser 1996.

E-mail addresses: pschnei@uni-muenster.de, csorensen@ucsd.edu

PETER SCHNEIDER, MATH. INSTITUT, UNIVERSITÄT MÜNSTER, MÜNSTER, GERMANY.

CLAUS SORENSEN, DEPT. OF MATHEMATICS, UC SAN DIEGO, LA JOLLA, USA.