

# Koszul duality and manifold calculus

## Advances in Homotopy Theory IV

---

Connor Malin

June 22, 2023

University of Notre Dame

# The rational story

In the 1970's, Quillen and Sullivan pioneered the study of rational homotopy theory, the study of spaces up to equivalence of their rational homotopy groups.

# The rational story

In the 1970's, Quillen and Sullivan pioneered the study of rational homotopy theory, the study of spaces up to equivalence of their rational homotopy groups.

$$\begin{array}{ccc} \mathrm{DGCA}^{f.t., \geq 2}(\mathbb{Q}) & \xrightarrow{K} & \mathrm{DGLA}^{f.t., \geq 1}(\mathbb{Q}) \\ & \nwarrow \Omega_{PL}^* \quad \nearrow & \\ & \mathrm{Top}_{*, \mathbb{Q}}^{f.t., \geq 2} & \end{array}$$

# The rational story

In the 1970's, Quillen and Sullivan pioneered the study of rational homotopy theory, the study of spaces up to equivalence of their rational homotopy groups.

$$\begin{array}{ccc} \mathrm{DGCA}^{f.t., \geq 2}(\mathbb{Q}) & \xrightarrow{K} & \mathrm{DGLA}^{f.t., \geq 1}(\mathbb{Q}) \\ & \nwarrow \Omega_{PL}^* \quad \nearrow & \\ & \mathrm{Top}_{*, \mathbb{Q}}^{f.t., \geq 2} & \end{array}$$

These are equivalences of homotopy theories.

The functor  $\bar{\Omega}_{PL}^*$  is a strictly commutative model of reduced singular cochains,

The functor  $\bar{\Omega}_{PL}^*$  is a strictly commutative model of reduced singular cochains, and the functor  $K$  is Koszul duality between nonunital commutative algebras and Lie algebras.

The functor  $\bar{\Omega}_{PL}^*$  is a strictly commutative model of reduced singular cochains, and the functor  $K$  is Koszul duality between nonunital commutative algebras and Lie algebras.

### Example

It is straightforward to check that

$$\bar{\Omega}_{PL}^*(S^{2k}) \simeq \Lambda\langle x, y \rangle$$

where  $|x| = 2k$ ,  $|y| = 4k - 1$  with  $dy = x^2$ .

The functor  $\bar{\Omega}_{PL}^*$  is a strictly commutative model of reduced singular cochains, and the functor  $K$  is Koszul duality between nonunital commutative algebras and Lie algebras.

### Example

It is straightforward to check that

$$\bar{\Omega}_{PL}^*(S^{2k}) \simeq \Lambda\langle x, y \rangle$$

where  $|x| = 2k$ ,  $|y| = 4k - 1$  with  $dy = x^2$ .

$$\bar{\Omega}_{PL}^*(S^{2k-1}) \simeq \Lambda\langle x \rangle$$

where  $|x| = 2k - 1$ .



The functor  $\bar{\Omega}_{PL}^*$  is a strictly commutative model of reduced singular cochains, and the functor  $K$  is Koszul duality between nonunital commutative algebras and Lie algebras.

### Example

It is straightforward to check that

$$\bar{\Omega}_{PL}^*(S^{2k}) \simeq \Lambda\langle x, y \rangle$$

where  $|x| = 2k$ ,  $|y| = 4k - 1$  with  $dy = x^2$ .

$$\bar{\Omega}_{PL}^*(S^{2k-1}) \simeq \Lambda\langle x \rangle$$

where  $|x| = 2k - 1$ .

The generators and differentials conspire to allow for a map  $S_{\mathbb{Q}}^{4k-1} \rightarrow S_{\mathbb{Q}}^{2k}$  which can be seen to generate  $\pi_{4k-1}(S^{2k}) \otimes \mathbb{Q} \cong \mathbb{Q}$ .

To what extent does this story extend to integral or  $\mathbb{F}_p$  valued cochains?

To what extent does this story extend to integral or  $\mathbb{F}_p$  valued cochains?

For finite sets  $I, J$  and  $a \in I$  let  $I \cup_a J = I - \{a\} \sqcup J$ .

To what extent does this story extend to integral or  $\mathbb{F}_p$  valued cochains?

For finite sets  $I, J$  and  $a \in I$  let  $I \cup_a J = I - \{a\} \sqcup J$ .

## Definition

An operad  $O$  in a symmetric monoidal category  $(C, \otimes)$  is a collection of objects  $O(I)$  with actions of  $\Sigma_I$ , for all nonempty finite sets  $I$ , together with maps for all  $I, J$  and  $a \in I$

$$O(I) \otimes O(J) \rightarrow O(I \cup_a J)$$

which satisfy associativity and equivariance conditions.

### Example

The commutative operad  $\text{com}$  in  $(\text{dgVect}_k, \otimes)$  is given by

$$\text{com}(I) = k$$

with all partial composites induced by  $k \otimes k = k$ .

### Example

The commutative operad  $\text{com}$  in  $(\text{dgVect}_k, \otimes)$  is given by

$$\text{com}(I) = k$$

with all partial composites induced by  $k \otimes k = k$ .

A cofibrant replacement of  $\text{com}$  is called an  $E_\infty$ -operad.

### Example

The commutative operad  $\text{com}$  in  $(\text{dgVect}_k, \otimes)$  is given by

$$\text{com}(I) = k$$

with all partial composites induced by  $k \otimes k = k$ .

A cofibrant replacement of  $\text{com}$  is called an  $E_\infty$ -operad.

### Definition

An algebra  $A$  over an operad  $O$  in  $(C, \otimes)$  is an object  $A \in C$  with maps

$$O(I) \otimes A^{\otimes I} \rightarrow A$$

satisfying equivariance and associativity conditions.

### Example

The commutative operad  $\text{com}$  in  $(\text{dgVect}_k, \otimes)$  is given by

$$\text{com}(I) = k$$

with all partial composites induced by  $k \otimes k = k$ .

A cofibrant replacement of  $\text{com}$  is called an  $E_\infty$ -operad.

### Definition

An algebra  $A$  over an operad  $O$  in  $(C, \otimes)$  is an object  $A \in C$  with maps

$$O(I) \otimes A^{\otimes I} \rightarrow A$$

satisfying equivariance and associativity conditions.

### Proposition

*The cup product on  $\bar{C}^*(X; k)$  naturally extends to an  $E_\infty$ -algebra structure.*



**Theorem (Mandell)**

*For simply connected spaces  $X, Y$  there is a homotopy equivalence  $X \simeq Y$ , if and only if, there is an equivalence of  $E_\infty$ -algebras  $C^*(X; \mathbb{Z}) \simeq C^*(Y; \mathbb{Z})$ .*

**Theorem (Mandell)**

*For simply connected spaces  $X, Y$  there is a homotopy equivalence  $X \simeq Y$ , if and only if, there is an equivalence of  $E_\infty$ -algebras  $C^*(X; \mathbb{Z}) \simeq C^*(Y; \mathbb{Z})$ .*

**Theorem (Mandell)**

*The composite*

$$\mathrm{Top}_*^{\geq 2} \xrightarrow{\bar{c}^*} \mathrm{Alg}_{E_\infty}(\mathrm{DGVect}_{\mathbb{F}_p}, \otimes) \xrightarrow{K} \mathrm{Alg}_{L_\infty}(\mathrm{DGVect}_{\mathbb{F}_p}, \otimes)$$

*takes values in contractible  $L_\infty$ -algebras.*

Consider the category  $(\mathrm{Spec}, \wedge)$ .

Consider the category  $(\mathrm{Spec}, \wedge)$ .

By setting  $\mathrm{com}(I) = S^0$ , we can define  $E_\infty$ -algebras as algebras over a cofibrant replacement of  $\mathrm{com}$ .

Consider the category  $(\mathrm{Spec}, \wedge)$ .

By setting  $\mathrm{com}(I) = S^0$ , we can define  $E_\infty$ -algebras as algebras over a cofibrant replacement of  $\mathrm{com}$ .

## Example

Given a pointed space  $X$ , the function spectrum  $F(\Sigma^\infty X, S^0)$  obtains an  $E_\infty$ -algebra structure from the diagonal of  $X$ .

Consider the category  $(\mathrm{Spec}, \wedge)$ .

By setting  $\mathrm{com}(I) = S^0$ , we can define  $E_\infty$ -algebras as algebras over a cofibrant replacement of  $\mathrm{com}$ .

## Example

Given a pointed space  $X$ , the function spectrum  $F(\Sigma^\infty X, S^0)$  obtains an  $E_\infty$ -algebra structure from the diagonal of  $X$ .

This  $E_\infty$ -structure is a lift of the  $E_\infty$ -structure on  $\bar{C}^*(X)$  to spectra.

A combination of results of Ching, Francis, Gaiitsgory, Harper, and Lurie establish that theories of Koszul duality for algebras and operads in spectra exist, and these lift algebraic Koszul duality in some sense.

A combination of results of Ching, Francis, Gaitsgory, Harper, and Lurie establish that theories of Koszul duality for algebras and operads in spectra exist, and these lift algebraic Koszul duality in some sense.

With this in mind, one defines the spectral lie operad  $\mathrm{lie} := K(\mathrm{com})$ , and one obtains a functor

$$\mathrm{Alg}_{E_\infty}(\mathrm{Spec}, \otimes) \xrightarrow{K} \mathrm{Alg}_{L_\infty}(\mathrm{Spec}, \otimes).$$



A combination of results of Ching, Francis, Gaitsgory, Harper, and Lurie establish that theories of Koszul duality for algebras and operads in spectra exist, and these lift algebraic Koszul duality in some sense.

With this in mind, one defines the spectral lie operad  $\mathrm{lie} := K(\mathrm{com})$ , and one obtains a functor

$$\mathrm{Alg}_{E_\infty}(\mathrm{Spec}, \otimes) \xrightarrow{K} \mathrm{Alg}_{L_\infty}(\mathrm{Spec}, \otimes).$$

Unfortunately, the negative results of Mandell still imply that this functor destroys all  $p$ -torsion information when applied to  $F(\Sigma^\infty X, S^0)$ .

## Finding the appropriate setting

We implicitly localize at  $p$  until stated otherwise.

## Finding the appropriate setting

We implicitly localize at  $p$  until stated otherwise. Chromatic homotopy theory studies a certain tower of localization of  $\mathrm{spec}$ :

# Finding the appropriate setting

We implicitly localize at  $p$  until stated otherwise. Chromatic homotopy theory studies a certain tower of localization of  $\text{spec}$ :

$$\begin{array}{ccccccc} L_0^f \text{Spec} = \text{Spec}_{\mathbb{Q}} & \longleftarrow & L_1^f \text{Spec} & \longleftarrow & \dots & \longleftarrow & L_n^f \text{Spec} \longleftarrow \dots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \text{Spec}_{v_0} = \text{Spec}_{\mathbb{Q}} & & \text{Spec}_{v_1} & & \dots & & \text{Spec}_{v_n} \end{array}$$

# Finding the appropriate setting

We implicitly localize at  $p$  until stated otherwise. Chromatic homotopy theory studies a certain tower of localization of  $\text{spec}$ :

$$\begin{array}{ccccccc} L_0^f \text{Spec} = \text{Spec}_{\mathbb{Q}} & \longleftarrow & L_1^f \text{Spec} & \longleftarrow & \dots & \longleftarrow & L_n^f \text{Spec} \longleftarrow \dots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \text{Spec}_{v_0} = \text{Spec}_{\mathbb{Q}} & & \text{Spec}_{v_1} & & \dots & & \text{Spec}_{v_n} \end{array}$$

The layers of this tower can be “destabilized” by work of Bousfield and Heuts to produce analogs of rational homotopy theory that detect  $p$ -torsion.

$$\text{Top}_{v_0} = \text{Top}_{\mathbb{Q}} \qquad \text{Top}_{v_1} \qquad \dots \qquad \text{Top}_{v_n} \qquad \dots$$

**Theorem (Heuts)**

*If  $n > 0$ , there is an equivalence  $\mathrm{Alg}_{\mathrm{lie}}(\mathrm{Spec}_{v_n}) \simeq \mathrm{Top}_{v_n}$ .*

**Theorem (Heuts)**

*If  $n > 0$ , there is an equivalence  $\mathrm{Alg}_{\mathrm{lie}}(\mathrm{Spec}_{v_n}) \simeq \mathrm{Top}_{v_n}$ .*

**Theorem (Behrens-Rezk, Heuts)**

*If the Goodwillie tower for the identity of  $X \in \mathrm{Top}_{v_n}$  converges, the Koszul dual of  $F(X, S_{v_n}^0)$  is the  $v_n$  lie algebra model of  $X$ . In particular, the Goodwillie tower converges for the spheres  $S_{v_n}^d$ .*

# Goodwillie calculus

We now drop all localizations of spaces and spectra. Consider a functor  $F : \mathbf{Top}_* \rightarrow \mathbf{Spec}$ . We have Goodwillie approximations



# Goodwillie calculus

We now drop all localizations of spaces and spectra. Consider a functor  $F : \mathbf{Top}_* \rightarrow \mathbf{Spec}$ . We have Goodwillie approximations

$$\begin{array}{c}
 P_\infty(F)(X) \\
 \downarrow \\
 \dots \\
 \downarrow \\
 P_n(F)(X) \longleftarrow \partial_n(F) \wedge_{h\Sigma_n} \Sigma^\infty X^{\wedge n} \\
 \downarrow \\
 \dots \\
 \downarrow \\
 P_1(F)(X) \longleftarrow \partial_1(F) \wedge \Sigma^\infty X \\
 \downarrow \\
 P_0(F)(X) \\
 \uparrow \swarrow \nearrow \searrow \\
 F(X) \longrightarrow P_0(F)(X)
 \end{array}$$

**Definition**

A right module  $R$  over an operad  $O$  is a collection of objects  $R(I)$  with actions of  $\Sigma_I$ , for all nonempty finite sets  $I$ , together with maps for all  $I, J$  and  $a \in I$

$$R(I) \otimes O(J) \rightarrow R(I \cup_a J)$$

which satisfy associativity and equivariance conditions.

**Definition**

A right module  $R$  over an operad  $O$  is a collection of objects  $R(I)$  with actions of  $\Sigma_I$ , for all nonempty finite sets  $I$ , together with maps for all  $I, J$  and  $a \in I$

$$R(I) \otimes O(J) \rightarrow R(I \cup_a J)$$

which satisfy associativity and equivariance conditions.

**Proposition (Arone-Ching)**

*There is a canonical right lie module structure on  $\partial_* F$ .*

**Definition**

A right module  $R$  over an operad  $O$  is a collection of objects  $R(I)$  with actions of  $\Sigma_I$ , for all nonempty finite sets  $I$ , together with maps for all  $I, J$  and  $a \in I$

$$R(I) \otimes O(J) \rightarrow R(I \cup_a J)$$

which satisfy associativity and equivariance conditions.

**Proposition (Arone-Ching)**

*There is a canonical right lie module structure on  $\partial_* F$ .*

**Theorem (Arone-Ching)**

*Let  $F : \text{Top}_* \rightarrow \text{Spec}$ . If  $\partial_* F$  is levelwise a finite, free  $\Sigma_n$ -spectrum, then*

$$P_i(F)(X) \simeq \text{Map}_{\text{lie}}^h((\partial_* \Sigma^\infty \text{Map}(X, -))^{\leq i}, (\partial_* F)^{\leq i}).$$

In fact, we can compute the derivatives of  $\Sigma^\infty \mathrm{Map}_*(X, -)$  explicitly.  
Define the comodule

$$X^\wedge(I) = X^{\wedge I}$$

In fact, we can compute the derivatives of  $\Sigma^\infty \mathrm{Map}_*(X, -)$  explicitly.

Define the com module

$$X^\wedge(I) = X^{\wedge I}$$

There is an equivalence of lie modules,  $K(\Sigma^\infty X^\wedge) \simeq \partial_*(\Sigma^\infty \mathrm{Map}(X, -))$ .

In fact, we can compute the derivatives of  $\Sigma^\infty \mathrm{Map}_*(X, -)$  explicitly.  
 Define the com module

$$X^\wedge(I) = X^{\wedge I}$$

There is an equivalence of lie modules,  $K(\Sigma^\infty X^\wedge) \simeq \partial_*(\Sigma^\infty \mathrm{Map}(X, -))$ .

So the Koszul duality of com and lie appears when we differentiate the Yoneda embedding

$$X \rightarrow \Sigma^\infty \mathrm{Map}_*(X, -).$$

In fact, we can compute the derivatives of  $\Sigma^\infty \mathrm{Map}_*(X, -)$  explicitly.  
 Define the com module

$$X^\wedge(I) = X^{\wedge I}$$

There is an equivalence of lie modules,  $K(\Sigma^\infty X^\wedge) \simeq \partial_*(\Sigma^\infty \mathrm{Map}(X, -))$ .

So the Koszul duality of com and lie appears when we differentiate the Yoneda embedding

$$X \rightarrow \Sigma^\infty \mathrm{Map}_*(X, -).$$

in order to form the “fake Goodwillie tower” of Arone-Ching.



# The little disks operad

Another classic example of an operad is the little  $n$ -disks operad  $E_n$ .

Another classic example of an operad is the little  $n$ -disks operad  $E_n$ .

**Definition**

The operad  $E_n$  has  $E_n(I)$  equal to the configuration space of  $n$ -disks, labeled by  $I$ , in an  $n$ -disk. Partial composites are determined by inserting configurations of disks into one another.

Another classic example of an operad is the little  $n$ -disks operad  $E_n$ .

**Definition**

The operad  $E_n$  has  $E_n(I)$  equal to the configuration space of  $n$ -disks, labeled by  $I$ , in an  $n$ -disk. Partial composites are determined by inserting configurations of disks into one another.

Examples of algebras over the  $E_n$  operad are  $n$ -fold loop spaces.

Another classic example of an operad is the little  $n$ -disks operad  $E_n$ .

**Definition**

The operad  $E_n$  has  $E_n(I)$  equal to the configuration space of  $n$ -disks, labeled by  $I$ , in an  $n$ -disk. Partial composites are determined by inserting configurations of disks into one another.

Examples of algebras over the  $E_n$  operad are  $n$ -fold loop spaces. If we apply  $\Sigma_+^\infty$  we obtain a family of operads of spectra whose algebras define increasing levels of commutativity for ring spectra.

## (framed) Manifold calculus

In the setting of (framed) manifolds, Weiss developed a calculus for presheaves  $F : (\mathcal{Mfld}_n^{\text{fr}})^{\text{op}} \rightarrow \mathcal{C}$ .

## (framed) Manifold calculus

In the setting of (framed) manifolds, Weiss developed a calculus for presheaves  $F : (\mathcal{Mfld}_n^{\text{fr}})^{\text{op}} \rightarrow \mathcal{C}$ .

$$\begin{array}{ccc} & T_{\infty}(F)(M) & \\ & \downarrow & \\ & \cdots & \\ & T_1(F)(M) & \\ & \downarrow & \\ F(M) & \longrightarrow & T_0(F)(M) \end{array}$$

The diagram illustrates the Taylor tower of a presheaf  $F$ . It shows a sequence of approximations  $T_0(F)(M), T_1(F)(M), \dots, T_{\infty}(F)(M)$  connected by vertical downward arrows. A horizontal arrow points from  $F(M)$  to  $T_0(F)(M)$ . Diagonal arrows point from  $F(M)$  to each  $T_i(F)(M)$  for  $i \geq 0$ .

## (framed) Manifold calculus

In the setting of (framed) manifolds, Weiss developed a calculus for presheaves  $F : (\mathcal{Mfld}_n^{\text{fr}})^{\text{op}} \rightarrow \mathcal{C}$ .

$$\begin{array}{ccc} & T_{\infty}(F)(M) & \\ & \downarrow & \\ & \dots & \\ & T_1(F)(M) & \\ & \downarrow & \\ F(M) & \longrightarrow & T_0(F)(M) \end{array}$$

By restricting  $F$  to the subcategory of disjoint unions of disks,  $F$  determines a right  $E_n$  module.

## (framed) Manifold calculus

In the setting of (framed) manifolds, Weiss developed a calculus for presheaves  $F : (\mathcal{Mfld}_n^{\text{fr}})^{\text{op}} \rightarrow \mathcal{C}$ .

$$\begin{array}{ccc} & T_{\infty}(F)(M) & \\ & \downarrow & \\ & \cdots & \\ & T_1(F)(M) & \\ & \downarrow & \\ F(M) & \longrightarrow & T_0(F)(M) \end{array}$$

By restricting  $F$  to the subcategory of disjoint unions of disks,  $F$  determines a right  $E_n$  module. Using some nonstandard notation, we call this right  $E_n$  module  $\partial_*(F)$ .



**Proposition (Boavida de Brito-Weiss)**

*At a framed  $n$ -manifold  $M$*

$$T_i(F)(M) \simeq \mathrm{Map}_{E_n}((\partial_* \mathrm{Emb}^{\mathrm{fr}}(-, M))^{\leq i}, (\partial_* F)^{\leq i}).$$

**Proposition (Boavida de Brito-Weiss)**

*At a framed  $n$ -manifold  $M$*

$$T_i(F)(M) \simeq \mathrm{Map}_{E_n}((\partial_* \mathrm{Emb}^{\mathrm{fr}}(-, M))^{\leq i}, (\partial_* F)^{\leq i}).$$

Let  $E_M$  denote  $\partial_* \mathrm{Emb}^{\mathrm{fr}}(-, M)$ . This can be identified with the collection of configuration spaces of disks in  $M$ .

**Proposition (Boavida de Brito-Weiss)**

*At a framed  $n$ -manifold  $M$*

$$T_i(F)(M) \simeq \mathrm{Map}_{E_n}((\partial_* \mathrm{Emb}^{\mathrm{fr}}(-, M))^{\leq i}, (\partial_* F)^{\leq i}).$$

Let  $E_M$  denote  $\partial_* \mathrm{Emb}^{\mathrm{fr}}(-, M)$ . This can be identified with the collection of configuration spaces of disks in  $M$ . By “collapsing disks”, one can show that the  $\mathrm{com}$  module  $(M_+)^{\wedge}$  that we saw earlier is the induction of  $(E_M)_+$  along  $(E_n)_+ \rightarrow \mathrm{com}$ .

# Comparison of manifold and Goodwillie calculus

By taking inductions and applying Koszul duality, one obtains a map of towers

$$\begin{array}{ccc}
 T_{\infty}(\Sigma_+^{\infty} \text{Emb}(-, M))(N) & \longrightarrow & P_{\infty}(\Sigma^{\infty} \text{Map}(N_+, -))(M_+) \\
 \downarrow & & \downarrow \\
 \dots & & \dots \\
 \downarrow & & \downarrow \\
 T_n(\Sigma_+^{\infty} \text{Emb}(-, M))(N) & \longrightarrow & P_n(\Sigma^{\infty} \text{Map}(N_+, -))(M_+) \\
 \downarrow & & \downarrow \\
 \dots & & \dots \\
 \downarrow & & \downarrow \\
 T_1(\Sigma_+^{\infty} \text{Emb}(-, M))(N) & \longrightarrow & P_1(\Sigma^{\infty} \text{Map}(N_+, -))(M_+)
 \end{array}$$

## Theorem (Arone-Ching)

*There is a commutative diagram of operads*

$$\begin{array}{ccccc} s_n \text{lie} & \longrightarrow & \Sigma_+^\infty E_n & \longrightarrow & \text{com} \\ \left| \simeq \right. & & \left| \simeq \right. & & \left| \simeq \right. \\ s_n K(\text{com}) & \longrightarrow & s_n K(\Sigma_+^\infty E_n) & \longrightarrow & K(\text{lie}) \end{array}$$

## Theorem (Arone-Ching)

*There is a commutative diagram of operads*

$$\begin{array}{ccccc} s_n \text{lie} & \longrightarrow & \Sigma_+^\infty E_n & \longrightarrow & \text{com} \\ \left| \simeq \right. & & \left| \simeq \right. & & \left| \simeq \right. \\ s_n K(\text{com}) & \longrightarrow & s_n K(\Sigma_+^\infty E_n) & \longrightarrow & K(\text{lie}) \end{array}$$

## Conjecture (Ching)

*There is an equivalence*

$$\text{res}_{\text{lie}} s_{(-n,-n)} \Sigma_+^\infty E_M \simeq \partial_*(\Sigma^\infty \text{Map}_*(M^+, -)).$$

# Koszul dual comparison

By instead restricting from  $E_n$  to  $s_n\text{lie}$ , subject to the conjecture, we obtain a **contravariant** comparison

$$\begin{array}{ccc}
 T_\infty(\Sigma_+^\infty \text{Emb}(-, M))(N) & \longrightarrow & P_\infty(\Sigma^\infty \text{Map}(M^+, -))(N^+) \\
 \downarrow & & \downarrow \\
 \dots & & \dots \\
 \downarrow & & \downarrow \\
 T_n(\Sigma_+^\infty \text{Emb}(-, M))(N) & \longrightarrow & P_n(\Sigma^\infty \text{Map}(M^+, -))(N^+) \\
 \downarrow & & \downarrow \\
 \dots & & \dots \\
 \downarrow & & \downarrow \\
 T_1(\Sigma_+^\infty \text{Emb}(-, M))(N) & \longrightarrow & P_1(\Sigma^\infty \text{Map}(M^+, -))(N^+)
 \end{array}$$

# Main Results

## Theorem (M.)

*For a framed  $n$ -manifold  $M$ , there is an equivalence*

$$\mathrm{res}_{\mathrm{lie} \mathcal{S}_{(-n,-n)}} \Sigma_+^\infty E_M \simeq \partial_*(\Sigma^\infty \mathrm{Map}_*(M^+, -)).$$



# Main Results

## Theorem (M.)

*For a framed  $n$ -manifold  $M$ , there is an equivalence*

$$\mathrm{res}_{\mathrm{lie}S_{(-n,-n)}} \Sigma_+^\infty E_M \simeq \partial_*(\Sigma^\infty \mathrm{Map}_*(M^+, -)).$$

In the process, we prove a stronger result extending the Koszul self duality of  $E_n$ . Let  $E_{M^+}$  denote the module of configurations of disks in  $M^+$ .

# Main Results

## Theorem (M.)

*For a framed  $n$ -manifold  $M$ , there is an equivalence*

$$\mathrm{res}_{\mathrm{lie}} S_{(-n, -n)} \Sigma_+^\infty E_M \simeq \partial_*(\Sigma^\infty \mathrm{Map}_*(M^+, -)).$$

In the process, we prove a stronger result extending the Koszul self duality of  $E_n$ . Let  $E_{M^+}$  denote the module of configurations of disks in  $M^+$ .

## Theorem (M.)

*There is a zigzag of equivalences of operads*

$$\Sigma_+^\infty E_n \simeq \cdots \simeq s_n K(\Sigma_+^\infty E_n)$$

*and a compatible zigzag of equivalences of modules*

$$\Sigma_+^\infty E_M \simeq \cdots \simeq s_{(n, n)} K(\Sigma^\infty E_{M^+}).$$

*These equivalences are natural with respect to framed embeddings.*

## Normal bundles of operads

A parametrized spectrum is a space  $X$  and a family of spectra indexed by the points of  $X$ .

## Normal bundles of operads

A parametrized spectrum is a space  $X$  and a family of spectra indexed by the points of  $X$ . This category has a smash product  $\bar{\wedge}$  by taking products of the base spaces and smash products of the fibers.

## Normal bundles of operads

A parametrized spectrum is a space  $X$  and a family of spectra indexed by the points of  $X$ . This category has a smash product  $\bar{\wedge}$  by taking products of the base spaces and smash products of the fibers. Call this category  $(\text{ParSp}, \bar{\wedge})$ .

# Normal bundles of operads

A parametrized spectrum is a space  $X$  and a family of spectra indexed by the points of  $X$ . This category has a smash product  $\bar{\wedge}$  by taking products of the base spaces and smash products of the fibers. Call this category  $(\text{ParSp}, \bar{\wedge})$ .

## Proposition (M.)

*There is a factorization*

$$\begin{array}{ccc} & & \text{Operad}(\text{ParSp}, \bar{\wedge}) \\ & \nearrow \xi(-) & \downarrow \text{Th}(-) \\ \text{Operad}(\text{Top}, \times) & \xrightarrow{K(\Sigma_+^\infty -)} & \text{Operad}(\text{Sp}, \wedge) \end{array}$$

*Here,  $\text{Th}(-)$  denotes the Thom complex of a parametrized spectrum, i.e. the spectrum obtained by collapsing the zero section.*

There is a similar construction of normal bundles  $\xi_{E_M}^c$ . Taking Thom complexes, we have

$$\mathrm{Th}(\xi_{E_M}^c) \simeq K(\Sigma^\infty E_{M+}).$$

There is a similar construction of normal bundles  $\xi_{E_M}^c$ . Taking Thom complexes, we have

$$\mathrm{Th}(\xi_{E_M}^c) \simeq K(\Sigma^\infty E_{M+}).$$

This assignment of the normal bundle can be seen to be covariant with respect to maps  $E_N \rightarrow E_M$  induced by  $N \subset M$ .



There is a similar construction of normal bundles  $\xi_{E_M}^c$ . Taking Thom complexes, we have

$$\mathrm{Th}(\xi_{E_M}^c) \simeq K(\Sigma^\infty E_{M+}).$$

This assignment of the normal bundle can be seen to be covariant with respect to maps  $E_N \rightarrow E_M$  induced by  $N \subset M$ .

## Corollary

*If  $\xi_{\mathbb{R}^n}^c$  has a trivialization as a module over  $\xi_{E_n}$ , then for any  $U \subset \mathbb{R}^n$  we may choose a module trivialization of  $\xi_U^c$ , natural with respect to inclusion.*

Using the Koszul duality of the operad  $E_n$ , it is not difficult to deduce the local case

$$\Sigma_+^\infty E_{\mathbb{R}^n} \simeq \cdots \simeq s_{(n,n)} K(\Sigma^\infty E_{(\mathbb{R}^n)_+}).$$

Using the Koszul duality of the operad  $E_n$ , it is not difficult to deduce the local case

$$\Sigma_+^\infty E_{\mathbb{R}^n} \simeq \cdots \simeq s_{(n,n)} K(\Sigma^\infty E_{(\mathbb{R}^n)_+}).$$

Such an equivalence implies that  $\xi_{E_{\mathbb{R}^n}}^c$  is trivial.

Using the Koszul duality of the operad  $E_n$ , it is not difficult to deduce the local case

$$\Sigma_+^\infty E_{\mathbb{R}^n} \simeq \cdots \simeq s_{(n,n)} K(\Sigma^\infty E_{(\mathbb{R}^n)_+}).$$

Such an equivalence implies that  $\xi_{E_{\mathbb{R}^n}}^c$  is trivial.

## Corollary

*If  $M$  is a codimension 0 submanifold of  $\mathbb{R}^n$ , there is a zigzag of equivalences*

$$\Sigma_+^\infty E_M \simeq \cdots \simeq s_{(n,n)} K(\Sigma^\infty E_{M_+}).$$

*These equivalences are natural with respect to inclusion.*

# Globalizing the result

It turns out that the functors

$$M \rightarrow \Sigma_+^\infty E_M$$

$$M \rightarrow s_{(n,n)} K(\Sigma^\infty E_{M+})$$

form Weiss cosheaves, i.e. are sheaves with respect to multilocal covers.

# Globalizing the result

It turns out that the functors

$$M \rightarrow \Sigma_+^\infty E_M$$

$$M \rightarrow s_{(n,n)} K(\Sigma^\infty E_{M^+})$$

form Weiss cosheaves, i.e. are sheaves with respect to multilocal covers. Using  $\infty$ -categorical techniques due to Ayala-Francis and Lurie,

# Globalizing the result

It turns out that the functors

$$M \rightarrow \Sigma_+^\infty E_M$$

$$M \rightarrow s_{(n,n)} K(\Sigma^\infty E_{M+})$$

form Weiss cosheaves, i.e. are sheaves with respect to multilocal covers. Using  $\infty$ -categorical techniques due to Ayala-Francis and Lurie, to show these functors agree on  $\mathcal{M}\mathrm{fld}_n^{\mathrm{fr}}$ , it suffices to check on  $\mathrm{Open}(\mathbb{R}^n)$ .

# Globalizing the result

It turns out that the functors

$$M \rightarrow \Sigma_+^\infty E_M$$

$$M \rightarrow s_{(n,n)} K(\Sigma^\infty E_{M+})$$

form Weiss cosheaves, i.e. are sheaves with respect to multilocal covers. Using  $\infty$ -categorical techniques due to Ayala-Francis and Lurie, to show these functors agree on  $\mathcal{Mfd}_n^{\text{fr}}$ , it suffices to check on  $\text{Open}(\mathbb{R}^n)$ .

## Theorem

*For a framed  $n$ -manifold  $M$ , there is a natural zigzag of equivalence of modules*

$$\Sigma_+^\infty E_M \simeq \cdots \simeq s_{(n,n)} K(\Sigma^\infty E_{M+}),$$

*compatible with the Koszul self duality of  $E_n$ .*



## Resolving Ching's conjecture

Recall there is an induction map  $E_{M^+} \rightarrow (M^+)^{\wedge}$ .

# Resolving Ching's conjecture

Recall there is an induction map  $E_{M^+} \rightarrow (M^+)^{\wedge}$ . By taking Koszul duals, we achieve a map of lie modules

$$K(\Sigma^{\infty}(M^+)^{\wedge}) \rightarrow \mathrm{res}_{\mathrm{lie}} K(\Sigma^{\infty} E_{M^+}).$$

# Resolving Ching's conjecture

Recall there is an induction map  $E_{M^+} \rightarrow (M^+)^{\wedge}$ . By taking Koszul duals, we achieve a map of lie modules

$$K(\Sigma^{\infty}(M^+)^{\wedge}) \rightarrow \mathrm{res}_{\mathrm{lie}} K(\Sigma^{\infty} E_{M^+}).$$

Since restriction is Koszul dual to induction, this map is an equivalence.

# Resolving Ching's conjecture

Recall there is an induction map  $E_{M^+} \rightarrow (M^+)^{\wedge}$ . By taking Koszul duals, we achieve a map of lie modules

$$K(\Sigma^{\infty}(M^+)^{\wedge}) \rightarrow \mathrm{res}_{\mathrm{lie}} K(\Sigma^{\infty} E_{M^+}).$$

Since restriction is Koszul dual to induction, this map is an equivalence. Appealing to the compactly supported self duality of  $E_M$ , we obtain a resolution to Ching's conjecture.

# Resolving Ching's conjecture

Recall there is an induction map  $E_{M^+} \rightarrow (M^+)^{\wedge}$ . By taking Koszul duals, we achieve a map of lie modules

$$K(\Sigma^{\infty}(M^+)^{\wedge}) \rightarrow \mathrm{res}_{\mathrm{lie}} K(\Sigma^{\infty} E_{M^+}).$$

Since restriction is Koszul dual to induction, this map is an equivalence. Appealing to the compactly supported self duality of  $E_M$ , we obtain a resolution to Ching's conjecture.

## Corollary

*There is an equivalence*

$$\mathrm{res}_{\mathrm{lie}} \mathcal{S}_{(-n, -n)} \Sigma_+^{\infty} E_M \simeq \partial_*(\Sigma^{\infty} \mathrm{Map}_*(M^+, -)).$$

**Questions?**

**Definition**

The Verdier dualizing parametrized spectrum  $P(X, A)$  of a pair  $(X, A)$  is the parametrized spectrum over  $X$  such that

$$P(X, A)|_x := \Gamma^c(\Sigma_X^\infty \text{Path}(x, -))$$

# Verdier duality

## Definition

The Verdier dualizing parametrized spectrum  $P(X, A)$  of a pair  $(X, A)$  is the parametrized spectrum over  $X$  such that

$$P(X, A)|_x := \Gamma^c(\Sigma_X^\infty \text{Path}(x, -))$$

## Theorem (Klein)

*Suppose  $A \rightarrow X$  is a cofibration such that  $A$  and  $X$  each have the homotopy type of a compact CW complex and  $X$  is connected, then for any  $q \in \text{Sp}_X$  there is a zigzag,*

$$\text{Th}(P(X, A) \wedge q) \leftarrow \text{Th}(P(X, A) \wedge F_X(P, q)) \rightarrow \Gamma^A(q)$$

*where the rightmost map is given on the fiber over  $x \in X$  by the composition*

$$F_X^A(\Sigma_X^\infty X, \Sigma_X^\infty \text{Path}(x, -)) \wedge F_X(\Sigma_X^\infty \text{Path}(x, -), q) \rightarrow F_X^A(\Sigma_X^\infty X, q) = \Gamma^A(q)$$

*If  $q$  is level-fibrant, then after deriving  $\wedge, \text{Th}$  these are equivalences*