The Slowly Fading Light Echo Around Type Ia Supernova 2009ig

Charlotte Wood, Peter Garnavich, Peter Milne, Dina Drozdov

April 17, 2019

University of Notre Dame - GPS Annual Conference

What are Type Ia Supernovae?

Type Ia Supernovae in a Nutshell

- Thermonuclear explosion of a white dwarf



Image Credit - Wikipedia

Type Ia Supernovae in a Nutshell

- Thermonuclear explosion of a white dwarf
- Have a "standard" luminosity



Image Credit - Wikipedia

Type Ia Supernovae in a Nutshell

- Thermonuclear explosion of a white dwarf
- Have a "standard" luminosity
- Spectroscopically classified by no hydrogen, no helium, strong silicon



Image Credit - Wikipedia

Single vs. Double Degenerate Progenitors

- One white dwarf, one main sequence/giant star
- Material accretes from companion onto white dwarf

- Two white dwarfs; combined mass $> 1.4 M_{\odot}$
- Close binary that eventually merges





Image credit - *Left:* D. A. Hardy/AstroArt.org; *Right:* GSFC/D. Berry

What are Light Echoes?

What is a light echo?



- Light is scattered by dust into our line of sight

Image credit - ESA/Hubble

What is a light echo?



- Light is scattered by dust into our line of sight
- NOT caused by emission

Image credit - ESA/Hubble

Geometry of Light Echoes

- Ellipsoid with event at one focus and observer at the other



Figure 1. Geometry of the problem in the single scattering approximation.

Image credit - Patat 2005

Geometry of Light Echoes

- Ellipsoid with event at one focus and observer at the other
- Approximated as a paraboloid (distance very large)



Figure 1. Geometry of the problem in the single scattering approximation.

Image credit - Patat 2005

Geometry of Light Echoes

- Ellipsoid with event at one focus and observer at the other
- Approximated as a paraboloid (distance very large)
- Light from all points on paraboloid take equal time to arrive



Figure 1. Geometry of the problem in the single scattering approximation.

Image credit - Patat 2005

What can light echoes tell us about supernovae?

- 3-D picture of the local dust distribution
- Distance
- Symmetry of explosion
- Type of supernova



Which supernovae have light echoes?

- Generally found around core-collapse supernovae
- Light echoes do occur around SNe Ia, but are rare
- Know SNe Ia light echoes: 1991T, 1995E, 1998bu, 2006X, 2007af, 2009ig, 2014J, 2012cg?



Image credit - Filippenko et al. 1992

Why do Type Ia Supernova Light Echoes Matter?

Implications?

Type Ia supernovae are used as standard candles for cosmology

Implications?

Type Ia supernovae are used as standard candles for cosmology

- Properties of SNe Ia are related to their environment

Implications?

Type Ia supernovae are used as standard candles for cosmology

- Properties of SNe Ia are related to their environment
- Hints at different progenitors

Single vs. Double Degenerate Progenitors

- One white dwarf, one main sequence/giant star
- Material accretes from companion onto white dwarf



- Two white dwarfs; combined mass $> 1.4 M_{\odot}$
- Close binary that eventually merges



Image credit - *Left:* D. A. Hardy/AstroArt.org; *Right:* GSFC/D. Berry

SN 2009ig Observations & Results

Why is SN2009ig interesting?



- Normal type Ia supernova
- Peak magnitude V $= 13.5$
- Nearby in NGC 1015 ($z = 0.0088$)
- Practically no host galaxy extinction ($A_V = 0.01 \pm 0.01$)
- Used in H_0 measurements!

Late-time Photometry



Image credit - C. M. Wood

Future Work

Future Work



- Resolved in archival Hubble data?

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)
  - Proposal for more Hubble time - light echo should be larger now

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)
  - Proposal for more Hubble time - light echo should be larger now
  - Determine actual geometry, distance to supernova

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)
  - Proposal for more Hubble time - light echo should be larger now
  - Determine actual geometry, distance to supernova
- Obtain spectra of SN2009ig?

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)
  - Proposal for more Hubble time - light echo should be larger now
  - Determine actual geometry, distance to supernova
- Obtain spectra of SN2009ig?
  - May be too faint

Image credit - ESA/Hubble & NASA, A. Riess

Future Work



- Resolved in archival Hubble data?
  - Barely (preliminary)
  - Proposal for more Hubble time - light echo should be larger now
  - Determine actual geometry, distance to supernova
- Obtain spectra of SN2009ig?
  - May be too faint
  - Alternative: analyze R-band data (2010-2018), early V, I data (2010-2012)

Image credit - ESA/Hubble & NASA, A. Riess

Questions?


SN 2009ig
Foley et al. 2012
KPNO
LBTO
B Magnitude
MJD
14
16
18
20
22
24
26
55000
55500
56000
56500
57000
57500
58000
58500