

# Chapter 11: Linear models with a factor variable with three or more levels

STATS 201/8

University of Auckland

# Learning Outcomes

In this chapter you will learn about:

- Explanatory factor with multiple levels—One-way analysis of variance
- The multiple comparisons problem
- Relevant `R`-code.

# Explanatory factor with multiple levels (One-way analysis of variance)

## Example—Fruit fly

In this case study we look at how the male fruit-fly's longevity is related to his reproductive activity.



Data from <http://www.cvgs.k12.va.us:81/digstats/Imain.html>.

## Fruit fly

Studies have shown that the longevity (life span) of female fruit flies decreases with an increase in reproduction, and this leads to a similar question related to males.

The hypothesis was that the males living alone or with uninterested females would live longer than the males living with the interested females. Since there are more than two group means to compare<sup>1</sup> an adjustment to how we interpret our model is used to determine if there is a significant difference between these group means.

How does one define “interest” in fruit-flies?

Here is this study’s definition of the question above:

Newly inseminated females will not usually mate again for at least two days. So, the males in the uninterested females groups were always living with newly inseminated females.

---

<sup>1</sup>If there were only two groups we could use a two sample two-sample  $t$ -test discussed in chapter 5

# Fruit fly...

The variable measured was:

**days** the number of days the male fly lived

The variables that were controlled were:

**group** the group they were allocated to where:

**G1** are males living alone,

**G2** are males living with one interested female,

**G3** are males living with eight interested females,

**G4** are males living with one uninterested female, and


**G5** are males living with eight uninterested females.

There were 25 flies in each group, for a total sample size of 125.

# Fruit fly...

Let us take a look at the data:

```
> Fruitfly.df = read.csv("Data/Fruitfly.csv", header = T)
> boxplot(days ~ group, data = Fruitfly.df)
```



## Fruit fly...

It seems male fruit flies do not live as long when they are in the presence of 'uninterested' females (G5). A result we were not expecting.

For females reproduction came at a cost (shorter lifespan), whereas for males, a lack of reproduction seems to cost them. Let us see if this effect is 'real' (or not).



# Fruit fly...

Explanatory factor with many levels ( $> 2$ )

If we believe this to be true then our underlying model is:

$$\text{days} = \beta_0 + \beta_1 \times \text{D2} + \beta_2 \times \text{D3} + \beta_3 \times \text{D4} + \beta_4 \times \text{D5} + \epsilon$$

where, as usual  $\epsilon \stackrel{iid}{\sim} N(0, \sigma^2)$ .


- **D2** is a dummy variable whereby: **D2**=1 if the fruit fly is in group 2—otherwise it is 0.
- **D3** is a dummy variable whereby: **D3**=1 if the fruit fly is in group 3—otherwise it is 0.
- ... And so on.

For example,  $\beta_1$  and  $\beta_2$  represent the differences in average longevity (**days**) when we compare groups 2 and 3 to group 1 (the baseline).

# Fruit fly...

## Assumption checks

```
> Fruitfly.fit = lm(days ~ group, data = Fruitfly.df)
> plot(Fruitfly.fit, which = 1)
```





The **EOV** assumption seem to be okay.

# Fruit fly...

## Assumption checks...

```
> normcheck(Fruitfly.fit)
```




The normality assumption seem to be okay.

# Fruit fly...

## Assumption checks...

```
> cooks20x(Fruitfly.fit)
```



No unduly influential data points.

# Fruit fly...

## $R^2$ and ANOVA table

We can trust this output. What is it telling us?

```
> anova(Fruitfly.fit)
Analysis of Variance Table

Response: days
          Df Sum Sq Mean Sq F value    Pr(>F)
group       4  11939  2984.82   13.612 3.516e-09 ***
Residuals 120   26314   219.28
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

This simply allows us to say that there is a clear difference in expected longevity between the five groups, which was fairly obvious from the first plot. The next step is to see where the differences lie.

Note that a significant result means you should investigate which groups are different from one another; there is more work to be done.

Now we can investigate whether female 'lack of interest' is 'killing' these wretched male fruit flies.

## Fruit fly...

$R^2$  and ANOVA table...

In the above output we see that the variability we observe for all our data can be broken down into two components **group** and **residual**.

The amount of variability that the variable **group** (as shown in the **Sum Sq** column) explains is **11939**. The residual variability (left over) is **26314**. The total variability is  $11939 + 26314 = 38253$ . The % of variability explained by **group** is therefore

$$100 \times \left( \frac{11939}{11939 + 26314} \right) = 100 \times \left( 1 - \frac{26314}{11939 + 26314} \right) = 31\%.$$

This is another way we can think of the % of variation explained or  $R^2$ .

It is the % of the data that is not residual or 'noise'.

# Fruit fly...

## One-way analysis of variance

Now we know that the variable `group` helps explain longevity, what can we say about these groups? Let us investigate. The `s20x` function `summary1way` provides the group means and their deviations from the overall (or 'grand') mean.

# Fruit fly...

## One-way analysis of variance...

```
> summary1way(Fruitfly.fit, draw.plot = FALSE)
```

ANOVA Table:

	Df	Sum Squares	Mean Square	F-statistic	p-value
Between Groups	4	11939.28	2984.82	13.61195	0
Within Groups	120	26313.52	219.27933		
Total	124	38252.8			

Numeric Summary:

	Sample size	Mean	Median	Std Dev	Midspread
All Data	125	57.44	58	17.56389	24
G1	25	63.56	62	16.45215	28
G2	25	64.80	65	15.65248	22
G3	25	63.36	65	14.53983	21
G4	25	56.76	56	14.92838	20
G5	25	38.72	40	12.10207	15

Table of Effects: (GrandMean and deviations from GM)

typ.val	G1	G2	G3	G4	G5
57.44	6.12	7.36	5.92	-0.68	-18.72



# Fruit fly...

## Interpreting the output

We see from above that the overall average longevity of the 125 male flies in the study is about 57.4 days.

We also see that group G5 has markedly lower longevity (18.75 fewer days) compared to the overall mean.

Note that if group does not explain any true underlying variation in longevity, then we expect all these group means to differ at most only moderately from the overall mean. This can be hard to judge informally, since we have to take into account the standard error of each group mean and how many groups there are.

That is why we have to rely on the  $P$ -value from the anova table.

# Fruit fly...

## Interpreting the output...

It is natural to ask which of the groups are different.

Here is our familiar `summary` output:

```
> summary(Fruitfly.fit)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	63.560	2.962	21.461	< 2e-16 ***
groupG2	1.240	4.188	0.296	0.768
groupG3	-0.200	4.188	-0.048	0.962
groupG4	-6.800	4.188	-1.624	0.107
groupG5	-24.840	4.188	-5.931	2.98e-08 ***

---

Residual standard error: 14.81 on 120 degrees of freedom

Multiple R-squared: 0.3121, Adjusted R-squared: 0.2892

F-statistic: 13.61 on 4 and 120 DF, p-value: 3.516e-09

Here we see that we have evidence to believe that  $\beta_4$  the parameter for group 5 is different from from zero.

We estimate that males with 8 uninterested females die, on average, 25 days earlier than males who are by themselves (our baseline group is G1).

# Fruit fly...

## Interpreting the output...

In the output above we are restricted to seeing how each of the groups, G2–G5, differs from the baseline group G1.

If we wish to see how the other groups differed from group G4, for example, then we could achieve this by changing the baseline group to group G4 by reordering the levels of the group factor variable.

This is very tedious, but here is how we make G4 the baseline level in R:

```
> Fruitfly.df$newgroup = factor(Fruitfly.df$group,  
+                               levels = c("G4", "G1", "G2", "G3", "G5"))
```

However, what if we wish to look at all pair-wise comparisons (i.e., G1 vs G2, G2 vs G3, ...)? Do we really have to do this re-ordering a bunch of times in order to find these out?

The answer is no: We can get R to do this 'heavy lifting' for us.

# Fruit fly...

## Multiple comparisons

Note that when we are looking at all pair-wise comparisons of 5 groups, we have a total of 10 different possibilities:

G1 vs G2, G1 vs G3, G1 vs G4, G1 vs G5, (4 comparisons)

G2 vs G3, G2 vs G4, G2 vs G5, (3 comparisons)

G3 vs G4, G3 vs G5, (2 comparisons)

G4 vs G5, (1 comparisons).

The number 10 comes from  $4 + 3 + 2 + 1 = 10$  or in fact  ${}^5C_2 = 10$  ways of choosing 2 objects from 5 (in no particular order).

Here we are asking 10 questions (comparisons) about our data, as we are looking to test 10 null hypotheses. Of all null hypotheses that are true, 5% are falsely rejected. Of all 95% confidence intervals, 5% of them do not contain the true parameter value.

# The multiple comparisons problem

# Erroneous evidence of an effect from multiple testing

The following **R** code fits a simple linear regression model to iid (independent and identically distributed) normal data.

**NOTE:** The null hypothesis  $H_0 : \text{slope} = 0$  is **true**.

```
> x = 1:30 ## Our explanatory variable
> x
[1] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25
[26] 26 27 28 29 30
> y = rnorm(30) ## y has NO relationship with x
> summary(lm(y ~ x))$coef ## Print only the coefficient table
              Estimate Std. Error    t value Pr(>|t|)
(Intercept) -0.107068614 0.35454802  -0.3019862 0.7648943
x            -0.004679368 0.01997124  -0.2343054 0.8164528
```

If this code is run many times over, then approximately 5% of the time the slope will have  $P\text{-value} < 0.05$ .<sup>2</sup>

That is, there will be erroneous evidence of an effect of **x** (i.e., evidence for a non-zero slope) about 1 time in 20!

---

<sup>2</sup>In fact, it can be shown that the  $P$ -value is uniformly distributed between 0 and 1 when  $H_0$  is true.

## Erroneous evidence of an effect from multiple testing...

When we do multiple tests (i.e., the 10 paired comparisons in this example) then we greatly increase the probability of obtaining at least one erroneous conclusion<sup>3</sup>.

This is known as the multiple comparison problem. It essentially says that if you look at enough things you will find something 'happening', even when there's nothing going on.

Remember, data always have variability, and if we are not careful we can 'discover' false structure that is not really there.

So, when we look at these 10 comparisons we need to adjust so that the overall error rate (the probability of any spurious significance) over all 10 comparison is no more the 5%. This can be done using a Tukey adjustment.

---

<sup>3</sup>Assuming independent comparisons, if we do 10 95% CIs we have an overall error rate of  $1 - (1 - .05)^{10} = 40\%$ , which is much higher than our original 5% error rate per comparison.

## Example—Fruit fly

### Tukey simultaneous confidence intervals

Let's get *simultaneous* 95% confidence intervals for all 10 comparisons via the `s20x` function `multipleComp`.

These confidence intervals are called “simultaneous” since we can be 95% confident that **they all** contain the true group difference simultaneously.

```
> multipleComp(Fruitfly.fit)
```

			Estimate	Tukey.L	Tukey.U	Tukey.p
G1	-	G2	-1.24	-12.8405	10.3605	0.9983
G1	-	G3	0.20	-11.4005	11.8005	1.0000
G1	-	G4	6.80	-4.8005	18.4005	0.4855
G1	-	G5	24.84	13.2395	36.4405	0.0000
G2	-	G3	1.44	-10.1605	13.0405	0.9970
G2	-	G4	8.04	-3.5605	19.6405	0.3127
G2	-	G5	26.08	14.4795	37.6805	0.0000
G3	-	G4	6.60	-5.0005	18.2005	0.5158
G3	-	G5	24.64	13.0395	36.2405	0.0000
G4	-	G5	18.04	6.4395	29.6405	0.0003

Here we see that most of these comparisons are not significantly different.



# Fruit fly

## Tukey simultaneous confidence intervals...

Lets extract the CIs where the Tukey adjusted  $P$ -value are less than 0.05.

```
> mc.fruitfly = multipleComp(Fruitfly.fit)
> ## Which entries have a P-value less than 0.05?
> mc.fruitfly[, "Tukey.p"] < 0.05
```

G1	-	G2	G1	-	G3	G1	-	G4	G1	-	G5	G2	-	G3	G2	-	G4	G2	-	G5	G3	-	G4
		FALSE			FALSE			FALSE			TRUE			FALSE			FALSE			TRUE			FALSE
G3	-	G5	G4	-	G5																		
		TRUE			TRUE																		

```
> ## Print entries which have a P-value less than 0.05
> mc.fruitfly[mc.fruitfly[, "Tukey.p"] < 0.05, ]
```

		Estimate	Tukey.L	Tukey.U	Tukey.p	
G1	-	G5	24.84	13.2395	36.4405	0e+00
G2	-	G5	26.08	14.4795	37.6805	0e+00
G3	-	G5	24.64	13.0395	36.2405	0e+00
G4	-	G5	18.04	6.4395	29.6405	3e-04

# Fruit fly...

Some conclusions:

- Our model explains 31% of variability in fruit fly longevity.
- We see that the effect of group 5 (males with 8 uninterested females) seems different from all the others.

On average, group 5 males live fewer days than:

- Group 1 (males living alone) by 13 to 36 fewer days.
- Group 2 (males living with one interested female) by 14 to 38 fewer days.
- Group 3 (males living with eight interested females) by 13 to 36 fewer days.
- Group 4 (males living with one uninterested female) by 6 to 30 fewer days.

## Fruit fly...

On a lighter note there is little evidence of a difference in longevity if no females or no more than one uninterested female is about, or if females are there and 'interested' in them — but in the presence of multiple uninterested females they die earlier (they 'drop like flies').

Recall also that in the original studies it was seen that females did not live as long if they reproduced.

It is tempting to make similar inference about the human species but that may be going too far!

## Relevant R-code

# Most of the R-code you need for this chapter

Note that this code comes with the usual code/checks discussed in chapters 1 and 2.

You do not need to create dummy variables - R does that for you. The baseline can be changed if you wish rather than having R choose it for you — see relevant R-code from chapter 9.

Use box plots to inspect the data for each level of the factor.

```
> ## Create the pairs plot of the five numeric variables  
> boxplot(days ~ group, data = Fruitfly.df)
```

Fit the model and use the ANOVA table to see if any of the means differ from one another (regardless of the baseline chosen).

```
> anova(Fruitfly.fit)
```

In order to see measure pair-wise differences between mean levels to adjust for multiple comparisons:

```
> multipleComp(Fruitfly.fit)
```