

Example:

Find least squares solution for the following system of nonlinear equations

Functional model:

$$x + y - 2y^{2} = -4$$
$$x^{2} + y^{2} = 8$$
$$3x^{2} - y^{2} = 7.7$$

Stochastic model for the observation:

$$p_1 = 1$$
, $p_2 = 1$, $p_3 = 1$



Observation equations:

$$-4 + v_1 = \hat{x} + \hat{y} - 2\hat{y}^2$$

$$8 + v_2 = \hat{x}^2 + \hat{y}^2$$

$$7.7 + v_3 = 3\hat{x}^2 - \hat{y}^2$$

Rearranging:

$$v_1 = \hat{x} + \hat{y} - 2\hat{y}^2 + 4$$

$$v_2 = \hat{x}^2 + \hat{y}^2 - 8$$

$$v_3 = 3\hat{x}^2 - \hat{y}^2 - 7.7$$

$$\sum_{i=1}^{n} p_i v_i^2 \to \min$$



$$\frac{\partial \Sigma p_i v_i^2}{\partial x} = 2(\hat{x} + \hat{y} - 2\hat{y}^2 + 4) + 2(\hat{x}^2 + \hat{y}^2 - 8)(2\hat{x}) + 2(3\hat{x}^2 - \hat{y}^2 - 7.7)(6\hat{x}) = 0$$

$$\frac{\partial \Sigma p_i v_i^2}{\partial y} = 2(\hat{x} + \hat{y} - 2\hat{y}^2 + 4)(1 - 4\hat{y}) + 2(\hat{x}^2 + \hat{y}^2 - 8)(2\hat{y}) + 2(3\hat{x}^2 - \hat{y}^2 - 7.7)(-2\hat{y}) = 0$$

Nonlinear normal equations:

$$\hat{x} + \hat{y} - 2\hat{y}^2 + 4 + 2\hat{x}^3 + 2\hat{x}\hat{y}^2 - 16\hat{x} + 18\hat{x}^3 - 6\hat{x}\hat{y}^2 - 46.2\hat{x} = 0$$

$$\hat{x} - 6\hat{y}^2 + 4 - 4\hat{x}\hat{y} + 8\hat{y}^3 - 15\hat{y} + 2\hat{x}^2\hat{y} + 2\hat{y}^3 - 16\hat{y} - 6\hat{x}^2\hat{y} + 2\hat{y}^3 + 15.4\hat{y} = 0$$

nonlinear normal equations



- Nonlinear functional model → nonlinear normal equations
 - → Question: How to solve nonlinear normal equations?
 - a) Direct solution (if possible, e.g. for quadratic equations), or
 - b) Linearisation (Taylor-Approximation) of normal equations and iterative solution, or
 - c) Heuristic Approaches, or
 - d) Replace the original functional model by a linearised functional model and iterative solution



Solution from linearisation of the original functional model

→ We replace our nonlinear functional model by an approximation (linear functional model)

Attention: We have to compute an iterative solution!



Nonlinear functional model:

$$-4.0 = x + y - 2y^{2}$$

$$8.0 = x^{2} + y^{2}$$

$$7.7 = 3x^{2} - y^{2}$$

$$\mathbf{L}_{n\times 1} = \mathbf{\Phi}_{n\times 1}(\mathbf{X}_{u\times 1}) = \begin{bmatrix} \boldsymbol{\varphi}_{1}(\mathbf{X}_{u\times 1}) \\ \boldsymbol{\varphi}_{2}(\mathbf{X}_{u\times 1}) \\ \vdots \\ \boldsymbol{\varphi}_{n}(\mathbf{X}_{u\times 1}) \end{bmatrix}$$

Question: How can we find a linear approximation?

→ Taylor series approximation for each equation



Taylor series approximation:

$$\mathbf{L}_{n\times 1} = \mathbf{\Phi}_{n\times 1} (\mathbf{X}_{u\times 1}) = \mathbf{\Phi}_{n\times 1} (\mathbf{X}_{u\times 1}^{0}) + \left(\frac{\partial \mathbf{\Phi}(\mathbf{X})}{\partial \mathbf{X}}\right)_{0 \le u} (\mathbf{X} - \mathbf{X}_{u\times 1}^{0})_{u\times 1} + \cdots$$

First-order Taylor series approximation at the place of the initial values \mathbf{X}^0



Start of iterative computing

 \mathbf{X}^0 : Initial values (starting vales, approximations) for the unknowns Initial values have to be close to the final solution, otherwise

- computation will not converge, or
- computation can converge to a local min for $\mathbf{v}^{\mathrm{T}}\mathbf{P}\mathbf{v} \to \min \to \text{worst case}$!



We introduce:

$$\mathbf{L}_{n\times 1}^0 = \mathbf{\Phi}_{n\times 1}(\mathbf{X}_{u\times 1}^0)$$

observations as functions of the unknowns

$$\mathbf{l}_{n\times 1} = \mathbf{L}_{n\times 1} - \mathbf{L}_{n\times 1}^0$$

vector of reduced observations

$$\Sigma_{ll} = \Sigma_{LL}$$

from variance covariance propagation with functional model $\mathbf{l} = \mathbf{L} - \mathbf{L}^0$

$$\mathbf{x}_{u \times 1} = \mathbf{X} - \mathbf{X}^0$$

vector of reduced unknowns vector of corrections for the unknowns

$$\mathbf{J}_{n \times u} = \left(\frac{\partial \Phi(\mathbf{X})}{\partial \mathbf{X}}\right)_{0 \ n \times u}$$

Jacobian matrix, contains the partial derivatives of each equation of the functional model



Linearised functional model:

$$\mathbf{L} - \underbrace{\mathbf{\Phi}(\mathbf{X}^0)}_{\mathbf{L}^0} = \left(\frac{\partial \mathbf{\Phi}(\mathbf{X})}{\partial \mathbf{X}}\right)_0 \underbrace{(\mathbf{X} - \mathbf{X}^0)}_{\mathbf{X}}$$

$$\mathbf{l}_{n\times 1}=\mathbf{J}_{n\times u}\mathbf{x}_{u\times 1}$$

with

$$\mathbf{I}_{n \times 1} = \begin{bmatrix} L_1 - \varphi_1(\mathbf{X}^0) \\ L_2 - \varphi_2(\mathbf{X}^0) \\ \vdots \\ L_n - \varphi_n(\mathbf{X}^0) \end{bmatrix}$$



and with

$$\mathbf{J}_{n \times u} = \begin{bmatrix} \frac{\partial \varphi_{1}(\mathbf{X}^{0})}{\partial X_{1}^{0}} & \frac{\partial \varphi_{1}(\mathbf{X}^{0})}{\partial X_{2}^{0}} & \cdots & \frac{\partial \varphi_{1}(\mathbf{X}^{0})}{\partial X_{u}^{0}} \\ \frac{\partial \varphi_{2}(\mathbf{X}^{0})}{\partial X_{1}^{0}} & \frac{\partial \varphi_{2}(\mathbf{X}^{0})}{\partial X_{2}^{0}} & \cdots & \frac{\partial \varphi_{2}(\mathbf{X}^{0})}{\partial X_{u}^{0}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \varphi_{n}(\mathbf{X}^{0})}{\partial X_{1}^{0}} & \frac{\partial \varphi_{n}(\mathbf{X}^{0})}{\partial X_{2}^{0}} & \cdots & \frac{\partial \varphi_{n}(\mathbf{X}^{0})}{\partial X_{u}^{0}} \end{bmatrix} = \begin{bmatrix} j_{11} & j_{12} & \cdots & j_{1u} \\ j_{21} & j_{22} & \cdots & j_{2u} \\ \vdots & \vdots & \ddots & \vdots \\ j_{n1} & j_{n2} & \cdots & j_{nu} \end{bmatrix}$$



Now: We take elements of matrix **J** and insert them into matrix **A**

$$\mathbf{A}_{n \times u} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1u} \\ a_{21} & a_{22} & \cdots & a_{2u} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nu} \end{bmatrix}$$

with



We obtain:

$$\mathbf{l}_{n\times 1} = \mathbf{A}_{n\times u} \; \mathbf{x}_{u\times 1}$$

Advantage of linearisation of the functional model: We can apply the simple algorithm for the linear case!

Observation equations:

$$\mathbf{l}_{n\times 1} + \mathbf{v}_{n\times 1} = \mathbf{A}_{n\times u} \,\, \hat{\mathbf{x}}_{u\times 1}$$

with



Vector of residuals:

$$\mathbf{v}_{n \times 1} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$
 here $\mathbf{v}_{3 \times 1} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$

Vector of adjusted reduced unknowns:

$$\hat{\mathbf{x}}_{u\times 1} = \begin{bmatrix} \hat{x}_1 \\ \hat{x}_2 \\ \vdots \\ \hat{x}_u \end{bmatrix} \qquad \text{here} \qquad \hat{\mathbf{x}}_{2\times 1} = \begin{bmatrix} d\hat{x} \\ d\hat{y} \end{bmatrix}$$



$$\Omega = \sum_{i=1}^{n} p_i v_i^2 \to \min$$

In matrix notation:

$$\mathbf{v}^{\mathrm{T}}\mathbf{P}\mathbf{v} \to \min$$
 with $\mathbf{v} = \mathbf{A}\hat{\mathbf{x}} - \mathbf{l}$

$$\mathbf{v} = \mathbf{A}\hat{\mathbf{x}} - \mathbf{I}$$

$$\Omega = (\mathbf{A}\hat{\mathbf{x}} - \mathbf{l})^{\mathrm{T}} \mathbf{P} (\mathbf{A}\hat{\mathbf{x}} - \mathbf{l})$$

$$\vdots$$

$$= \hat{\mathbf{x}}^{\mathrm{T}} \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{A} \hat{\mathbf{x}} - 2 \cdot \hat{\mathbf{x}}^{\mathrm{T}} \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{l} + \mathbf{l}^{\mathrm{T}} \mathbf{P} \mathbf{l}$$

Minimum:

$$\frac{\partial \Omega}{\partial \hat{\mathbf{x}}^{T}} = 2 \cdot \mathbf{A}^{T} \mathbf{P} \mathbf{A} \hat{\mathbf{x}} - 2 \cdot \mathbf{A}^{T} \mathbf{P} \mathbf{l} = 0$$
$$\mathbf{A}^{T} \mathbf{P} \mathbf{A} \hat{\mathbf{x}} = \mathbf{A}^{T} \mathbf{P} \mathbf{l}$$



Normal Equations:

$$\underbrace{\mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{A}}_{\mathbf{N}} \hat{\mathbf{x}} = \underbrace{\mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{I}}_{\mathbf{n}}$$

- $\mathbf{N}_{u \times u} = \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{A}$: Normal matrix
- $\mathbf{n}_{u \times 1} = \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{l}$: Right hand side of normal equations

If **N** regular \rightarrow we can compute N^{-1}

Solution of normal equations:

$$\hat{\mathbf{x}} = (\mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{A})^{-1} \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{I}$$

$$\hat{\mathbf{x}} = \mathbf{N}^{-1} \mathbf{n}$$

What is $\hat{\mathbf{x}}$? $\rightarrow \hat{\mathbf{x}}$ is a correction for the initial values \mathbf{X}^0



Adjusted unknowns:


$$\hat{\mathbf{X}} = \mathbf{X}^0 + \hat{\mathbf{x}}$$

Residuals:

$$\mathbf{v} = \mathbf{A}\hat{\mathbf{x}} - \mathbf{l}$$

Adjusted observations:

$$\hat{\mathbf{L}} = \mathbf{L} + \mathbf{v}$$





Attention: $\widehat{\mathbf{X}}$ is solution for the linearised problem (approximation)

- → We have to perform iterative computation to reach final solution
- \rightarrow We have to introduce $\hat{\mathbf{X}}$ as new approximation \mathbf{X}^0 and perform the adjustment again and again!

Question: How often do we have to repeat the computation?

Answer: Until

- a) the corrections for the unknowns $\hat{\mathbf{x}}$ become sufficiently small
- b) final check $\hat{\mathbf{L}} \Phi(\hat{\mathbf{X}}) \stackrel{!}{\approx} \mathbf{0}$ Mereoring



a) "sufficiently small"

If our unknowns are coordinates and we want to obtain results with mm $(10^{-3}\text{m}) \rightarrow \max |\hat{x}_i|$ for $i=1,\ldots,u \leq 10^{-5}$

Rule: $\max |\hat{x}_i|$ should be two orders of magnitude smaller than the resolution of the result

b)
$$\hat{\mathbf{L}} - \mathbf{\Phi}(\widehat{\mathbf{X}}) \stackrel{!}{pprox} \mathbf{0}$$

 $\Phi(\widehat{X})$: Original <u>nonlinear</u> functional model

Result contains:

- Computation error
- Linearisation error
 - → "How far away is my solution for the linearized problem from the solution for the nonlinear problem"



Rule:

$$\max |\hat{L}_i - \Phi_i(\widehat{\mathbf{X}})|$$
 should be smaller than 10^{-8}

- → If these break-off conditions are not met, continue with iterative computation
- \rightarrow Attention: Introduce \widehat{X} as new starting values X^0 and start iterative computing at step "

6.7.5 Variance-Covariance Matrices (VCM) for the results



Precision measures (see also Section 6.7.2):

• Empirical reference standard deviation:
$$s_0 = \sqrt{\frac{\mathbf{v}^\mathrm{T}\mathbf{P}\mathbf{v}}{n-u}}$$

• $\mathbf{Q}_{\widehat{X}\widehat{X}} = \mathbf{Q}_{\widehat{x}\widehat{x}}$ from variance covariance propagation with functional model $\widehat{\mathbf{X}} = \mathbf{X}^0 + \widehat{\mathbf{x}}$

• VCM of adjusted unknowns:
$$\Sigma_{\hat{X}\hat{X}} = s_0^2 \mathbf{Q}_{\hat{X}\hat{X}}$$
 with $\mathbf{Q}_{\hat{X}\hat{X}} = \mathbf{N}^{-1}$

• Cofactor matrix of adjusted observations:
$$\mathbf{Q}_{\hat{L}\hat{L}} = \mathbf{A}\mathbf{Q}_{\hat{X}\hat{X}}\mathbf{A}^{\mathrm{T}}$$

• VCM of adjusted observations:
$$\Sigma_{\hat{L}\hat{L}} = s_0^2 \mathbf{Q}_{\hat{L}\hat{L}}$$

• Cofactor matrix of residuals:
$$\mathbf{Q}_{vv} = \mathbf{Q}_{LL} - \mathbf{Q}_{\hat{L}\hat{L}}$$

• VCM of residuals:
$$\Sigma_{vv} = s_0^2 \mathbf{Q}_{vv}$$



<u>Example</u>: Find least squares solution for the unknown parameters and compute their standard deviation

Functional model:

$$\varphi_1: 4.0 = x + y - 2y^2$$
 (1)
 $\varphi_2: 8.0 = x^2 + y^2$ (2)
 $\varphi_3: 7.7 = 3x^2 - y^2$ (3)

Values 4.0, 8.0, 7.7 are equally weighted and uncorrelated observations Parameters x, y are unknowns

Observation vector:

$$\mathbf{L} = \begin{bmatrix} 4.0 \\ 8.0 \\ 7.7 \end{bmatrix}$$



Stochastic model of the observations:

$$p_1 = 1$$
, $p_2 = 1$, $p_3 = 1$

$$\mathbf{P} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \mathbf{I}$$

Vector of adjusted unknowns:

$$\widehat{\mathbf{X}} = \begin{bmatrix} \widehat{x} \\ \widehat{y} \end{bmatrix}$$

Nonlinear functional model \rightarrow solution from iterative computing with linearised functional model \rightarrow introduction of approximate values x^0 , y^0



→ Vector of starting values:

$$\mathbf{X}^0 = \begin{bmatrix} x^0 \\ y^0 \end{bmatrix}$$

Vector of adjusted reduced unknowns:

$$\hat{\mathbf{x}} = \begin{bmatrix} d\hat{x} \\ d\hat{y} \end{bmatrix} = \hat{\mathbf{X}} - \mathbf{X}^0 = \begin{bmatrix} \hat{x} - x^0 \\ \hat{y} - y^0 \end{bmatrix}$$

Vector of reduced observations:

Prof. Dr.-Ing. Frank Neitzel | Adjustment Theory I



Jacobian matrix:

$$\mathbf{J} = \begin{bmatrix} \varphi_1 & \frac{\partial \varphi_1}{\partial x^0} & \frac{\partial \varphi_1}{\partial y^0} \\ \frac{\partial \varphi_2}{\partial x^0} & \frac{\partial \varphi_2}{\partial y^0} \\ \frac{\partial \varphi_3}{\partial x^0} & \frac{\partial \varphi_3}{\partial y^0} \end{bmatrix} = \begin{bmatrix} x^0 & y^0 \\ y^0 & y^0 \\ \varphi_2 & y^0 \\ \varphi_3 & \frac{\partial \varphi_3}{\partial x^0} & \frac{\partial \varphi_3}{\partial y^0} \end{bmatrix}$$

Design matrix:

$$\mathbf{A} = \begin{bmatrix} 1 & (1 - 4y^0) \\ 2x^0 & 2y^0 \\ 6x^0 & -2y^0 \end{bmatrix}$$



Normal equations:

$$\mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{A}\hat{\mathbf{x}} = \mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{l}$$

Solutions of normal equations:

$$\hat{\mathbf{x}} = \left(\mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{A}\right)^{-1}\mathbf{A}^{\mathrm{T}}\mathbf{P}\mathbf{I}$$

Adjusted unknowns:

$$\widehat{\mathbf{X}} = \mathbf{X}^0 + \widehat{\mathbf{x}}$$


Residuals:

$$\mathbf{v} = \mathbf{A}\hat{\mathbf{x}} - \mathbf{l}$$

Adjusted observations:

$$\hat{\mathbf{L}} = \mathbf{L} + \mathbf{v}$$







Empirical reference standard deviation:

$$s_0 = \sqrt{\frac{\mathbf{v}^{\mathrm{T}} \mathbf{P} \mathbf{v}}{n - u}}$$

VCM of adjusted unknowns:

$$\mathbf{\Sigma}_{\hat{X}\hat{X}} = s_0^2 \cdot \mathbf{Q}_{\hat{X}\hat{X}}$$
 with $\mathbf{Q}_{\hat{X}\hat{X}} = \mathbf{N}^{-1}$



Standard deviation of unknowns:

$$\mathbf{\Sigma}_{\hat{X}\hat{X}} = s_0^2 \cdot \begin{bmatrix} q_{\hat{X}\hat{X}} & q_{\hat{X}\hat{Y}} \\ q_{\hat{Y}\hat{X}} & q_{\hat{Y}\hat{Y}} \end{bmatrix}$$

- Cofactor for unknown value x: $q_{\hat{x}\hat{x}}$
- Cofactor for unknown value y: $q_{\hat{y}\hat{y}}$

$$s_{\hat{x}} = \sqrt{s_0^2 \cdot q_{\hat{x}\hat{x}}} = s_0 \cdot \sqrt{q_{\hat{x}\hat{x}}}$$
$$s_{\hat{y}} = \sqrt{s_0^2 \cdot q_{\hat{y}\hat{y}}} = s_0 \cdot \sqrt{q_{\hat{y}\hat{y}}}$$



Least-squares Adjustment for Nonlinear Adjustment Problems

- Iterative solution with linearized functional model -

$$L_{1} = \varphi_{1}(X_{1}, X_{2}, \dots, X_{u})$$

$$L_{2} = \varphi_{2}(X_{1}, X_{2}, \dots, X_{u})$$

$$\vdots$$

 $L_n = \varphi_n \left(X_1, X_2, \dots, X_u \right)$

Nonlinear functional model in matrix notation:
$$\mathbf{L} = \mathbf{\Phi}(\mathbf{X}) = \begin{bmatrix} \varphi_1(\mathbf{X}) \\ \psi_2(\mathbf{X}) \\ \vdots \\ \varphi_n(\mathbf{X}) \end{bmatrix}$$

$$\vdots$$

$$\varphi_n(\mathbf{X})$$

Vector of observations: $\mathbf{L}_{n+1} = \begin{bmatrix} L_1 & L_2 & \cdots & L_n \end{bmatrix}^{\mathrm{T}}$

Variance covariance matrix of the observations: $\boldsymbol{\Sigma_{LL}}_{n,n} = \begin{bmatrix} \sigma_{L_1}^2 & \sigma_{L_1L_2} & \cdots & \sigma_{L_1L_n} \\ \sigma_{L_2L_1} & \sigma_{L_2}^2 & \cdots & \sigma_{L_2L_n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{L_LL_1} & \sigma_{L_LL_2} & \cdots & \sigma_{L_L}^2 \end{bmatrix} \text{ with theoretical values } \sigma_i$

$$\mathbf{S_{LL}} = \begin{bmatrix} s_{L_1}^2 & s_{L_1L_2} & \cdots & s_{L_1L_n} \\ s_{L_2L_1} & s_{L_2}^2 & \cdots & s_{L_2L_n} \\ \vdots & \vdots & \ddots & \vdots \\ s_{L_nL_1} & s_{L_nL_2} & \cdots & s_{L_n}^2 \end{bmatrix}$$
 with empirical values s_i

Nonlinear functional model for the unknowns:



VCM of the reduced observations from variance covariance propagation with the functional model $\mathbf{l} = \mathbf{L} - \mathbf{L}^0$

$$\sum_{\substack{\mathbf{l} \mathbf{l} \\ n,n}} = \sum_{\substack{\mathbf{l} \mathbf{L} \mathbf{L} \\ n,n}}$$

with theoretical values σ_i

$$\mathbf{S}_{\mathbf{ll}} = \mathbf{S}_{\mathbf{LL}}$$

with empirical values s_i

Theoretical reference standard deviation:

 σ_0 (or theoretical reference variance σ_0^2)

Cofactor matrix of the observations and reduced observations:

$$\mathbf{Q}_{\mathbf{LL}} = \frac{1}{\sigma_0^2} \mathbf{\Sigma}_{\mathbf{LL}}$$
 respectively $\mathbf{Q}_{\mathbf{LL}} = \frac{1}{\sigma_0^2} \mathbf{S}_{\mathbf{LL}}$

Weight matrix of the observations and reduced observations:

$$\mathbf{P}_{n,n} = \mathbf{Q}_{\mathbf{LL}}^{-1}$$

Vector of adjusted unknowns:

$$\hat{\mathbf{X}}_{u,1} = \begin{bmatrix} \hat{X}_1 & \hat{X}_2 & \cdots & \hat{X}_u \end{bmatrix}^{\mathrm{T}}$$

Vector of { initial values starting values for the unknowns: approximations

$$\mathbf{X}_{u,1}^0 = \begin{bmatrix} X_1^0 & X_2^0 & \cdots & X_u^0 \end{bmatrix}^{\mathrm{T}}$$

Vector of adjusted reduced unknowns:

$$\hat{\mathbf{x}} = \hat{\mathbf{X}} - \mathbf{X}^0$$



→ Observations as functions of the approximations for the unknowns:

$$\mathbf{L}_{n,1}^0 = \mathbf{\Phi}_{n,1} \left(\mathbf{X}_{n,1}^0 \right)$$

Vector of reduced observations:

$$\mathbf{l}_{n,1} = \mathbf{L} - \mathbf{L}_{n,1}^0$$

Jacobian matrix:

$$\mathbf{J}_{n,u} = \left(\frac{\partial \mathbf{\Phi}(\mathbf{X})}{\partial \mathbf{X}}\right)_{\mathbf{X} = \mathbf{X}^{0}} = \begin{bmatrix} \frac{\partial \mathbf{\Phi}(\mathbf{X})}{\partial X_{1}} & \frac{\partial \varphi_{2}(\mathbf{X})}{\partial X_{2}} & \cdots & \frac{\partial \varphi_{2}(\mathbf{X})}{\partial X_{u}} \\ \frac{\partial \varphi_{2}(\mathbf{X})}{\partial X_{1}} & \frac{\partial \varphi_{2}(\mathbf{X})}{\partial X_{2}} & \cdots & \frac{\partial \varphi_{2}(\mathbf{X})}{\partial X_{u}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \varphi_{n}(\mathbf{X})}{\partial X_{1}} & \frac{\partial \varphi_{n}(\mathbf{X})}{\partial X_{2}} & \cdots & \frac{\partial \varphi_{n}(\mathbf{X})}{\partial X_{u}} \end{bmatrix}_{\mathbf{X} = \mathbf{X}^{0}}$$

Coefficient matrix of the linearized functional model: $\mathbf{A} = \mathbf{J}$ "Design Matrix"

Vector of residuals:

$$\mathbf{v}_{n,1} = \begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}^{\mathrm{T}}$$

Observation equations:

$$\mathbf{l} + \mathbf{v} = \mathbf{A} \, \hat{\mathbf{x}} \\ _{n,1} = \mathbf{k} \, \hat{\mathbf{x}}$$

Normal equations:

$$\mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{A} \hat{\mathbf{x}} = \mathbf{A}^{\mathrm{T}} \mathbf{P} \mathbf{I}$$

Normal matrix:

$$\mathbf{N}_{u,u} = \mathbf{A}^{\mathrm{T}}_{u,n} \mathbf{P}_{n,n} \mathbf{A}_{n,u}$$



Right hand side of normal equations:	$\mathbf{n} = \mathbf{n}$	\mathbf{A}^{T}	PI
•	u,1	u,n	n, n, n, 1

Normal equations:
$$N \hat{\mathbf{x}} = \mathbf{n}$$

$$u, u, u, 1$$

$$u, u, 1$$

Inversion of normal matrix:
$$\mathbf{Q}_{\hat{\mathbf{x}}\hat{\mathbf{x}}} = \mathbf{N}^{-1}_{u,u}$$

Solution of normal equations:
$$\hat{\mathbf{x}} = \mathbf{Q}_{\hat{\mathbf{x}}\hat{\mathbf{x}}} \mathbf{n}_{u,u}$$

Solution for the unknowns:
$$\hat{\mathbf{X}} = \hat{\mathbf{X}}^0 + \hat{\mathbf{X}}_{u,1}$$

Vector of residuals:
$$\mathbf{v} = \mathbf{A} \hat{\mathbf{x}} - \mathbf{l}_{n,1}$$

Vector of adjusted observations:
$$\hat{\mathbf{L}} = \mathbf{L} + \mathbf{v}_{n,1}$$

Check 1:
$$\max |\hat{x}_i| \le \varepsilon \quad \forall i = 1, ..., u \quad \text{MatLab: } \max (\text{abs}(x \text{ hat}))$$

Check 2:
$$\max \left| \hat{L}_i - \varphi_i(\hat{\mathbf{X}}) \right| \le \delta \quad \forall i = 1, ..., n$$

If
$$\left\{ \left(\max |\hat{x}_i| \le \varepsilon \quad \forall i = 1, ..., u \right) \land \left(\max |\hat{L}_i - \varphi_i(\hat{\mathbf{X}})| \le \delta \quad \forall i = 1, ..., n \right) \right\}$$

 $\hat{\mathbf{X}}$ is the solution for the nonlinear adjustment problem

Else

Use $\hat{\mathbf{X}}$ as new approximation for the unknowns \mathbf{X}^0 and continue with step " \rightarrow "



Empirical reference standard deviation:

$$S_0 = \sqrt{\frac{\mathbf{v}^\mathsf{T} \mathbf{P} \mathbf{v}}{n - u}}$$

(or empirical reference variance s_0^2)

Cofactor matrix of adjusted unknowns from variance covariance propagation with the functional model $\hat{\mathbf{X}} = \mathbf{X}^0 + \hat{\mathbf{x}}$

$$\mathbf{Q}_{\hat{\mathbf{X}}\hat{\mathbf{X}}} = \mathbf{Q}_{\hat{\mathbf{x}}\hat{\mathbf{X}}}_{n,n}$$

VCM of adjusted unknowns:

$$\sum_{\substack{\hat{\mathbf{X}}\hat{\mathbf{X}}\\u,u}} = s_0^2 \, \mathbf{Q}_{\substack{\hat{\mathbf{X}}\hat{\mathbf{X}}\\u,u}}$$

Cofactor matrix of adjusted observations:

$$\mathbf{Q}_{\hat{\mathbf{L}}\hat{\mathbf{L}}} = \mathbf{A}_{n,u} \mathbf{Q}_{\hat{\mathbf{X}}\hat{\mathbf{X}}} \mathbf{A}^{\mathrm{T}}_{u,n}$$

VCM of adjusted observations:


$$\sum_{\substack{\hat{\mathbf{L}}\hat{\mathbf{L}}\\n,n}} = S_0^2 \mathbf{Q}_{\substack{\hat{\mathbf{L}}\hat{\mathbf{L}}\\n,n}}$$

Cofactor matrix of the residuals:

$$\mathbf{Q}_{\mathbf{v}\mathbf{v}} = \mathbf{Q}_{\mathbf{L}\mathbf{L}} - \mathbf{Q}_{\hat{\mathbf{L}}\hat{\mathbf{L}}}$$

VCM of the residuals:

$$\sum_{\substack{\mathbf{v}\mathbf{v}\\n,n}} = s_0^2 \, \mathbf{Q}_{\substack{\mathbf{v}\mathbf{v}\\n,n}}$$





Copyright notice:

This provided digital course material is protected by International copyright laws. All rights reserved.

Any reproduction and redistribution of the material without written permission is prohibited other than the following: You may print or download to a local hard disk extracts for your personal use only while attending the Geodesy and Geoinformation Science master's programme.

You may not distribute or commercially exploit the content. You may not transmit it or store it in any other website.

Adjustment Theory I

Chapter 6 – Introduction to Least Squares Adjustment

Prof. Dr.-Ing Frank Neitzel | Institute of Geodesy and Geoinformation Science