

*Institute of Geodesy and
Geoinformation Science*
*Chair of Geodesy and
Adjustment Theory*



Adjustment Theory I

Chapter 5: The Gaussian or Normal Distribution

Prof. Dr.-Ing. Frank Neitzel

Copyright notice:

This provided digital course material is protected by International copyright laws. All rights reserved.

Any reproduction and redistribution of the material without written permission is prohibited other than the following:
You may print or download to a local hard disk extracts for your personal use only while attending the Geodesy and Geoinformation Science master's programme.

You may not distribute or commercially exploit the content.
You may not transmit it or store it in any other website.

1. Definitions
2. Random variables
3. The random vector
4. Propagation of observation errors
5. The Gaussian or Normal Distribution
6. Introduction to least squares adjustment
7. Applications of least squares adjustment
8. Least squares adjustment with constraints
for the unknowns parameters
9. Least squares adjustment with constant values
in the functional model

5. The Gaussian or Normal Distribution

- In previous sections we have discussed random variables with arbitrary distribution
- Now: we consider random variables with certain distribution functions
- Amongst all distribution functions the Normal Distribution plays a special role because random variables in geodesy (almost) follow this distribution
- For the derivation of normal distribution we apply „theory of elementary errors“.

5.1 Theory of Elementary Errors

Theory: Was developed by BESSEL and HAGEN in 1837

Random deviation ε of an observation L can be described by a sum of q very small „elementary errors“ Δ_i .

$$\varepsilon = \Delta_1 + \Delta_2 + \Delta_3 + \dots + \Delta_q \quad ; \quad \varepsilon = \sum_{i=1}^q \Delta_i$$

Under the following conditions:

- All values Δ_i have the same absolute value $\Delta_i = |\delta| \quad \forall_i$
- All values Δ_i differ only in their sign $\varepsilon = \pm\delta \pm \delta \pm \delta + \dots$
- Positive and negative signs have same probability

$$P(\Delta_i = +\delta) = P(\Delta_i = -\delta) = 0.5$$

5.1 Theory of Elementary Errors

For $q \rightarrow \infty \Rightarrow \delta \rightarrow 0$

(for ∞ number of elementary errors the magnitude of the errors is almost zero)

Expectation $E(\Delta_h) = 0$

$$\Rightarrow E(\varepsilon) = \sum_{h=1}^q E(\Delta_h) = 0$$

$$\sigma^2 = E(\varepsilon^2) = \sum_{h=1}^q E(\Delta_h^2) = q \cdot \delta^2$$

5.1 Theory of Elementary Errors

Examples: $\varepsilon = \Delta_1 + \Delta_2 + \Delta_3 + \dots + \Delta_q$
 $\Rightarrow q = 1 : \varepsilon = \Delta_1$

Probability: $P_i(\varepsilon) = \frac{k_i}{\sum k_i}$

Case number	Δ_1	ε	Frequency (k)	Relative Frequency = Probability P
1	+	$+\delta$	1	$\frac{1}{2} = 0.5$
2	-	$-\delta$	1	$\frac{1}{2} = 0.5$

$\Rightarrow q = 2 : \varepsilon = \Delta_1 + \Delta_2$

Case number	Δ_1	Δ_2	ε	Frequency (k)	Relative Frequency = Probability P
1	+	+	$+2\delta$	1	$\frac{1}{4} = 0.25$
2	+	-	0	2	$\frac{2}{4} = 0.5$
3	-	+	0		
4	-	-	-2δ	1	$\frac{1}{4} = 0.25$

5.1 Theory of Elementary Errors

$$\Rightarrow q = 4 : \varepsilon = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4$$

Case number	Δ_1	Δ_2	Δ_3	Δ_4	ε	Frequency (k)	Relative Frequency = Probability P
1	+	+	+	+	4δ	1	$1/16 = 0.0625$
2	+	+	+	-	2δ	4	$4/16 = 0.25$
3	+	+	-	+			
4	+	-	+	+	0	6	$6/16 = 0.375$
5	-	+	+	+			
6	+	+	-	-			
7	+	-	-	+			
8	-	-	+	+			
9	-	+	+	-			
10	-	+	-	+			
11	+	-	+	-			

5.1 Theory of Elementary Errors

$$\Rightarrow q = 4 : \varepsilon = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4$$

Case number	Δ_1	Δ_2	Δ_3	Δ_4	ε	Frequency (k)	Relative Frequency = Probability P
12	-	-	-	+			
13	-	-	+	-	-2δ	4	$4/16 = 0.25$
14	-	+	-	-			
15	+	-	-	-			
16	-	-	-	-	-4δ	1	$1/16 = 0.0625$

→ The larger the deviation (“error”) the smaller the probability

5.1 Theory of Elementary Errors

Pascal Triangle

$\varepsilon = \sum \Delta_i$	-6 δ	-5 δ	-4 δ	-3 δ	-2 δ	- δ	0	+ δ	+2 δ	+3 δ	+4 δ	+5 δ	+6 δ	$\sum k$
$q = 1$						1		1						2
$q = 2$					1		2		1					4
$q = 3$			1		3	+	3		1					8
$q = 4$		1		4	+	6	+	4		1				16
$q = 5$	1		5		10		10		5		1			32
$q = 6$	1		6		15		20		15		6		1	64

5.1 Theory of Elementary Errors

Formula for all cases

$$\varepsilon = (q - 2 \cdot w) \cdot \delta$$

with $w = 0, 1, 2, \dots, q$

$q = 1$

$w = 0$	$\varepsilon = (1 - 2 \cdot 0) \cdot \delta$	$+ \delta$
$w = 1$	$\varepsilon = (1 - 2 \cdot 1) \cdot \delta$	$- \delta$

$q = 2$

$w = 0$	$\varepsilon = (2 - 2 \cdot 0) \cdot \delta$	$+2\delta$
$w = 1$	$\varepsilon = (2 - 2 \cdot 1) \cdot \delta$	0
$w = 2$	$\varepsilon = (2 - 2 \cdot 2) \cdot \delta$	-2δ

5.1 Theory of Elementary Errors

General Law


$$P(\varepsilon) = P\{\varepsilon = (q - 2w)\delta\} = \frac{1}{2^q} \binom{q}{w}, \quad \text{with } \binom{q}{w} = \frac{q!}{(q-w)!w!}$$

Please note: $\sum_{w=0}^q \binom{q}{w} = 2^q$ and $\sum_{w=0}^q P(\varepsilon) = \frac{1}{2^q} \sum_{w=0}^q \binom{q}{w} = 1$

$q = 10$


We can now directly compute ε_i and $P(\varepsilon_i)$, $w = 0, 1, 2, \dots, 10$

w	0	1	2	3	4	5	6	7	8	9	10
ε	10δ	8δ	6δ	4δ	2δ	0	-2δ	-4δ	-6δ	-8δ	-10δ
$P(\varepsilon)$	0.001	0.010	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.010	0.001



5.1 Theory of Elementary Errors

Graphical representation for $q = 10$.




5.1 Theory of Elementary Errors

getting big \checkmark *& getting small*
Now: $q \rightarrow \infty \Rightarrow \delta \rightarrow 0$

& Continuous bell curve function

$$P(\xi \leq \varepsilon < \xi + d\xi) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{\xi^2}{2\sigma^2}\right) d\xi = f(\xi) d\xi$$



Definition

A random variable is normally distributed if it can be described by a sum of a large amount of “elementary errors” with (more or less) same magnitude.

It is: $\varepsilon = \Delta_1 + \Delta_2 + \dots + \Delta_n$

With: ε random deviation

Δ_i elementary components of the deviation

$$E(\Delta_i) = 0 \quad \forall_i \leftarrow \text{expectation}$$

$$n \rightarrow \infty$$

no Δ_i is dominating

5.2 Central Limit Theorem (CLT)

Comparison with BESSEL and HAGEN:

- All values Δ_i have the same absolute value $\Delta_i = |\delta| \quad \forall_i$
- All values Δ_i differ only in their sign $\varepsilon = \pm\delta \pm \delta \pm \delta \pm \dots$
- Positive and negative signs have same probability $P(\Delta_i = +\delta) = P(\Delta_i = -\delta) = 0.5$

For $n \rightarrow \infty \Rightarrow \delta \rightarrow 0$

\Rightarrow Then it holds:

$$P(\xi \leq \varepsilon < \xi + d\xi) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{\xi^2}{2 \cdot \sigma^2}\right) d\xi = f(\xi) d\xi$$

With σ = Constant value, to be defined (tbd)

$f(\xi)$ = Density of normal distribution

History of Central Limit Theorem (CLT)

de Moivre 1730

Laplace 1892

}

First ideas, no proof

Gauss 1820

introduction into error theory (\rightarrow “Gauss Distribution”)

Lyapunov 1901

Proof!

In practice:

If many “elementary errors” have (more or less) the same magnitude,
we can **assume** a normal distribution.

5.2 Expectation and Variance of a normally distributed random variable

Given: random variable


$$L = \mu + \varepsilon \rightarrow \varepsilon = L - \mu$$

what we want

with: μ expectation $E(L)$

ε random deviation

$f(x)$ density of L



Shifting the coordinate system:

$$x = \mu + \xi \rightarrow \xi = \underbrace{x - \mu}_{d\xi} \rightarrow d\xi = dx \rightarrow f(x) = f(\xi)$$

Substitution yields

$$P(\xi \leq \varepsilon < \xi + d\xi) = \underbrace{f(\xi)}_{f(x)} d\xi$$

$$P(x - \cancel{\mu} \leq L - \cancel{\mu} < x - \cancel{\mu} + dx)$$

$$P(x \leq L < x + dx) = f(x)dx$$

5.2 Expectation and Variance of a normally distributed random variable

Normal distribution from Central Limit Theorem: $f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2}\frac{\xi^2}{\sigma^2}\right)$

Substitution $\frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}\right) = f(x)$

$$P(x \leq L < x + dx) = \underbrace{\frac{1}{\sigma\sqrt{2\pi}}}_{f(x)} \exp\left(-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}\right) dx$$

Probability density and thus the distribution function of a normally distributed random variable is completely defined by two parameters: Expectation μ and variance σ^2 !!

$$L \sim N(\mu, \sigma^2)$$


follows := ?

Expectation

Variance

5.2 Expectation and Variance of a normally distributed random variable

" L follows the normal distribution with expectation μ and variance σ^2 "



μ : defines the location

σ : defines the shape

σ never reaches x -axis

sigma defines shape

3 sigma rule
if G exceeds $3\sigma = 99.7\%$ = it's a
rare error

Adjustment Theory I

Chapter 5: The Gaussian or Normal Distribution

Prof. Dr.-Ing. Frank Neitzel

Copyright notice:

This provided digital course material is protected by International copyright laws. All rights reserved.

Any reproduction and redistribution of the material without written permission is prohibited other than the following:
You may print or download to a local hard disk extracts for your personal use only while attending the Geodesy and Geoinformation Science master's programme.

You may not distribute or commercially exploit the content.
You may not transmit it or store it in any other website.