
#### 3.2 The *m*-dimensional random vector



#### 3.2.1 Theoretical expectation and theoretical covariance matrix

**Example:** Measurement of directions and distances in a geodetic network



# 3.2.1 Theoretical expectation and theoretical covariance matrix



- ▶ Given: m random variables  $L_1, L_2, ..., L_m$
- Random vector

$$\mathbf{L}_{m \times 1} = \begin{bmatrix} L_1 \\ L_2 \\ \vdots \\ L_m \end{bmatrix}$$

► Vector of expectations

$$\mathbf{\mu}_{m \times 1} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_m \end{bmatrix} = E\{\mathbf{L}_{m \times 1}\} = \begin{bmatrix} E(L_1) \\ E(L_2) \\ \vdots \\ E(L_m) \end{bmatrix}$$

# 3.2.1 Theoretical expectation and theoretical covariance matrix



► Vector of random deviations

v Bepletation

$$\mathbf{\varepsilon}_{m\times 1} = \begin{bmatrix} \varepsilon_{1} \\ \varepsilon_{2} \\ \vdots \\ \varepsilon_{m} \end{bmatrix} = \mathbf{L}_{m\times 1} - \mathbf{\mu}_{m\times 1} = \begin{bmatrix} L_{1} - \mu_{1} \\ L_{2} - \mu_{2} \\ \vdots \\ L_{m} - \mu_{m} \end{bmatrix}$$

$$5 \times \mathcal{A} \text{ Additive } \mathcal{A} \text{ Ad$$

▶ as in 2.4.1

$$E(\mathbf{\varepsilon}_{m\times 1}) = E\{\mathbf{L}_{m\times 1} - \mathbf{\mu}_{L_{m\times 1}}\} = \underbrace{E(\mathbf{L}_{m\times 1})}_{\mathbf{L}_{m\times 1}} - \mathbf{\mu}_{L_{m\times 1}} = \mathbf{0}$$

$$E\{\boldsymbol{\varepsilon}_{L_{m\times 1}}\cdot\boldsymbol{\varepsilon}_{L_{1\times m}}^{\mathrm{T}}\} = \begin{bmatrix} E(\varepsilon_{1}^{2}) & E(\varepsilon_{1}\cdot\varepsilon_{2}) & \cdots & E(\varepsilon_{1}\cdot\varepsilon_{m}) \\ E(\varepsilon_{2}\cdot\varepsilon_{1}) & E(\varepsilon_{2}^{2}) & \cdots & E(\varepsilon_{2}\cdot\varepsilon_{m}) \\ \vdots & \vdots & \ddots & \vdots \\ E(\varepsilon_{m}\cdot\varepsilon_{1}) & E(\varepsilon_{m}\cdot\varepsilon_{2}) & \cdots & E(\varepsilon_{m}\cdot\varepsilon_{m}) \end{bmatrix} = \boldsymbol{\Sigma}_{LL_{m\times m}}$$

Prof. Dr.-Ing. Frank Neitzel | Adjustment Theory I

Varme.

# 3.2.1 Theoretical expectation and theoretical covariance matrix



► Theoretical VCM of L

$$\mathbf{\Sigma}_{LL_{m \times m}} = \begin{bmatrix} \sigma_{1}^{2} & \sigma_{12} & \cdots & \sigma_{1m} \\ \sigma_{21} & \sigma_{2}^{2} & \cdots & \sigma_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{m1} & \sigma_{m2} & \cdots & \sigma_{m}^{2} \end{bmatrix} = \begin{bmatrix} \sigma_{1}^{2} & \rho_{12} \cdot \sigma_{1} \cdot \sigma_{2} & \cdots & \sigma_{1m} \\ \rho_{21} \cdot \sigma_{2} \cdot \sigma_{1} & \sigma_{2}^{2} & \cdots & \sigma_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ \rho_{m1} \cdot \sigma_{m} \cdot \sigma_{1} & \rho_{m2} \cdot \sigma_{m} \cdot \sigma_{2} & \cdots & \sigma_{m}^{2} \end{bmatrix}$$

ightharpoonup Theoretical Variance of  $L_i$ 

# 3.2.2 Empirical expectation and empirical covariance matrix



- ▶ Given: m random variables  $\mathbf{L}_{m \times 1} = \begin{bmatrix} L_1 \\ L_2 \\ \vdots \\ L_m \end{bmatrix}$
- $\rightarrow m \times n$ -dimensional observation matrix

$$\mathbf{l}_{m \times n} = \begin{bmatrix} l_{11} & l_{21} & \cdots & l_{m1} \\ l_{12} & l_{22} & \cdots & l_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ l_{1n} & l_{2n} & \cdots & l_{mn} \end{bmatrix}$$

m different random variables were measured n times

 $\blacktriangleright$  m-dimensional vector of mean values respective expectations

$$\bar{\mathbf{I}} = \begin{bmatrix} \bar{l}_1 \\ \bar{l}_2 \\ \vdots \\ \bar{l}_m \end{bmatrix} \qquad \text{resp.} \qquad \mathbf{\mu} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_m \end{bmatrix}$$

# 3.2.2 Empirical expectation and empirical covariance matrix




► m-dimensional VCM

$$\mathbf{S}_{ll_{m \times m}} = \begin{bmatrix} s_1^2 & s_{12} & \cdots & s_{1m} \\ s_{21} & s_2^2 & \cdots & s_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ s_{m1} & s_{m2} & \cdots & s_m^2 \end{bmatrix}$$

Correlation coefficient for the m-dimensional case

$$ho_{ij} = rac{\sigma_{ij}}{\sigma_i \cdot \sigma_j}$$
 respective  $r_{ij} = rac{s_{ij}}{s_i \cdot s_j}$  for  $i, j = 1, 2, ..., m$  and  $i \neq j$ 

→ All computations (e.g. residuals, empirical variances and covariances and correlation coefficients as in 2-dimensional case)





#### **Copyright notice:**

This provided digital course material is protected by International copyright laws. All rights reserved.

Any reproduction and redistribution of the material without written permission is prohibited other than the following: You may print or download to a local hard disk extracts for your personal use only while attending the Geodesy and Geoinformation Science master's programme.

You may not distribute or commercially exploit the content. You may not transmit it or store it in any other website.

#### Adjustment Theory I

Chapter 3 - The Random Vector

Prof. Dr.-Ing Frank Neitzel | Institute of Geodesy and Geoinformation Science