

geospatialsuite: Comprehensive Geospatiotemporal Analysis and Multimodal Integration Toolkit for R

28 December 2025

Summary

Geospatialsuite is a comprehensive R package for geospatiotemporal analysis that provides researchers with a unified toolkit for multimodal spatial data integration. The package features over 60 vegetation indices with automatic satellite band detection, universal spatial join operations that work with any raster-vector combination, robust raster visualization with built-in error handling, and rapid mapping capabilities through the `quick_map()` function. Designed for reliability and universal applicability, geospatialsuite addresses critical needs in agricultural research, environmental monitoring, and remote sensing applications while maintaining computational efficiency for large datasets.

The package seamlessly integrates with the modern R spatial ecosystem through `terra` (Hijmans 2022) and `sf` (Pebesma 2018), providing enhanced NDVI calculations with quality filtering, comprehensive water quality analysis using multiple indices (NDWI (McFeeters 1996), MNDWI (Xu 2006), NDMI), crop data layer (CDL) analysis (USDA National Agricultural Statistics Service 2024), spatial interpolation techniques (Cressie 1993), and terrain analysis. Unlike existing solutions that are often limited by complex dependencies or geographic scope, geospatialsuite emphasizes universal functionality—working seamlessly across different regions, satellite platforms (Landsat, Sentinel-2, MODIS), and data types with comprehensive error handling throughout. The package has enabled research in agricultural systems monitoring (Olatunde D. Akanbi, Bhuvanagiri, et al. 2024) and multimodal data integration (Olatunde D. Akanbi, Li, et al. 2024).

Statement of Need

Geospatial analysis in environmental and agricultural research requires reliable, standardized tools that can handle diverse datasets across different geographic regions and satellite platforms. While foundational packages like `terra` (Hijmans 2022) and `sf` (Pebesma 2018) provide essential spatial data handling capabilities, researchers often face significant challenges when attempting to integrate multiple data sources, calculate specialized indices, or create reliable visualizations from large raster datasets.

Current solutions in the R ecosystem present several critical limitations: vegetation index packages typically cover only a subset of available indices and lack automatic band detection across different satellite platforms; spatial join operations often fail with edge cases or require extensive preprocessing; visualization tools frequently encounter memory issues with large rasters or produce inconsistent results; and multimodal data integration requires custom workflows that are difficult to reproduce, error-prone, and time-consuming to implement.

geospatialsuite addresses these fundamental challenges through several key innovations:

Universal spatial operations with robust error handling: The package provides a universal spatial join function that reliably handles any raster-vector combination with comprehensive error checking, automatic coordinate system handling, and graceful failure recovery. This eliminates the common frustrations researchers face when working with diverse spatial datasets.

Comprehensive vegetation analysis with automatic detection: Over 60 vegetation indices including NDVI (Rouse Jr et al. 1974), EVI (Huete et al. 2002), SAVI (Rondeaux, Steven, and Baret 1996), and ARVI (Kaufman and Tanré 1992) with automatic band detection that works seamlessly across Landsat, Sentinel-2, and MODIS platforms. Quality filtering and standardized output formats ensure consistent results regardless of input data source.

Reliable visualization for large datasets: Built on `terra`'s efficient raster handling, the package provides publication-quality mapping through functions like `quick_map()` that handle large rasters without memory issues, automatic legend generation, and consistent color schemes. The visualization system includes robust error handling that prevents common plotting failures.




Figure 1: Example output from `geospatialsuite`'s `quick_map()` function demonstrating universal spatial mapping capabilities. The function automatically detects data types, coordinate systems, and optimal visualization parameters, requiring only a single line of code to produce publication-quality maps from any spatial data format.

Multimodal integration workflows: Streamlined functions for combining satellite imagery, weather data, crop data layers, and administrative boundaries with built-in spatial and temporal alignment, quality checks, and standardized output formats. This addresses the type of complex geospatial analysis challenges found across diverse research domains (Gordon et al. 2025).

The package architecture prioritizes reproducibility, computational efficiency, and ease of use through comprehensive documentation, extensive testing, and a consistent API design. This approach has proven essential in research funded by the National Science Foundation Engineering Research Center for Advancing Sustainable and Distributed Fertilizer Production (CASFER), where reliable geospatial analysis tools are critical for understanding agricultural systems at multiple scales (Olatunde D. Akanbi, Bhuvanagiri, et al. 2024; Olatunde David Akanbi 2024).

`geospatialsuite` fills a critical gap by providing a single, well-documented package that handles the most common geospatial analysis challenges with the reliability and consistency required for reproducible science, while maintaining the flexibility needed for specialized research applications.

Software Architecture

geospatialsuite is organized into 10 functional categories comprising 165 functions (Figure 2), designed to provide comprehensive geospatial analysis capabilities while maintaining ease of use and reliability. The package architecture emphasizes modularity, with each functional category serving specific analytical needs while integrating seamlessly with other components.



Figure 2: geospatialsuite functional organization showing the 10 major categories of functions: Core Spatial Operations, Water Quality and Indices, Spatial Interpolation, Testing & Validation, Utility Functions, Temporal Analysis, Data Processing, Visualization & Mapping, Comprehensive Workflows, and Vegetation Analysis.

The core design philosophy centers on universal compatibility and robust error handling. Functions automatically detect data types, coordinate systems, and optimal processing methods, reducing the technical barrier for researchers while maintaining the flexibility needed for advanced applications. This architecture enables both novice users to quickly generate results and experienced researchers to implement sophisticated analytical workflows.

Availability

geospatialsuite is available on the Comprehensive R Archive Network (CRAN) at <https://cran.r-project.org/web/packages/geospatialsuite/> and can be installed using `install.packages("geospatialsuite")`. The source code is actively maintained on GitHub with comprehensive documentation, examples, and issue tracking to support the research community.

Acknowledgements

This material is based upon financial support by the National Science Foundation, EEC Division of Engineering Education and Centers, NSF Engineering Research Center for Advancing Sustainable and Distributed Fertilizer production (CASFER), NSF 20-553 Gen-4 Engineering Research Centers award 2133576. The authors thank the broader CASFER research community for feedback and testing that improved the package’s functionality and usability.

References

- Akanbi, Olatunde David. 2024. “Leveraging Multimodal Data for Geospatiotemporal Analytics.” Master’s thesis, Case Western Reserve University.
- Akanbi, Olatunde D, Deepa C Bhuvanagiri, Erika I Barcelos, Arafath Nihar, Brian Gonzalez Hernandez, Jeffrey M Yarus, and Roger H French. 2024. “Integrating Multiscale Geospatial Analysis for Monitoring Crop Growth, Nutrient Distribution, and Hydrological Dynamics in Large-Scale Agricultural Systems.” *Journal of Geovisualization and Spatial Analysis* 8 (1): 9. <https://doi.org/10.1007/s41651-023-00164-y>.
- Akanbi, Olatunde D, Jiaqi Li, Vibha Mandayam, Arafath Nihar, Yinghui Wu, Laura S Bruckman, Jeffrey M Yarus, Erika I Barcelos, and Roger H French. 2024. “Integrating Multimodal Geospatiotemporal Data for Societal, Economic, and Environmental (SEE) Analysis of Large Agricultural Systems.” In *2024 IEEE International Conference on Big Data (BigData)*, 1–1. IEEE Computer Society. <https://doi.org/10.1109/BigData62323.2024.10825456>.
- Cressie, Noel AC. 1993. *Statistics for Spatial Data*. New York: John Wiley & Sons.
- Gordon, Jonathan E, Olatunde D Akanbi, Deepa C Bhuvanagiri, Hope E Omodolor, Vibha Mandayam, Roger H French, Jeffrey M Yarus, and Erika I Barcelos. 2025. “Geospatial Modeling of Near Subsurface Temperatures of the Contiguous United States for Assessment of Materials Degradation.” *Scientific Reports* 15 (1): 1053. <https://doi.org/10.1038/s41598-024-85050-3>.
- Hijmans, Robert J. 2022. “Terra: Spatial Data Analysis.” *R Package Version 1.6-17*. <https://CRAN.R-project.org/package=terra>.
- Huete, Alfredo, Kamel Didan, Tomoaki Miura, E Patricia Rodriguez, Xiang Gao, and Laerte G Ferreira. 2002. “Overview of the Radiometric and Biophysical Performance of the MODIS Vegetation Indices.” *Remote Sensing of Environment* 83 (1-2): 195–213. [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2).
- Kaufman, Yoram J, and Didier Tanré. 1992. “Atmospherically Resistant Vegetation Index (ARVI) for EOS-MODIS.” *IEEE Transactions on Geoscience and Remote Sensing* 30 (2): 261–70. <https://doi.org/10.1109/36.134076>.
- McFeeters, Stuart K. 1996. “The Use of the Normalized Difference Water Index (NDWI) in the Delineation of Open Water Features.” *International Journal of Remote Sensing* 17 (7): 1425–32. <https://doi.org/10.1080/01431169608948714>.
- Pebesma, Edzer. 2018. “Simple Features for R: Standardized Support for Spatial Vector Data.” *The R Journal* 10 (1): 439–46. <https://doi.org/10.32614/RJ-2018-009>.
- Rondeaux, Gilles, Michael Steven, and Frédéric Baret. 1996. “Optimization of Soil-Adjusted Vegetation Indices.” *Remote Sensing of Environment* 55 (2): 95–107. [https://doi.org/10.1016/0034-4257\(95\)00186-7](https://doi.org/10.1016/0034-4257(95)00186-7).
- Rouse Jr, John W, RH Haas, JA Schell, and DW Deering. 1974. “Monitoring Vegetation Systems in the Great Plains with ERTS.” *NASA Special Publication* 351: 309.
- USDA National Agricultural Statistics Service. 2024. *Cropland Data Layer*. https://www.nass.usda.gov/Research_and_Science/Cropland/.
- Xu, Hanqiu. 2006. “Modification of Normalised Difference Water Index (NDWI) to Enhance Open Water

Features in Remotely Sensed Imagery.” *International Journal of Remote Sensing* 27 (14): 3025–33.
<https://doi.org/10.1080/01431160600589179>.