

# Homework 3

Quantum Mechanics

Sept 15th, 2022

CLAYTON SEITZ

**Problem 1.** *Problem 2.1 from Sakurai*

**Solution.** The Heisenberg equation of motion reads

$$\frac{dA}{dt} = \frac{1}{i\hbar} [A, H]$$

For the spin precession problem, we have the Hamiltonian

$$H = - \left( \frac{eB}{mc} \right) S_z = \omega S_z$$

For  $A = S_x, S_y, S_z$ , the time evolution is given by

$$\begin{aligned} \frac{dS_x}{dt} &= \frac{\omega}{i\hbar} [S_x, S_z] = -\omega S_y \\ \frac{dS_y}{dt} &= \frac{\omega}{i\hbar} [S_y, S_z] = \omega S_x \\ \frac{dS_z}{dt} &= \frac{\omega}{i\hbar} [S_z, S_z] = 0 \end{aligned}$$

The above system has a straightforward solution:

$$\begin{aligned} S_x(t) &= \cos(\omega t) \\ S_y(t) &= \sin(\omega t) \\ S_z(t) &= S_z(0) \end{aligned}$$

■

**Problem 2.** *Problem 2.3 from Sakurai*

**Solution.** We are given that  $\vec{B} = B\hat{z}$  and that we are in the eigenstate  $|\psi(0)\rangle = |\mathbf{S} \cdot \hat{\mathbf{n}}\rangle_+$ , which reads

$$\begin{aligned} |\psi(0)\rangle &= \psi_+ |+\rangle + \psi_- |-\rangle \\ &= \cos \frac{\beta}{2} |+\rangle + \sin \frac{\beta}{2} |-\rangle \end{aligned}$$

where we have set  $\alpha = 0$  since the ket is in the x-z plane. This state will evolve according to a Hamiltonian

$$H = - \left( \frac{eB}{m_e c} \right) S_z$$

Let  $\omega = |e|B/m_e c$  giving  $H = \omega S_z$ . We have the energies

$$E_{\pm} = \mp \frac{e\hbar B}{2m_e c} = \mp \hbar \omega$$

$$\begin{aligned} |\psi(t)\rangle &= \psi_+(0) \exp \left( \frac{-iE_+ t}{\hbar} \right) |+\rangle + \psi_-(0) \exp \left( \frac{-iE_- t}{\hbar} \right) |-\rangle \\ &= \cos \frac{\beta}{2} \exp \left( \frac{-i\omega t}{2} \right) |+\rangle + \sin \frac{\beta}{2} \exp \left( \frac{i\omega t}{2} \right) |-\rangle \end{aligned}$$

In general, the probability of measuring  $|+\rangle_x = \frac{1}{\sqrt{2}} |+\rangle + \frac{1}{\sqrt{2}} |-\rangle$  is given by the inner product

$$\begin{aligned} |\langle S_x; + | \psi; t \rangle|^2 &= \left| \left( \frac{1}{\sqrt{2}} \langle + | + \frac{1}{\sqrt{2}} \langle - | \right) \cdot \right. \\ &\quad \left. \left( \psi_+ \exp \left( \frac{-i\omega t}{2} \right) |+\rangle + \psi_- \exp \left( \frac{i\omega t}{2} \right) |-\rangle \right) \right|^2 \\ &= \left| \frac{1}{\sqrt{2}} \cos \frac{\beta}{2} \exp \left( \frac{-i\omega t}{2} \right) + \frac{1}{\sqrt{2}} \sin \frac{\beta}{2} \exp \left( \frac{i\omega t}{2} \right) \right|^2 \end{aligned}$$

Using the half-angle identity for  $\sin \theta$  and some straightforward arithmetic gives

$$|\langle S_x; + | \psi; t \rangle|^2 = \frac{1 + \sin \beta \cos \omega t}{2}$$

For the time-dependence of  $\langle S_x \rangle$ , we have

$$\begin{aligned} \langle S_x \rangle(t) &= \langle \psi; t | S_x | \psi; t \rangle \\ &= \left( \psi_+ \exp \left( \frac{i\omega t}{2} \right) \langle + | + \psi_- \exp \left( \frac{-i\omega t}{2} \right) \langle - | \right) \\ &\quad \cdot \frac{\hbar}{2} \left( \psi_+ \exp \left( -\frac{i\omega t}{2} \right) | - \rangle + \psi_- \exp \left( \frac{i\omega t}{2} \right) | + \rangle \right) \end{aligned}$$

Substituting  $\psi_+$  and  $\psi_-$  with the same values as above, we get

$$\langle S_x \rangle(t) = \frac{\hbar}{2} \sin \beta \cos \omega t$$

When  $\beta = \pi/2$  the probability oscillates between 0 and 1 with frequency  $\omega$  and when  $\beta = 0$  then the probability is always 1/2, as expected. The expectation value also makes sense because when  $\beta = 0$ , we can get  $\pm\hbar/2$  with equal probability, giving zero on average. When  $\beta = \pi/2$  the expectation value oscillates between  $\hbar/2$  and  $-\hbar/2$ . ■

**Problem 3.** *Problem 2.9 from Sakurai*

**Solution.** ■

**Problem 4.** *Problem 2.10 from Sakurai*

**Solution.** Let  $|\psi\rangle = \alpha|a'\rangle + \beta|a''\rangle$  be an eigenvector of the Hamiltonian. Note that this must be real for the eigenvalue to be real. That means that

$$\begin{aligned} H|\psi\rangle &= (|a'\rangle\delta\langle a''| + |a''\rangle\delta\langle a'|)(\alpha|a'\rangle + \beta|a''\rangle) \\ &= \delta(\alpha|a''\rangle + \beta|a'\rangle) \end{aligned}$$

Therefore  $\alpha = \beta = \frac{1}{\sqrt{2}}$  or  $\alpha = \frac{1}{\sqrt{2}}$  and  $\beta = -\frac{1}{\sqrt{2}}$ . Giving eigenvalues  $\pm\delta$ . To get the time evolution of the state, we need to express these in the basis of  $H$ . Just based on inspection of the two bases, we can tell that

$$|a'\rangle = \frac{1}{\sqrt{2}} (|\psi_1\rangle - |\psi_2\rangle)$$

$$|a''\rangle = \frac{1}{\sqrt{2}} (|\psi_1\rangle + |\psi_2\rangle)$$

and, since the Hamiltonian is time-independent, a state prepared in  $|a'\rangle$  will evolve according to

$$|\alpha(t)\rangle = \frac{1}{\sqrt{2}} \exp\left(\frac{-i\delta t}{\hbar}\right) |\psi_1\rangle - \frac{1}{\sqrt{2}} \exp\left(\frac{i\delta t}{\hbar}\right) |\psi_2\rangle$$

The probability of finding the system in the state  $|a''\rangle$  at a later time is

$$\begin{aligned} |\langle a''|\alpha(t)\rangle|^2 &= \left| \frac{1}{\sqrt{2}} (\langle\psi_1| + \langle\psi_2|) \right. \\ &\quad \cdot \left( \frac{1}{\sqrt{2}} \exp\left(\frac{-i\delta t}{\hbar}\right) |\psi_1\rangle - \frac{1}{\sqrt{2}} \exp\left(\frac{i\delta t}{\hbar}\right) |\psi_2\rangle \right) \Big|^2 \\ &= \frac{1}{4} \sin^2 \frac{\delta t}{\hbar} \end{aligned}$$

This could describe a system in which the eigenvectors of the Hamiltonian are simultaneous with the eigenvectors of  $S_x$ , however the states  $|a'\rangle$  and  $|a''\rangle$  are expressed in the  $S_z$  basis. ■

**Problem 5.** *Problem 2.12 from Sakurai*

**Solution.** The state is prepared in

$$|\alpha; t=0\rangle = \frac{1}{\sqrt{2}} |0\rangle + \frac{\exp(i\delta)}{\sqrt{2}} |1\rangle$$

In general, the energies of  $|n\rangle$  are  $E_n = (n + \frac{1}{2}) \hbar\omega$ . Therefore, the time dependence of the state can be evaluated as

$$\begin{aligned} |\alpha; t\rangle &= \exp\left(-\frac{iHt}{\hbar}\right) |\alpha; t=0\rangle \\ &= \frac{1}{\sqrt{2}} \exp\left(-\frac{i\omega t}{2}\right) |0\rangle + \frac{1}{\sqrt{2}} \exp(i\delta) \exp\left(-\frac{3i\omega t}{2}\right) |1\rangle \end{aligned}$$

$$\langle x'|\alpha;t\rangle = \frac{1}{\sqrt{2}} \exp \frac{-i\omega t}{2} \langle x'|0\rangle + \frac{1}{\sqrt{2}} \exp(i\delta) \exp \frac{-3i\omega t}{2} \langle x'|1\rangle$$

and we know in general that the position representation of  $|n\rangle$  i.e.,  $\langle x'|n\rangle$  are

$$\langle x'|n\rangle = \psi_n(x) = \frac{1}{\sqrt{2^n n!}} \left( \frac{m\omega}{\pi \hbar} \right)^{1/4} e^{-\frac{m\omega x^2}{2\hbar}} H_n \left( \sqrt{\frac{m\omega}{\hbar}} x \right)$$

$$\begin{aligned} \langle x\rangle(t) &= \langle \alpha;t|x|\alpha;t\rangle \\ &= \left( \frac{1}{\sqrt{2}} \exp \frac{i\omega t}{2} \langle 0| + \frac{1}{\sqrt{2}} \exp(-i\delta) \exp \frac{3i\omega t}{2} \langle 1| \right) \\ &\quad x \left( \frac{1}{\sqrt{2}} \exp \frac{-i\omega t}{2} |0\rangle + \frac{1}{\sqrt{2}} \exp(i\delta) \exp \frac{-3i\omega t}{2} |1\rangle \right) \\ &= \frac{1}{2} \langle 0|x|0\rangle + \frac{1}{2} \langle 1|x|1\rangle \\ &\quad + \frac{1}{2} \exp(i\delta) \exp(-i\omega t) \langle 0|x|1\rangle + \frac{1}{2} \exp(-i\delta) \exp(i\omega t) \langle 1|x|0\rangle \end{aligned}$$

Now recall the general expression for the matrix element of  $x$

$$\langle n'|x|n\rangle = \sqrt{\frac{\hbar}{2m\omega}} \left( \sqrt{n} \delta_{n',n-1} + \sqrt{n+1} \delta_{n',n+1} \right)$$

which means that the above expression simplifies to

$$\begin{aligned} \langle x\rangle(t) &= \frac{1}{2} \exp(i\delta) \exp(-i\omega t) \langle 0|x|1\rangle + \frac{1}{2} \exp(-i\delta) \exp(i\omega t) \langle 1|x|0\rangle \\ &= \frac{1}{2} \sqrt{\frac{\hbar}{2m\omega}} (\exp(i\delta) \exp(-i\omega t) + \exp(-i\delta) \exp(i\omega t)) \\ &= \sqrt{\frac{\hbar}{2m\omega}} \cos(\delta - \omega t) \\ &= \sqrt{\frac{\hbar}{2m\omega}} \cos(\omega t - \delta) \end{aligned}$$

For momentum, we can just replace the operator  $x$  with  $p$  in the expressions above:

$$\begin{aligned}
\langle p \rangle(t) &= \langle \alpha; t | x | \alpha; t \rangle \\
&= \left( \frac{1}{\sqrt{2}} \exp \frac{i\omega t}{2} \langle 0 | + \frac{1}{\sqrt{2}} \exp(-i\delta) \exp \frac{3i\omega t}{2} \langle 1 | \right) \\
&\hat{p} \left( \frac{1}{\sqrt{2}} \exp \frac{-i\omega t}{2} | 0 \rangle + \frac{1}{\sqrt{2}} \exp(i\delta) \exp \frac{-3i\omega t}{2} | 1 \rangle \right) \\
&= \frac{1}{2} \langle 0 | p | 0 \rangle + \frac{1}{2} \langle 1 | p | 1 \rangle \\
&\quad + \frac{1}{2} \exp(i\delta) \exp(-i\omega t) \langle 0 | p | 1 \rangle + \frac{1}{2} \exp(-i\delta) \exp(i\omega t) \langle 1 | p | 0 \rangle
\end{aligned}$$

and we have another general expression for the matrix element of  $p$

$$\langle n' | p | n \rangle = i \sqrt{\frac{m\hbar\omega}{2}} \left( -\sqrt{n} \delta_{n',n-1} + \sqrt{n+1} \delta_{n',n+1} \right)$$

which again means that the above expression simplifies to

$$\begin{aligned}
\langle p \rangle(t) &= \frac{1}{2} \exp(i\delta) \exp(-i\omega t) \langle 0 | p | 1 \rangle + \frac{1}{2} \exp(-i\delta) \exp(i\omega t) \langle 1 | p | 0 \rangle \\
&= \frac{i}{2} \sqrt{\frac{m\hbar\omega}{2}} \left( -\exp(i\delta) \exp(-i\omega t) + \exp(-i\delta) \exp(i\omega t) \right) \\
&= -\sqrt{\frac{m\hbar\omega}{2}} \sin(\omega t - \delta)
\end{aligned}$$

■

**Problem 6.** *Problem 2.13 from Sakurai*

**Solution.**

■