

Synaptic Plasticity

“the only game in town”

-R. Malenka

Short Term Plasticity

Paired pulse facilitation / depression

Presynaptic modulation

Presynaptic receptors

Postsynaptic modulation

Long Term Plasticity

LTP, LTD

Synapses vary!

Some important questions...

Synapses are modified in response to a wide range of stimuli, activity patterns, drugs, behavioral states, etc...

When changes in synaptic strength occur, what caused the change?

- ...change in neurotransmitter release?

 - Mechanism?

- ...change in postsynaptic sensitivity to neurotransmitter?

 - Mechanism?


- ...change in synapse number?

 - Mechanism?


“Necessary”, “Sufficient”, “Mediator”, “Modulator”....

Paired Pulse Ratio

ISI = 50 msec




ISI = 200 msec




Paired Pulse Ratio

Facilitation




Ahmad et al 2012

Depression





Shypshyna & Veselovsk 2015



Post-Tetanic Potentiation at the NMJ




Modulation of Synaptic Strength via Presynaptic Receptors






Nicotine Enhances Excitatory Synaptic Transmission




Nicotine Enhances Excitatory Synaptic Transmission via $\alpha 7$ nAChRs




$\beta 2^*$ nAChRs enhance inhibitory drive to spinal dorsal horn neurons




Presynaptic NMDA receptors in spinal cord are increased in a chronic pain state




Deng et al., 2019




Presynaptic NMDA Receptors and Neuropathic Pain




Hippocampus






www.BrainConnection.com
© 1999 Scientific Learning Corporation



Hippocampus









LONG-LASTING POTENTIATION OF SYNAPTIC TRANSMISSION IN THE DENTATE AREA OF THE ANAESTHETIZED RABBIT FOLLOWING STIMULATION OF THE PERFORANT PATH

By T. V. P. BLISS AND T. LØMO

*From the National Institute for Medical Research, Mill Hill,
 London NW7 1AA and the Institute of Neurophysiology,
 University of Oslo, Norway*

(Received 12 February 1973)





Intracellular injections of EGTA block induction of hippocampal long-term potentiation

Gary Lynch, John Larson, Stephen Kelso,
German Barrionuevo & Frank Schottler

Center for the Neurobiology of Learning and Memory,
University of California, Irvine, California 92717, USA

Hippocampal long-term potentiation (LTP) is a remarkably stable facilitation of synaptic responses resulting from very brief trains of high-frequency stimulation^{1,2}. Because of its persistence and modest induction conditions, LTP represents a promising candidate for a substrate of memory. Some progress has been made in localizing the changes responsible for the effect; for example, it has been shown that LTP is not accompanied by changes in the fibre volleys of the test afferents³ or by generalized alterations of the dendrites of their target cells⁴. However, it is unknown whether the potentiation is due to pre- or postsynaptic changes and there is evidence in favour of each (for example, see refs 5, 6). We now report that intracellular injections of the calcium chelator EGTA block the development of LTP. These results strongly suggest that LTP is caused by a modification of the postsynaptic neurone and that its induction depends on the level of free calcium.




Table 1 e.p.s.p. amplitudes of cells which did not spike to single pulse stimulation after induction of extracellular long-term potentiation

	e.p.s.p. amplitude before high-frequency stimulation (mV)	Time (min)		
		5	10	15
Control	7.0 ± 0.6	+30 ± 5%	+30 ± 4%	+28 ± 4%
EGTA	7.1 ± 0.6	+8 ± 3%*	+1 ± 3%*	+5 ± 4%*

The e.p.s.p. amplitudes (mean ± s.e.m.) for 28 of 33 cells recorded with regular electrodes (control) and 24 of 26 cells recorded with EGTA-filled electrodes.


* $P < 0.001$.

Frequency-dependent involvement of NMDA receptors in the hippocampus: a novel synaptic mechanism


Caroline E. Herron, Robin A. J. Lester,
Elizabeth J. Coan & Graham L. Collingridge

Department of Pharmacology, University of Bristol, Medical School,
University Walk, Bristol BS8 1TD, UK


Fig. 2 Effect of 50 μ M APV on synaptic responses recorded intracellularly in CA1 neurones in response to stimulation of the Schaffer collateral-commissural pathway. *a*, Averages of five successive records of synaptic potentials evoked at 30-s intervals before (control), during and after washout of APV. *b*, Single records of responses of the same cell to high-frequency stimulation (20 shocks at 10-ms intervals) in the presence and following washout of APV. The e.p.s.p.s shown in *a* (centre and right-hand records) were obtained immediately before the two respective periods of high-frequency stimulation. *c*, Single records of responses to high-frequency stimulation in another cell. The times of stimulation are indicated by arrowheads and stimulus artefacts have been blanked out for clarity. Stimulus intensity and membrane potentials were constant throughout. Action potentials in *b* are truncated.




Long Term Potentiation



Long Term Depression





Long Term Potentiation, Cont'd



Long Term Depression, Cont'd

Endocannabinoid-mediated presynaptic inhibition



Long Term Potentiation, Cont'd

