

Homework 1

Quantum Mechanics

August 29th, 2022

CLAYTON SEITZ

Problem 1. For the spin $1/2$ state $|+\rangle_x$, evaluate both sides of the inequality

$$\langle(\Delta A)^2\rangle\langle(\Delta B)^2\rangle \geq \frac{1}{4}|\langle[A, B]\rangle|^2$$

for the operators $A = S_x$ and $B = S_y$, and show that the inequality is satisfied. Repeat for the operators $A = S_z$ and $B = S_y$

Solution.

Let $A = S_x$ and $B = S_y$. The variance $\langle(\Delta S_x)^2\rangle$ in state $|+\rangle_x$ must be zero since $|+\rangle_x$ is an eigenvector of S_x

$$\langle(\Delta S_x)^2\rangle = \langle S_x^2\rangle - \langle S_x\rangle^2 = 0$$

Therefore, the LHS of the above inequality is zero. The commutator $[S_x, S_y] = i\hbar S_z$ and

$$\langle S_z\rangle = \langle + |_x S_z | + \rangle_x = 0$$

Clearly the inequality is satisfied since both sides are zero. Now let $A = S_z$ and $B = S_y$. Since the state is prepared in $|+\rangle_x$, the variances $\langle(\Delta S_x)^2\rangle$ and $\langle(\Delta S_y)^2\rangle$ must be $1/4$ (this is just a fair coin toss).

The commutator $[S_z, S_y] = -i\hbar S_x$ and $\langle S_x\rangle = \frac{\hbar}{2}$. The inequality then reads

$$\frac{1}{16} \geq \frac{\hbar^2}{16}$$

which is satisfied given that $\hbar \approx 10^{-34} \text{ J} \cdot \text{s}$

■

Problem 2. Suppose a 2×2 matrix X (not necessarily Hermitian, nor unitary) is written as

Solution.

$$\begin{aligned}\mathrm{Tr}(X) &= \mathrm{Tr}(a_0) + \mathrm{Tr}\left(\sum_k a_k \sigma_k\right) \\ &= 2a_0\end{aligned}$$

$$\begin{aligned}\mathrm{Tr}(\sigma_k X) &= \mathrm{Tr}\left(\sigma_k a_0 + \sigma_k \sum_j a_j \sigma_j\right) \\ &= \mathrm{Tr}\left(\sigma_k a_0 + \sum_j a_j \sigma_k \sigma_j\right) \\ &= \mathrm{Tr}\left(\sum_j a_j \sigma_k \sigma_j\right)\end{aligned}$$

We can write out the equation $X = a_0 + \sigma \cdot a$ explicitly

$$X = \begin{pmatrix} a_0 + a_3 & a_1 - ia_3 \\ a_1 + ia_2 & a_0 - a_3 \end{pmatrix}$$

Thus we have four equations involving X_{ij} 's and a_k for $k = (1, 2, 3)$. We can manipulate those four equations to show that

$$\begin{aligned}a_0 &= \frac{X_{11} + X_{22}}{2} \\ a_1 &= \frac{X_{12} + X_{21}}{2} \\ a_2 &= \frac{X_{21} - X_{12}}{2} \\ a_3 &= \frac{X_{11} - X_{22}}{2}\end{aligned}$$

■

Problem 3.

Solution.



Problem 4.

Solution.



Problem 5.

Solution.



Problem 6.

Solution.

