

# Advancing super resolution microscopy for quantitative in-vivo imaging of chromatin nanodomains

Clayton W. Seitz

July 9, 2024

# Outline

Introduction to fluorescence nanoscopy

Method 1: Enhanced nanoscopy with single photon avalanche diode (SPAD) cameras


Method 2: Enhanced nanoscopy with deep generative models

Super-resolution of nucleosome nanodomains *in-vivo*


## Introduction to fluorescence nanoscopy

# Fluorescence microscopy and the diffraction limit


Minimal resolvable distance  $d \sim \lambda$




(a)



(b)




# Stochastic optical reconstruction microscopy (STORM)




Lakadamyali, M. et al. Nature Methods 17, (2020).

# Stochastic optical reconstruction microscopy (STORM)

## a Photoswitching





## b Temporal separation





- ▶ STORM and similar nanoscopy techniques are diffraction-unlimited
- ▶ Photoswitching enables resolution of emitters below the diffraction limit

# Stochastic optical reconstruction microscopy (STORM)

## a Photoswitching



## b Temporal separation



- ▶ STORM and similar nanoscopy techniques are diffraction-unlimited
- ▶ Photoswitching enables resolution of emitters below the diffraction limit

# Vanilla SMLM by localizing isolated fluorescent emitters

Modeling the point spread function permits sub-pixel localization

$$\mu_k = i_0 \int \int O(u, v) du dv$$


$$i_0 = g_k \eta N_0 \Delta$$

$g_k$  – pixel gain

$\eta$  – quantum efficiency

$N_0$  – photon emission rate


$\Delta$  – exposure time



Assume  $N_0$  is constant over  $\Delta$  (homogeneous Poisson)

$$\theta^* = \operatorname{argmax}_{\theta} \prod_k p(\mathbf{x}_k | \theta) = \operatorname{argmin}_{\theta} - \sum_k \log p(\mathbf{x}_k | \theta)$$

# Applications of single molecule localization microscopy




Wu et al. Trends in Cell Biology. 30 (2020)

## Method 1: Enhanced nanoscopy with single photon avalanche diode (SPAD) cameras

# Enhanced nanoscopy with single photon avalanche diode (SPAD) cameras



SPAD512




a



- ▶ Exposures as low as 1 microsecond
- ▶ Fast gated imaging for widefield fluorescence lifetime imaging
- ▶ Zero readout noise (imaging weak fluorescent signals)

## Method 2: Enhanced nanoscopy with deep generative models

## How to pack more localizations in a single frame?



$$p(\mathbf{x}_k|\theta) = A \sum_{q=0}^{\infty} \frac{1}{q!} e^{-\omega_k} \omega_k^q \frac{1}{\sqrt{2\pi w_k}} e^{-\frac{(\mathbf{x}_k - g_k q - o_k)^2}{2w_k^2}} \approx \text{Poisson}(\omega'_k)$$

- ▶ Would like to estimate a high-resolution  $\mathbf{y}$  from low-resolution  $\mathbf{x}$ , but it is many to one
- ▶ Must then model a *distribution* over  $\mathbf{y}$  i.e.,  $p_\theta(\mathbf{y}|\mathbf{x})$



## Trivial example of sampling from a mixture of Gaussians

Consider a two-component Gaussian mixture  $p(\mathbf{x}) = \sum_{k=1}^2 \pi_k \mathcal{N}(\mu_k, \sigma_k^2)$

If we know  $\nabla \log p(\mathbf{x})$  we can sample from  $p(\mathbf{x})$  with Langevin dynamics:

$$\mathbf{x}_i = \mathbf{x}_{i-1} + \epsilon \nabla \log p(\mathbf{x}) + \sqrt{2\epsilon} \eta \quad \eta \sim \mathcal{N}(0, I)$$

## Bayesian image restoration with diffusion models




Need to approximate the gradient  $s_\theta(\mathbf{y}_t) \approx \nabla \log p(\mathbf{x})$  and sample:

$$\mathbf{y}_{t-1} = \frac{1}{\sqrt{1 - \beta_t}} (\mathbf{y}_t + \beta_t s_\theta(\mathbf{y}_t)) + \sqrt{\beta_t} \xi \quad \xi \sim \mathcal{N}(0, I)$$


## Super-resolution of nucleosome nanodomains *in-vivo*

# Hierarchical structure of chromatin




Fyodorov, D. et al. Nat Rev Mol Cell Biol **19**, (2018).

# Bromodomain protein 4 (BRD4) binds acetylated chromatin









Zheng, B. et al. Molecular Cell **16**, (2023).




Fillapakoulos, P. et al. Nature **468**, (2010).

# BET inhibitor (+)-JQ1 decondenses chromatin and increases mobility




Blots: top - BRD4, bottom - GAPDH

# Bromodomain protein 4 (BRD4) binds acetylated chromatin




# BRD4 phosphorylation state is necessary for maintenance of chromatin structure




# Coarse grained molecular dynamics of chromatin binders at 310K

a



b




100kb chromatin chains interact with binders via the potential

$$U_{ij} = \epsilon \left( 1 - \left( \frac{|r_{ij}|}{R_0} \right)^2 \right)^3$$

- ▶ A (B) type particles represent unacetylated (acetylated) chromatin beads
- ▶ BRD4-like C particles bind B type particles with variable energies





# Multivalent chromatin binding reduces chromatin mobility



Integrate Brownian dynamics:  $\dot{r} = \gamma^{-1} \nabla U + \sqrt{2k_B T} \gamma^{-1/2} \xi$   $\gamma = 10^{-6}$

Stochastic forcing is a delta-correlated white-noise  
 $\xi \sim \mathcal{N}(0, 1)$ ,  $\langle \xi(t) \xi(t + \tau) \rangle = \delta(\tau)$

# Multivalent chromatin binding reduces chromatin mobility



Experiment:  $D_{WT} - D_{7D} \approx 10^{-3} \mu\text{m}^2/\text{s}, \gamma = 10^{-6}$

## **Temporary page!**

$\text{\LaTeX}$  was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away because  $\text{\LaTeX}$  now knows how many pages to expect for this document.