

Interferon- γ induction of GBP5 in HeLa cells

Clayton W. Seitz

October 24, 2022

The principle of Interferon- γ induced transcriptional memory





Figure 1


Siwek et al. *Activation of Clustered IFN γ Target Genes Drives Cohesin-Controlled Transcriptional Memory*. Molecular Cell 2020

Rare HeLa cell GBP5 expression @ 24h after reinduction with IFN- γ


Single max intensity projection from a 10x10x9 (2mm x 2mm x 8um) tiled acquisition



Rare HeLa cell GBP5 expression @ 24h after reinduction with IFN- γ



Intensity histogram for rare GBP5 expression



- ▶ Very few (1 – 5%) reinduced cells express GBP5, but those that do express at levels comparable to GAPDH
- ▶ Control sample shows little to no GBP5 expression

Comments on ergodicity of transcription

1. Priming leads to more transcription




2. Priming leads to more cells transcribing




- ▶ RNA flow cannot apply to non-ergodic systems (yet ergodicity is often assumed)
- ▶ Previous work suggests that IFN- γ induces epigenetic changes at the GBP5 locus
- ▶ What is the epigenetic change? Is the epigenetic change all or nothing?

Epigenetic changes at GBP genes after IFN- γ treatment

Siwek et al. *Activation of Clustered IFN γ Target Genes Drives Cohesin-Controlled Transcriptional Memory*. Molecular Cell 2020




STORM imaging of H2B




Lakadamyali et al. *Visualizing the genome in high resolution challenges our textbook understanding*. Nature Methods 2020

Working principle of dSTORM



Details on STORM timing setup



Remaining questions on the application of STORM

If STORM works with the H2B-HaloTag

- ▶ How do we interpret H2B density maps?
- ▶ Can you correlate ChIP-seq/ATAC-seq data with H2B density?